



HAL
open science

Some geometric applications of the discrete heat flow

Alexandros Eskenazis

► **To cite this version:**

| Alexandros Eskenazis. Some geometric applications of the discrete heat flow. 2023. hal-04272902

HAL Id: hal-04272902

<https://hal.science/hal-04272902>

Preprint submitted on 6 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SOME GEOMETRIC APPLICATIONS OF THE DISCRETE HEAT FLOW

ALEXANDROS ESKENAZIS

ABSTRACT. We present two geometric applications of heat flow methods on the discrete hypercube $\{-1, 1\}^n$. First, we prove that if X is a finite-dimensional normed space, then the bi-Lipschitz distortion required to embed $\{-1, 1\}^n$ equipped with the Hamming metric into X satisfies

$$c_X(\{-1, 1\}^n) \gtrsim \sup_{p \in [1, 2]} \frac{n}{T_p(X) \min\{n, \dim(X)\}^{1/p}},$$

where $T_p(X)$ is the Rademacher type p constant of X . This estimate yields a mutual refinement of distortion lower bounds which follow from works of Oleszkiewicz (1996) and Ivanisvili, van Handel and Volberg (2020) for low-dimensional spaces X . The proof relies on an extension of an important inequality of Pisier (1986) on the biased hypercube combined with an application of the Borsuk–Ulam theorem from algebraic topology. Secondly, we introduce a new metric invariant called metric stable type as a functional inequality on the discrete hypercube and prove that it coincides with the classical linear notion of stable type for normed spaces. We also show that metric stable type yields bi-Lipschitz nonembeddability estimates for weighted hypercubes.

2020 Mathematics Subject Classification. Primary: 46B85; Secondary: 30L15, 42C10, 46B07.

Key words. Hamming cube, Rademacher type, metric embeddings, Borsuk–Ulam theorem, stable type.

1. INTRODUCTION

Let $\{-1, 1\}^n$ be the n -dimensional discrete hypercube equipped with the Hamming metric

$$\forall x, y \in \{-1, 1\}^n, \quad \rho(x, y) = \frac{1}{2} \sum_{i=1}^n |x(i) - y(i)|, \quad (1)$$

where $x = (x(1), \dots, x(n))$ and $y = (y(1), \dots, y(n))$. The purpose of the present paper is to investigate certain metric properties of the Hamming cube via heat flow methods.

1.1. Dimensionality of Hamming metrics. If (M, d_M) is a metric space and $(Y, \|\cdot\|_Y)$ is a normed space, we say that M embeds into Y with bi-Lipschitz distortion at most $D \in [1, \infty)$, if there exists a mapping $f : M \rightarrow Y$ satisfying

$$\forall p, q \in M, \quad d_M(p, q) \leq \|f(p) - f(q)\|_Y \leq D d_M(p, q). \quad (2)$$

The least $D \geq 1$ for which such an embedding exists will be denoted by $c_Y(M)$. The rapidly growing field of metric dimension reduction aims to uncover conditions under which given families of metric spaces admit (or do not admit) embeddings into low-dimensional normed spaces with prescribed properties. Without attempting to survey this vast area, we note that important contributions have been made on low-dimensional embeddings of finite subsets of Hilbert space [29], arbitrary finite metric spaces [30, 3, 41, 42], discrete hypercubes [56, 34], diamond graphs [14, 35, 52], Laakso graphs [24, 34], ultrametric spaces [8], series-parallel graphs [15], recursive cycle graphs [1], Heisenberg-type metrics [32, 54], ℓ_p variants of thin Laakso structures [7, 9] and expander graphs [48, 50]. We refer to the survey [49] for more bibliographic information and to [26, 38, 69, 2] for a sample of algorithmic applications.

The first to study the bi-Lipschitz embeddability of hypercubes into normed spaces was Enflo. In the seminal work [20], he introduced the notion of *roundness* of a metric space and used it to show that any embedding of the Hamming cube $\{-1, 1\}^n$ into an $L_p(\mu)$ space, where $p \in [1, 2]$, incurs bi-Lipschitz distortion at least $n^{1-1/p}$ (see also [19, 21] for additional early

results along these lines). More specifically, Enflo proved that if $p \in [1, 2]$, then any mapping of the form $f : \{-1, 1\}^n \rightarrow L_p(\mu)$ satisfies the estimate

$$\int_{\{-1, 1\}^n} \|f(x) - f(-x)\|_{L_p(\mu)}^p d\sigma_n(x) \leq \sum_{i=1}^n \int_{\{-1, 1\}^n} \|f(x) - f(x(1), \dots, -x(i), \dots, x(n))\|_{L_p(\mu)}^p d\sigma_n(x), \quad (3)$$

where σ_n is the uniform probability on $\{-1, 1\}^n$. This readily implies that if f has bi-Lipschitz distortion D , then $D \geq n^{1-1/p}$. In the follow-up work [21], he raised an influential problem by asking for which normed spaces $(X, \|\cdot\|_X)$, inequality (3) is satisfied for X -valued functions f up to a multiplicative constant T , independent of the choice of f or the dimension n . Restricting this requirement to linear functions $f(x) = \sum_{i=1}^n x_i v_i$, one recovers the necessary condition

$$\int_{\{-1, 1\}^n} \left\| \sum_{i=1}^n x_i v_i \right\|_X^p d\sigma_n(x) \leq T^p \sum_{i=1}^n \|v_i\|_X^p, \quad (4)$$

which must be satisfied for every $n \in \mathbb{N}$ and vectors $v_1, \dots, v_n \in X$. If a normed space X satisfies (4), we say that X has *Rademacher type p* and the least constant T is denoted by $T_p(X)$. After decades of substantial efforts (see [13, 64, 53, 25, 22]), Ivanisvili, van Handel and Volberg resolved Enflo's problem in the breakthrough work [27] by proving the sufficiency of this condition, namely that any normed space of Rademacher type p also has Enflo's nonlinear type p . Consequently, any bi-Lipschitz embedding of $\{-1, 1\}^n$ into a normed space X of Rademacher type p incurs distortion at least a constant multiple of $T_p(X)^{-1} n^{1-1/p}$. We note in passing that, conversely, a classical theorem of Pisier [57] implies that if X does not have type p for any $p > 1$, then $\{-1, 1\}^n$ embeds into X with bi-Lipschitz distortion at most $1 + \varepsilon$, for any $\varepsilon > 0$.

Independently of this line of research, the beautiful (but perhaps overlooked) work [56] of Oleszkiewicz established a nonembeddability result for discrete hypercubes in the context of dimensionality reduction. Following Ball, Carlen and Lieb [5], we say that a normed space is *p -uniformly smooth*, where $p \in [1, 2]$, if there exists a constant $S > 0$ such that

$$\forall x, y \in X, \quad \frac{\|x\|_X^p + \|y\|_X^p}{2} \leq \left\| \frac{x+y}{2} \right\|_X^p + S^p \left\| \frac{x-y}{2} \right\|_X^p; \quad (5)$$

the least such constant S is denoted by $S_p(X)$. A well-known tensorization argument due to Pisier [60] shows that $T_p(X) \leq S_p(X)$, yet there exist examples of normed spaces X for which $T_p(X) < \infty$ whereas $S_p(X) = \infty$ for $p \in (1, 2]$, see [61, 28, 65]. The main result of Oleszkiewicz's paper [56] asserts that the distortion required to embed $\{-1, 1\}^n$ into a finite-dimensional normed space X satisfies

$$c_X(\{-1, 1\}^n) \geq \sup_{p \in [1, 2]} \frac{n}{S_p(X) \min\{n, \dim(X)\}^{1/p}} \quad (6)$$

(see also [6] for a precursor of this result for linear embeddings). This substantially improves the bound $c_X(\{-1, 1\}^n) \gtrsim \sup_{p \in [1, 2]} T_p(X)^{-1} n^{1-1/p}$ which follows from [27], at least for spaces X with p -smoothness constant $S_p(X) \asymp 1$ and dimension $\dim(X) \ll n$.

The first goal of the present paper is to revisit the technique used for Oleszkiewicz's nonembeddability theorem [56], in particular proving the following mutual refinement of his result and of the recent work of Ivanisvili, van Handel and Volberg [27].

Theorem 1. *Let $(X, \|\cdot\|_X)$ be a finite-dimensional normed space. Then, for any $n \geq 1$, we have*

$$c_X(\{-1, 1\}^n) \gtrsim \sup_{p \in [1, 2]} \frac{n}{T_p(X) \min\{n, \dim(X)\}^{1/p}} \quad (7)$$

We emphasize that, in contrast to Oleszkiewicz's bound (6), Theorem 1 captures more accurately the nonembeddability of the hypercube into (finite-dimensional subspaces of) normed spaces which have Rademacher type p but are not r -smooth for any $r \in (1, 2]$, see [28, 65].

About the proof. Theorem 1 is proven by a combination of semigroup tools with a clever topological trick of [56]. More specifically, let $f : \{-1, 1\}^n \rightarrow X$ be a function, where X is a d -dimensional normed space and $d < n$. An application of the Borsuk–Ulam theorem [43] for the unique multilinear extension of f implies that there exists a subset $\sigma \subseteq \{1, \dots, n\}$ with $|\sigma| = d$, a product measure ν on $\{-1, 1\}^\sigma$ and a point $w \in \{-1, 1\}^{\sigma^c}$ such that

$$\int_{\{-1, 1\}^\sigma} f(x, w) \, d\nu(x) = \int_{\{-1, 1\}^\sigma} f(-x, -w) \, d\nu(x). \quad (8)$$

Then, a Poincaré inequality à la Enflo for the product measure ν (instead of the uniform measure σ_d) on the d -dimensional subcubes $\{-1, 1\}^\sigma \times \{w\}$ and $\{-1, 1\}^\sigma \times \{-w\}$ yields the distortion bounds of Theorem 1 (see Theorem 14 and also equation (78) below).

In the case of p -uniformly smooth spaces, Oleszkiewicz [56] used (8) and a bootstrap argument for the Lipschitz constant of f , based on the two-point inequality (5), to obtain (6). In our case, the biased Poincaré inequality which will yield (7) is an extension of an inequality for the uniform measure that was proven in [27]. The key technical contribution of [27] was a novel representation of the time derivative of the heat flow on $\{-1, 1\}^n$. Instead, we consider a Markov process having the product measure ν as stationary measure (see Section 2) and prove a formula for the time derivative of the corresponding semigroup (see Proposition 13) which extends the formula of [27] (see also (47) below). Due to the fact that our product measure ν is no longer the stationary measure of the random walk on a group, the resulting identity lacks some homogeneity properties that were used in [27], but nevertheless it is sufficient for the proof of the biased Poincaré inequality which is needed for our geometric application.

1.2. A Pisier–Talagrand inequality on the biased cube. For $\alpha \in (0, 1)$ we denote by μ_α the α -biased probability measure on $\{-1, 1\}$ with $\mu_\alpha\{1\} = \alpha$ and $\mu_\alpha\{-1\} = 1 - \alpha$. The proof of Theorem 1 yields as a consequence the following extension and refinement of Pisier’s inequality [64].

Theorem 2. *For every $p \in [1, \infty)$ and $\alpha \in (0, 1)$, there exist $K_{p,\alpha}, C_\alpha \in (0, \infty)$ such that the following holds. For any normed space $(X, \|\cdot\|_X)$ and any $n \in \mathbb{N}$, every function $f : \{-1, 1\}^n \rightarrow X$ satisfies*

$$\begin{aligned} \left\| f - \int_{\{-1, 1\}^n} f \, d\mu_\alpha^n \right\|_{L_p(\log L)^{p/2}(\mu_\alpha^n; X)} &\leq K_{p,\alpha} \left(\int_{\{-1, 1\}^n} \left\| \sum_{i=1}^n \delta_i \partial_i^\alpha f \right\|_{L_p(\mu_\alpha^n; X)}^p \, d\sigma_n(\delta) \right)^{1/p} \\ &\quad + C_\alpha (\log n + 1) \int_{\{-1, 1\}^n} \left\| \sum_{i=1}^n \delta_i \partial_i^\alpha f \right\|_{L_1(\mu_\alpha^n; X)} \, d\sigma_n(\delta). \end{aligned} \quad (9)$$

If additionally X is assumed to be of finite cotype, then there exists $K_{p,\alpha}(X) \in (0, \infty)$ such that

$$\left\| f - \int_{\{-1, 1\}^n} f \, d\mu_\alpha^n \right\|_{L_p(\log L)^{p/2}(\mu_\alpha^n; X)} \leq K_{p,\alpha}(X) \left(\int_{\{-1, 1\}^n} \left\| \sum_{i=1}^n \delta_i \partial_i^\alpha f \right\|_{L_p(\mu_\alpha^n; X)}^p \, d\sigma_n(\delta) \right)^{1/p}. \quad (10)$$

The (standard) definitions of Orlicz norms, cotype and discrete derivatives will be given at the main part of the article. Theorem 2 is an optimal vector-valued version of a deep logarithmic Sobolev inequality of Talagrand [67]. In the case of the uniform measure which corresponds to $\alpha = \frac{1}{2}$, Theorem 2 was proven recently in [16].

1.3. Metric stable type. Besides its interest in the context of embedding theory, the work [27] of Ivanisvili, van Handel and Volberg was a major contribution to a long-standing research program in nonlinear functional analysis called the *Ribe program*. Put simply, the foundational idea of the program (as put forth by Bourgain [12], who was inspired by a landmark rigidity theorem of Ribe [66]) is that isomorphic local properties of normed spaces can be equivalently reformulated using only distances between pairs of points, with no reference to the spaces’ linear structure (see also [46, 4]). In this sense, the main result of [27] completes the Ribe program for Rademacher type by proving that it is equivalent to nonlinear Enflo type.

Among the various isomorphic properties studied in the local theory of normed spaces, *stable type* (see [58, 59]) plays a prominent role, in particular due to its relation to the problem of identifying ℓ_p^n subspaces of Banach spaces [63]. Recall that a symmetric random variable θ is distributed according to the standard p -stable law, where $p \in (0, 2]$, if it satisfies $\mathbb{E}\exp(it\theta) = \exp(-|t|^p)$ for every $t \in \mathbb{R}$. A Banach space $(X, \|\cdot\|_X)$ is said to have stable type p if for every $r < p$, every $n \in \mathbb{N}$ and every vectors $v_1, \dots, v_n \in X$, we have

$$\left(\mathbb{E}\left\|\sum_{i=1}^n \theta_i v_i\right\|_X^r\right)^{1/r} \leq C_{p,r}(X) \left(\sum_{i=1}^n \|v_i\|_X^p\right)^{1/p}, \quad (11)$$

where the constant $C_{p,r}(X)$ depends only on p, r and X , but not on n or the choice of v_1, \dots, v_n , and $\theta_1, \theta_2, \dots$ are independent standard symmetric p -stable random variables.

In the special case $p = 2$, p -stable random variables are Gaussians and thus stable type 2 coincides with Rademacher type 2. Moreover, any space with stable type $p \in [1, 2)$ also has Rademacher type p but the converse does not necessarily hold. In fact, early results of Maurey and Pisier in the theory of type (see [58, Proposition 3] and [59, Théorème 1]), show that a normed space X has stable type $p \in [1, 2)$ if and only if X has Rademacher type $p + \varepsilon$ for some $\varepsilon > 0$. Therefore, formally, the Ribe program for stable type $p \in [1, 2)$ has been completed by [27]: a normed space X has stable type p if and only if X has Enflo type $p + \varepsilon$ for some $\varepsilon > 0$. In other words, stable type p of a normed space X can be characterized by the validity of some, a priori unknown, X -valued Poincaré-type inequality from a given family.

To amend this, we shall introduce a new nonlinear invariant called *metric stable type* which is again a Poincaré-type inequality for functions defined on the discrete hypercube and will serve as a metric characterization of stable type for normed spaces. If (M, d_M) is a metric space and $f : \{-1, 1\}^n \rightarrow M$ is an M -valued function, we denote

$$\forall x \in \{-1, 1\}^n, \quad \mathfrak{d}_i f(x) \stackrel{\text{def}}{=} \frac{1}{2} d_M(f(x), f(x(1), \dots, -x(i), \dots, x(n))). \quad (12)$$

Moreover, we shall denote the weak ℓ_p norm on \mathbb{R}^n by

$$\forall w \in \mathbb{R}^n, \quad \|w\|_{\ell_{p,\infty}^n} \stackrel{\text{def}}{=} \sup_{r \geq 0} \{r \cdot \#\{i : |w_i| \geq r\}^{1/p}\}, \quad (13)$$

where $\#S$ denotes the cardinality of a finite set S .

Definition 3. A metric space (M, d_M) has metric stable type $p \in (0, 2)$ with constant $S \in (0, \infty)$ if for any $n \in \mathbb{N}$, every function $f : \{-1, 1\}^n \rightarrow M$ satisfies

$$\int_{\{-1, 1\}^n} d_M(f(x), f(-x))^p d\sigma_n(x) \leq S^p \int_{\{-1, 1\}^n} \left\|(\mathfrak{d}_1 f(x), \dots, \mathfrak{d}_n f(x))\right\|_{\ell_{p,\infty}^n}^p d\sigma_n(x). \quad (14)$$

Since $\|\cdot\|_{\ell_{p,\infty}^n} \leq \|\cdot\|_{\ell_p^n}$ by Markov's inequality, any space with metric stable type p also has Enflo type p with the same constant. We shall prove the following metric characterization.

Theorem 4. A normed space X has metric stable type $p \in [1, 2)$ if and only if X has stable type p .

Pisier's K -convexity theorem [62], asserts that a Banach space X is K -convex if and only if it has Rademacher type p for some $p > 1$. In view of the aforementioned characterization of stable type [58, 59], this is equivalent to X having stable type 1 and thus we derive the following metric characterization of K -convexity by a Poincaré inequality.

Corollary 5. A normed space X is K -convex if and only if X has metric stable type 1.

Embeddings. The fact that stable type of Banach spaces is a refinement of Rademacher type can also be seen at the level of (linear) embeddings. Indeed, if X is a space of Rademacher type p , then standard considerations show that the Banach–Mazur distance between ℓ_1^n and any n -dimensional subspace of X is at least a constant multiple of $n^{1-1/p}$. This estimate is sharp for $X = \ell_p$. On the other hand, if X has stable type p , then the characterization of [58, 59]

implies that X has Rademacher type r for some $r > p$. Hence, there exists $\varepsilon(X) > 0$ such that the Banach–Mazur distance between ℓ_1^n and any n -dimensional subspace of X is at least $n^{1-1/p+\varepsilon(X)}$.

In the metric setting, Enflo type gives distortion lower bounds for embeddings of $\{-1, 1\}^n$. More generally, given a vector $\mathbf{w} = (w_1, \dots, w_n) \in \mathbb{R}_+^n$, consider the metric space $\{-1, 1\}_{\mathbf{w}}^n$ which is the hypercube $\{-1, 1\}^n$ equipped with the *weighted* Hamming metric

$$\forall x, y \in \{-1, 1\}^n, \quad \rho_{\mathbf{w}}(x, y) = \frac{1}{2} \sum_{i=1}^n w_i |x(i) - y(i)|. \quad (15)$$

It follows readily from the definitions that if M has Enflo type p , then any embedding of $\{-1, 1\}_{\mathbf{w}}^n$ into M incurs distortion at least a constant multiple of $\|\mathbf{w}\|_{\ell_1^n} / \|\mathbf{w}\|_{\ell_p^n}$. For spaces of metric stable type p , we have the following improvement for the distortion of weighted hypercubes.

Proposition 6. *If a metric space (M, d_M) has metric stable type p with constant S , then*

$$\forall \mathbf{w} \in \mathbb{R}_+^n, \quad c_M(\{-1, 1\}_{\mathbf{w}}^n) \geq \frac{\|\mathbf{w}\|_{\ell_1^n}}{S \|\mathbf{w}\|_{\ell_{p,\infty}^n}}. \quad (16)$$

Concretely, choosing the weight vector \mathbf{w} with $w_i = i^{-1/p}$, we have that $\|\mathbf{w}\|_{\ell_{p,\infty}^n} \asymp 1$, whereas $\|\mathbf{w}\|_{\ell_p^n} \asymp (\log n)^{1/p}$, therefore any embedding of $\{-1, 1\}_{\mathbf{w}}^n$ into a metric space of Enflo type p incurs distortion at least $n^{1-1/p}/(\log n)^{1/p}$ instead of the asymptotically stronger lower bound $n^{1-1/p}$ which one gets for target spaces of metric stable type p .

On the nonlinear Maurey–Pisier problem for type. A landmark theorem of Maurey and Pisier [44] asserts that if $p_X \in [1, 2]$ is the supremal Rademacher type of an infinite-dimensional Banach space X , then X contains the spaces $\ell_{p_X}^n$ uniformly. In [63], Pisier used stable type to give a much simpler proof of this important theorem, while simultaneously obtaining the following quantitative refinement: for any $\varepsilon > 0$ and $p \in [1, 2)$, every normed space X contains a subspace which is $(1 + \varepsilon)$ -isomorphic to ℓ_p^d provided that d is smaller than some explicit (unbounded) function of ε and the stable type p constant of X . When X is infinite-dimensional, this yields the result of [44] since X does not have stable type p_X by the results of [58, 59].

In [13], Bourgain, Milman and Wolfson introduced a notion of nonlinear type, which is now referred to as BMW type, and showed that if a metric space M does not have BMW type p for any $p > 1$, then $c_M(\{-1, 1\}^n) = 1$ for any $n \in \mathbb{N}$. Finding a satisfactory nonlinear Maurey–Pisier theorem for metric spaces of supremal nonlinear type $p > 1$ remains an open problem (see also [47, Section 6]). It would be interesting to understand whether the newly introduced notion of metric stable type can lead to a nonlinear version of the result of [63]. We refer to Section 6 for further remarks and open problems which naturally arise from this work.

Acknowledgements. I wish to thank Florent Baudier, Paata Ivanisvili and Assaf Naor for their constructive feedback on this work.

2. PRELIMINARIES

2.1. Probability. In this section, we outline the basics of analysis on the biased hypercube, with an emphasis on the underlying semigroup structure.

The biased measure. Recall that, for $\alpha \in (0, 1)$, the α -biased probability measure μ_α on $\{-1, 1\}$ is given by $\mu_\alpha\{1\} = \alpha$ and $\mu_\alpha\{-1\} = 1 - \alpha$. Moreover, if $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_n) \in (0, 1)^n$, then we shall denote by $\mu_{\boldsymbol{\alpha}}$ the product measure $\mu_{\alpha_1} \otimes \dots \otimes \mu_{\alpha_n}$ on the hypercube $\{-1, 1\}^n$.

The Markov process. For $\alpha \in (0, 1)$, consider the transition matrices $\{p_t^\alpha\}_{t \geq 0}$ on $\{-1, 1\}$ given by

$$\forall t \geq 0, \quad \begin{pmatrix} p_t^\alpha(1, 1) & p_t^\alpha(1, -1) \\ p_t^\alpha(-1, 1) & p_t^\alpha(-1, -1) \end{pmatrix} = \begin{pmatrix} 1 - (1 - e^{-t})(1 - \alpha) & (1 - e^{-t})(1 - \alpha) \\ (1 - e^{-t})\alpha & 1 - (1 - e^{-t})\alpha \end{pmatrix} \quad (17)$$

Moreover, for $\alpha = (\alpha_1, \dots, \alpha_n) \in (0, 1)^n$ consider the corresponding tensor products $\{p_t^\alpha\}_{t \geq 0}$ on $\{-1, 1\}^n$ given by

$$\forall x, y \in \{-1, 1\}^n, \quad p_t^\alpha(x, y) = \prod_{i=1}^n p_t^{\alpha_i}(x(i), y(i)). \quad (18)$$

As each $p_t^{\alpha_i}$ is a row-stochastic 2×2 matrix with nonnegative entries, the same holds also for the $2^n \times 2^n$ matrices p_t^α . Therefore, $\{p_t^\alpha\}_{t \geq 0}$ is the transition kernel of a time-homogeneous Markov chain $\{X_t^\alpha\}_{t \geq 0}$ on $\{-1, 1\}^n$, that is

$$\forall t, s \geq 0, \quad \mathbb{P}\{X_{t+s}^\alpha = y \mid X_s^\alpha = x\} = p_t^\alpha(x, y), \quad (19)$$

where $x, y \in \{-1, 1\}^n$. We shall need the following simple facts for this process.

Lemma 7. Fix $\alpha \in (0, 1)^n$ and let $\{X_t^\alpha\}_{t \geq 0}$ be a Markov process on $\{-1, 1\}^n$ with transition kernels $\{p_t^\alpha\}_{t \geq 0}$. Then, $\{X_t^\alpha\}_{t \geq 0}$ is stationary and reversible with respect to μ_α .

Proof. Due to the product structure of the Markov chain, it suffices to consider the case $n = 1$, that is, to prove that for $\alpha \in (0, 1)$,

$$\forall x, y \in \{-1, 1\}, \quad \mu_\alpha(x)p_t^\alpha(x, y) = \mu_\alpha(y)p_t^\alpha(y, x). \quad (20)$$

This follows automatically by the expression (17) for the transition matrix. The simple fact that reversibility implies stationarity is well-known [37, Proposition 1.20]. \square

The stationary Markov process $\{X_t^\alpha\}_{t \geq 0}$ has a simple probabilistic interpretation which we shall now describe. For $i = 1, \dots, n$, let $\{N_t(i)\}_{t \geq 0}$ be n independent Poisson processes of unit rate and suppose that X_0^α is sampled from μ_α independently of $\{N_t(i)\}_{t \geq 0}$. Then, at any time $t > 0$ for which the process $N_t(i)$ jumps for some $i \in \{1, \dots, n\}$, the corresponding value $X_t^\alpha(i)$ is updated independently from μ_{α_i} . An explicit calculation shows that this probabilistic construction gives rise exactly to the transition kernel of (17) and (18).

The corresponding semigroup. Fix $\alpha \in (0, 1)^n$ and let $\{P_t^\alpha\}_{t \geq 0}$ be the Markov semigroup associated to the process $\{X_t^\alpha\}_{t \geq 0}$. Concretely, if X is a vector space, then for every function $f : \{-1, 1\}^n \rightarrow X$ and $t \geq 0$, we denote by

$$\forall x \in \{-1, 1\}^n, \quad P_t^\alpha f(x) = \mathbb{E}[f(X_t^\alpha) \mid X_0^\alpha = x]. \quad (21)$$

In view of the above interpretation of $\{X_t^\alpha\}_{t \geq 0}$ by means of a Poisson process, the action of the semigroup $\{P_t^\alpha\}_{t \geq 0}$ can be computed via the identity

$$\begin{aligned} P_t^\alpha f &= \sum_{S \subseteq \{1, \dots, n\}} \mathbb{P}\{N_t(i) > 0 \text{ for } i \in S \text{ and } N_t(i) = 0 \text{ for } i \notin S\} \int_{\{-1, 1\}^S} f \, d \prod_{i \in S} \mu_{\alpha_i} \\ &= \sum_{S \subseteq \{1, \dots, n\}} (1 - e^{-t})^{|S|} e^{-t(n-|S|)} \int_{\{-1, 1\}^S} f \, d \prod_{i \in S} \mu_{\alpha_i}. \end{aligned} \quad (22)$$

Lemma 8. Fix $\alpha = (\alpha_1, \dots, \alpha_n) \in (0, 1)^n$ and let $\{P_t^\alpha\}_{t \geq 0}$ be the semigroup (21). Then, the action of its generator \mathcal{L}_α on a function $f : \{-1, 1\}^n \rightarrow X$, where X is a vector space, is given by

$$\forall x \in \{-1, 1\}^n, \quad \mathcal{L}_\alpha f(x) = - \sum_{i=1}^n \partial_i^{\alpha_i} f(x) \quad (23)$$

where $\partial_i^\beta f(x) = f(x) - \int_{\{-1, 1\}} f(x_1, \dots, x_{i-1}, y, x_{i+1}, \dots, x_n) \, d\mu_\beta(y)$ for $\beta \in (0, 1)$.

Proof. The claim follows from the expression (22) of the semigroup and the definition

$$\forall x \in \{-1, 1\}^n, \quad \mathcal{L}_\alpha f(x) = \left. \frac{d}{dt} \right|_{t=0} P_t^\alpha f(x). \quad \square$$

2.2. Topology. Apart from the probabilistic elements from analysis on biased hypercubes, the proof of Theorem 1 also has a crucial topological component, following an idea of [56].

The Borsuk–Ulam theorem. While the Poincaré-type inequality of Enflo for X -valued functions on $\{-1, 1\}^n$ cannot capture the dimension of the target space X , a key part of the argument towards Theorem 1 is to show that there exists a $\dim(X)$ -dimensional subcube of $\{-1, 1\}^n$ along with a bias vector α for which the α -biased Poincaré inequalities (see Theorem 14 below) on this subcube and its antipodal yield much better distortion lower bounds. This will be proven using the Borsuk–Ulam theorem from algebraic topology, see [43].

Theorem 9 (Borsuk–Ulam). *For every continuous function $g : \mathbb{S}^d \rightarrow \mathbb{R}^d$, where $d \in \mathbb{N}$, there exists a point $w \in \mathbb{S}^d$ such that $g(w) = g(-w)$.*

Multilinear extension and low-dimensional faces of the cube. Every function $f : \{-1, 1\}^n \rightarrow X$ admits a unique multilinear extension on the solid cube $[-1, 1]^n$, given by

$$\forall y \in [-1, 1]^n, \quad F(y) \stackrel{\text{def}}{=} \sum_{S \subseteq \{-1, 1\}^n} \left(\frac{1}{2^n} \sum_{x \in \{-1, 1\}^n} f(x) w_S(x) \right) w_S(y), \quad (24)$$

where $w_S(a) = \prod_{i \in S} a_i$, which is usually referred to as the Fourier–Walsh expansion of f . Extending f to the continuous cube allows for the use of topological methods. In what follows, we will exploit the fact that the cube $[-1, 1]^n$ is equipped with a canonical CW complex structure. Concretely, for $d \in \{1, \dots, n\}$, consider the subsets

$$\mathcal{C}_d^n = \left\{ x \in [-1, 1]^n : \text{there exists } \sigma \subseteq \{1, \dots, n\} \text{ with } |\sigma| \geq n - d \text{ and } |x(i)| = 1, \forall i \in \sigma \right\} \quad (25)$$

consisting of all ℓ -dimensional faces of $[-1, 1]^n$ for $\ell \leq d$, so that $\mathcal{C}_n^n = [-1, 1]^n$ and $\mathcal{C}_0^n = \{-1, 1\}^n$. We shall use the following elementary topological fact (see [56, Lemma 1]).

Lemma 10. *If $d < n$, there exists a continuous map $h_d : \mathbb{S}^d \rightarrow \mathcal{C}_d^n$ with $h_d(-x) = -h_d(x)$, $\forall x \in \mathbb{S}^d$.*

Combining this and the Borsuk–Ulam theorem, we deduce the following useful lemma.

Lemma 11. *If $n, d \in \mathbb{N}$ with $d < n$, then for every continuous function $F : \mathcal{C}_d^n \rightarrow \mathbb{R}^d$ there exists a point $z \in \mathcal{C}_d^n$ such that $F(z) = F(-z)$.*

Proof. Consider the function $g \stackrel{\text{def}}{=} F \circ h_d : \mathbb{S}^d \rightarrow \mathbb{R}^d$, where h_d is the function of Lemma 10. By the Borsuk–Ulam theorem and the oddness of h_d , there exists a point $w \in \mathbb{S}^d$ such that

$$F(h_d(w)) = g(w) = g(-w) = F(h_d(-w)) = F(-h_d(w)) \quad (26)$$

and the conclusion follows by choosing $z = h_d(w) \in \mathcal{C}_d^n$. \square

3. PROOF OF THEOREM 1

We are now ready to proceed to the main part of the proof. The main analytic component is a biased version of the key formula of [27] for the time derivative of the heat flow on $\{-1, 1\}^n$. Given $t \geq 0$, $\alpha \in (0, 1)$ and an auxiliary parameter $\theta \in \mathbb{R}$, consider the matrix $\eta_t^\alpha(\cdot, \cdot; \theta)$ given by

$$\forall t \geq 0, \quad \begin{pmatrix} \eta_t^\alpha(1, 1; \theta) & \eta_t^\alpha(1, -1; \theta) \\ \eta_t^\alpha(-1, 1; \theta) & \eta_t^\alpha(-1, -1; \theta) \end{pmatrix} = \begin{pmatrix} \frac{e^{-t} - \theta}{p_t^\alpha(1, 1)} & \frac{-\theta}{p_t^\alpha(-1, 1)} \\ \frac{\theta - e^{-t}}{p_t^\alpha(1, -1)} & \frac{\theta}{p_t^\alpha(-1, -1)} \end{pmatrix}. \quad (27)$$

For future reference, we record the following straightforward properties of $\eta_t^\alpha(\cdot, \cdot; \theta)$.

Lemma 12. *Fix $t \geq 0$ and $\alpha \in (0, 1)$. Then,*

$$\forall x \in \{-1, 1\}, \theta \in \mathbb{R}, \quad p_t^\alpha(x, 1) \eta_t^\alpha(1, x; \theta) + p_t^\alpha(x, -1) \eta_t^\alpha(-1, x; \theta) = 0 \quad (28)$$

and

$$\begin{aligned} & \min_{\theta \in \mathbb{R}} \max_{x \in \{-1, 1\}} \left\{ p_t^\alpha(x, 1) \eta_t^\alpha(1, x; \theta)^2 + p_t^\alpha(x, -1) \eta_t^\alpha(-1, x; \theta)^2 \right\} \\ &= \frac{e^{-t}}{(e^t - 1)(\sqrt{\alpha p_t^\alpha(-1, -1)} + \sqrt{(1 - \alpha) p_t^\alpha(1, 1)})^2} \leq \frac{1}{e^t - 1}. \end{aligned} \quad (29)$$

Proof. The centering condition (28) can be checked easily using the explicit formulas (17) and (27) of the matrices. For (29), we compute that for any $\theta \in \mathbb{R}$,

$$\begin{aligned} \max_{x \in \{-1,1\}} \left\{ p_t^\alpha(x,1)\eta_t^\alpha(1,x;\theta)^2 + p_t^\alpha(x,-1)\eta_t^\alpha(-1,x;\theta)^2 \right\} \\ = \max \left\{ \frac{(e^{-t} - \theta)^2}{p_t^\alpha(1,1)p_t^\alpha(1,-1)}, \frac{\theta^2}{p_t^\alpha(-1,1)p_t^\alpha(-1,-1)} \right\}. \end{aligned} \quad (30)$$

As this is the maximum of two quadratic functions in θ , its minimum is attained at the point θ^* where they intersect in the interval $(0, e^{-t})$, namely at

$$\theta^* = \frac{e^{-t} \sqrt{\alpha p_t^\alpha(-1,-1)}}{\sqrt{\alpha p_t^\alpha(-1,-1)} + \sqrt{(1-\alpha) p_t^\alpha(1,1)}}. \quad (31)$$

The first equality in (29) is immediate, whereas for the inequality we compute

$$\begin{aligned} \frac{e^{-t}}{(e^t - 1)(\sqrt{\alpha p_t^\alpha(-1,-1)} + \sqrt{(1-\alpha) p_t^\alpha(1,1)})^2} &\leq \frac{e^{-t}}{(e^t - 1)(\alpha p_t^\alpha(-1,-1) + (1-\alpha) p_t^\alpha(1,1))} \\ &= \frac{e^{-t}}{(e^t - 1)(1 - (1 - e^{-t})(\alpha^2 + (1-\alpha)^2))} \leq \frac{1}{e^t - 1}, \end{aligned} \quad (32)$$

where both inequalities follow from the convexity of $x \mapsto x^2$. \square

The key technical ingredient in the proof of Theorem 1 is the following identity.

Proposition 13. Fix $n \in \mathbb{N}$, $\alpha = (\alpha_1, \dots, \alpha_n) \in (0, 1)^n$, $t \geq 0$ and $\theta_1, \dots, \theta_n \in \mathbb{R}$. Then, for every function $f : \{-1, 1\}^n \rightarrow X$, where X is a vector space, we have

$$\forall x \in \{-1, 1\}^n, \quad \mathcal{L}_\alpha P_t^\alpha f(x) = -\mathbb{E} \left[\sum_{i=1}^n \eta_t^{\alpha_i}(x(i), X_t^\alpha(i); \theta_i) \partial_i^{\alpha_i} f(X_t^\alpha) \mid X_0^\alpha = x \right]. \quad (33)$$

Proof. In view of (23) and the product structure of the process $\{X_t^\alpha\}_{t \geq 0}$, it suffices to check the claim for $n = 1$, namely that for every $\beta \in (0, 1)$, $\theta \in \mathbb{R}$ and $f : \{-1, 1\} \rightarrow X$,

$$\forall x \in \{-1, 1\}, \quad e^{-t} \partial^\beta f(x) = P_t^\beta \partial^\beta f(x) = \mathbb{E} \left[\eta_t^\beta(x, X_t^\beta; \theta) \partial^\beta f(X_t^\beta) \mid X_0^\beta = x \right], \quad (34)$$

where the first equality follows from the probabilistic representation (22). Taking into account that $\beta \partial^\beta f(1) + (1 - \beta) \partial^\beta f(-1) = 0$, this amounts to the system of equations

$$\begin{cases} e^{-t} = p_t^\beta(1,1)\eta_t^\beta(1,1;\theta) - \frac{\beta}{1-\beta} p_t^\beta(1,-1)\eta_t^\beta(1,-1;\theta) \\ e^{-t} = -\frac{1-\beta}{\beta} p_t^\beta(-1,1)\eta_t^\beta(-1,1;\theta) + p_t^\beta(-1,-1)\eta_t^\beta(-1,-1;\theta) \end{cases} \quad (35)$$

which can be easily verified by direct computation. \square

Theorem 14. Fix $p \in [1, 2]$ and let $(X, \|\cdot\|_X)$ be a normed space of Rademacher type p . Then, for any $n \in \mathbb{N}$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in (0, 1)^n$, every function $f : \{-1, 1\}^n \rightarrow X$ satisfies

$$\int_{\{-1,1\}^n} \left\| f(x) - \int_{\{-1,1\}^n} f \, d\mu_\alpha \right\|_X^p \, d\mu_\alpha(x) \leq (2\pi \Gamma_p(X))^p \sum_{i=1}^n \int_{\{-1,1\}^n} \left\| \partial_i^{\alpha_i} f(x) \right\|_X^p \, d\mu_\alpha(x). \quad (36)$$

Proof. Writing

$$f(x) - \int_{\{-1,1\}^n} f \, d\mu_\alpha = P_0^\alpha f(x) - P_\infty^\alpha f(x) = - \int_0^\infty \mathcal{L}_\alpha P_t^\alpha f(x) \, dt \quad (37)$$

and using Jensen's inequality and Proposition 13, we see that for $\theta_1(t), \dots, \theta_n(t) \in \mathbb{R}$,

$$\begin{aligned} & \left(\int_{\{-1,1\}^n} \left\| f(x) - \int_{\{-1,1\}^n} f \, d\mu_\alpha \right\|_X^p d\mu_\alpha(x) \right)^{1/p} \leq \int_0^\infty \left(\int_{\{-1,1\}^n} \left\| \mathcal{L}_\alpha P_t^\alpha f(x) \right\|_X^p d\mu_\alpha(x) \right)^{1/p} dt \\ & = \int_0^\infty \left(\int_{\{-1,1\}^n} \left\| \mathbb{E} \left[\sum_{i=1}^n \eta_t^{\alpha_i}(x(i), X_t^\alpha(i); \theta_i(t)) \partial_i^{\alpha_i} f(X_t^\alpha) \mid X_0^\alpha = x \right] \right\|_X^p d\mu_\alpha(x) \right)^{1/p} dt \quad (38) \\ & \leq \int_0^\infty \left(\mathbb{E} \left\| \sum_{i=1}^n \eta_t^{\alpha_i}(X_0^\alpha(i), X_t^\alpha(i); \theta_i(t)) \partial_i^{\alpha_i} f(X_t^\alpha) \right\|_X^p \right)^{1/p} dt, \end{aligned}$$

where in the last expectation X_0^α is distributed according to μ_α . Now, by the reversibility of the chain, this expectation can be written as

$$\begin{aligned} & \mathbb{E} \left\| \sum_{i=1}^n \eta_t^{\alpha_i}(X_0^\alpha(i), X_t^\alpha(i); \theta_i(t)) \partial_i^{\alpha_i} f(X_t^\alpha) \right\|_X^p \\ & = \int_{\{-1,1\}^n} \sum_{y \in \{-1,1\}^n} p_t^\alpha(x, y) \left\| \sum_{i=1}^n \eta_t^{\alpha_i}(y(i), x(i); \theta_i(t)) \partial_i^{\alpha_i} f(x) \right\|_X^p d\mu_\alpha(x). \end{aligned} \quad (39)$$

Fixing $x \in \{-1,1\}^n$, equation (28) asserts that each $\eta_t^{\alpha_i}(y(i), x(i); \theta_i(t))$ is a centered random variable when $y(i)$ is distributed according to $p_t^{\alpha_i}(x(i), \cdot)$. Therefore, as $p_t^\alpha(x, \cdot)$ is a product measure, the Rademacher type condition for sums of centered independent random vectors (see [33, Proposition 9.11]) yields the bound

$$\begin{aligned} & \int_{\{-1,1\}^n} \sum_{y \in \{-1,1\}^n} p_t^\alpha(x, y) \left\| \sum_{i=1}^n \eta_t^{\alpha_i}(y(i), x(i); \theta_i(t)) \partial_i^{\alpha_i} f(x) \right\|_X^p d\mu_\alpha(x) \\ & \leq (2T_p(X))^p \int_{\{-1,1\}^n} \sum_{i=1}^n \sum_{y(i) \in \{-1,1\}} p_t^{\alpha_i}(x(i), y(i)) \left| \eta_t^{\alpha_i}(y(i), x(i); \theta_i(t)) \right|^p \left\| \partial_i^{\alpha_i} f(x) \right\|_X^p d\mu_\alpha(x) \\ & \leq (2T_p(X))^p \sum_{i=1}^n \int_{\{-1,1\}^n} \left(\sum_{y(i) \in \{-1,1\}} p_t^{\alpha_i}(x(i), y(i)) \left| \eta_t^{\alpha_i}(y(i), x(i); \theta_i(t)) \right|^2 \right)^{p/2} \left\| \partial_i^{\alpha_i} f(x) \right\|_X^p d\mu_\alpha(x), \end{aligned} \quad (40)$$

where we also used that $p \leq 2$. Now, choosing the $\theta_i(t)$ which minimize the quantity in the left-hand side of (29) with bias α_i , and combining (38), (39) and (40), we conclude that

$$\begin{aligned} & \left(\int_{\{-1,1\}^n} \left\| f(x) - \int_{\{-1,1\}^n} f \, d\mu_\alpha \right\|_X^p d\mu_\alpha(x) \right)^{1/p} \\ & \leq 2T_p(X) \int_0^\infty \left(\sum_{i=1}^n \int_{\{-1,1\}^n} \left\| \partial_i^{\alpha_i} f(x) \right\|_X^p d\mu_\alpha(x) \right)^{1/p} \frac{dt}{\sqrt{e^t - 1}}, \end{aligned} \quad (41)$$

which is precisely the desired estimate. \square

Equipped with the biased Poincaré inequality of Theorem 14, we can conclude the proof.

Proof of Theorem 1. Let $X = (\mathbb{R}^d, \|\cdot\|_X)$ be a d -dimensional normed space and suppose that $f : \{-1,1\}^n \rightarrow X$ is a function such that

$$\forall x, y \in \{-1,1\}^n, \quad \rho(x, y) \leq \|f(x) - f(y)\|_X \leq D\rho(x, y) \quad (42)$$

for some $D \geq 1$. The conclusion of the theorem follows from [27] when $d \geq n$ so we shall assume that $d < n$. Let $F : [-1,1]^n \rightarrow X$ be the multilinear extension of f given by (24). Then, F is clearly continuous as a polynomial and therefore, by Lemma 11, there exists a point $z \in \mathcal{C}_d^n$ such that $F(z) = F(-z)$. As z has at least $n - d$ coordinates equal to 1 in absolute value we shall

assume without loss of generality that $|z(d+1)| = \dots = |z(n)| = 1$ and consider the functions $h_+, h_- : \{-1, 1\}^d \rightarrow X$ which are defined as

$$\forall x \in \{-1, 1\}^d, \quad h_{\pm}(x) = f(\pm x(1), \dots, \pm x(d), \pm z(d+1), \dots, \pm z(n)). \quad (43)$$

Consider also the bias vector $\alpha_z = \left(\frac{1+z(1)}{2}, \dots, \frac{1+z(d)}{2}\right) \in (0, 1)^d$ and notice that, by the multilinearity of F , we have the identity

$$\int_{\{-1, 1\}^d} h_+(x) \, d\mu_{\alpha_z}(x) = F(z) = F(-z) = \int_{\{-1, 1\}^d} h_-(x) \, d\mu_{\alpha_z}(x). \quad (44)$$

Therefore, by the triangle inequality and Theorem 14 we get

$$\begin{aligned} & \int_{\{-1, 1\}^d} \|h_+(x) - h_-(x)\|_X^p \, d\mu_{\alpha_z}(x) \\ & \leq 2^{p-1} \int_{\{-1, 1\}^d} \|h_+(x) - F(z)\|_X^p + \|h_-(x) - F(-z)\|_X^p \, d\mu_{\alpha_z}(x) \\ & \leq 2^{2p-1} (\pi \mathbb{T}_p(X))^p \sum_{i=1}^d \int_{\{-1, 1\}^d} \|\partial_i^{\frac{1+z(i)}{2}} h_+(x)\|_X^p + \|\partial_i^{\frac{1+z(i)}{2}} h_-(x)\|_X^p \, d\mu_{\alpha_z}(x). \end{aligned} \quad (45)$$

Now, in view of the lower Lipschitz condition (42), we clearly have

$$\|h_+(x) - h_-(x)\|_X = \|f(x(1), \dots, x(d), z(d+1), \dots, z(n)) - f(-x(1), \dots, -x(d), -z(d+1), \dots, -z(n))\|_X \geq n$$

for every $x \in \{-1, 1\}^d$. On the other hand, for a fixed $i \in \{1, \dots, d\}$ and $\beta = \frac{1+z(i)}{2}$, we have

$$\begin{aligned} & \int_{\{-1, 1\}^d} \|\partial_i^{\beta} h_+(x)\|_X^p \, d\mu_{\beta}(x(i)) = \beta \|\partial_i^{\beta} h_+(x(1), \dots, 1, \dots, x(d))\|_X^p + (1-\beta) \|\partial_i^{\beta} h_+(x(1), \dots, -1, \dots, x(d))\|_X^p \\ & = (\beta(1-\beta)^p + (1-\beta)\beta^p) \|h_+(x(1), \dots, 1, \dots, x(d)) - h_+(x(1), \dots, -1, \dots, x(d))\|_X^p \leq \frac{D^p}{2^p}, \end{aligned}$$

where in the last equality we used that $p \leq 2$ along with the upper Lipschitz condition (42). The same bound also holds for h_- . Integrating the last two inequalities and combining them with (45), we deduce that

$$n^p \leq (2\pi \mathbb{T}_p(X))^p d D^p, \quad (46)$$

which completes the proof of the theorem. \square

Remark 15. *The identity of Proposition 13 in the case of the uniform measure σ_n (which was obtained in [27]) is simpler. Let $\xi_1(t), \dots, \xi_n(t)$ be i.i.d. random variables distributed according to $\mu_{\beta(t)}$, where $\beta(t) = \frac{1+e^{-t}}{2}$. Then, for any point $x \in \{-1, 1\}^n$, the corresponding unbiased process $\{X_t(i)\}_{t \geq 0}$ with $X_0 = x$ has distribution equal to $x(i)\xi_i(t)$ at time t . Thus applying formula (33) with $\alpha_i = \frac{1}{2}$ and $\theta_i(t) = \frac{e^{-t}}{2}$, we recover the usual identity*

$$\forall x \in \{-1, 1\}^n, \quad \mathcal{L}P_t f(x) = -\mathbb{E} \left[\sum_{i=1}^n \frac{\xi_i(t) - e^{-t}}{e^t - e^{-t}} \cdot \partial_i f(x\xi(t)) \right], \quad (47)$$

where $x\xi(t) = (x(1)\xi_1(t), \dots, x(n)\xi_n(t))$, as was proven in [27].

4. PROOF OF THEOREM 2

Recall that a normed space $(X, \|\cdot\|_X)$ has cotype $q \in [2, \infty)$ with constant $C \in (0, \infty)$ if for every $n \in \mathbb{N}$ and $v_1, \dots, v_n \in X$, we have

$$\int_{\{-1, 1\}^n} \left\| \sum_{i=1}^n x_i v_i \right\|_X^q \, d\sigma_n(x) \geq \frac{1}{C^q} \sum_{i=1}^n \|v_i\|_X^q. \quad (48)$$

We say that X has finite cotype if it has cotype q for some $q \in [2, \infty)$.

The proof of Theorem 2 is similar to the arguments of [27, 16] using as input the new identity (33) for the time derivative of the biased semigroup $\{P_t^\alpha\}_{t \geq 0}$. In view of that, we shall omit various simple details and we will be less attentive with the values of the implicit constants.

We start by proving the (weaker) biased Pisier inequality, in which the Orlicz norm on the left hand side of the conclusions of Theorem 2 is replaced by an L_p norm.

Theorem 16. *For every $\alpha \in (0, 1)$, there exists $K_\alpha \in (0, \infty)$ such that the following holds when $p \in [1, \infty)$. For any normed space $(X, \|\cdot\|_X)$ and any $n \in \mathbb{N}$, every function $f : \{-1, 1\}^n \rightarrow X$ satisfies*

$$\left\| f - \int_{\{-1, 1\}^n} f \, d\mu_\alpha^n \right\|_{L_p(\mu_\alpha^n; X)} \leq K_\alpha (\log n + 1) \left(\int_{\{-1, 1\}^n} \left\| \sum_{i=1}^n \delta_i \partial_i^\alpha f \right\|_{L_p(\mu_\alpha^n; X)}^p \, d\sigma_n(\delta) \right)^{1/p}. \quad (49)$$

If additionally X is assumed to be of finite cotype, then there exists $K_{p, \alpha}(X)$ such that

$$\left\| f - \int_{\{-1, 1\}^n} f \, d\mu_\alpha^n \right\|_{L_p(\mu_\alpha^n; X)} \leq K_{p, \alpha}(X) \left(\int_{\{-1, 1\}^n} \left\| \sum_{i=1}^n \delta_i \partial_i^\alpha f \right\|_{L_p(\mu_\alpha^n; X)}^p \, d\sigma_n(\delta) \right)^{1/p}. \quad (50)$$

Proof. Let $\alpha = (\alpha, \dots, \alpha)$ and without loss of generality assume that $\int f \, d\mu_\alpha = 0$. We start by proving the general inequality (49). Notice that, by convexity,

$$\|\mathcal{L}_\alpha f\|_{L_p(\mu_\alpha; X)} \stackrel{(23)}{\leq} \sum_{i=1}^n \|\partial_i^\alpha f\|_{L_p(\mu_\alpha; X)} \leq 2n \|f\|_{L_p(\mu_\alpha; X)}, \quad (51)$$

where the last inequality follows from the definitions of ∂_i^α . Therefore,

$$\forall t \geq 0, \quad \|e^{-t\mathcal{L}_\alpha} f\|_{L_p(\mu_\alpha; X)} \leq \sum_{m=0}^{\infty} \frac{t^m}{m!} \|\mathcal{L}_\alpha^m f\|_{L_p(\mu_\alpha; X)} \leq \sum_{m=0}^{\infty} \frac{(2nt)^m}{m!} \|f\|_{L_p(\mu_\alpha; X)} = e^{2nt} \|f\|_{L_p(\mu_\alpha; X)}.$$

Since $P_t^\alpha = e^{t\mathcal{L}_\alpha}$ for $t \geq 0$, this inequality applied to $P_t^\alpha f$ with $t = \frac{1}{n}$ implies that

$$\|f\|_{L_p(\mu_\alpha; X)} \leq e^2 \|P_{1/n}^\alpha f\|_{L_p(\mu_\alpha; X)}. \quad (52)$$

Similarly to (38) and (39), we thus have

$$\begin{aligned} \|f\|_{L_p(\mu_\alpha; X)} &\leq e^2 \|P_{1/n}^\alpha f\|_{L_p(\mu_\alpha; X)} \\ &\leq e^2 \int_{1/n}^{\infty} \left(\int_{\{-1, 1\}^n} \sum_{y \in \{-1, 1\}^n} p_t^\alpha(x, y) \left\| \sum_{i=1}^n \eta_i^\alpha(y(i), x(i); \theta_i(t)) \partial_i^\alpha f(x) \right\|_X^p \, d\mu_\alpha(x) \right)^{1/p} dt \end{aligned} \quad (53)$$

Fix $t \geq 0$ and $x \in \{-1, 1\}^n$. Since $p_t^\alpha(x, \cdot)$ is a product measure, the centering condition (28) gives

$$\begin{aligned} \sum_{y \in \{-1, 1\}^n} p_t^\alpha(x, y) \left\| \sum_{i=1}^n \eta_i^\alpha(y(i), x(i); \theta_i(t)) \partial_i^\alpha f(x) \right\|_X^p \\ \leq 2^p \int_{\{-1, 1\}^n} \sum_{y \in \{-1, 1\}^n} p_t^\alpha(x, y) \left\| \sum_{i=1}^n \delta_i |\eta_i^\alpha(y(i), x(i); \theta_i(t))| \partial_i^\alpha f(x) \right\|_X^p \, d\sigma_n(\delta). \end{aligned} \quad (54)$$

To prove (49), set $\theta_i(t) = \frac{e^{-t}}{2}$ and observe that by the contraction principle [33, Theorem 4.4], we can further bound this quantity by

$$\begin{aligned} 2^p \sum_{y \in \{-1, 1\}^n} p_t^\alpha(x, y) \int_{\{-1, 1\}^n} \left\| \sum_{i=1}^n \delta_i |\eta_i^\alpha(y(i), x(i); \frac{e^{-t}}{2})| \partial_i^\alpha f(x) \right\|_X^p \, d\sigma_n(\delta) \\ \leq 2^p \max_{\chi, \psi \in \{-1, 1\}} |\eta_t^\alpha(\psi, \chi; \frac{e^{-t}}{2})|^p \int_{\{-1, 1\}^n} \left\| \sum_{i=1}^n \delta_i \partial_i^\alpha f(x) \right\|_X^p \, d\sigma_n(\delta). \end{aligned} \quad (55)$$

It is now elementary to check that we have

$$\max_{\chi, \psi \in \{-1, 1\}} \left| \eta_t^\alpha \left(\psi, \chi; \frac{e^{-t}}{2} \right) \right| \lesssim_\alpha \begin{cases} \frac{1}{t}, & \text{for } t \in (0, 1) \\ e^{-t}, & \text{for } t \geq 1 \end{cases} \quad (56)$$

and thus combining all the above we get

$$\|f\|_{L_p(\mu_\alpha; X)} \lesssim_\alpha \left(\int_{1/n}^1 \frac{dt}{t} + \int_1^\infty e^{-t} dt \right) \cdot \left(\int_{\{-1, 1\}^n} \left\| \sum_{i=1}^n \delta_i \partial_i f \right\|_{L_p(\mu_\alpha; X)}^p d\sigma_n(\delta) \right)^{1/p}, \quad (57)$$

which readily concludes the proof of (49).

For the proof of (50), we combine (38) and (39) with (54) to get

$$\|f\|_{L_p(\mu_\alpha; X)} \leq 2 \int_0^\infty \left(\int_{\{-1, 1\}^{2n}} \sum_{y \in \{-1, 1\}^n} p_t^\alpha(x, y) \left\| \sum_{i=1}^n \delta_i \left| \eta_t^\alpha \left(y(i), x(i); \frac{e^{-t}}{2} \right) \right| \partial_i^\alpha f(x) \right\|_X^p d\sigma_n(\delta) d\mu_\alpha(x) \right)^{1/p} dt.$$

Fix $x \in \{-1, 1\}^n$ and $t \geq 0$. Denoting by $I(x) = \{i : x(i) = 1\}$, the inner term is bounded above by

$$\begin{aligned} & \int_{\{-1, 1\}^n} \sum_{y \in \{-1, 1\}^n} p_t^\alpha(x, y) \left\| \sum_{i=1}^n \delta_i \left| \eta_t^\alpha \left(y(i), x(i); \frac{e^{-t}}{2} \right) \right| \partial_i^\alpha f(x) \right\|_X^p d\sigma_n(\delta) \\ & \leq 2^{p-1} \int_{\{-1, 1\}^{I(x)}} \sum_{y \in \{-1, 1\}^{I(x)}} p_t^\alpha(x, y) \left\| \sum_{i \in I(x)} \delta_i \left| \eta_t^\alpha \left(y(i), 1; \frac{e^{-t}}{2} \right) \right| \partial_i^\alpha f(x) \right\|_X^p d\sigma_{I(x)}(\delta) \\ & \quad + 2^{p-1} \int_{\{-1, 1\}^{I(x)^c}} \sum_{y \in \{-1, 1\}^{I(x)^c}} p_t^\alpha(x, y) \left\| \sum_{i \notin I(x)} \delta_i \left| \eta_t^\alpha \left(y(i), -1; \frac{e^{-t}}{2} \right) \right| \partial_i^\alpha f(x) \right\|_X^p d\sigma_{I(x)^c}(\delta). \end{aligned} \quad (58)$$

When $y \in \{-1, 1\}^{I(x)}$ is distributed according to $p_t^\alpha(x, \cdot)$ and δ is a uniformly random sign, the random variables $\delta_i \left| \eta_t^\alpha \left(y(i), 1; \frac{e^{-t}}{2} \right) \right|$, $i \in I(x)$, are independent and identically distributed. Therefore, standard comparison principles going back to works of Maurey and Pisier (see, e.g., [33, Proposition 9.14]) show that if X has cotype $q < \infty$ and $r > \max\{p, q\}$, then

$$\begin{aligned} & \int_{\{-1, 1\}^{I(x)}} \sum_{y \in \{-1, 1\}^{I(x)}} p_t^\alpha(x, y) \left\| \sum_{i \in I(x)} \delta_i \left| \eta_t^\alpha \left(y(i), 1; \frac{e^{-t}}{2} \right) \right| \partial_i^\alpha f(x) \right\|_X^p d\sigma_{I(x)}(\delta) \\ & \lesssim_{r, X} \left(\sum_{\psi \in \{-1, 1\}} p_t^\alpha \left(1, \psi; \frac{e^{-t}}{2} \right) \left| \eta_t^\alpha \left(\psi, 1; \frac{e^{-t}}{2} \right) \right|^r \right)^{p/r} \int_{\{-1, 1\}^{I(x)}} \left\| \sum_{i \in I(x)} \delta_i \partial_i^\alpha f(x) \right\|_X^p d\sigma_{I(x)}(\delta). \end{aligned} \quad (59)$$

It is again elementary to show that

$$\max_{\chi \in \{-1, 1\}} \left(\sum_{\psi \in \{-1, 1\}} p_t^\alpha \left(\chi, \psi; \frac{e^{-t}}{2} \right) \left| \eta_t^\alpha \left(\psi, \chi; \frac{e^{-t}}{2} \right) \right|^r \right)^{1/r} \lesssim_{r, \alpha} \begin{cases} \frac{1}{t^{1-\frac{1}{r}}}, & \text{for } t \in (0, 1) \\ e^{-t}, & \text{for } t \geq 1 \end{cases}. \quad (60)$$

In view of the convergence of the integrals

$$\int_0^1 \frac{dt}{t^{1-\frac{1}{r}}} + \int_1^\infty e^{-t} dt \lesssim_r 1, \quad (61)$$

combining (59), the corresponding estimate on $I(x)^c$ and (58), we finally get

$$\|f\|_{L_p(\mu_\alpha; X)} \lesssim_{r, \alpha, X} \left(\int_{\{-1, 1\}^n} \int_{\{-1, 1\}^n} \left\| \sum_{i \in I(x)} \delta_i \partial_i^\alpha f(x) \right\|_X^p + \left\| \sum_{i \notin I(x)} \delta_i \partial_i^\alpha f(x) \right\|_X^p d\sigma_n(\delta) d\mu_\alpha(x) \right)^{1/p}.$$

Now, if F, G are two independent centered X -valued random vectors, then

$$\mathbb{E}\|F + G\|_X^p \geq \max\{\mathbb{E}\|F\|_X^p + \mathbb{E}\|G\|_X^p, \mathbb{E}_G\|\mathbb{E}_F[F] + G\|_X^p\} = \max\{\mathbb{E}\|F\|_X^p, \mathbb{E}\|G\|_X^p\}. \quad (62)$$

Therefore,

$$\begin{aligned} & \left(\int_{\{-1,1\}^n} \int_{\{-1,1\}^n} \left\| \sum_{i \in I(x)} \delta_i \partial_i^\alpha f(x) \right\|_X^p + \left\| \sum_{i \notin I(x)} \delta_i \partial_i^\alpha f(x) \right\|_X^p d\sigma_n(\delta) d\mu_{\mathbf{a}}(x) \right)^{1/p} \\ & \lesssim \left(\int_{\{-1,1\}^n} \left\| \sum_{i=1}^n \delta_i \partial_i^\alpha f(x) \right\|_{L_p(\mu_{\mathbf{a}}; X)}^p d\sigma_n(\delta) \right)^{1/p} \end{aligned} \quad (63)$$

and this concludes the proof of the theorem. \square

We now proceed to discuss the vector-valued L_p logarithmic Sobolev inequality of Theorem 2. Recall that given a function $f : (\Omega, \mu) \rightarrow X$, we define the $L_p(\log L)^a$ Orlicz norm of f as

$$\|f\|_{L_p(\log L)^a(\mu; X)} = \inf \left\{ \gamma > 0 : \int_{\Omega} \frac{\|f(\omega)\|_X^p}{\gamma^p} \log^a \left(e + \frac{\|f(\omega)\|_X^p}{\gamma^p} \right) d\mu(\omega) \leq 1 \right\}. \quad (64)$$

In the case of the unbiased cube, Theorem 2 was proven recently in [16] and the proof used as a black box the unbiased analogue of Theorem 16 which is due to [64, 27]. As the modifications which are required to derive Theorem 2 from Theorem 16 in the biased case are almost mechanical, we shall only offer a high-level description of the proof.

Sketch of the proof of Theorem 2. Given a scalar-valued function $h : \{-1, 1\}^n \rightarrow \mathbb{R}$, we denote by $Mh : \{-1, 1\}^n \rightarrow \mathbb{R}_+$ the asymmetric gradient of h , given by

$$\forall x \in \{-1, 1\}^n, \quad Mh(x) = \left(\sum_{i=1}^n \partial_i h(x)_+^2 \right)^{1/2}, \quad (65)$$

where $a_+ = \max\{a, 0\}$ for $a \in \mathbb{R}$. A combination of the triangle inequality with a technical result of Talagrand [67, Proposition 5.1] on the biased cube implies that if a vector-valued function $f : \{-1, 1\}^n \rightarrow X$ satisfies $\int f d\mu_{\mathbf{a}}^n = 0$, then

$$\|f\|_{L_p(\log L)^{p/2}(\mu_{\mathbf{a}}^n; X)} \lesssim_{p, \alpha} \|M\|f\|_X\|_{L_p(\mu_{\mathbf{a}}^n; X)} + \|f\|_{L_1(\mu_{\mathbf{a}}^n; X)}. \quad (66)$$

The second term can be controlled by the right hand side of (9) or (10) using the results of Theorem 16. For the first term, we use the key pointwise inequality of [16], asserting that

$$\forall x \in \{-1, 1\}^n, \quad M\|f\|_X(x) \leq \sqrt{2} \left(\int_{\{-1,1\}^n} \left\| \sum_{i=1}^n \delta_i \partial_i f(x) \right\|_X^p d\sigma_n(\delta) \right)^{1/p}. \quad (67)$$

To further upper bound this by the square function where ∂_i is replaced by ∂_i^α , observe that

$$\partial_i^\alpha f(x) \in \{2\alpha \partial_i f(x), 2(1-\alpha) \partial_i f(x)\} \quad (68)$$

and thus the contraction principle and integration in x yield the desired conclusions. \square

5. PROOF OF THEOREM 4

In this section we shall prove the equivalence of stable type and metric stable type for norms. The key ingredient in the proof is the following characterization of stable type (see, e.g., [33, Proposition 9.12 (iii)]) which goes back at least to [40].

Lemma 17. *A Banach space $(X, \|\cdot\|_X)$ has stable type $p \in [1, 2)$ if and only if there exists a constant $ST_p(X) \in (0, \infty)$ such that for every $n \in \mathbb{N}$ and every vectors $v_1, \dots, v_n \in X$, we have*

$$\int_{\{-1,1\}^n} \left\| \sum_{i=1}^n x_i v_i \right\|_X^p d\sigma_n(x) \leq ST_p(X)^p \left(\|v_1\|_X, \dots, \|v_n\|_X \right)_{\ell_{p, \infty}^n}^p. \quad (69)$$

Proof of Theorem 4. It is clear from Lemma 17 that any normed space with metric stable type p also has stable type p . For the converse implication, we treat the case $p = 1$ separately. Assuming that X has stable type 1, it follows from [59] that it also has nontrivial Rademacher type and thus (in view of [57]) finite cotype. Therefore, by the X -valued Pisier inequality with a dimension-free constant of [27], we get

$$\begin{aligned} \int_{\{-1,1\}^n} \|f(x) - f(-x)\|_X \, d\sigma_n(x) &\leq 2 \int_{\{-1,1\}^n} \left\| f(x) - \int_{\{-1,1\}^n} f \, d\sigma_n \right\|_X \, d\sigma_n(x) \\ &\lesssim_X \int_{\{-1,1\}^n} \int_{\{-1,1\}^n} \mathbb{E} \left\| \sum_{i=1}^n \delta_i \partial_i f(x) \right\|_X \, d\sigma_n(\delta) \, d\sigma_n(x). \end{aligned} \quad (70)$$

Applying Lemma 17 conditionally on $x \in \{-1, 1\}^n$, we can further bound this quantity as

$$\begin{aligned} \int_{\{-1,1\}^n} \int_{\{-1,1\}^n} \mathbb{E} \left\| \sum_{i=1}^n \delta_i \partial_i f(x) \right\|_X \, d\sigma_n(\delta) \, d\sigma_n(x) \\ \leq \text{ST}_1(X) \int_{\{-1,1\}^n} \left\| (\|\partial_1 f(x)\|_X, \dots, \|\partial_n f(x)\|_X) \right\|_{\ell_{1,\infty}^n} \, d\sigma_n(x) \end{aligned} \quad (71)$$

and this proves the converse implication since $\mathfrak{d}_i f(x) = \|\partial_i f(x)\|_X$.

While this proof extends to all values of p , in the case $p > 1$ we present a more cumbersome argument which avoids the X -valued Pisier inequality and thus gives better dependence on parameters of X . For $t \geq 0$, let $\xi_1(t), \dots, \xi_n(t)$ be i.i.d. random variables distributed according to $\mu_{\beta(t)}$, where $\beta(t) = \frac{1+e^{-t}}{2}$, and denote by $\eta_i(t) = \frac{\xi_i(t) - e^{-t}}{e^t - e^{-t}}$. Then, it follows from the semigroup argument leading to (38) along with identity (47) of [27] and Jensen's inequality that

$$\begin{aligned} \left(\int_{\{-1,1\}^n} \|f(x) - f(-x)\|_X^p \, d\sigma_n(x) \right)^{1/p} &\leq 2 \left(\int_{\{-1,1\}^n} \left\| f(x) - \int_{\{-1,1\}^n} f \, d\sigma_n \right\|_X^p \, d\sigma_n(x) \right)^{1/p} \\ &\leq 2 \int_0^\infty \left(\int_{\{-1,1\}^n} \mathbb{E} \left\| \sum_{i=1}^n \eta_i(t) \partial_i f(x) \right\|_X^p \, d\sigma_n(x) \right)^{1/p} \, dt \end{aligned} \quad (72)$$

Fixing $x \in \{-1, 1\}^n$, the independent random vectors $\eta_1(t) \partial_1 f(x), \dots, \eta_n(t) \partial_n f(x)$ are centered. Thus, by standard symmetrization estimates, we can further bound the last term by

$$\begin{aligned} 2 \int_0^\infty \left(\int_{\{-1,1\}^n} \mathbb{E} \left\| \sum_{i=1}^n \eta_i(t) \partial_i f(x) \right\|_X^p \, d\sigma_n(x) \right)^{1/p} \, dt \\ \leq 4 \int_0^\infty \left(\int_{\{-1,1\}^n} \mathbb{E} \left\| \sum_{i=1}^n \delta_i \eta_i(t) \partial_i f(x) \right\|_X^p \, d\sigma_n(x) \right)^{1/p} \, dt, \end{aligned} \quad (73)$$

where the expectation on the right hand side is with respect to $(\delta_i, \eta_i(t))$, where the δ_i are uniformly random signs, independent of the $\eta_i(t)$. Therefore, conditioning first on the values of $\eta_1(t), \dots, \eta_n(t)$ and using Lemma 17 for the integrand, we get

$$\mathbb{E} \left\| \sum_{i=1}^n \delta_i \eta_i(t) \partial_i f(x) \right\|_X^p \leq \text{ST}_p(X)^p \mathbb{E} \left\| \sum_{i=1}^n \eta_i(t) \|\partial_i f(x)\|_X e_i \right\|_{\ell_{p,\infty}^n}^p, \quad (74)$$

where $\{e_1, \dots, e_n\}$ is the standard basis of \mathbb{R}^n . It is well-known that, since $p > 1$, $\ell_{p,\infty}^n$ is isomorphic to a normed space up to constants depending only on p . Moreover, it has finite cotype with a constant independent of n (for instance because it can be realized as a real interpolation space between ℓ_1^n and ℓ_2^n , see [11, Section 5.3]). Therefore, using again the comparison

principle [33, Proposition 9.14], we have

$$\begin{aligned} \mathbb{E} \left\| \sum_{i=1}^n \eta_i(t) \|\partial_i f(x)\|_X e_i \right\|_{\ell_{p,\infty}^n}^p &\lesssim_p \|\eta_1(t)\|_{L_r}^p \mathbb{E} \left\| \sum_{i=1}^n \delta_i \|\partial_i f(x)\|_X e_i \right\|_{\ell_{p,\infty}^n}^p \\ &= \|\eta_1(t)\|_{L_r}^p \left\| (\|\partial_1 f(x)\|_X, \dots, \|\partial_n f(x)\|_X) \right\|_{\ell_{p,\infty}^n}^p \end{aligned} \quad (75)$$

for any $r > \max\{r', p\}$, where r' is the cotype of $\ell_{p,\infty}^n$. Combining all the above along with the fact that $t \mapsto \|\eta_1(t)\|_{L_r}$ is integrable for any $r < \infty$, we conclude the proof. It is worth emphasizing that this proof shows additionally that the metric stable type constant of X is proportional to the parameter $\text{ST}_p(X)$ of (69) up to constants depending only on p , provided that $p > 1$. \square

Finally, we present the simple proof of the distortion bound (16).

Proof of Proposition 6. Suppose that (M, d_M) has metric stable type p with constant S and suppose that a function $f : \{-1, 1\}^n \rightarrow M$ satisfies

$$\forall x, y \in \{-1, 1\}^n, \quad s\rho_{\mathbf{w}}(x, y) \leq d_M(f(x), f(y)) \leq sD\rho_{\mathbf{w}}(x, y) \quad (76)$$

for some constants $s > 0$ and $D \geq 1$. Combined with the stable type assumption, this gives

$$\begin{aligned} s^p \|\mathbf{w}\|_{\ell_1^n}^p &\stackrel{(76)}{\leq} \int_{\{-1, 1\}^n} d_M(f(x), f(-x))^p d\sigma_n(x) \leq S^p \int_{\{-1, 1\}^n} \left\| (\partial_1 f(x), \dots, \partial_n f(x)) \right\|_{\ell_{p,\infty}^n}^p d\sigma_n(x) \\ &\stackrel{(76)}{\leq} (sSD)^p \int_{\{-1, 1\}^n} \|(w_1, \dots, w_n)\|_{\ell_{p,\infty}^n}^p d\sigma_n(x) = (sSD)^p \|\mathbf{w}\|_{\ell_{p,\infty}^n}^p. \end{aligned} \quad (77)$$

Rearranging gives the desired lower bound (16) for the distortion D . \square

6. DISCUSSION AND OPEN PROBLEMS

1. The proof of Theorem 1 in fact implies a Poincaré-type inequality for *restrictions* of functions $f : \{-1, 1\}^n \rightarrow X$ if $\dim(X) < n$, which in turn yields the refined distortion lower bounds. An inspection of the argument reveals that for every such f there exists a subset $\sigma \subseteq \{1, \dots, n\}$ with $|\sigma| \leq \dim(X)$, a point $w \in \{-1, 1\}^{\sigma^c}$ and a bias vector $\alpha = (\alpha_i)_{i \in \sigma} \in (0, 1)^\sigma$ such that

$$\begin{aligned} \int_{\{-1, 1\}^\sigma} \|f(x, w) - f(-x, -w)\|_X^p d\mu_\alpha(x) \\ \leq 2^{2p-1} (\pi T_p(X))^p \sum_{i \in \sigma} \int_{\{-1, 1\}^\sigma} \|\partial_i^{\alpha_i} f(x, w)\|_X^p + \|\partial_i^{\alpha_i} f(-x, -w)\|_X^p d\mu_\alpha(x). \end{aligned} \quad (78)$$

2. Such refinements of Poincaré-type inequalities for topological reasons had not been exploited since Oleszkiewicz's original work [56]. The last decades have seen the development of many metric inequalities on graphs which yield nonembeddability results into normed spaces. We believe that investigating whether the distortion estimates which one obtains this way can be further improved assuming upper bounds for the dimension of the target space is a very worthwhile research program. As examples, we mention the nonembeddability of graphs with large girth into uniformly smooth spaces [39, 51], of ℓ_∞ -grids into spaces of finite cotype [45] and of trees [12, 36] and diamond graphs [31, 23] into uniformly convex spaces.

3. The results of [27] in fact imply that any Lipschitz embedding of $\{-1, 1\}^n$ into a normed space of Rademacher type p incurs p -average distortion at least a constant multiple of $T_p(X)^{-1} n^{1-1/p}$. It would be interesting to understand whether the bound of Theorem 1 can be extended to average distortion embeddings beyond bi-Lipschitz ones.

4. The Poincaré inequality (78) implies that any $f : \{-1, 1\}^n \rightarrow X = (\mathbb{R}^d, \|\cdot\|_X)$ satisfies

$$\sup_{(x,y): \rho(x,y)=n} \frac{n}{\|f(x) - f(y)\|_X} \sup_{(x,y): \rho(x,y)=1} \|f(x) - f(y)\|_X \gtrsim \frac{n}{T_p(X) d^{1/p}}, \quad (79)$$

provided that $d < n$. This is stronger than the lower bound of Theorem 1 as it implies that f incurs large distortion only on *specific* pairs of points, namely those which are either antipodal or connected by an edge. Moreover though, (79) is sharp for any value of d and p . Indeed, assume without loss of generality that $\frac{n}{d}$ is an odd integer and consider a partition I_1, \dots, I_d of $\{1, \dots, n\}$ in d parts of size $\frac{n}{d}$. Then, the mapping $f : \{-1, 1\}^n \rightarrow \ell_p^d$, where $p \in [1, 2]$, given by

$$\forall x \in \{-1, 1\}^n, \quad f(x) \stackrel{\text{def}}{=} \left(\sum_{i \in I_1} x_i, \dots, \sum_{i \in I_d} x_i \right) \quad (80)$$

satisfies

$$\inf_{(x,y): \rho(x,y)=n} \|f(x) - f(y)\|_{\ell_p^d} = \inf_{x \in \{-1,1\}^n} \left(\sum_{k=1}^d \left| \sum_{i \in I_k} x_i \right|^p \right)^{1/p} = d^{1/p} \quad (81)$$

and $\|f(x) - f(y)\|_{\ell_p^d} = 1$ when $\rho(x, y) = 1$.

5. The functional inequality (78) in fact implies that if $\theta \in (0, 1)$, then the bi-Lipschitz distortion of the θ -snowflake of $\{-1, 1\}^n$ into a finite dimensional normed space X satisfies

$$c_X(\{-1, 1\}^n, \rho^\theta) \gtrsim \frac{n^\theta}{T_p(X) \min\{n, \dim(X)\}^{1/p}}. \quad (82)$$

6. As the reasons behind the impossibility of dimension reduction of Theorem 1 are partly topological, it is natural to ask in what other settings can one deduce similar conclusions. For instance, do d -dimensional manifolds of nonpositive or nonnegative curvature (which have Enflo type 2, see [55]) admit the same distortion bounds as normed spaces of type 2?

7. In this paper we extended the semigroup machinery developed in [27] to the biased cube, driven by the geometric application obtained in Theorem 1. As an aside, this led to biased versions of the vector-valued Poincaré and logarithmic Sobolev inequalities of [27, 16]. The same reasoning also yields biased extensions of the vector-valued versions of Talagrand's influence inequality [68] which were studied in [17] for the uniform probability measure on the hypercube. Moreover, combining the biased semigroup machinery with the arguments of [10, Section 2.5], one can derive biased versions of the isoperimetric-type inequalities of Eldan and Gross [18]. As these extensions are mostly mechanical given the material of this paper and in lack of a specific application which may follow from them, we omit them.

REFERENCES

- [1] Alexandr Andoni, Moses S. Charikar, Ofer Neiman, and Huy L. Nguyen. Near linear lower bound for dimension reduction in ℓ_1 . In *2011 IEEE 52nd Annual Symposium on Foundations of Computer Science—FOCS 2011*, pages 315–323. IEEE Computer Soc., Los Alamitos, CA, 2011.
- [2] Alexandr Andoni, Piotr Indyk, and Ilya Razenshteyn. Approximate nearest neighbor search in high dimensions. In *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. IV. Invited lectures*, pages 3287–3318. World Sci. Publ., Hackensack, NJ, 2018.
- [3] Juan Arias-de-Reyna and Luis Rodríguez-Piazza. Finite metric spaces needing high dimension for Lipschitz embeddings in Banach spaces. *Israel J. Math.*, 79(1):103–111, 1992.
- [4] Keith Ball. The Ribe programme. *Astérisque*, (352):Exp. No. 1047, viii, 147–159, 2013. Séminaire Bourbaki. Vol. 2011/2012. Exposés 1043–1058.
- [5] Keith Ball, Eric A. Carlen, and Elliott H. Lieb. Sharp uniform convexity and smoothness inequalities for trace norms. *Invent. Math.*, 115(3):463–482, 1994.
- [6] Imre Bárány and Victor S. Grinberg. On some combinatorial questions in finite-dimensional spaces. *Linear Algebra Appl.*, 41:1–9, 1981.
- [7] Yair Bartal, Lee-Ad Gottlieb, and Ofer Neiman. On the impossibility of dimension reduction for doubling subsets of ℓ_p . *SIAM J. Discrete Math.*, 29(3):1207–1222, 2015.
- [8] Yair Bartal and Manor Mendel. Dimension reduction for ultrametrics. In *Proceedings of the Fifteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 664–665. ACM, New York, 2004.
- [9] Florent Baudier, Krzysztof Świąćki, and Andrew Swift. No dimension reduction for doubling subsets of ℓ_q when $q > 2$ revisited. *J. Math. Anal. Appl.*, 504(2):Paper No. 125407, 18, 2021.

- [10] David Beltran, Paata Ivanisvili, and José Madrid. On sharp isoperimetric inequalities on the hypercube. Preprint available at <https://arxiv.org/abs/2303.06738>, 2023.
- [11] Jöran Bergh and Jörgen Löfström. *Interpolation spaces. An introduction*, volume No. 223 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin-New York, 1976.
- [12] Jean Bourgain. The metrical interpretation of superreflexivity in Banach spaces. *Israel J. Math.*, 56(2):222–230, 1986.
- [13] Jean Bourgain, Vitali Milman, and Haim Wolfson. On type of metric spaces. *Trans. Amer. Math. Soc.*, 294(1):295–317, 1986.
- [14] Bo Brinkman and Moses Charikar. On the impossibility of dimension reduction in l_1 . *J. ACM*, 52(5):766–788, 2005.
- [15] Bo Brinkman, Adriana Karagiozova, and James R. Lee. Vertex cuts, random walks, and dimension reduction in series-parallel graphs. In *STOC'07—Proceedings of the 39th Annual ACM Symposium on Theory of Computing*, pages 621–630. ACM, New York, 2007.
- [16] Dario Cordero-Erausquin and Alexandros Eskenazis. Discrete logarithmic Sobolev inequalities in Banach spaces. Preprint available at <https://arxiv.org/abs/2304.03878>, 2023.
- [17] Dario Cordero-Erausquin and Alexandros Eskenazis. Talagrand’s influence inequality revisited. *Anal. PDE*, 16(2):571–612, 2023.
- [18] Ronen Eldan and Renan Gross. Concentration on the Boolean hypercube via pathwise stochastic analysis. *Invent. Math.*, 230(3):935–994, 2022.
- [19] Per Enflo. On a problem of Smirnov. *Ark. Mat.*, 8:107–109, 1969.
- [20] Per Enflo. On the nonexistence of uniform homeomorphisms between L_p -spaces. *Ark. Mat.*, 8:103–105, 1969.
- [21] Per Enflo. On infinite-dimensional topological groups. In *Séminaire sur la Géométrie des Espaces de Banach (1977–1978)*, pages Exp. No. 10–11, 11. École Polytech., Palaiseau, 1978.
- [22] Alexandros Eskenazis. On Pisier’s inequality for UMD targets. *Canad. Math. Bull.*, 64(2):282–291, 2021.
- [23] Alexandros Eskenazis, Manor Mendel, and Assaf Naor. Diamond convexity: a bifurcation in the Ribe program. Preprint, 2023.
- [24] Anupam Gupta, Robert Krauthgamer, and James R. Lee. Bounded geometries, fractals, and low-distortion embeddings. In *44th Annual IEEE Symposium on Foundations of Computer Science, 2003. Proceedings.*, pages 534–543, 2003.
- [25] Tuomas Hytönen and Assaf Naor. Pisier’s inequality revisited. *Studia Math.*, 215(3):221–235, 2013.
- [26] Piotr Indyk. Algorithmic applications of low-distortion geometric embeddings. In *42nd IEEE Symposium on Foundations of Computer Science (Las Vegas, NV, 2001)*, pages 10–33. IEEE Computer Soc., Los Alamitos, CA, 2001.
- [27] Paata Ivanisvili, Ramon van Handel, and Alexander Volberg. Rademacher type and Enflo type coincide. *Ann. of Math. (2)*, 192(2):665–678, 2020.
- [28] Robert C. James. Nonreflexive spaces of type 2. *Israel J. Math.*, 30(1-2):1–13, 1978.
- [29] William B. Johnson and Joram Lindenstrauss. Extensions of Lipschitz mappings into a Hilbert space. In *Conference in modern analysis and probability (New Haven, Conn., 1982)*, volume 26 of *Contemp. Math.*, pages 189–206. Amer. Math. Soc., Providence, RI, 1984.
- [30] William B. Johnson, Joram Lindenstrauss, and Gideon Schechtman. On Lipschitz embedding of finite metric spaces in low-dimensional normed spaces. In *Geometrical aspects of functional analysis (1985/86)*, volume 1267 of *Lecture Notes in Math.*, pages 177–184. Springer, Berlin, 1987.
- [31] William B. Johnson and Gideon Schechtman. Diamond graphs and super-reflexivity. *J. Topol. Anal.*, 1(2):177–189, 2009.
- [32] Vincent Lafforgue and Assaf Naor. A doubling subset of L_p for $p > 2$ that is inherently infinite dimensional. *Geom. Dedicata*, 172:387–398, 2014.
- [33] Michel Ledoux and Michel Talagrand. *Probability in Banach spaces*, volume 23 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1991. Isoperimetry and processes.
- [34] James R. Lee, Manor Mendel, and Assaf Naor. Metric structures in L_1 : dimension, snowflakes, and average distortion. *European J. Combin.*, 26(8):1180–1190, 2005.
- [35] James R. Lee and Assaf Naor. Embedding the diamond graph in L_p and dimension reduction in L_1 . *Geom. Funct. Anal.*, 14(4):745–747, 2004.
- [36] James R. Lee, Assaf Naor, and Yuval Peres. Trees and Markov convexity. *Geom. Funct. Anal.*, 18(5):1609–1659, 2009.
- [37] David A. Levin and Yuval Peres. *Markov chains and mixing times*. American Mathematical Society, Providence, RI, second edition, 2017. With contributions by Elizabeth L. Wilmer, With a chapter on “Coupling from the past” by James G. Propp and David B. Wilson.
- [38] Nathan Linial. Finite metric-spaces—combinatorics, geometry and algorithms. In *Proceedings of the International Congress of Mathematicians, Vol. III (Beijing, 2002)*, pages 573–586. Higher Ed. Press, Beijing, 2002.
- [39] Nathan Linial, Avner Magen, and Assaf Naor. Girth and Euclidean distortion. *Geom. Funct. Anal.*, 12(2):380–394, 2002.

- [40] Michael B. Marcus and Gilles Pisier. Characterizations of almost surely continuous p -stable random Fourier series and strongly stationary processes. *Acta Math.*, 152(3-4):245–301, 1984.
- [41] Jiří Matoušek. Note on bi-Lipschitz embeddings into normed spaces. *Comment. Math. Univ. Carolin.*, 33(1):51–55, 1992.
- [42] Jiří Matoušek. On the distortion required for embedding finite metric spaces into normed spaces. *Israel J. Math.*, 93:333–344, 1996.
- [43] Jiří Matoušek. *Using the Borsuk-Ulam theorem*. Universitext. Springer-Verlag, Berlin, 2003. Lectures on topological methods in combinatorics and geometry, Written in cooperation with Anders Björner and Günter M. Ziegler.
- [44] Bernard Maurey and Gilles Pisier. Séries de variables aléatoires vectorielles indépendantes et propriétés géométriques des espaces de Banach. *Studia Math.*, 58(1):45–90, 1976.
- [45] Manor Mendel and Assaf Naor. Metric cotype. *Ann. of Math. (2)*, 168(1):247–298, 2008.
- [46] Assaf Naor. An introduction to the Ribe program. *Jpn. J. Math.*, 7(2):167–233, 2012.
- [47] Assaf Naor. Comparison of metric spectral gaps. *Anal. Geom. Metr. Spaces*, 2(1):1–52, 2014.
- [48] Assaf Naor. A spectral gap precludes low-dimensional embeddings. In *33rd International Symposium on Computational Geometry*, volume 77 of *LIPICs. Leibniz Int. Proc. Inform.*, pages Art. No. 50, 16. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2017.
- [49] Assaf Naor. Metric dimension reduction: a snapshot of the Ribe program. In *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. I. Plenary lectures*, pages 759–837. World Sci. Publ., Hackensack, NJ, 2018.
- [50] Assaf Naor. An average John theorem. *Geom. Topol.*, 25(4):1631–1717, 2021.
- [51] Assaf Naor, Yuval Peres, Oded Schramm, and Scott Sheffield. Markov chains in smooth Banach spaces and Gromov-hyperbolic metric spaces. *Duke Math. J.*, 134(1):165–197, 2006.
- [52] Assaf Naor, Gilles Pisier, and Gideon Schechtman. Impossibility of dimension reduction in the nuclear norm. *Discrete Comput. Geom.*, 63(2):319–345, 2020.
- [53] Assaf Naor and Gideon Schechtman. Remarks on non linear type and Pisier’s inequality. *J. Reine Angew. Math.*, 552:213–236, 2002.
- [54] Assaf Naor and Robert Young. Foliated corona decompositions. *Acta Math.*, 229(1):55–200, 2022.
- [55] Shin-Ichi Ohta. Markov type of Alexandrov spaces of non-negative curvature. *Mathematika*, 55(1-2):177–189, 2009.
- [56] Krzysztof Oleszkiewicz. On a discrete version of the antipodal theorem. *Fund. Math.*, 151(2):189–194, 1996.
- [57] Gilles Pisier. Sur les espaces de Banach qui ne contiennent pas uniformément de l_n^1 . *C. R. Acad. Sci. Paris Sér. A-B*, 277:A991–A994, 1973.
- [58] Gilles Pisier. “Type” des espaces normés. In *Séminaire Maurey-Schwartz 1973–1974: Espaces L^p , applications radonifiantes et géométrie des espaces de Banach, Exp. No. 3,* pages 12 pp. (errata, p. E.1). „ 1974.
- [59] Gilles Pisier. Une propriété du type p -stable. In *Séminaire Maurey-Schwartz 1973–1974: Espaces L^p , applications radonifiantes et géométrie des espaces de Banach,* pages Exp. No. 8, 10 pp. (errata, p. E.1). École Polytech., Paris, 1974.
- [60] Gilles Pisier. Martingales with values in uniformly convex spaces. *Israel J. Math.*, 20(3-4):326–350, 1975.
- [61] Gilles Pisier. Un exemple concernant la super-réflexivité. In *Séminaire Maurey-Schwartz 1974–1975: Espaces L^p applications radonifiantes et géométrie des espaces de Banach, Annexe No. 2,* page 12. „ 1975.
- [62] Gilles Pisier. Holomorphic semigroups and the geometry of Banach spaces. *Ann. of Math. (2)*, 115(2):375–392, 1982.
- [63] Gilles Pisier. On the dimension of the l_p^n -subspaces of Banach spaces, for $1 \leq p < 2$. *Trans. Amer. Math. Soc.*, 276(1):201–211, 1983.
- [64] Gilles Pisier. Probabilistic methods in the geometry of Banach spaces. In *Probability and analysis (Varenna, 1985)*, volume 1206 of *Lecture Notes in Math.*, pages 167–241. Springer, Berlin, 1986.
- [65] Gilles Pisier and Quanhua Xu. Random series in the real interpolation spaces between the spaces v_p . In *Geometrical aspects of functional analysis (1985/86)*, volume 1267 of *Lecture Notes in Math.*, pages 185–209. Springer, Berlin, 1987.
- [66] Martin Ribe. On uniformly homeomorphic normed spaces. *Ark. Mat.*, 14(2):237–244, 1976.
- [67] Michel Talagrand. Isoperimetry, logarithmic Sobolev inequalities on the discrete cube, and Margulis’ graph connectivity theorem. *Geom. Funct. Anal.*, 3(3):295–314, 1993.
- [68] Michel Talagrand. On Russo’s approximate zero-one law. *Ann. Probab.*, 22(3):1576–1587, 1994.
- [69] Santosh S. Vempala. *The random projection method*, volume 65 of *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*. American Mathematical Society, Providence, RI, 2004. With a foreword by Christos H. Papadimitriou.

CNRS, INSTITUT DE MATHÉMATIQUES DE JUSSIEU, SORBONNE UNIVERSITÉ, FRANCE AND TRINITY COLLEGE, UNIVERSITY OF CAMBRIDGE, UK.

Email address: alexandros.eskenazis@imj-prg.fr, ae466@cam.ac.uk