

Age, sex, and ancestry-related variation in cortical bone and dentine thickness in modern humans and relevance to evolutionary anthropology

Mathilde Augoyard, Clément Zanolli, Frédéric Santos, Anna C Oettlé, Ericka l'Abbé, Antonio Profico, Mona Le Luyer, Marine Cazenave, Thomas Colard, Jakobus W Hoffman, et al.

▶ To cite this version:

Mathilde Augoyard, Clément Zanolli, Frédéric Santos, Anna C Oettlé, Ericka l'Abbé, et al.. Age, sex, and ancestry-related variation in cortical bone and dentine thickness in modern humans and relevance to evolutionary anthropology. European Society for the study of Human Evolution meeting, Sep 2023, Aarhus (Danemark), Denmark. 10.48738/2023.iss2.1093. hal-04280541

HAL Id: hal-04280541 https://hal.science/hal-04280541v1

Submitted on 14 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Age, sex, and ancestry-related variation in cortical bone and dentine thickness in modern humans and relevance to evolutionary anthropology

Mathilde Augoyardı, Clément Zanolliı,2, Frédéric Santosı, Anna C. Oettlé3, Ericka N. L'Abbé4, Antonio Profico5, Mona Le Luyerı,6,7, Marine Cazenave8,9,10, Thomas Colardı,11, Jakobus Hoffmanı2, Priscilla Bayle1

1 - University of Bordeaux, CNRS, MCC, PACEA, UMR 5199, F-33600 Pessac, France · 2 - Department of Maxillofacial and Oral Surgery, Sefako Makgatho Health Sciences University, Ga-Rankuwa, Pretoria, South Africa · 3 - Department of Anatomy and Histology, Sefako Makgatho Health Sciences University, Ga-Rankuwa, Pretoria, South Africa · 4 - Forensic Anthropology Research Centre, Department of Anatomy, Faculty of Health Sciences, University of Pretoria, Pretoria 0031, South Africa · 5 - Department of Biology, University of Pisa, Pisa, Italy · 6 - Center for Genomic Medicine, Massachusetts General Hospital, Boston, USA · 7 - Department of Psychiatry, Harvard Medical School, Boston, USA · 8 - Division of Anthropology, American Museum of Natural History, New York, NY, USA · 9 - Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent, Canterbury, UK · 10 - Department of Anatomy, Faculty of Health Sciences, University of Pretoria, South Africa · 11 - Department of Oral Radiology, University of Lille, Lille University Hospital, France · 12 - South African Nuclear Energy Corporation SOC Ltd., Pelindaba, South Africa

Cortical bone and dentine share an embryological origin, a development, and a genetic background, that are not shared with enamel [1]. Knowing how these tissues relate in individuals is of great importance to decipher the factors acting on their morphogenesis, and ultimately to understand the mechanisms responsible for the different patterns of tissue proportions shown by hominins. However, only a few studies have analyzed bone and tooth structures in combination [2], and no study has investigated a possible correlation between the volumes of cortical bone and dentine relative to endogenous and exogenous factors. The aims of our study are to examine age-, sex-, and ancestry-related variation in cortical bone and dentine volumes, and to preliminary assess the possible covariation between these tissues in modern humans. The analytical sample includes 12 immature individuals and 17 adults from French medieval cemeteries, as well as 32 adults of African and European ancestries from South African contemporary osteological collections. Variables describing three-dimensional tissue proportions were assessed from microtomographic records of radii and permanent maxillary canines. Results suggest an age-related pattern in bone development and important sex differences in cortical bone and dentine thickness. The developmental pattern of dentine also seems to vary according to individual's ancestry. In addition, preliminary results have outlined a covariation trend between volumes of cortical bone and dentine that does not exist with enamel nor pulp. However, when splitting the sample by age, sex, and ancestry, no signal of covariation is found in the immature subsample, while the adults of South African and European ancestry show a covariation between cortical bone and either radicular or crown dentine, respectively. Thus, accessing the genetic-related morphological variation may be complexified by the impact of environmental factors, particularly mechanical ones, which play an important role in structuring the mechanosensitive tissues throughout life. To identify variations in degree of bone and dentine functional sensitivity, and thus disentangle genetic and environmental factors, we also documented the local variation of cortical bone and dentine distribution along the bone shaft and the dental remains by using morphometric maps [3,4]. More research on the cortical bone-dentine covariation would provide a better understanding of the impact of endogenous and exogenous factors on the development of the mineralized tissues. Ultimately, this may shed light on the pattern of joint tissue thickening observed in the palaeoanthropological record, and especially in the majority of Neandertal specimens [5].

Work supported by the "Ministère de l'Enseignement supérieur et de la Recherche" (2021-24; doctoral grant received by M. Augoyard, University of Bordeaux), the CNRS, the "Investments for the future" program IdEx Bordeaux (3Dent'in project, ANR-10-IDEX-03-02, coord.: P. Bayle; STEP ITAP project, OPE-2018-0181, coord.: P. Bayle and J.-G. Bordes), the Région Nouvelle-Aquitaine (NATCH project, 2016-1R40204, coord.: J.-Ph. Faivre), the Département des Sciences Archéologiques de Bordeaux, and the Bakeng se Afrika project 597924-EPP-1-2018-1-ZA-EPPKA2-CBHE-JP, an Erasmus plus project of the EU (coord.: E. L'Abbé). This research benefited from the scientific framework of the University of Bordeaux's IdEx "Investments for the Future" program / GPR "Human Past". For access to the specimens and technical support, we thank C. Beauval, D. Castex, NL Japhta, F. de Beer, E. de Jager, A. Masses, N. Vanderesse, and R. Ledevin.

References:[1] Burr, D.B., Allen, M.R., 2019. Basic and Applied Bone Biology (2nd ed). Academic Press, Amsterdam, The Netherlands. [2] Coquerelle, M., Prados-Frutos, J.C., Benazzi, S., Bookstein, F.L., Senck, S., Mitteroecker, P., Weber, G.W., 2012. Infant growth patterns of the mandible in modern humans: a closer exploration of the developmental interactions between the symphyseal bone, the teeth, and the suprahyoid and tongue muscle insertion sites. Journal of Anatomy 222(2), 178-192. [3] Bondioli, L., Bayle, P., Dean, C., Mazurier, A., Puymerail, L., Ruff, C., Stock, J.T., Volpato, V., Zanolli, C., Macchiarelli, R., 2010. Technical note: Morphometric maps of long bone shafts and dental roots for imaging topographic thickness variation. American Journal of Physical Anthropology 142(2), 328-334. [4] Profico, A., Bondioli, L., Raia, P., O'Higgins, P., Marchi, D., 2021. morphomap: An R package for long bone landmarking, cortical thickness, and cross-sectional geometry mapping. American Journal of Physical Anthropology 174(1), 129-139. [5] Churchill, S.E., 1998. Cold adaptation, heterochrony, and neandertals. Evolutionary Anthropology: Issues, News, and Reviews 7(2), 46-60.