

## Preliminary palaeoproteomic insights into the Plio-Pleistocene australopiths from South Africa

Palesa Madupe, Patrick Leopold Rüther, Nomawethu Hlazo, Meaghan Mackie, Ioannis Patramanis, Alberto Taurozzi, Mirriam Tawane, Lauren Schroeder, Fernando Racimo, Clément Zanolli, et al.

#### ▶ To cite this version:

Palesa Madupe, Patrick Leopold Rüther, Nomawethu Hlazo, Meaghan Mackie, Ioannis Patramanis, et al.. Preliminary palaeoproteomic insights into the Plio-Pleistocene australopiths from South Africa. 12th European Society for the study of Human Evolution meeting (ESHE 2022), Sep 2022, Tübingen, Germany. 10.48738/2022.iss2.809. hal-04280759

### HAL Id: hal-04280759 https://hal.science/hal-04280759

Submitted on 14 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#### Preliminary palaeoproteomic insights into the Plio-Pleistocene australopiths from South Africa

# Palesa Madupe1,2, Patrick L. Rüther3, Nomawethu Hlaz02,4, Meaghan Mackie1,3, Ioannis Patramanis5, Alberto Taur02zi1, Mirriam Tawane6, Lauren Schroeder2,7, Fernando Racim05, Clément Zanolli8, Jesper V. Olsen3, Rebecca R. Ackermann2,4, Enrico Cappellini1

1 - Evolutionary Genomics Section, Globe Institute, University of Copenhagen, Denmark · 2 - Human Evolution Research Institute, University of Cape Town, South Africa · 3 - Proteomics Program, Novo Nordisk Foundation Center for Protein Research, University of Copenhagen, Denmark · 4 - Department of Archaeology, University of Cape Town, South Africa · 5 - Section for Molecular Ecology and Evolution, Globe Institute, Faculty of Health and Medical Sciences, University of Copenhagen, Denmark · 6 - Ditsong National Museum of Natural History, Pretoria, South Africa · 7 - Department of Anthropology, University of Toronto Mississauga, Canada · 8 - Univ. Bordeaux, CNRS, MCC, PACEA, UMR 5199, 33600 Pessac, France

Palaeoproteomics allows for the recovery of genetic information beyond the limits of ancient DNA preservation. Dental enamel proteins older than 900 thousand years from temperate and subtropical areas can be phylogenetically informative about phylogeny, as recently demonstrated by the analysis of fossil teeth from *Gigantopithecus blacki* (1.9 Ma) [1]. Recent work on *Homo antecessor* (900 ka) [2] has demonstrated the viability of applying these approaches to ancient fossil hominins. Here we recovered dental enamel proteins from australopith remains from South Africa: in particular, four *Paranthropus robustus* specimens from Swartkrans, dated to 2.2-1.8 Ma, and 1 specimen of *Australopithecus africanus* from Sterkfontein, dated to 2.8-2.2 Ma. This enabled us to determine the sex of these individuals and learn about their phylogenetic placement relative to present-day and ancient hominids.

We employed a digestion-free enamel extraction protocol [3] to recover ancient hominin proteins from the South African specimens SK 830, SK 835, SK 850, SK 14132, and Sts 63, and leveraged tandem mass spectrometry for deep sequencing. To confidently assign the retrieved MS/MS spectra to peptide sequences, we used reference database-dependent and open searches, using MaxQuant [4] and Peaks [5] software. The reference database included enamel and dentine protein sequences from publicly available repositories (Uniprot and NCBI) taxonomically restricted to hominids. For those taxa whose enamel and dentine protein sequences were not available, we translated protein sequences from publicly available genomic archives

Our results show that it is possible to retrieve enamel proteins from these ancient South African hominins. This is an encouraging step forward in ancient protein studies and is the first ancient proteome recovered from African Plio-Pleistocene fossils. Specimens SK 835, SK 850 and Sts 63 are identified as biologically male individuals. This opens the possibility of reassessing our understanding of sexual dimorphism in these and other early hominins and differentiating variation due to dimorphism from that resulting from taxonomic distinctiveness. Although the recovered enamel proteome is small, we also identified phylogenetically informative single amino acid polymorphisms. Future work will focus on expanding our sample, to better understand variation within and between early hominin taxa in Africa.

This project has received funding from the European Union's Horizon 2020 Research and Innovation Programme under (i) the Marie Sklodowska-Curie grant agreement No. 861389 "PUSHH", and (ii) the European Research Council (ERC) grant agreement No 101021361 "BACKWARD". We thank the South African Heritage Resources Agency (SAHRA), the Ditsong National Museum of Natural History for giving us access to the fossils under their care. Permit identification numbers: 3079 and 3132.

References: [1] Welker, F., Ramos-Madrigal, J., Kuhlwilm, M., Liao, W., Gutenbrunner, P., de Manuel, M., Samodova, D., Mackie, M., Allentoft, M.E., Bacon, A.M., Collins, M.J., 2019. Enamel proteome shows that *Gigantopithecus* was an early diverging pongine. Nature 576 (7786), 262-265. [2] Welker, F., Ramos-Madrigal, J., Gutenbrunner, P., Mackie, M., Tiwary, S., Rakownikow Jersie-Christensen, R., Chiva, C., Dickinson, M.R., Kuhlwilm, M., de Manuel, M., Gelabert, P., 2020. The dental proteome of *Homo antecessor*. Nature 580(7802), 235-238 [3] Cappellini, E., Welker, F., Pandolfi, L., Ramos-Madrigal, J., Samodova, D., Rüther, P.L., Fotakis, A.K., Lyon, D., Moreno-Mayar, J.V., Bukhsianidze, M., Rakownikow Jersie-Christensen, R., 2019. Early Pleistocene enamel proteome from Dmanisi resolves *Stephanorhinus* phylogeny. Nature 574 (7776), 103-107. [4] Cox, J. and Mann, M., 2008. MaxQuant enables high peptide identification rates, individualized ppb-range mass accuracies and proteome-wide protein quantification. Nature Biotechnology 26(12), 1367-1372.[5] Li, D., Fu, Y., Sun, R., Ling, C.X., Wei, Y., Zhou, H., Zeng, R., Yang, Q., He, S., Gao, W., 2005. pFind: a novel database-searching software system for automated peptide and protein identification via tandem mass spectrometry. Bioinformatics 21(13), 3049-3050.