

# Harbour geoarchaeology of Lechaion (Corinth area, Greece) sheds new light on economics during the Late Bronze Age/Early Iron Age transition

Antoine Chabrol, Hugo Delile, Sandrine Baron, Catherine Bouras, Panagiotis Athanasopoulos, Bjørn Lovén

## ► To cite this version:

Antoine Chabrol, Hugo Delile, Sandrine Baron, Catherine Bouras, Panagiotis Athanasopoulos, et al.. Harbour geoarchaeology of Lechaion (Corinth area, Greece) sheds new light on economics during the Late Bronze Age/Early Iron Age transition. Marine Geology, 2023, 465, pp.107167. 10.1016/j.margeo.2023.107167. hal-04282063

# HAL Id: hal-04282063 https://hal.science/hal-04282063

Submitted on 16 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

- 1 Harbour geoarchaeology of Lechaion (Corinth area, Greece)
- 2 sheds new light on economics during the Late Bronze Age/Early
- <sup>3</sup> Iron Age transition
- 4

# Antoine Chabrol<sup>1</sup>, Hugo Delile<sup>2</sup>, Sandrine Baron<sup>3</sup>, Catherine Bouras<sup>4</sup>, Panagiotis Athanasopoulos<sup>5</sup>, Bjørn Lovén<sup>5</sup>

- 8 <sup>1</sup> University of Paris-Sorbonne, UR Médiations, Paris, France (<u>chabrol.antoine@yahoo.com</u>)
- 9 <sup>2</sup> CNRS, Archéorient, UMR 5133, Maison de l'Orient et de la Méditerranée, University of Lyon
- 10 2, Lyon, France, (<u>hugo.delile@cnrs.fr</u>)
- <sup>3</sup> University of Toulouse, CNRS, Traces, UMR 5608, Maison de la Recherche, Toulouse, France
   (sandrine.baron@univ-tlse2.fr)
- <sup>4</sup> French School of Archaeology at Athens, Athens, Greece (<u>catherine.bouras@efa.gr</u>)
- <sup>5</sup>Danish Institute at Athens, Athens, Greece (panagiotis.athanasopoulos@diathens.gr;
- 15 <u>bjoern.loven@gmail.com</u>)
- 16
- 17 18

#### 19 ABSTRACT

20 Lechaion in Corinth, Greece, is the largest ancient port in Greece. Harbour geoarchaeological 21 investigations, based on <sup>14</sup>C-dated palaeoenvironmental archives taken from a well-studied inner basin (Basin 3) and a still unknown outer basin (Basin 4), revealed anthropogenic lead excesses 22 starting from the 12<sup>th</sup> century BCE, associated with brown coal fragments, for the first time 23 discovered in such an ancient archaeological context. Given that historical sources trace the 24 foundation of the port back to the 7<sup>th</sup> century BCE, these results attest to protohistoric industrial 25 use of the site and push back its chronology by over five centuries. The existence of such ancient 26 27 port activity not only extends the chronological horizon for harbour activity in the Corinthian area, 28 but also provides new insights for the Late Bronze Age/Early Iron Age (LBA/EIA) transition, 29 including potential trading routes that may have transited through Lechaion, likely spanning across 30 the Gulf of Corinth and possibly even beyond, to western Mediterranean urban centers.

31

#### 32 **1. INTRODUCTION**

The Late Bronze Age/Early Iron Age (LBA/EIA) transition (12<sup>th</sup>-8<sup>th</sup> century BCE) in Greece is a pivotal period that laid the foundations for the emergence of Greek civilization in its "classical"

form. Up to recently considered a "dark age", this transition had been seen as a period of decline

in the Aegean world. Research tends to revisit this view (Lemos, 2006; Ben Yosef, 2019) and

37 Greek sites from the Early Iron Age show that commercial exchanges, particularly with the Levant,

38 did not disappear (**Bintliff et al., 2012 ; Mazarakis-Ainian et al., 2012**). During these centuries,

39 port activities have certainly played a central role in the expansion of the Phoenicians and Greek 40 colonies in the Mediterranean. However, no sedimentary evidence of such ancient port activity has vet been discovered in Greece. Located in an ancient river outlet, Lechaion was the primary 41 harbour of ancient Corinth, spanning an area of approximately 0.7 km<sup>2</sup>. It is the largest ancient 42 43 port in Greece and its strategic location played a vital role in the economic and military growth of the ancient city (Rothaus, 1995) (Fig. 1A&1B). While historical sources suggest that the port was 44 established in the 7<sup>th</sup> or 6<sup>th</sup> centuries BCE, the presence of an older protohistoric, i.e. Bronze Age, 45 port in the Corinthian region has long been suspected by archaeologists (Blegen, 1920; Tzonou 46 47 and Morgan, 2019). Evidence of commerce across the Gulf of Corinth during Bronze Age is 48 provided by remains and settlements established near the Lechaion coastline, such as the 49 prehistoric site of Korakou (Fig. 1A), which demonstrates contacts with Achaea and Italy until it 50 was abandoned around 1050 BCE (Tzonou and Morgan, 2019). Although regional archaeological 51 evidence attests to the continuity of settlement over the centuries of the Late Bronze Age/Early 52 Iron Age (LBA/EIA) transition (Bintliff, 2012), the question of an active protohistoric port that 53 could have been open to the Gulf of Corinth and the western Mediterranean remains unresolved 54 (Tzonou and Morgan, 2019). To address this topic and to shed new light on the "muted" topic of the LBA/EIA transition (Ben-Yosef, 2019), geoarchaeological investigations took place in the 55 56 harbour area of Lechaion in which two sediment cores, Tar11, and Drill2, were taken in harbour 57 basins 3 and 4, respectively (Fig. 1A). In combination with multivariate analysis (MVA) we 58 conducted Attenuated total reflectance Fourier transform infrared spectroscopy (FTIR-ATR), 59 portable X-ray fluorescence spectrometry (pXRF) and sedimentological analyses on core sediment samples at high resolution. These tools are used to reconstruct palaeo-environmental changes and 60 61 Pb palaeo-pollution signal likely to have trapped traces of early port activities (Delile et al., 2017 62 ; 2018). We also conducted SEM-EDS analyses on brown coal materials found in Drill2 (Fig. 3). 63

64 65

#### 2. PHYSICAL SETTING

66 The geographical and geomorphological framework of the port of Lechaion has been the subject of numerous studies since the 1990s. Lechaion is located in the eastern part of the Gulf of Corinth, 67 68 less than 3 kilometers north of the ancient city. The Gulf of Corinth is an active asymmetric rift 69 zone (with some of the highest extension rates in the world), and seismic activity is significant in 70 the area (Armijo et al., 1996; Charalampakis et al., 2014; Moretti et al., 2003). At this location, 71 the northern shore of the Peloponnese experiences ongoing uplift during the Quaternary period 72 (numerous uplifted marine terraces are present in the Corinth region), and the coastline of 73 Lechaion, as well as the harbour itself, have experienced multiple co-seismic uplift events during 74 historical periods (Apostolopoulos et al., 2015; Morhange et al., 2012; Stiros et al., 1996). 75 Marine dynamics such as longshore drift, swells, and waves are powerful and occasionally 76 morphogenic in this area (Hadler et al., 2013; Vött et al., 2018; Mourtzas et al., 2014). 77

78 Nowadays, the area corresponding to the ancient harbour extends both on land and underwater. 79 Recent excavations (Lechaion Harbour Project) have allowed for the identification and naming of visible port structures (Loven et al., 2018 ; Athanasopoulos, 2020) (Fig. 1A). The inner port 80 81 consists of three basins interconnected by canals. The connection to the sea is also facilitated by a 82 canal. The outer port comprises 5 moles (L-M1 to L-M5) that are mostly submerged. The existence 83 of a fourth basin (Basin 4) located between moles L-M1 and L-M2 was first revealed in this study. 84 The meandering shape of the inner basins suggests that they were constructed downstream of a 85 former river channel, perhaps in a small estuary or pre-existing delta or coastal marsh (Hadler et al., 2013; Vött et al., 2018; Morhange et al., 2012; Kolaiti et al., 2017) 86 87 88 3. MATERIALS AND METHODS

88

89 <u>3.1 Coring and <sup>14</sup>C sampling</u>

90

91 Tar11 core was performed in Basin 3 using a hand auger in order to minimize sediment compaction 92 and Drill2 using a geotechnical drilling rig. In the field, the sediments were photographically 93 documented into stratigraphic units based on texture and colour (using the Munsell code). For each 94 <sup>14</sup>C sample, 20 g of sediment were collected and sieved (500 µm mesh) with distilled water. 95 Organic remains were then individually identified (micro-charcoals, wood and plant remains) 96 under a binocular microscope, extracted and sent for AMS dating.

97

99

98 <u>3.2 Sedimentology and harbour units identification</u>

100 110 samples were analyzed (60 from Tar11 and 50 from Drill2). Grain size-analyses were 101 conducted by Laser Diffraction Spectroscopy and the organic content of the sediment was 102 measured by LOI (Loss On Ignition) (**Supplemental Material A and B**).

103 Based on pXRF and FTIR-ATR analyses, we followed the methods of (Delile et al., 2014) and 104 (Martinez et al., 2021), respectively, to decipher geochemical and mineralogical compositions 105 involved in the formation of sedimentary deposits of the ancient harbour basins of Lechaion 106 (Supplemental Material D, Tab. S1 and Fig. S1). We also used Pb enrichment factors ( $EF_{Pb}$ ) to 107 identify harbour deposit levels from pre-harbour environment in the stratigraphy of cores 108 (Supplemental Material C). Indeed, it has been established that ancient port sediments are 109 characterized by traces of human activities, such as Trace Metal Element (TME) and lead (Pb) 110 pollution seen as an urban change proxy of ancient cities (Delile et al., 2014, 2016, 2017).

- 111
- 112

## 113 <u>3.3 Brown coal fragments analyses</u>

114

After ultrasonic cleaning for 10 minutes, followed by drying during 24 hours, the materials (n=5)
 were analyzed using SEM-EDS (Scanning Electron Microscopy with Energy Dispersive X-ray
 Spectroscopy). Analyses are expressed in X-ray spectrum as well as weight percentage of oxides

118 (Fig. 3). The method is only semi-quantitative, identifying the major elements present in the object.

119 The mineral fraction, composed by Na, Mg, Al, Si, S, Cl, K, Ca and Fe remains below 5wt.% for

120 all the EDS analyses. It should be noted that, due to the extremely high carbon content, the analyses

121 were carried out under non-ideal conditions. The mass balances are therefore only overestimated

122 (for O content) and in no way represent the precise content of the elements from the mineral

- 123 fraction. Nonetheless, these results provide ample information on the present elements and their 124 relative proportions.
- 125
- 126

#### 127 **4. RESULTS**

128 The port facilities that are visible today mainly date back to the Roman period and consist of three 129 interior basins (Harbour basins 1, 2, 3) and an outer basin open to the sea (Harbour basin 4) (**Fig.** 

130 **1A**). The morphology of the port during pre-roman periods is widely unknown. In both Tar11 and

131 Drill2 cores, the radiocarbon datings, Pb enrichment factors ( $EF_{Pb} > 1.5$ ) and geochemical

compositions (redox conditions) of the sediments allow us to subdivide the stratigraphy into three
 main sedimentary units: pre-harbour (unit A), harbour (unit B), and post-harbour (unit C) (Fig. 2).

133 134

135 Pre-harbour deposits (Unit A) in Tar11 and Drill2 cores, ranged at a depth of 2.5 to 3.1 m below

sea level (b.s.l.) and 2.3 to 9.3 m b.s.l., respectively, show lowest EF<sub>Pb</sub> values ranging from 0.71.3 (mean value of 0.9) in Tar11 core, and 0.5-2.5 in Drill2 core (mean value of 1.2). These

sediments of sandy muds and gravely muddy sands refer to detrital inputs dominated by silicate

minerals (and carbonates in Drill2 core between 7.6 and 2.3 m b.s.l.) under weak anoxic water

140 column conditions (**Fig. 2**). These alluvial deposits corroborate the hypothesis that the ancient port

141 of Lechaion was built at the mouth of a coastal temporary stream (Stiros, 1996; Morhange et al.,

- 142 **2012 ; Kolaiti et al., 2017**).
- 143

Both for the inner and the outer harbour basins, a surge of anthropogenic lead excesses (consistent

- 145  $EF_{Pb} > 1.5$ ) and an oxygen depletion in the water column reveal unambiguously harbour deposits
- 146 at 0.6-2.5 m b.s.l. depth in Tar11 core and 0.3-2.3 m b.s.l. depth in Drill2 core (**Fig. 2**) (Unit B).
- 147 According to the radiocarbon dates (**Table 1**), harbour basins were in use from Late Bronze Age
- 148 (1381-1056 cal BCE and 1122-927 cal BCE in Tar11 and Drill2 cores, respectively) to Late
- 149 Antiquity (333-753 CE in Drill2 core). Five medium size pieces of brown coal (B-axe 2 to 5 cm;
- radiocarbon age >45000 Cal BP) were discovered at 0.9 m b.s.l. depth in Drill2 core. According
  to the C content, these materials are situated between lignite (C content of 66wt.%; n=6 analyses)
- and bituminous coal (C content circa 81wt.%; n=2) (Fig. 3).
- 153 As in many ancient Mediterranean ports, the post-harbour unit (0.6 to 0.1 m b.s.l. in Tar11 core
- and 0.2 to 0 m b.s.l. Drill2 core) is characterized by a major environmental change with an oxygen-
- 155 deficient eutrophic regime evolving towards a submerged aquatic environment (Unit C) (**Fig. 2**).
- 156 Post-harbour deposits record the highest EF<sub>Pb</sub> values ranging from 3.9-335.2 (mean value of 58.1)
- 157 in Tar11 core, and 5.9-12.4 in Drill2 core (mean value of 8.1). This sharp increasing trend of Pb

158 contamination in the post-harbour unit has been recorded in numerous fillings of ancient harbour

basins, especially during Medieval times (e.g. Rome, **Delile et al., 2017**; Neapolis, **Delile et al.,** 

160 **2016**; Ephesus, **Delile et al.**, **2015**), and may correspond to other inland local or regional industrial

161 activities related to the reassignment of harbour activities.

162

#### 163 **5. DISCUSSION**

164 The dated stratigraphic connection of the brown coal materials with high Pb excess in Drill2 core seem to highlight, for the first time, a palaeo-metallurgical activity in a harbour area during the 165 166 LBA/EIA transition and/or a regional trade of metals (as ingots shipping or local metallurgical 167 transformation close to the harbour) and/or coal. Even if the use of brown coal in LBA/EIA Greek 168 metallurgy has not yet been demonstrated by mining and metallurgical archaeology, its high C 169 content (> 65wt.%) coupled with an alkaline mineral fraction with little sulfur content (Fig. 3) 170 make it relevant for metallurgical use favoring slag formation directly from the lead-silver ores 171 loaded in furnaces. The presence of these fragments of brown coal in a level dating from the end 172 of the Bronze Age completes recent discoveries. : the exploitation and the use of brown coal as a 173 combustion source during LBA has recently been indirectly identified in mycenaean Tiryns 174 (Peloponnese) (Buckley et al., 2021) and there are lignite sources in the Peloponnese (Kavvadas 175 et al., 2020) (Fig. 1B).

On a broader regional scale, this study raises questions about the potential connections that may have existed between local metallurgical activities (attested during the LBA by recent excavations around Lechaion, **Xanthopoulou et al., 2020**), regional mining operations, metal trade and the development of extensive maritime commercial exchanges. Indeed, there is evidence of silver-lead mining as early as the 3<sup>rd</sup> millennium BCE at Laurion (Attic peninsula) (**Morin and Photiades**, **2005**), Sifnos, Syros, Milos (Cyclades) and South Euboea (**Vaxevanopoulos et al., 2022**) (**Fig. 1B**). Recent studies highlight the continuity of metal exploitation in the Eastern Mediterranean and

183 their diffusion towards the west after the collapse of Bronze Age civilizations at the beginning of

184 the Iron Age, primarily through maritime trade (Eshel et al., 2019; 2023).

185

186 This ongoing harbour activity during the centuries of the LBA/EIA transition would have made 187 Lechaion a central site in Eastern Mediterranean, whose economic significance led to its likely 188 expansion and the construction of monumental port structures from the 7<sup>th</sup> century BCE onwards. 189 Up to now, however, the only visible harbour remains at Lechaion date to historical periods and 190 any protohistoric port structures, if they exist, are yet to be discovered.

- 191
- 192
- 193
- 194
- 195
- 196

#### 197 Data availability statement :

198 Datasets related to this article (XRF and laser granulometry raw data) can be found at :

199 https://data.mendeley.com/datasets/rkd5vpbtv5/1

200 Chabrol, Antoine; Delile, Hugo (2023), "Harbour geoarchaeology of Lechaion sheds new light on 201 economics during the LBA/EIA transition. XRF and granulometry datasets", Mendeley Data, V1, doi: 202 10.17632/rkd5vpbtv5.1

203

#### 204 ACKNOWLEDGMENTS

We would like to thank the Carlsberg and Augustinus Foundations for funding the Lechaion Harbour Project and this study, as well as the Danish Institute at Athens. This work would also not

have been possible without the first official drilling permits in Lechaion provided by the Greek

208 Ministry of Culture, the Greek Ephorates of Underwater Antiquities and of Antiquities of Corinth,

and without the help of D. Kourkoumelis and G. Sanders. We also thank the ARTEMIS program

210 for carrying out the SMA radiocarbon datings. This work was conducted with the analytical

- support of the OMEAA technology platform (UMR 5600 & UMR 5133 CNRS) based at the
- 212 University Lumière Lyon 2 and with the support of the LABEX IMU (ANR-10-LABX-0088) of
- 213 the University of Lyon, within the framework of the "Investissements d'Avenir" program managed
- by the Agence Nationale de la Recherche (ANR). . We are grateful to Isabelle Théry-Parisot for
- her initial support on the identification of the brown coal materials and Luc Robbiola (TRACES,

216 CNRS, "ArchéoSciences platform) for helping us to perform the SEM-EDS analyses. Finally, we

- 217 acknowledge the material assistance provided by the Ecole française d'Athènes during the
- fieldwork in 2017.
- 219

## 220 **REFERENCES CITED**

22Apostolopoulos, G., Minos-Minopoulos, D., Pavlopoulos, K., 2015. Geophysical investigation for the
detection of liquefaction phenomena in an archaeological site, Lechaion, Greece. GEOPHYSICS
80, EN105–EN117. <u>https://doi.org/10.1190/geo2014-0109.1</u>

22**A**rmijo, R., Meyer, B., King, G.C.P., Rigo, A., Papanastassiou, D., 1996. Quaternary evolution of the 225 Corinth Rift and its implications for the Late Cenozoic evolution of the Aegean. Geophysical

226 Journal International 126, 11–53. <u>https://doi.org/10.1111/j.1365-246X.1996.tb05264.x</u>

- 227Athanasopoulos, P., 2020. Inde i en middelhavshavn: Lechaions havneanlaeg i Korinth, *SFINX 32 årgang 2020, Nr. 2*, pp. 50-55 (in Danish).
- 22**B**en-Yosef, E., 2019. Archaeological science brightens Mediterranean dark age. Proceedings of the National Academy of Sciences 116, 5843–5845. <u>https://doi.org/10.1073/pnas.1901885116</u>

23Bintliff, J., 2012. The Greek Early Iron Age and the Concept of a "Dark Age," in: The Complete
232 Archaeology of Greece. John Wiley & Sons, Ltd, pp. 207–233.
233 <u>https://doi.org/10.1002/9781118255179.ch8</u>

23**B**legen, C.W., 1920. Corinth in Prehistoric Times. American Journal of Archaeology 24, 1–13. <u>https://doi.org/10.2307/497547</u>

23**B**uckley, S., Power, R.C., Andreadaki-Vlazaki, M., Akar, M., Becher, J., Belser, M., Cafisso, S., 237 Eisenmann, S., Fletcher, J., Francken, M., Hallager, B., Harvati, K., Ingman, T., Kataki, E., Maran,

- 238 J., Martin, M.A.S., McGeorge, P.J.P., Milevski, I., Papadimitriou, A., Protopapadaki, E., Salazar-
- 239 García, D.C., Schmidt-Schultz, T., Schuenemann, V.J., Shafiq, R., Stuijts, I., Yegorov, D., Yener,
- 240 K.A., Schultz, M., Spiteri, C., Stockhammer, P.W., 2021. Archaeometric evidence for the earliest

exploitation of lignite from the bronze age Eastern Mediterranean. Sci Rep 11, 24185.
 <u>https://doi.org/10.1038/s41598-021-03544-w</u>

24C haralampakis, M., Lykousis, V., Sakellariou, D., Papatheodorou, G., Ferentinos, G., 2014. The
tectono-sedimentary evolution of the Lechaion Gulf, the south eastern branch of the Corinth
graben, Greece. Marine Geology 351, 58–75. <a href="https://doi.org/10.1016/j.margeo.2014.03.014">https://doi.org/10.1016/j.margeo.2014.03.014</a>

24Delile, H., Blichert-Toft, J., Goiran, J.-P., Stock, F., Arnaud-Godet, F., Bravard, J.-P., Brückner, H.,
Albarède, F., 2015. Demise of a harbor: a geochemical chronicle from Ephesus. Journal of
Archaeological Science 53, 202–213. <u>https://doi.org/10.1016/j.jas.2014.10.002</u>

24Delile, H., Goiran, J.P., Blichert-Toft, J., 2018. The contribution of geochemistry to ancient harbor
geoarcheology: The example of Ostia Antica. Quaternary Science Reviews 193, 170–187.
https://doi.org/10.1016/j.quascirev.2018.06.019

25Delile, H., Keenan-Jones, D., Blichert-Toft, J., Goiran, J.-P., Arnaud-Godet, F., Albarède, F., 2017.

- Rome's urban history inferred from Pb-contaminated waters trapped in its ancient harbor basins.
  Proceedings of the National Academy of Sciences 114, 10059–10064.
  https://doi.org/10.1073/pnas.1706334114
- 25Delile, H., Keenan-Jones, D., Blichert-Toft, J., Goiran, J.-P., Arnaud-Godet, F., Romano, P., Albarède,
  F., 2016. A lead isotope perspective on urban development in ancient Naples. Proceedings of the
  National Academy of Sciences 113, 6148–6153. https://doi.org/10.1073/pnas.1600893113

25Delile, H., Mazzini, I., Blichert-Toft, J., Goiran, J.P., Arnaud-Godet, F., Salomon, F., Albarède, F.,
260 2014. Geochemical investigation of a sediment core from the Trajan basin at Portus, the harbor of
261 ancient Rome. Quaternary Science Reviews 87, 34–45.
262 <u>https://doi.org/10.1016/j.quascirev.2014.01.002</u>

26Eshel, T., Erel, Y., Yahalom-Mack, N., Tirosh, O., Gilboa, A., 2019. Lead isotopes in silver reveal
earliest Phoenician quest for metals in the west Mediterranean. Proceedings of the National
Academy of Sciences 116, 6007–6012. <u>https://doi.org/10.1073/pnas.1817951116</u>

26Eshel, T., Gilboa, A., Tirosh, O., Erel, Y., Yahalom-Mack, N., 2023. The earliest silver currency hoards
in the Southern Levant: Metal trade in the transition from the Middle to the Late Bronze Age.
Journal of Archaeological Science 149, 105705. https://doi.org/10.1016/j.jas.2022.105705

26 Pladler, H., Vött, A., Koster, B., Mathes-Schmidt, M., Mattern, T., Ntageretzis, A.K., Reicherter, K.,

270 Willershäuser, T., 2013. Multiple late-Holocene tsunami landfall i in the eastern Gulf of Corinth

recorded in the palaeotsunami geo-archive at Lechaion, harbour of ancient Corinth (Peloponnese,
Greece). Zeitschrift für Geomorphologie, Supplementary Issues 57, 139–180.

273 https://doi.org/10.1127/0372-8854/2013/S-00138

27Kavvadas, M., Roumpos, C., Schilizzi, P., 2020. Stability of Deep Excavation Slopes in Continuous
Surface Lignite Mining Systems. Geotech Geol Eng 38, 791–812. <u>https://doi.org/10.1007/s10706-</u>
019-01066-x

- 27Kolaiti, E., Papadopoulos, G.A., Morhange, C., Vacchi, M., Triantafyllou, I., Mourtzas, N.D., 2017.
- Palaeoenvironmental evolution of the ancient harbor of Lechaion (Corinth Gulf, Greece): Were
   changes driven by human impacts and gradual coastal processes or catastrophic tsunamis? Marine
- 280 Geology 392, 105–121. <u>https://doi.org/10.1016/j.margeo.2017.08.004</u>

28Lemos, I., 2006. The 'Dark Age' of Greece, in : Bispham, Edward, Harrison Thomas, Sparkes, Brian

(eds.), *The Edinburgh Companion to Ancient Greece and Rome*, Edinburgh: Edinburgh University
 Press

28Longman, J., Veres, D., Finsinger, W., Ersek, V., 2018. Exceptionally high levels of lead pollution in
the Balkans from the Early Bronze Age to the Industrial Revolution. Proceedings of the National
Academy of Sciences 115, E5661–E5668. <u>https://doi.org/10.1073/pnas.1721546115</u>

- 28 Loven B., Kourkoumelis D., Micha P., Athanasopoulos P., 2018. Lechaion Harbour Project 2013-2014.
- 288 In A. Simosi (ed.), Βουτιά στα Περασμένα. Η υποβρύχια αρχαιολογική έρευνα 1974-2014. Αθήνα:

289 ТАПА, pp. 85-90. (in Modern Greek)

- 29Martínez Cortizas, A., Sjöström, J.K., Ryberg, E.E., Kylander, M.E., Kaal, J., López-Costas, O.,
- Álvarez Fernández, N., Bindler, R., 2021. 9000 years of changes in peat organic matter
   composition in Store Mosse (Sweden) traced using FTIR-ATR. Boreas 50, 1161–1178.
   https://doi.org/10.1111/bor.12527
- 29Mazarakis-Ainian, A.I., Coulson, W.D.E., 2011. The "dark ages" revisited : acts of an international
  symposium in memory of William D.E. Coulson, University of Thessaly, Volos, 14-17 June 2007.
  University of Thessaly Press.
- 29 Moretti, I., Sakellariou, D., Lykousis, V., Micarelli, L., 2003. The Gulf of Corinth: an active half
- graben? Journal of Geodynamics, Active Faults: Analysis, Processes and Monitoring 36, 323–340.
   <u>https://doi.org/10.1016/S0264-3707(03)00053-X</u>
- 30 Morhange, C., Pirazzoli, P.A., Evelpidou, N., Marriner, N., 2012. Late Holocene Tectonic Uplift and
- the Silting Up of Lechaion, the Western Harbor of Ancient Corinth, Greece. Geoarchaeology 27,
   278–283. https://doi.org/10.1002/gea.21388
- 30Morin, D., Photiades, A., 2005. Nouvelles recherches sur les mines antiques du Laurion (Grèce). Pallas 304 327–358.
- 30 Mourtzas, N.D., Kissas, C., Kolaiti, E., 2014. Archaeological and geomorphological indicators of the
- historical sea level changes and the related palaeogeographical reconstruction of the ancient
- 307 foreharbour of Lechaion, East Corinth Gulf (Greece). Quaternary International, Relative sea level
- 308 changes: Signature on continental shelves, changing coastlines, and implications for coastal 309 morphodynamics 332, 151–171. https://doi.org/10.1016/j.quaint.2012.12.037
- 31**R**othaus, R., 1995. Lechaion, Western Port of Corinth: A Preliminary Archaeology and History. Oxford
- 311 Journal of Archaeology 14, 293–306. <u>https://doi.org/10.1111/j.1468-0092.1995.tb00065.x</u>
- 31Stiros, S., Pirazzoli, P., Rothaus, R., Papageorgiou, S., Laborel, J., Arnold, M., 1996. On the date of
  construction of Lechaion, western harbor of ancient Corinth, Greece. Geoarchaeology 11, 251–
  263. https://doi.org/10.1002/(SICI)1520-6548(199605)11:3<251::AID-GEA4>3.0.CO;2-2
- 31**T**zonou, I., Morgan, C., 2019. The Corinthia, in: A Companion to the Archaeology of Early Greece and 316 the Mediterranean. John Wiley & Sons, Ltd, pp. 719–741. 317 https://doi.org/10.1002/9781118769966.ch29
- 318/axevanopoulos, M., Blichert-Toft, J., Davis, G., Albarède, F., 2022. New findings of ancient Greek 319 silver sources. Journal of Archaeological Science 137, 105474. 320 https://doi.org/10.1016/j.jas.2021.105474
- 32Vött, A., Hadler, H., Koster, B., Matthes-Schmidt, M., Röbke, B.R., Willershäuser, T., Reicherter, K.,
- 322 2018. Returning to the facts: Response to the refusal of tsunami traces in the ancient harbour of
- Lechaion (Gulf of Corinth, Greece) by 'non-catastrophists' Reaffirmed evidence of harbour
- destruction by historical earthquakes and tsunamis in AD 69–79 and the 6th cent. AD and a preceding pre-historical event in the early 8th cent. BC. Zeitschrift für Geomorphologie 28.
- 326 preceding pre-instorical event in the early still cent. BC. Zent
- 327Xanthopoulou M., Banou A., Zymi E., Giannouli E., Karapanagiotou A. and Koumousi A., (eds).
- 328 2020. Το αρχαιολογικό έργο στην Πελοπόννησο 2. Πρακτικά της Β΄ Επιστημονικής Συνάντησης
- 329 Καλαμάτα, 1-4 Νοεμβρίου 2017. pp. 105–14.

#### 330 FIGURES AND TABLE CAPTIONS



Figure 1. A. The Lechaion harbour area and the drilling locations. The pre- and protohistoric site of Korakou, which was occupied until the end of the Bronze Age, is located on top of a hill, west of the harbour area. **B.** Location of the sites and pre- and protohistoric silver-lead mines mentioned

in the text (Vaxevanopoulos et al., 2022; Morin and Photiades, 2005). The brown coal (lignite)

336 sources in the Peloponnese (Kavvadas et al., 2020)

337



338

Figure 2. Stratigraphic log of the Tar11 and Drill2 cores showing granulometry, Principal
 Component Analysis (PCA) of major and trace element concentrations, fractionation of Fourier
 transform infrared spectroscopy (FTIR-ATR) communalities, and Enrichment Factor of Pb
 (normalized to Al). Right panels show correlation circles of the PCA.









Sample code	Sample #	Elevation (cm bmsl)	Material	$\Delta^{13}C$	<sup>14</sup> C Age (BP)	$2\sigma$ Max and Min Cal. Age		Median prob. age
Poz#2-141434	Tar11-1	-127,8	charcoal	-16	2085 ± 30	194 BCE	5 BCE	94, BCE
Poz-140728	Tar11-2	-151,8	charcoal	-26	1785 ± 35	204 CE	381 CE	292,5 CE
Poz-140677	Tar11-4	-223,8	charcoal	-26,1	2225 ± 35	388 BCE	197 BCE	292,5 BCE
Poz-140741	Tar11-5	-246,8	plant remains	-17,8	2985 ± 35	1381 BCE	1056 BCE	1218,5 BCE
Poz#2-141023	Drill2-2	-8,1	marine shell	3,4	2135 ± 30	333 CE	753 CE	543 CE
Poz-157392	Drill2-2'	-63,1	charcoal	-30,2	2860 ± 30	1122 BCE	927 BCE	1024,5 BCE
Poz-140727	Drill2-5	-88,1	lignite	-24,1	>46 000 ± 6180	46990 BCE	45685 BCE	46337,5 BCE

**Table 1.** Radiocarbon dating. AMS  $^{14}$ C radiocarbon dates calibrated using OxCal (v4.4.4) withthe IntCal20 and MarineCal20 calibration curves