



## The diversification of the antimicrobial peptides from marine worms is driven by environmental conditions

Renato Bruno, Céline Boidin-Wichlacz, Oleg Melnyk, Daniela Zeppilli, Celine Landon, Frédéric Thomas, Marie-Anne Cambon-Bonavita, Mickaël Lafond, Kamel Mabrouk, François Massol, et al.

### ► To cite this version:

Renato Bruno, Céline Boidin-Wichlacz, Oleg Melnyk, Daniela Zeppilli, Celine Landon, et al.. The diversification of the antimicrobial peptides from marine worms is driven by environmental conditions. Science of the Total Environment, 2023, 879, pp.162875. 10.1016/j.scitotenv.2023.162875 . hal-04283278

**HAL Id: hal-04283278**

**<https://hal.science/hal-04283278>**

Submitted on 13 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# **The diversification of the antimicrobial peptides from marine worms is driven by environmental conditions**

Renato Bruno<sup>1</sup>, Céline Boidin-Wichlacz<sup>1</sup>, Oleg Melnyk<sup>1</sup>, Daniela Zeppilli<sup>2</sup>, Céline Landon<sup>3</sup>,  
Frédéric Thomas<sup>4</sup>, Marie-Anne Cambon<sup>2</sup>, Mickael Lafond<sup>5</sup>, Kamel Mabrouk<sup>6</sup>, François  
Massol<sup>1</sup>, Stéphane Hourdez<sup>7</sup>, Marc Maresca<sup>5</sup>, Didier Jollivet<sup>8</sup>, Aurélie Tasiemski<sup>1\*</sup>

<sup>1</sup> Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 9017 -  
CIIL - Center for Infection and Immunity of Lille, F-59000 Lille, France

<sup>2</sup> IFREMER Centre Brest REM/EEP/LEP, ZI de la Pointe du Diable, CS10070,  
Plouzané F-29280, France

<sup>3</sup> Center for Molecular Biophysics, CNRS, UPR 4301, Orleans, France

<sup>4</sup> CREEC/(CREES), MIVEGEC, Unité Mixte de Recherches, IRD 224-CNRS 5290-  
Université de Montpellier, Montpellier, France

<sup>5</sup> Aix-Marseille Univ, CNRS, Centrale Marseille, iSm2, Marseille F-13013, France

<sup>6</sup> Aix-Marseille Univ, CNRS, UMR 7273, ICR, Marseille F-13013, France

<sup>7</sup> Sorbonne Université, LECOB, UMR 8222, Observatoire Océanologique de Banyuls,  
1 Avenue Pierre Fabre, 66650 Banyuls-sur-Mer, France

<sup>8</sup> Sorbonne Université, CNRS, UMR 7144 AD2M, Station Biologique de Roscoff, Place  
Georges Teissier CS90074, Roscoff F-29688, France

\*Corresponding author: Email: aurelie.tasiemski@univ-lille.fr, address: Center for  
Infection & Immunity of Lille (CIIL), CGIM Team, Institut Pasteur de Lille, Bâtiment IBL, 1  
rue du Professeur Calmette, CS 50447, F-59021 Lille Cedex, France

## **Orcid#**

Renato Bruno 0000-0002-5195-4192

Celine Boidin-Wichlacz 0000-0002-0242-868X

Daniela Zeppilli 0000-0002-0809-2566

Oleg Melnyk 0000-0002-6030-3666

Céline Landon 0000-0002-0248-1990

Marie-Anne Cambon 0000-0002-4076-0472

Mickael Lafond: 0000-0003-2646-0286

34 Kamel Mabrouk  
35 Frederic Thomas 0000-0003-2238-1978  
36 François Massol 0000-0002-4098-955X  
37 Stéphane Hourdez 0000-0001-6418-3887  
38 Marc Maresca 0000-0002-3585-4765  
39 Didier Jollivet 0000-0001-8042-1857  
40 Aurélie Tasiemski 0000-0003-3559-5115

41

42 **Keywords:** BRICHOS-AMPs, disulfide bridges, annelids, polar, hydrothermal, external  
43 immunity, immunoecology

44

45 **Credit author statement:**

46 Renato Bruno: Writing the initial draft, performing the experiments, data presentation,  
47 investigation. Céline Boidin-Wichlacz: Design of methodology, performing the  
48 experiments, data presentation. Daniela Zeppilli: Critical review and co PI of the Pioneer  
49 Project. Oleg Melnyk: Mass spectrometry experiments and analyses. Céline Landon,  
50 Frederic Thomas and Mickael Lafond: Critical review, commentary. Marie-Anne Cambon:  
51 Provision of biological materials. Kamel Mabrouk: Provision and design of the ABU  
52 peptides. François Massol: Statistical analyses, creation of models, conceptualization.  
53 Stéphane Hourdez: Design and creation of the polar annelid databank, critical review.  
54 Marc Maresca: Design of methodology, performing the experiments, critical review. Didier  
55 Jollivet: Design and creation of the polar annelid databank, identification of POL from  
56 transcriptomic analyses, conceptualization, critical review. Aurélie Tasiemski: Data  
57 presentation, writing the initial draft, review and final editing of the published work,  
58 conceptualization, supervision and project administration.

59

**Abstract:** Antimicrobial peptides (AMPs) play a key role in the external immunity of animals, offering an interesting model for studying the influence of the environment on the diversification and evolution of immune effectors. Arenicin (ARE), alvinellacin (ALV) and polaricin (POL, a novel AMP identified here), characterized from three marine worms inhabiting contrasted habitats (temperate, 'hot' vents and polar respectively), possess a well conserved BRICHOS domain in their precursor molecule despite a profound amino acid and structural diversification of the C-terminal part containing the core peptide. Data not only showed that ARE, ALV and POL display an optimal bactericidal activity against the bacteria typical of the habitat where each worm species is living but also that this killing efficacy is optimal under the thermochemical conditions encountered by their producers *in situ*. Moreover, the correlation between species habitat and the cysteine content of POL, ARE and ALV led us to investigate the importance of disulfide bridges in their biological efficacy as a function of abiotic pressures (pH and temperature). The construction of variants using non-proteinogenic residues instead of cysteines ( $\alpha$ -aminobutyrate chemical variants) leading to AMPs devoid of disulfide bridges, provided evidence that the disulfide pattern of the three AMPs allows for a better bactericidal activity and constitutes an adaptive way to sustain the fluctuations of the worm's environment. This work shows that the external immune effectors exemplified here by BRICHOS AMPs are evolving under strong diversifying environmental pressures to be structurally shaped and more efficient/specific under the ecological niche of their producer.

## 1. INTRODUCTION

External immunity constitutes the first defense of metazoans to face pathogens by handling the surrounding microbial communities to both avoid infection and establish symbiosis [1]. As such, it can be viewed as an extended arm of the immune system to prevent infection [1]. Many organisms including plants, invertebrates, vertebrates and, even bacteria, secrete antimicrobial peptides (AMPs) as an extrinsic protective shield against the surrounding microbiome that is usually rich in potentially pathogenic agents [2–4]. AMPs are predominantly short cationic peptides (up to 100 amino acids), with a broad-spectrum of activities against diverse microbes (bacteria, virus or fungi) with a rapid and selective killing processing mode [4]. Lacking antibodies, invertebrates nevertheless benefit from a powerful innate immune system, characterized, among other processes, by a great variety of AMPs [5].

Marine organisms constitute almost half of the biodiversity of the Earth, and after a long-term diversification (Cambrian explosion: 541 to 485 mya ago) and co-evolution with fungi, bacteria and archaea, acquired the capacity of producing many unique antimicrobial substances [6–8]. Marine AMPs were found to be structurally different from their counterparts produced by terrestrial species, usually displaying novel structures, taxa-specific or even species-specific [6,8,9]. A long evolutionary story of marine organisms (including bacteria) under diversifying environmental pressures, such as the microbial community composition and/or highly fluctuating abiotic factors (temperature, pH, salinity, or hydrostatic pressure, which can vary to extreme values from polar regions to the Equator and the intertidal to deep sea) have probably driven the evolution of the physiological adaptation of most marine invertebrates, including on their immune functions [10].

In particular, worms have colonized and dominated in biomass most of marine and freshwater habitats, coping with a wide number of environmental conditions and bacterial assemblages, especially in extreme environments such as polar regions, deep-sea hydrothermal vents, or highly anthropized shallow areas [11,12]. In the case of annelids (ringed worms), the majority of AMPs is species-specific, probably as the result of a long parallel evolution of their defense arsenal to a wide variety of habitats on Earth (aquatic and terrestrial), mimicking their highly diverse life styles and ecology [13]. In this context, the BRICHOS-domain AMP family of marine annelids living under extreme

conditions, which is specifically involved in the worms' external immune defense constitutes a unique and attractive model to study the adaptive evolution of the immune system to environmental changes.

BRICHOS (initially found in Bri2, chondromodulin, and prosurfactant protein C in human) is a 100 amino acids domain, present in several protein precursors, acting as an intramolecular chaperone [17–21]. Despite their very old phylogenetic separation (at the end of the Cambrian diversification), BRICHOS precursors all display the same structure consisting of: (i) a highly hydrophobic N-terminal signal sequence, (ii) a pro-region containing a BRICHOS domain, and (iii) a C-terminal region with  $\beta$ -sheet propensities. In the unique case of marine worms, this latter corresponds to an AMP which exerts its antimicrobial activities once cleaved from the precursor [6,14–16]. Among the BRICHOS-domain AMPs, arenicin (ARE) was the first identified from *Arenicola marina*, an annelid inhabiting the intertidal zone of the European temperate shore [14]. Later, the AMP alvinellacin (ALV) was purified from the deep-sea hydrothermal vent annelid *Alvinella pompejana*, which inhabits the hottest part of this peculiar environment [15]. Finally, another annelid BRICHOS-domain AMP, the nicomicin, was identified from *Nicomache minor*, a polychaete colonizing both temperate and arctic habitats [16].

Interestingly, precursors of BRICHOS-AMPs show a well-conserved BRICHOS domain but highly divergent primary AMP sequences, suggesting either a strong diversifying selection at the interspecific level of the antibiotic peptides to face the diverse microbial communities, and/or abiotic conditions, or some exon shuffling between the well-conserved BRICHOS domain and AMPs of different genomic origins [22]. In order to enlarge our panel of BRICHOS-AMPs from extreme habitats, we report here a novel AMP, named polaricin (POL), from an undescribed terebellid polychaete belonging to the genus *Amphitritides* endemic to subtidal polar waters of the Antarctic habitat. POL together with ALV and ARE, represent relevant molecules to study AMP adaptation to environmental stressors as they have evolved under highly contrasted (polar, hot vents and temperate respectively) environmental conditions from a shared but old ancestor [14,15] (Fig. 1).

In this work, we focused our attention on how selective pressures associated with biotic and abiotic factors have affected the structure and bioactivity of the three POL, ALV and ARE peptides. Through combined analyses of their bioactivities and structural characteristics under various set of environmental and bacterial conditions, we

deciphered whether these immune peptides better suited to the environmental conditions typifying the worms' habitats.

## **2. MATERIAL AND METHODS**

### **2.1. AMPs and habitat specificities of the worms from which they derived from**

Arenicin (ARE) was purified from the lugworm *Arenicola marina* (Annelid, Polychaeta, Scolecida), a polychaete that inhabits temperate near-shore (intertidal) sediments, digging tunnels in the sand [60,61]. The burrows of these worms are continuously exposed to periodic tidal emersion, which imposes severe thermal variations (-5 °C to 25 °C) and temporal hypoxia with mild acidic conditions (pH from 5.6 to 8.2) [24–32].

Alvinellacin (ALV) was isolated from *Alvinella pompejana* (Annelid, Polychaeta, Terebellida), a polychaete that inhabits parchment-like tubes along the outer walls of active hydrothermal vents chimneys of East Pacific Rise [62]. Its habitat is characterized by frequent and unpredictable mixing of hot and anoxic fluids with cold seawater (2-3 °C), which expose the worm to extreme temperatures (from 2 °C to upper 100 °C) that greatly vary inside the tube (14 °C to 80 °C with a thermal optimum around 42 °C) and acidic gradients (pH from 5.33 to 6.4) due to the mixing of bottom seawater with the acidic vent fluid [33–38].

Polaricin (POL) was identified by analogy from the transcriptome of *Amphitritides* sp. (Annelid, Polychaeta, Terebellida), a marine annelid inhabiting marine sediments at the base of sponges or bryozoans of the Antarctic plateau at depth of 100-150 m. This “spaghetti worm” is sedentary and lives inside a slightly consolidated tube made with sand and coral debris on the Antarctic continental plateau where it faces extremely cold but stable coastal waters (about -1.8 °C), with very small fluctuations of temperature and pH [63–66].

### **2.2. Molecular identification of the AMP precursors**

The molecular characterization of the preproarenicin and preproalvinellacin has been described in previous studies done by Ovchinnikova, T. V. *et al* (2004) [14] and by Tasiemski A. and colleagues [15].

The complete cDNA sequence of prepropolaricin was obtained by homology with the preproalvinellacin gene (Tasiemski *et al.* 2014, accession N° KJ489380) from a tblastx query on the transcriptome assembly previously obtained for the Antarctic *Amphitritides* sp. under scrutiny (blast database) using the software Geneious (blast cut-off p-value =

10-15). The *Amphitritides* RNAseq dataset used for the assembly was obtained from an Illumina 150 bp paired-ends sequencing of a stranded mRNA library on a HiSeq 2500 machine at Genome Québec (half a line: 150 M PE reads). Transcripts (Ngenes=99970, N50=1440 bp, median contig length=488 bp, GC%=36.2) were then obtained with the Trinity 2.4.0 software from R1 and R2 pairs after trimming adapters and regions of low PHRED scores (Trimmomatic 0.36: ILLUMINACLIP: illumina. fa: 2:30:10, LEADING:5, TRAILING:5, SLIDINGWINDOW:4:15 MINLEN:36) and a subsequent normalization of the reads.

### **2.3. Alignments of the precursor amino acid sequences and identification of the BRICHOS and AMP domains**

BRICHOS domain sequence was identified by using MyHits Motif Scan ([https://myhits.isb-sib.ch/cgi-bin/motif\\_scan](https://myhits.isb-sib.ch/cgi-bin/motif_scan)) and the SignalP 5.0 software (<https://www.cbs.dtu.dk/services/SignalP/>) was used to detect the presence of signal peptide.

The amino acid sequence alignments of peptides precursors of ALV (GenBank accession number KJ489380), ARE (GenBank accession number AY684856), nicomicin (GenBank accession number MH898866) and POL, was performed by using CLC Sequence Viewer software (version 8.0). The computation of homology/identity percentage was performed at the SIB (ExPASy software, <https://www.expasy.org/>) using the BLAST network service (BLASTP, version 2.2.31+). The amino acid sequences of the three AMPs presented in this alignment were used for the chemical synthesis of native and ABU peptides as described further on.

### **2.4. Chemical synthesis of AMPs with (ALV, ARE, POL) and without (ABU ALV, ABU ARE, ABU POL) disulfide bridges**

The amino acid sequences used for the peptide synthesis are presented in tables S1 and S2. The chemically synthesized peptides (purity of 98%), *i.e.*, ALV, ARE and POL, were produced by GenScript (Leiden, Netherlands) with the cysteine (Cys) pattern involved in disulfide bridges accordingly to their native forms:

- ALV: Four cysteine residues engaged in two intramolecular disulfide bonds (Cys3-Cys21; Cys7-Cys7);
- ARE: Two cysteine residues engaged in one intramolecular disulfide bond (Cys3-Cys20);
- POL: One cysteine (Cys10) engaged in one intermolecular disulfide bond (Cys10-Cys10).



The  $\alpha$ -aminobutyrate (ABU) modified peptides were chemically produced using the solid-phase method with an automated peptide synthesizer (Model 433A, Applied Biosystems Inc.) in the Radical Chemistry Institute by the Radical Organic Chemistry and Specialty Polymers Team (CROPS) laboratory (France, Marseilles) [67,68]. The disulfide bonds were removed by replacing all the cysteine residues by ABU residues (hydrophobic non-proteinogenic alpha amino acid, also known as homoalanine) which do not associate covalently and thus do not form disulfide bridges [69,70]. The identity and degree of homogeneity of the peptides were verified by: (1) analytical C18 reversed-phase HPLC(NUCLEODUR 100-5 C18, 250  $\times$  10 mm, MACHERY NAGEL) using a 60-min linear gradient from 0 to 60% of buffer B [0.08% (v/v) TFA/acetonitrile] in buffer A [0.1% (v/v) TFA/H<sub>2</sub>O], at a flow rate of 1 mL/min; (2) amino acid analysis after acidolysis [6N HCl/2% (w/v) phenol, 20 h, 118°C, N<sub>2</sub> atmosphere] and (3) molecular mass analysis by MALDI-TOF mass spectrometry. The physicochemical properties of all the synthesized peptides (molecular weight, isoelectric point, net charge at neutral pH) were calculated by using Innovagen Pepcalc.com server (Innovagen AB, SE-22370 Lund, SWEDEN).

### **2.5. Comparison of the three-dimensional (3D) structures of ALV, ARE and POL**

The 3D structure of ARE and ALV was gained from previous studies [14,15]. 3D peptide models of the homooligomer 1:1 of POL and of the sequence monomer were predicted using the Alpha Fold software. The confidence in the accuracy of the predicted structures was measured by the predicted local distance difference test (pLDDT): score 0 to 100. The peptides 3D images were generated using ChimeraX.

### **2.6. Mass spectrometry of POL**

A sample of POL (about 2  $\mu$ L, at concentration of 2 mg/mL) was characterized by analytical UPLC–MS using a System Ultimate 3000 UPLC (ThermoFisher) equipped with an Acquity peptide BEH300 C18 column (1.7  $\mu$ m, 2.1  $\times$  100 mm, 300 Å), a diode array detector and a mass spectrometer (Ion trap LCQfleet). Analysis was performed at 70 °C using a linear gradient of 0-70% of eluent B in eluent A over 20 min at a flow rate of 0.4 mL min<sup>-1</sup> (eluent A = 0.1% TFA in H<sub>2</sub>O; eluent B = 0.1% TFA in CH<sub>3</sub>CN). The column eluate was monitored by UV at 215 nm. The peptide masses were measured by on-line UPLC–MS (LCQ Fleet Ion Trap Mass Spectrometer, ThermoFisherScientific): heat temperature 450 °C, spray voltage 2.8 kV, capillary temperature 400 °C, capillary voltage 10 V, tube lens voltage 75 V.

### **2.7. Bacterial strains used for the antimicrobial assays**

A series of bacterial strains was used to estimate the killing efficiency of the different AMPs and test whether they are more efficient against bacteria living in the same environment of the worms. The bacterial strains used in this study are all Gram-negative from different marine environments:

--*Pseudomonas sp.* AT1238 and *Vibrio diabolicus* HE800 provided by Ifremer LMEE culture collection (Mission HERO'1991) were isolated from a mix of tubes of *A. pompejana* and small pieces of the surrounding polymetallic sulfides of deep-sea hydrothermal chimneys (East Pacific Rise) [72].

--*Shewanella algae*, *Oceanisphaera donghaensis*, and *Vibrio fluvialis* were isolated from intertidal mud sediment near to the burrow of *Arenicola marina*. Raw sediment samples were incubated overnight in a liquid Marine Broth media at 28 °C to stimulate bacterial growth. Well-growing colonies were isolated by multiple pricking out onto agar plates containing marine seawater to identify different bacterial strains. Pure colonies were identified with the Universally applicable bacterial primer set 16S-F and 16S-R (forward primer: 5'-TCCTACGGGAGGCAGCAGT -3'; reverse primer: 5'-GGACTACCAGGGTATCTAATCCTGTT -3') as described in *Cuvillier et al* [73] . The sequences were compared against nucleotide sequences from the Eztaxon database using BLASTn UPGMA trees were constructed using the Mega-X software package (<http://www.megasoftware.net>) (Fig. S4).

--*Vibrio alginolyticus* provided by the Ifremer is worldwide distributed in coastal waters of temperate and also polar areas [74,75]. *V. alginolyticus* and *V. diabolicus* are genetically closely related (Fig. S4).

## **2.8. Determination of the spectrum of antimicrobial activities of ALV, ARE and POL**

Peptides Minimal Inhibitory Concentration (MIC) and Minimal Bactericidal Concentration (MBC), were determined by the broth microdilution method against bacterial growth in microtiter plates as previously described by Bulet [76]. The AMPs were resuspended in sterile pure water. Briefly, 100 µL of bacterial dilutions (10<sup>6</sup> Colony Forming Unit/mL) were added to each well into sterile 96-well flat bottom plates (CELLSTAR, Greiner bio-one) containing serial dilutions (from 40 to 0.00195 µg/mL) of peptides (10 µL per well). The plates were incubated overnight at 28 °C on a rotary shaker (140 rpm). The MIC was taken as the lowest peptide concentration at which observable microbial growth was inhibited, the measurements were realized using a microtiter plate reader (Tecan Sunrise

Microplate Reader) at A600nm. MBC was measured by streaking on proper agar (MHB with 15 g/L bacteriological agar, Euromedex) petri dishes, the entire volume of each well (110 µL) with no bacterial growth in the previously obtained MIC plates. After incubation for about 24 hours at 28°C, the MBC was defined as the peptide concentration where no colony growth was observed. All tests were conducted in triplicate.

A 'home versus away' graph was constructed from antimicrobial assays against the whole set of environmental marine bacteria with log-transformed MIC data organized in two groups, depending on the habitat (polar/temperate intertidal vs deep-sea hot vents) to highlight patterns of AMP efficiency in both native and transplanted habitats.

### **2.9. Kinetics of bactericidal activities of ALV, ARE and POL against *Vibrio* strains depending on the thermal range**

Kinetics of bactericidal activity of the three AMPs were assessed on two bacteria belonging to the same genus but colonizing two contrasted habitats: *V. alginolyticus* (polar and temperate) and *V. diabolicus* ("hot" hydrothermal vent) according to the protocol described in Tasiemski et al [77].

Cultures were incubated with peptides (at a concentration of 5-fold MICs) for 5, 10, 15, 30, 60, 120 minutes and 24 h at two temperatures (10 °C and 42 °C) to mimic the worms' thermal environment. Viabilities of *Vibrio* strains were assessed by plating the bacterial suspensions on MH agar plates (sterile petri dish, 90 mm). The plates were then incubated overnight at 28 °C and the resulting bacterial colonies were counted (CFU/mL). Three independent experiments were run in parallel and data were log-transformed (log CFU/mL).

### **2.10. Thermostability assays on ALV, ARE, POL and their ABU forms against *Vibrio* strains**

The biological activities of AMPs were evaluated after their exposition to a thermal stress (high and low temperatures). Two *Vibrio* species genetically closely related, *V. alginolyticus* and *V. diabolicus* were chosen for this assay, as representatives of the two marine environments. The 10<sup>6</sup> CFU/mL diluted culture was added (100 µL per well) into sterile 96-well flat bottom plates (CELLSTAR, Greiner bio-one) containing a volume of 10 µL/well of peptides (at a concentration of 5-fold MIC values) previously incubated for 30 minutes and 1 hour at 4, 10, 42 and 90°C. The plates were then incubated overnight at 28°C at 140 rpm. The antimicrobial activity was evaluated by measuring bacteria growth on a microtiter plate reader (Tecan Sunrise Microplate Reader) at 600 nm. The results

were expressed as percentage of bacterial growth, in comparison with the control (no AMP added to the MHB medium). The tests were conducted in triplicate.

The same experiment was also conducted in triplicate with the ABU forms to determine the role of the disulfide bonds in the thermal stability of the three native AMPs

### **2.11. pH-stability assay of ALV, ARE, POL and their ABU forms against *V. alginolyticus***

The biological activities of the three AMPs were estimated after their exposition to an acid/basic stress following the method developed by Yang *et al.* 2017 [46]. For comparative purposes, only a strain sensitive to the three AMPs was used: *V. diabolica* was excluded and *V. alginolyticus* was chosen because of its lowest MICs to all three AMPs. One colony of *V. alginolyticus* was grown in MHB medium overnight at 28°C (140 rpm). Diluted bacterial cultures ( $1 \times 10^6$  CFU/mL) were added (100 µL per well) into sterile 96-well flat bottom plates (CELLSTAR, Greiner bio-one) containing serial dilutions of peptides (from 20 to 0.00195 µg/mL), previously incubated for 3 hours at 10°C in a pH buffered solution. Four acid-to-basic 100 mM solutions (sodium acetate buffer (pH 4.0), sodium phosphate buffer (pH 6.0), Tris-HCL buffer (pH 8.0), and glycine-NaOH buffer (pH 10.0)) were tested against control samples incubated in MHB medium (pH 7.4) at 10°C. The plates were then incubated overnight at 28°C and 140 rpm. The antimicrobial activity of peptides was determined by the broth microdilution method [78]. All tests were conducted in triplicate and the results were expressed as MIC values (in µM).

The same experiment was then conducted in triplicate with the ABU forms to determine the role of the disulfide bonds in the pH stability of the three native AMPs.

## **3. RESULTS**

### **3.1. The newly identified Polaricin (POL) supports a wider sequence and structural diversification of AMPs from the BRICHOS-AMP family**

The molecular identification of prepropolaricin was performed following a tblastx on the transcriptome assembly of the Antarctic terebellid worm *Amphitritides* sp. using the sequence of the preproalvinellacin as query (Fig. 2 and Fig. S1). Pfam analysis revealed the presence of a conserved BRICHOS domain, in line with the previously described members of BRICHOS-AMP family [14–16]. Data highlighted a higher percentage of sequence identity of the BRICHOS domain compared to the AMP itself, which seems to

have evolved at a much faster rate (Table S1). As opposed to the AMP itself, the BRICHOS domain keeps a clear phylogenetic signal in our set of polychaetes with *Alvinella* and *Amphitritides* being closer one to each other when compared to other annelid species even if they inhabit opposite habitat.

POL is processed from the C-terminal part of prepropolaricin (Fig. 2). It is a cationic 19 amino acids peptide (Table S2) containing an unique cysteine residue by contrast with ARE and ALV, which respectively possess two and four cysteine residues involved in the formation of disulfide bonds that stabilize a  $\beta$ -hairpin conformation (Fig. 3).

As already evidenced by Nuclear Magnetic Resonance (NMR) [14,15], ALV and ARE share the same monomeric organization despite great differences in primary structure: two twisted antiparallel  $\beta$ -strands, forming a  $\beta$ -hairpin conformation stabilized by two or one disulfide bond(s) respectively. By contrast with ARE and ALV, MALDI-TOF mass spectrometry analyses of POL revealed that 3/4 of the polar AMP molecules dimerize presumably through an intermolecular bond engaging the cysteine residue of POL (Fig. S2). Depending on the dimerization state, the predicted 3D structure for POL, based on an *in silico* approach (see Methods) is different: The monomer adopts an alpha-helix conformation while the homodimer forms two parallel  $\beta$ -strands exposing face to face each cysteine residue. The best predicted models are presented in Figure 3.

### **3.2. BRICHOS-AMPs co-evolved with the worms' microbial community from which they are derived from**

Crossed antimicrobial assays (Table 1) were performed between bacterial strains (cultivable under MIC assay conditions) typical of the polar/temperate shallow water or hot vents, and the three BRICHOS-AMPs (*i.e.*, POL, ARE and ALV). AMPs specificities for microbial communities characterizing thermal habitats encountered by the worms were then analyzed *via* a "home *versus* away" diagram, a method commonly used in ecology to identify a local adaptation (see Material and methods) [23]. To do so, MIC values of the three AMPs against the different strains were plotted together to build the diagram (Fig. 4). The peptides which are the best locally adapted to the home site of their producers display the lowest values of MIC (higher fitness level) and *vice versa*. Analysis showed that ALV displayed a strong antimicrobial activity against all the bacteria tested with a better fitness against hydrothermal vent bacteria than against temperate/polar bacteria. ARE was also active against all the strains tested including *S. algae*, showing unlike ALV, a

better fitness against temperate-coastal bacteria than against hydrothermal vent bacteria. Among the three peptides, POL was not active against the hot vent *V. diabolicus* and has the lowest efficacy against all the other bacterial strains tested. However, its fitness was the best against *V. alginolyticus* a strain that can also be found in polar habitat. Consequently, BRICHOS-AMPs are locally adapted to (*i.e.* specifically target) the microbial community associated with the worm habitat from which they derive from.

### **3.3. BRICHOS-AMPs are fitted to handle the thermal and pH ranges sustained by the worms**

In a first step, the bactericidal activity, which reflects the requested time for an AMP to kill bacteria, was investigated through killing-time kinetics assays against the two *Vibrio* strains typical of the deep-sea 'hot' vents (*V. diabolicus*) and of the intertidal temperate/polar environment (*V. alginolyticus*) (Fig. 5). The antimicrobial efficacy was tested either at 10°C, which corresponds to the optimal temperature of *Arenicola marina* [24–32] or at 42°C, which corresponds to the optimal temperature of *A. pompejana* [33–38]. The assay was first adjusted according to the growth curve of the bacteria previously determined at the different temperatures (Fig. S3). The observed effects are consequently not reflecting a direct impact of these temperatures on the growth of these bacterial strains but rather differential AMP activities on the bacteria. Figure 5 shows that POL was slightly active against *V. alginolyticus* at 10°C and inactive against *V. diabolicus* at both 10°C and 42°C. Figure 5 shows also that ARE and ALV are bactericidal against both strains with a better efficacy against their respective surrounding *Vibrio* communities. When mimicking local temperatures, the efficiency of ARE and ALV is even more pronounced against the local strain (*i.e.*, 15 min to kill 100% of *V. diabolicus* at 42°C versus >24 h at 10°C for ALV and 30 min to kill 100% of *V. alginolyticus* at 10°C versus 24 h at 42°C for ARE).

In a second step, the thermal stabilities of the three AMPs were tested by comparing their bactericidal properties after being incubated for 30 min to 1 hour at 4 °C, 10 °C, 42 °C and even 90 °C (Fig. 6A). The “polar” AMP (POL) exerted its antimicrobial activity against *V. alginolyticus* at 4 °C and 10 °C only and lost all its activity following incubations at 42 °C and 90 °C. By contrast, the “vent” AMP, ALV remained active against both bacteria following incubations at 42 °C and 90 °C with a 50% decrease of its activity after 1 h of incubation for the highest temperature tested. Data confirmed the higher efficiency of ALV

toward *V. diabolicus*. The “temperate” ARE displayed an intermediate pattern in which the activity started to decrease by 80% at 42 °C and almost disappeared after 1 h at 90 °C with only 20% of residual activity.

The effect of the environmental pH on the three AMPs was evaluated by comparing their antimicrobial efficacies at various pH (4, 6, 7.4, 8 and 10) corresponding to the pH range of the habitat of each worm (Fig. 3B), the narrowest range being the polar environment (7.8-8.6 versus 3.9-8.5 for the hot vent). Because POL is not active against *V. diabolicus*, only *V. alginolyticus* was used for the assays (Fig. 7A). The peptides remained soluble whenever the pH. Antibacterial activities of ARE and ALV were not affected by their incubation under acid-to-basic pH conditions although POL exhibited a reduced efficiency (4-fold in the most extreme conditions pH 4 and 10) as soon as the pH is not neutral.

#### **3.4. Importance of disulphide bridges for the antibacterial activities of BRICHOS-AMPs under different thermal and pH conditions**

The more the marine worms are exposed to high temperatures and extreme pH values, the more their BRICHOS-AMP exhibits a higher content of cysteine residues engaged in disulfide bridges (Fig. 3). Cysteine replacement by alpha amino butyric acid (ABU) in the three AMPs enabled removing the intermolecular (ABU-POL) or intramolecular disulfide bonds (ABU-ALV, ABU-ARE) without modifying their main biochemical features (summarized in Table S3).

When comparing both the MIC and MBC values (Table 2) of the three AMPs and their ABU forms, ABU ALV and ABU ARE were slightly less efficient against *V. alginolyticus* while ABU POL has become slightly more active against this bacterium. Against *V. diabolicus*, the replacement of cysteines had a strong negative effect on the killing efficiency of the three AMPs. The disulfide bridges are likely to play a crucial role in the mode of action against *V. diabolicus*.

The importance of the cysteines in the thermostability of the BRICHOS-AMPs was also investigated. The ABU forms of the three BRICHOS-AMPs were exposed for 30 min to 1 hour to the three temperatures (4 °C, 10 °C and 42 °C) considered as optimal for their respective worms (Fig. 6B) before being used in antibacterial assays against *V. alginolyticus* and *V. diabolicus*. ABU-POL kept the same activity as POL (Fig. 6A) against *V. alginolyticus* at 4 °C and 10 °C. However, when incubated at higher temperatures (42 °C and 90 °C), ABU-POL still inhibited partly the growth of *V. alginolyticus* unlike the native

peptide POL, which is inactive in such conditions. ABU-POL even showed a reduced but still existing activity against this bacterium after being heated 30 min at 90 °C. Both POL and its ABU form were inactive against *V. diabollicus* whatever the AMP thermal incubations tested. ABU-ARE was as active as ARE against *V. alginolyticus* and *V. diabollicus* at 4 °C and 10 °C. At 42 °C and above, ABU-ARE became almost inactive against the two tested strains. Without disulfide bridges, ALV was only active against *V. alginolyticus* at 4 °C and 10 °C. No activity was observed at 42 °C and above.

Because POL, ABU-POL and ABU-ALV were not active against *V. diabollicus*, the importance of the disulfide bridges in the pH stability of the ABU-AMPs versus the AMPs was only investigated against *V. alginolyticus* (Fig. 7). Upon acid/basic conditions, ABU-variants kept their antibacterial activity against *V. alginolyticus* (Fig. 7B), displaying a slight decrease of activity (2-4-fold) especially under the most extreme pH conditions (4 and 10). ABU-POL was more resistant to pH changes than the native form, with an antimicrobial activity that slightly increased compared to POL.

#### 4. DISCUSSION

The members of the BRICHOS AMP family constitute a remarkably attractive model to study the evolution of AMPs, as actors of worm's immune defense in extreme and fluctuating environmental conditions. In a previous study, genetic analyses of the *ALV precursor* encoding gene in two sister and syntopic species of alvinellid worms (*A. pompejana* and *A. caudata*) sharing the same hydrothermal vent conditions, showed no differences of the AMP sequence over a geographic range spanning the 6000 km of East Pacific Rise [39][22]. Despite multiple genetic duplications the *ALV precursor* gene in both sister species [22], sharing the same biotic and abiotic environmental conditions constitutes a strong genetic purifying way of maintaining exactly the same AMP to interact with common microbiome during the evolution race [40,41].

Here, an “*in silico* approach” consisting in blasting the sequence of preproalvinellacin on the Antarctic *Amphitritides sp.* transcriptome [15] allowed us to identify a novel member of the BRICHOS-AMP family, named polaricin (POL). POL does not show any similarities with other known molecules. Focusing on BRICHOS-AMP precursors (ARE, POL and ALV) from annelids living in opposite habitats (temperate, polar, hot vent), we provided evidence for almost no similarity in AMP primary sequences and conformations, while exhibiting a nearly conserved BRICHOS domain.



477 According to the mass spectrometry analysis, the Antarctic AMP seems to adopt a  
478 dimerized form presumably through a disulfide bridge engaging each unique cysteine  
479 residue of POL, what is unusual for an AMP. The difference between the predicted  
480 structural conformations of the homodimer (two parallel beta strands) *versus* the  
481 monomeric form (alpha helix) of POL underlines puzzling mechanisms that require  
482 further investigations. The best predicted 3D structures of POL do not show any  $\beta$ -hairpin  
483 formation with two antiparallel strands unlike the solved structures of ARE and ALV.  
484 These data suggest a common origin of the precursor molecule with a clear divergent  
485 evolution of the AMP itself, which seems to be driven by strong disruptive ecological  
486 constraints associated with the worm's lifestyle.

487 According to common-garden approaches, the manipulation of environmental factors  
488 (microbial community, temperature and pH) is a powerful method to test for the role of  
489 the species *preferenda* in the evolution of their external immune defenses [1]. Using the  
490 "home *versus* away" approach, we showed that ARE and ALV are locally adapted to the  
491 bacterial communities of the worm's environment they belong to, while POL is  
492 maladapted to interact with both the shallow temperate and the deep hydrothermal-vent  
493 microbiota. According to the MIC data, ALV and ARE indeed kill faster and more efficiently  
494 the bacterial strains encountered in the habitat of their own worm. Interestingly, POL was  
495 inefficient against the hydrothermal vent *V. diabolicus* but displayed the best efficacy  
496 against *V. alginolyticus*, a bacterium that colonizes both polar and cold temperate habitats,  
497 what supports a local adaptation of this AMP. The demonstration would have been even  
498 more convincing by observing a strong activity of POL against a strictly polar marine  
499 bacterium but this requires the use of a strain cultivable under the conditions of a MIC  
500 assay. Such a strain was not available so far. Unlike *Vibrio*, *Pseudomonas* sp. isolated from  
501 hydrothermal vents was easily killed by the three AMPs regardless of their habitat of  
502 origin. The *Pseudomonas* genus is cosmopolitan to marine waters and terrestrial habitats  
503 (e.g., freshwater, soil...) and can be considered as one of the most successful bacterial  
504 groups on Earth. Worms studied here likely have encountered species from the  
505 *Pseudomonas* genus close to the species used here. Such AMP specificities support the role  
506 of AMPs in the invertebrate adaptation to environmental changes through selective  
507 constraints associated with the surrounding bacterial communities as previously  
508 exemplified in studies demonstrating their involvement in the immune defense and in the  
509 control of the vital ectosymbiosis of *A. pompejana* and of the shrimp *Rimicaris exoculata*

[15,43]. Data confirmed not only that ARE, ALV and POL display an optimal killing activity against the *Vibrio* bacterium typical of the habitat where each worm species is living, but also that this killing efficacy is optimal/kept under the thermo-chemical conditions under which the worms are living.

The correlation between species habitat and the cysteine content of the BRICHOS-AMPs led us to investigate the importance of disulfide bridges in the biological efficacy of the three molecules as a function of abiotic pressures (pH and temperature). The use of ABU-chemical variants, provided evidence of the strong involvement of internal disulfide bridges in the stability of BRICHOS-AMPs when exposed to thermal and/or pH stresses. This allowed us to conclude that the disulfide bridge patterns of ALV and ARE are not only designed for a better bacterial killing but also a way to sustain the highly fluctuations of the worm's environment (tidal emersion and seasonality for *A. marina* and chaotic mixing of hot fluid with cold seawater for *A. pompejana*) [24,32,35,44]. These results are consistent with previous data on chemically synthesized fragments (*i.e.* not naturally existing) designed from an ARE isoform [45]: the synthetic fragment containing disulfide bridges were not affected by pH variations (4.0 to 10.0) and more thermostable (range 20-80 °C) than alternative linear fragments (devoid of disulfide bridges) which also exhibited lowest antibacterial activities [46]. By ensuring the  $\beta$ -hairpin structure, the disulfide bridges confer both a higher stability and a much more efficient bactericidal activity to ALV and ARE [47–51].

Unexpectedly, ABU-POL was more effective against both mesophilic and thermophilic bacteria, displaying a more stable structure at higher temperatures and more acid or basic waters than the native dimeric form. To date, the dimerization of AMPs was mainly performed with the aim of improving the performance of synthetic peptides for therapeutic use (reviewed in Lorenzon et al., 2019 [52]). Chemical dimerization *via* the creation of disulfide bonds of some AMPs such as magainin and histatin conducted to an enhanced antimicrobial potency and a reinforced resistance to proteases while the dimerization of others such as the alpha helical aurein, reduced the antimicrobial activities by changing the mechanism of action of the peptide [53]. In the specific case of POL, the formation of homodimers might have been selected during the evolution race, probably as a response to the progressive cooling of the Antarctic continent, about 25 Mya [54,55]. Although less efficient against temperate to thermophilic bacterial strains,

peptide dimerization might be advantageous against polar psychrophilic microbes and such hypothesis as well as the structural conformation of POL still need to be investigated.

## **Conclusion and perspectives**

Based on POL, a novel BRICHOS-AMP sequence discovered in a new terebellid polychaete strictly inhabiting Antarctica, we have performed a comparative analysis between different BRICHOS-domain AMPs, specific of marine polychaetes inhabiting contrasted thermal habitats. Due to its unusual dimerization, further investigations are however required to elucidate both the structure and the mode of action of this novel polar AMP. The presence of a BRICHOS domain linked to a potent AMP and its successful evolution in polychaetes raised the matter of fully investigating on their specific molecular interaction notably by investigating the role of the BRICHOS domain as a molecular chaperone to the folding/unfolding of the host defense factors of these worms.

For the first time, we provided evidence that BRICHOS-domain AMPs evolved under strong diversifying environmental pressures to be more efficient under the ecological preferences of the worm they belong to. They are structurally shaped and adapted to exert their activities under specific conditions and against specific microbial targets. Considering the multiple interacting and changing nature of the biotic and abiotic conditions of the worms, these surveys might provide guidelines of the evolutionary response of organisms to future scenarios, like ocean warming and acidification [56–58]. The present data also support the importance of considering the optimal/natural environment of active substances when investigating and optimizing their use in therapeutic treatment of microbial diseases known to modify/alter the cellular microenvironment [59].

**Acknowledgments:** This research was funded by the Université de Lille, the Total Foundation-Ifremer (PIONEER projects) and the French Polar Institute (KREVET IPEV project: Coord. J.Y. Toullec). Renato Bruno benefitted from a Université de Lille-Total Foundation fellowship for his PhD. The French government is also warmly acknowledged for supporting this work.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## TABLES

**Table 1:** Antibacterial activities against environmental bacteria of the three BRICHOS-AMPs: the “cold” POL, “temperate” ARE and “hot” ALV. MIC: Minimal Inhibitory Concentration; MBC: Minimal Bactericidal Concentration; > not active at this concentration and beyond

HABITATS	BACTERIA	MIC (μM)			MBC (μM)		
		ALV	ARE	POL	ALV	ARE	POL
Temperate, polar Intertidal	<i>Vibrio alginolyticus</i>	0.31	0.15	0.625	0.625	0.31	5
Temperate Intertidal	<i>Vibrio fluvialis</i>	0.625	0.31	40	10	1.25	>40
Temperate Intertidal	<i>Shewanella algae</i>	>40	40	>40	>40	>40	>40
Temperate Intertidal	<i>Oceanisphaera donghaensis</i>	0.625	0.15	5	1.25	0.625	20
Hot vent Deep sea	<i>Vibrio diabolicus</i>	2.5	5	>40	20	20	>20
Hot vent Deep sea	<i>Pseudomonas sp.</i>	0.07	0.15	0.31	0.31	1.25	5

**Table 2:** Comparison of the antibacterial activities against the hot vent *Vibrio diabolicus* and the polar/temperate *Vibrio alginolyticus* of the three BRICHOS-AMPs (the “cold” POL, “temperate” ARE and “hot” ALV) versus their respective ABU variants devoid of disulfide bridges. MIC: Minimal Inhibitory Concentration; MBC: Minimal Bactericidal Concentration; > not active at this concentration and beyond

BACTERIA	MIC (μM)						MBC (μM)					
	ALV	ABU-ALV	ARE	ABU-ARE	POL	ABU-POL	ALV	ABU-ALV	ARE	ABU-ARE	POL	ABU-POL
<i>V. alginolyticus</i>	0.31	0.62	0.15	0.31	0.625	0.31	0.625	1.25	0.31	1.25	5	2.5
<i>V. diabolicus</i>	2.5	>40	5	>40	>40	>40	20	>40	20	>40	>20	>40

## References

1. Otti, O.; Tragust, S.; Feldhaar, H. Unifying External and Internal Immune Defences. *Trends Ecol. Evol.* **2014**, *29*, 625–634, doi:10.1016/j.tree.2014.09.002.
2. Rakers, S.; Niklasson, L.; Steinhagen, D.; Kruse, C.; Schaubert, J.; Sundell, K.;

- Paus, R. Antimicrobial Peptides (AMPs) from Fish Epidermis: Perspectives for Investigative Dermatology. *J. Invest. Dermatol.* **2013**, *133*, 1140–1149, doi:10.1038/jid.2012.503.
3. Wang, G. Human Antimicrobial Peptides and Proteins. **2014**, 545–594, doi:10.3390/ph7050545.
4. Zasloff, M. Antimicrobial Peptides of Multicellular Organisms. *Nature* **2002**, *415*, 389–395, doi:10.1038/415389a.
5. Stensvåg, K.; Haug, T.; Sperstad, S. V.; Rekdal, Ø.; Indrevoll, B.; Styrvold, O.B. Arasin 1, a Proline-Arginine-Rich Antimicrobial Peptide Isolated from the Spider Crab, *Hias Araneus*. *Dev. Comp. Immunol.* **2008**, *32*, 275–285, doi:10.1016/j.dci.2007.06.002.
6. Bruno, R.; Maresca, M.; Canaan, S.; Cavalier, J.; Kamel, M.; Boidin-Wichlacz, C.; Olleik, H.; Zeppilli, D.; Broidin, P.; Massol, F.; et al. Worms ' Antimicrobial Peptides. *Mar. Drugs* **2019**, *17*, doi:10.3390/md17090512.
7. Kang, H.K.; Seo, C.H.; Park, Y. Marine Peptides and Their Anti-Infective Activities. *Mar. Drugs* **2015**, *13*, 618–654, doi:10.3390/md13010618.
8. Bruno, R.; Zeppilli, D.; Boidin-wichlacz, C.; Sandulli, R. Screening for Antibacterial Molecules in Meiobenthic Nematodes Belonging to the Oncholaimidae Family. **2020**, 129–142, doi:10.21411/CBM.A.D87636D.
9. Cheung, R.C.F.; Ng, T.B.; Wong, J.H. *Marine Peptides: Bioactivities and Applications*; 2015; Vol. 13; ISBN 8523943803.
10. Nalini, S.; Sandy Richard, D.; Mohammed Riyaz, S.U.; Kavitha, G.; Inbakandan, D. Antibacterial Macro Molecules from Marine Organisms. *Int. J. Biol. Macromol.* **2018**, *115*, 696–710, doi:10.1016/j.ijbiomac.2018.04.110.
11. Bulgheresi, S. Calling the Roll on Laxus Oneistus Immune Defense Molecules. *Symbiosis* **2011**, *55*, 127–135, doi:10.1007/s13199-012-0157-3.
12. Heip, C.; Vincx, M.; Vranken, G. The Ecology of Marine Nematodes. Pdf 1985, 399–489.
13. A.Tasiemski Antimicrobial Peptides in Annelids. *Lab. Neuroimmunol. des Annelides* . **2008**, 75–82.
14. Ovchinnikova, T. V.; Aleshina, G.M.; Balandin, S. V.; Krasnodembetskaya, A.D.; Markelov, M.L.; Frolova, E.I.; Leonova, Y.F.; Tagaev, A.A.; Krasnodembsky, E.G.; Kokryakov, V.N. Purification and Primary Structure of Two Isoforms of Arenicin, a Novel Antimicrobial Peptide from Marine Polychaeta *Arenicola Marina*. *FEBS Lett.* **2004**, *577*, 209–214, doi:10.1016/j.febslet.2004.10.012.
15. Tasiemski, A.; Jung, S.; Boidin-Wichlacz, C.; Jollivet, D.; Cuvillier-Hot, V.; Pradillon, F.; Vetriani, C.; Hecht, O.; Sönnichsen, F.D.; Gelhaus, C.; et al. Characterization and Function of the First Antibiotic Isolated from a Vent Organism: The Extremophile Metazoan *Alvinella Pompejana*. *PLoS One* **2014**, *9*, 1–10, doi:10.1371/journal.pone.0095737.
16. Panteleev, P. V.; Tsarev, A. V.; Bolosov, I.A.; Paramonov, A.S.; Marggraf, M.B.; Sychev, S. V.; Shenkarev, Z.O.; Ovchinnikova, T. V. Novel Antimicrobial Peptides from the Arctic Polychaeta *Nicomache Minor* Provide New Molecular Insight into Biological Role of the BRICHOS Domain. *Mar. Drugs* **2018**, *16*, doi:10.3390/md16110401.
17. Sánchez-Pulido, L.; Devos, D.; Valencia, A. BRICHOS: A Conserved Domain in Proteins Associated with Dementia, Respiratory Distress and Cancer. *Trends Biochem. Sci.* **2002**, *27*, 329–332, doi:10.1016/S0968-0004(02)02134-5.
18. Johansson, H.; Nordling, K.; Weaver, T.E.; Johansson, J. The Brichos Domain-

- Containing C-Terminal Part of pro-Surfactant Protein C Binds to an Unfolded Poly-Val Transmembrane Segment. *J. Biol. Chem.* **2006**, *281*, 21032–21039, doi:10.1074/jbc.M603001200.
19. Hedlund, J.; Johansson, J.; Persson, B. BRICHOS - A Superfamily of Multidomain Proteins with Diverse Functions. *BMC Res. Notes* **2009**, *2*, 1–10, doi:10.1186/1756-0500-2-180.
  20. Kim, Y.; De Zoysa, M.; Lee, Y.; Whang, I.; Lee, J. BRICHOS Domain-Containing Leukocyte Cell-Derived Chemotaxin 1-like CDNA from Disk Abalone *Haliotis Discus*. *Fish Shellfish Immunol.* **2010**, *29*, 899–902, doi:10.1016/j.fsi.2010.07.021.
  21. Willander, H.; Hermansson, E.; Johansson, J.; Presto, J. BRICHOS Domain Associated with Lung Fibrosis, Dementia and Cancer - A Chaperone That Prevents Amyloid Fibril Formation? *FEBS J.* **2011**, *278*, 3893–3904, doi:10.1111/j.1742-4658.2011.08209.x.
  22. Papot, C.; Massol, F.; Jollivet, D.; Tasiemski, A. Antagonistic Evolution of an Antibiotic and Its Molecular Chaperone: How to Maintain a Vital Ectosymbiosis in a Highly Fluctuating Habitat. *Sci. Rep.* **2017**, *7*, 1454, doi:10.1038/s41598-017-01626-2.
  23. Savolainen, O.; Lascoux, M.; Merilä, J. Ecological Genomics of Local Adaptation. *Nat. Rev. Genet.* **2013**, *14*, 807–820, doi:10.1038/nrg3522.
  24. Alyakrinskaya, I.O. Some Ecological Features of the Lugworm *Arenicola Marina* L. (Annelida, Polychaeta) and Its Morphological and Biochemical Adaptations to Burrowing. **2003**, *30*, 411–418.
  25. Bat, L.; Raffaelli, D. Sediment Toxicity Testing: A Bioassay Approach Using the Amphipod *Corophium Volutator* and the Polychaete *Arenicola Marina*. **1998**, *226*, 217–239.
  26. A. Sommer, H.O.P. Exposure of *Arenicola Marina* to Extreme Temperatures: Adaptative Flexibility of a Boreal and a Subpolar Population. Pdf 1999.
  27. Sommer, A.M.; Pörtner, H.O. Metabolic Cold Adaptation in the Lugworm *Arenicola Marina*: Comparison of a North Sea and a White Sea Population. *Mar. Ecol. Prog. Ser.* **2002**, *240*, 171–182.
  28. Toulmond, A. Blood Oxygen Transport and Metabolism of the Confined Lugworm *Arenicola Marina* (L.). *J. Exp. Biol.* **1975**, *63*, 647–660.
  29. Sommer, A.; Klein, B.; Portner, H.O. Temperature Induced Anaerobiosis in Two Populations of the Polychaete Worm *Arenicola Marina* (L.). **1997**, 25–35.
  30. Juretschke, H. -P; Kamp, G. Influence of Intracellular PH on Reduction of Energy Metabolism during Hypoxia in the Lugworm *Arenicola Marina*. *J. Exp. Zool.* **1990**, *256*, 255–263, doi:10.1002/jez.1402560304.
  31. Weber, E. Respiratory Properties of Erythrocrucorin (Extracellular Hemoglobin) in the Blood of the Annelid *Arenicola marina* with Special Reference to the Salinity and Temperature. **1979**, *Ophelia*, *18*(2), 151–170.
  32. Shumway, S.E.; Davenport, J. Some Aspects of the Physiology of *Arenicola Marina* (Polychaeta) Exposed to Fluctuating Salinities. *J. mar. biol. ass. U.K.* **1977**, 907–924.
  33. Von Damm, K.L. Chemistry of Hydrothermal Vent Fluids from 9°-10°N, East Pacific Rise: "Time Zero," the Immediate Post-eruptive Period. *J. Geophys. Res. Solid Earth* **2000**, *105*, 11203–11222, doi:10.1029/1999jb900414.
  34. Le Bris, N.; Gaill, F. How Does the Annelid *Alvinella Pompejana* Deal with an Extreme Hydrothermal Environment? *Life Extrem. Environ.* **2007**, 315–339, doi:10.1007/978-1-4020-6285-8-20.

35. Desbruyères, D.; Chevaldonné, P.; Alayse, A.M.; Jollivet, D.; Lallier, F.H.; Jouin-Toulmond, C.; Zal, F.; Sarradin, P.M.; Cosson, R.; Caprais, J.C.; et al. Biology and Ecology of the “Pompeii Worm” (*Alvinella Pompejana* Desbruyères and Laubier), a Normal Dweller of an Extreme Deep-Sea Environment: A Synthesis of Current Knowledge and Recent Developments. *Deep. Res. Part II Top. Stud. Oceanogr.* **1998**, *45*, 383–422, doi:10.1016/S0967-0645(97)00083-0.
36. Le Bris, N.; Zbinden, M.; Gaill, F. Processes Controlling the Physico-Chemical Micro-Environments Associated with Pompeii Worms. *Deep. Res. Part I Oceanogr. Res. Pap.* **2005**, *52*, 1071–1083, doi:10.1016/j.dsr.2005.01.003.
37. Di Meo-Savoie, C.; Luther, G.W.; Cary, S.C. Physicochemical Characterization of the Microhabitat of the Epibionts Associated with *Alvinella Pompejana*, a Hydrothermal Vent Annelid. **2004**, *68*, 2055–2066, doi:10.1016/j.gca.2003.10.039.
38. Ravaux, J.; Hamel, G.; Zbinden, M.; Tasiemski, A.A.; Boutet, I.; Léger, N.; Tanguy, A.; Jollivet, D.; Shillito, B. Thermal Limit for Metazoan Life in Question: In Vivo Heat Tolerance of the Pompeii Worm. *PLoS One* **2013**, *8*, 4–9, doi:10.1371/journal.pone.0064074.
39. Bulle, C.T.; Bertrand, D.; Nagarajan, N.; Copley, R.R.; Corre, E. Genomic Patterns of Divergence in the Early and Late Steps of Speciation of the Deep - Sea Vent Thermophilic Worms of the Genus *Alvinella*. *BMC Ecol. Evol.* **2022**, 1–17, doi:10.1186/s12862-022-02057-y.
40. Bulmer, M.S.; Crozier, R.H. Variation in Positive Selection in Termite GNBPs and Relish. **2004**, doi:10.1093/molbev/msj037.
41. Fuller, C.A.; Postava-Davignon, M.A.; West, A.; Rosengaus, R.B. Environmental Conditions and Their Impact on Immunocompetence and Pathogen Susceptibility of the Caribbean Termite *Nasutitermes Acajutlae*. *Ecol. Entomol.* **2011**, *36*, 459–470, doi:10.1111/j.1365-2311.2011.01289.x.
42. Erler, S.; Lhomme, P.; Rasmont, P.; Lattorff, H.M.G. Infection , Genetics and Evolution Rapid Evolution of Antimicrobial Peptide Genes in an Insect Host – Social Parasite System. *Infect. Genet. Evol.* **2014**, *23*, 129–137, doi:10.1016/j.meegid.2014.02.002.
43. Bloa, S. Le; Boidin-wichlacz, C.; Cueff-gauchard, V.; Pradillon, F.; Tasiemski, A. Antimicrobial Peptides and Ectosymbiotic Relationships : Involvement of a Novel Type IIa Crustin in the Life Cycle of a Deep-Sea Vent Shrimp. **2020**, *11*, 1–18, doi:10.3389/fimmu.2020.01511.
44. Pradillon, F.; Bris, N. Le; Shillito, B.; Young, C.M.; Gaill, F. Influence of Environmental Conditions on Early Development of the Hydrothermal Vent Polychaete *Alvinella Pompejana*. **2005**, 1551–1561, doi:10.1242/jeb.01567.
45. Yang, N.; Wang, X.; Teng, D.; Mao, R.; Hao, Y.; Feng, X.; Wang, J. Deleting the First Disulphide Bond in an Arenicin Derivative Enhances Its Expression in *Pichia Pastoris*. *Lett. Appl. Microbiol.* **2017**, doi:10.1111/lam.12770.
46. Yang, N.; Liu, X.; Teng, D.; Li, Z.; Wang, X.; Mao, R. Antibacterial and Detoxifying Activity of NZ17074 Analogues with Multi-Layers of Selective Antimicrobial Actions against *Escherichia Coli* and *Salmonella Enteritidis*. *Sci. Rep.* **2017**, 1–19, doi:10.1038/s41598-017-03664-2.
47. Ju-Un Lee Dong-Il Kang, Wan Long Zhu, Song Yub Shin, Kyung-Soo Hahm, Y.K. Solution Structures and Biological Functions of the Antimicrobial Arenicin-1, and Its Linear Derivative. *Biopolym. - Pept. Sci. Sect.* **2007**, *88*, 208–216, doi:10.1002/bip.
48. Andrä, J.; Hammer, M.U.; Grötzinger, J.; Jakovkin, I.; Lindner, B.; Vollmer, E.;

- Fedders, H.; Leippe, M.; Gutschmann, T. Significance of the Cyclic Structure and of Arginine Residues for the Antibacterial Activity of Arenicin-1 and Its Interaction with Phospholipid and Lipopolysaccharide Model Membranes. *Biol. Chem.* **2009**, *390*, 337–349, doi:10.1515/BC.2009.039.
49. Lai, J.R.; Huck, B.R.; Weisblum, B.; Gellman, S.H. Design of Non-Cysteine-Containing Antimicrobial -Hairpins : Structure - Activity Relationship Studies with Linear Protegrin-1 Analogues †. **2002**, 12835–12842.
50. Nan, Y.H.; Jacob, B.; Yub, S. Linear Bactenecin Analogs with Cell Selectivity and Anti-Endotoxic Activity. **2012**, 740–747, doi:10.1002/psc.2460.
51. Wu, M.; Hancock, R.E.W.; Bacs, L. Interaction of the Cyclic Antimicrobial Cationic Peptide Bactenecin with the Outer and Cytoplasmic Membrane \* Showed Activity against the Gram-Positive Bacteria. **1999**, 274, 29–35.
52. Lorenzon, E.N.; Piccoli, J.P.; Santos-filho, N.A.; Cilli, E.M. Dimerization of Antimicrobial Peptides: A Promising Strategy to Enhance Antimicrobial Peptide Activity. **2019**, 98–107, doi:10.2174/0929866526666190102125304.
53. Lorenzon, E.N.; Sanches, P.R.S.; Nogueira, L.G.; Bauab, T.M.; Cilli, E.M. Dimerization of Aurein 1.2 : Effects in Structure , Antimicrobial Activity and Aggregation of Candida Albicans Cells. **2013**, 1521–1528, doi:10.1007/s00726-013-1475-3.
54. Rogers, A.D.; Murphy, E. j.; Johnston, N.M.; Clarke, A. Introduction . Antarctic Ecology : From Genes to Ecosystems . Part 2 . Evolution , Diversity and Functional Ecology. **2007**, 2187–2189, doi:10.1098/rstb.2007.2135.
55. Cowart, D.A.; Schiaparelli, S.; Alvaro, M.C.; Cecchetto, M.; Le, S.; Didier, P.; Stephane, J. Origin , Diversity , and Biogeography of Antarctic Scale Worms ( Polychaeta : Polynoidae ) : A Wide- - Scale Barcoding Approach. **2022**, 1–17, doi:10.1002/ece3.9093.
56. Mcculloch, M.; Falter, J.; Trotter, J.; Montagna, P. Coral Resilience to Ocean Acidification and Global Warming through PH Up-Regulation. *Nat. Clim. Chang.* **2012**, 2, 1–5, doi:10.1038/nclimate1473.
57. Wilkins, L.G.E.; Leray, M.; Yuen, B.; Peixoto, R. Host-Associated Microbiomes and Their Roles in Marine Ecosystem Functions. **2019**, 1–27.
58. Sternberg, E.D.; Thomas, M.B. Local Adaptation to Temperature and the Implications for Vector-Borne Diseases. *Trends Parasitol.* **2014**, 1–8, doi:10.1016/j.pt.2013.12.010.
59. Crabbé, A.; Ledesma, M.A.; Nickerson, C.A. Mimicking the Host and Its Microenvironment in Vitro for Studying Mucosal Infections by Pseudomonas Aeruginosa. *Pathog. Dis.* **2014**, 1–19, doi:10.1111/2049-632X.12180.
60. Bulletin, B. The Feeding and Ecology of Two North Pacific Abarenicola Species ( Arenicolidae , Polychaeta ) Author ( s ) : Katharine D . Hobson Reviewed Work ( s ) : Published by : Marine Biological Laboratory. **1967**, 133, 343–354.
61. Riisgard, H.U.; Banta, G.T. Irrigation and Deposit Feeding by the Lugworm Arenicola Marina, Characteristics and Secondary Effects on the Environment. A Review of Current Knowledge 1998, 243–257.
62. Desbruyères, D.; Laubier, L. *Alvinella Pompejana* Gen. Sp. Nov., Ampharetidae Aberrant Des Sources Hydrothermales de La Ride Est-Pacifique. *Oceanol. Acta* **1980**, 3, 267–274.
63. Thiel, A.; Portner, H.O.; Arntz, W.E. Marine Life at Low Temperatures - a Comparison of Polar and Deep-Sea Characteristics 1996, 183–219.
64. Aoki, S.; Rintoul, S.R.; Ushio, S.; Watanabe, S.; Bindoff, N.L. Freshening of the



- Adélie Land Bottom Water near 140 ° E. *Geophys. Res. Lett.* **2005**, 32, 2–5, doi:10.1029/2005GL024246.
65. Matson, P.G.; Martz, T.R.; Hofmann, G.E. High-Frequency Observations of PH under Antarctic Sea Ice in the Southern Ross Sea. **2011**, 7, 1–7, doi:10.1017/S0954102011000551.
66. Mcneil, B.I.; Tagliabue, A.; Sweeney, C. A Multi - Decadal Delay in the Onset of Corrosive ‘ Acidified ’ Waters in the Ross Sea of Antarctica Due to Strong Air - Sea CO 2 Disequilibrium. **2010**, 37, 1–5, doi:10.1029/2010GL044597.
67. Mabrouk, K.; Ram, N.; Boisseau, S.; Strappazzon, F.; Rehaïm, A. Critical Amino Acid Residues of Maurocalcine Involved in Pharmacology , Lipid Interaction and Cell Penetration. **2007**, 1768, 2528–2540, doi:10.1016/j.bbamem.2007.06.030.
68. Merrifield, B. Solid Phase Synthesis. *Science (80-. )*. **1986**, 232, 341–347.
69. Fajloun, Z.; Ferrat, G.; Carlier, E.; Fathallah, M.; Lecomte, C.; Sandoz, G.; Luccio, E.; Mabrouk, K.; Legros, C.; Darbon, H.; et al. Synthesis , 1 H NMR Structure , and Activity of a Three-Disulfide- Bridged Maurotoxin Analog Designed to Restore the Consensus Motif of Scorpion Toxins \*. **2000**, 275, 13605–13612.
70. Sabatier, J.; Mabrouk, K.; Rochat, H. Maurotoxin, pi1 and hstx1 derivatives 2009, 1, 1–6.
71. Lamiable, A.; Th, P.; Rey, J.; Vavrusa, M.; Derreumaux, P.; Tuff, P. PEP-FOLD3 : Faster Denovo Structure Prediction for Linear Peptides in Solution and in Complex. **2016**, 44, 449–454, doi:10.1093/nar/gkw329.
72. Raguénès, G.; Christen, R.; Guezennec, J.; Pignet, P.; Barbier, G. *Vibrio Diabolicus* Sp. Nov., a New Polysaccharide-Secreting Organism Isolated from a Deep-Sea Hydrothermal Vent Polychaete Annelid, *Alvinella Pompejana*. *Int. J. Syst. Bacteriol.* **1997**, 47, 989–995, doi:10.1099/00207713-47-4-989.
73. Cuvillier-hot, V.; Marylène, S.; Massol, F.; Boidin-wichlacz, C.; Pennel, T.; Lesven, L.; Net, S.; Papot, C.; Ravaux, J.; Vekemans, X.; et al. Science of the Total Environment Immune Failure Reveals Vulnerability of Populations Exposed to Pollution in the Bioindicator Species Hediste Diversicolor. *Sci. Total Environ.* **2017**, doi:10.1016/j.scitotenv.2017.08.259.
74. Urakawa, H.; Rivera, I.N.G. The Biology of Vibrios 2006.
75. Holt, H.M.; Bruun, B. *Shewanella* Algae and *Shewanella* Putrefaciens : Clinical and Microbiological Characteristics. *Eur. Soc. Clin. Infect. Dis.* **2005**, 11, 347–352, doi:10.1111/j.1469-0691.2005.01108.x.
76. Bulet, P.; Cociancich, S.; Dimarcq, J.; Lambert, J.; Reichhart, J.; Hoffmann, D.; Hetru, C.; Hoffmanns, J.A. Isolation From a Coleopteran Insect of a Novel Inducible Antibacterial Peptide and of New Members of the Insect Defensin Family. **1991**, 266, 24520–24525.
77. Tasiemski, A.; Vandenbulcke, F.; Mitta, G.; Lemoine, J.; Lefebvre, C.; Sautière, P.E.; Salzert, M. Molecular Characterization of Two Novel Antibacterial Peptides Inducible upon Bacterial Challenge in an Annelid, the Leech *Theromyzon Tessulatum*. *J. Biol. Chem.* **2004**, 279, 30973–30982, doi:10.1074/jbc.M312156200.
78. Wiegand, I.; Hilpert, K.; Hancock, R.E.W. Agar and Broth Dilution Methods to Determine the Minimal Inhibitory Concentration ( MIC ) of Antimicrobial Substances. **2008**, 3, 163–175, doi:10.1038/nprot.2007.521.

## FIGURE LEGENDS

**Figure 1:** Geographical distribution of the three polychaetes studied together with the description of their marine habitats.

**Figure 2:** Alignment of the precursors of the BRICHOS-AMP family members: polaricin (POL), nicomicin, alvinellacin (ALV) and arenicin (ARE). BRICHOS and AMP regions are respectively in black and dotted black rectangles. In the AMP part, cysteine residues involved in disulfide bridges are in black circles. The number of residues relative to the N-terminal segment of the precursor is shown on the right side of each sequence. The color of amino-acids background underlines the higher level of conservation in the BRICHOS part as compared to the AMP part itself. The alignment was generated using CLC Sequence Viewer software (version 8.0).

**Figure 3:** Three-dimensional structures of ALV, ARE and the predicted ones of POL generated by Alpha Fold software. AlphaFold produces a per-residue confidence score (pLDDT) between 0 and 100: Very high confidence (pLDDT > 90); Confident (90 > pLDDT > 70); Low (70 > pLDDT > 50); Very low (pLDDT < 50). Number of cysteines in the three AMPs as a function of thermal and pH encountered in the habitat of each worm.


**Figure 4:** A representation of BRICHOS-AMPs local adaptation (home vs. away diagram) to temperate-coastal and hydrothermal vents-deep sea habitats, using their MIC values against environmental bacterial strains. Y-axis, representative of fitness factor, stands for peptides MIC log-transformed data: higher MIC values correspond to lower fitness levels. The colored areas gather the values belonging to the same peptide. Dashed lines, connecting the areas across the two habitats, show the molecular adaptation of ARE and ALV to their respective habitats, displaying lower MIC values against “home strains” and *vice versa*. Missing polar bacterial strains, POL displays the highest MIC values against the investigated “away strains”. Peptides and bacteria sharing a common environment are displayed by the same color: red for hot vent, blue for temperate and purple for polar. The cases of not-active peptide are represented with – on the top of the diagram, as leaning to infinity.

**Figure 5:** Kinetics profiles of the bactericidal activities of ALV, ARE and POL against *V. alginolyticus* and *V. diabolicus*, at 10 °C and 42 °C. The results are shown as time-plot of *Vibrio* spp. viability ( $\log_{10}$  of CFU/mL), without (control) and with peptides. Temperatures, peptides and strains belonging to the temperate, hydrothermal vent and polar environments are in blue, red and purple respectively. The dotted grey lines are the negative controls with no peptides.

**Figure 6:** Effect of temperature on biological activity of (A) native peptides and (B) ABU-peptides, against *V. alginolyticus* and *V. diabolicus*, after 30 minutes to 1 hour of peptide incubation at different temperatures (4, 10, 42 and 90 °C). Each bar is expressed as the average with error bars, representing  $\pm$ SD from 3 independent experiments (n = 3). The Y-axis (in A) displays a break in the range of values to improve its readability.


**Figure 7:** Effect of pH on the biological activity of (A) native peptides and (B) ABU-peptides activities, against *V. alginolyticus*. The tests were performed after peptide incubation (3 hours) under acid/basic conditions (at pH 4, 6, 8 and 10). The control is represented by peptide MIC values in MHB medium (pH 7.4) for each peptide.

## Graphical abstract



- Local adaptation of AMPs to the ecological niche of the producers

Figure 1



Annelid species	<i>Amphitritides</i> <i>sp.</i>	<i>Arenicola</i> <i>marina</i>	<i>Alvinella</i> <i>pompejana</i>
AMP name	Polaricin (POL)	Arenicin (ARE)	Alvinellacin (ALV)
Environment	Polar, coastal	Temperate, coastal	Hot chimneys of hydrothermal vents, deep sea,
Distribution	Antarctica (Adelie Land)	North Sea	East Pacific Rise 9 North



Figure 2







Figure 3

	POL	ARE	ALV
Structure	 PDB : 2JSB	 PDB : 2LLR	 PDB : 2LLR
Number of Cys	1	2	4
Number of intramolecular disulfide bridges	0	1	2

Thermal range of the worms' habitat

			<i>Amphitritides sp.</i>	<i>Arenicola marina</i>	<i>Alvinella pompejana</i>
HABITAT	T°	Range	-1.8 °C – 1 °C	5 °C – 25 °C	2 °C - >100 °C
		Optimum	~0 °C	~10 °C	~42 °C

Figure 4






Figure 5




Figure 6





Figure 7



pH range of the worms' habitat

HABITAT	pH	range	<i>Amphitritides sp.</i>	<i>Arenicola marina</i>	<i>Alvinella pompejana</i>
			7.8 – 8.6	5.4 – 8.4	3.9 – 8.5
HABITAT	pH	optimum	7.5 - 8	6.5 - 8	5.7 - 8