



HAL
open science

SL(2,C) Scheme Processing of Singularities in Quantum Computing and Genetics

Michel Planat, Marcelo M Amaral, David Chester, Klee Irwin

► **To cite this version:**

Michel Planat, Marcelo M Amaral, David Chester, Klee Irwin. SL(2,C) Scheme Processing of Singularities in Quantum Computing and Genetics. *Axioms*, 2023, 12 (3), pp.12030233 (12). hal-04283936

HAL Id: hal-04283936

<https://hal.science/hal-04283936>

Submitted on 14 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

$SL(2, \mathbb{C})$ Scheme Processing of Singularities in Quantum Computing and Genetics

Michel Planat ^{1,*}, Marcelo M. Amaral ^{2,†}, David Chester ^{2,†} and Klee Irwin ^{2,†}

¹ Institut FEMTO-ST CNRS UMR 6174, Université de Bourgogne-Franche-Comté, F-25044 Besançon, France

² Quantum Gravity Research, Los Angeles, CA 90290, USA; Marcelo@QuantumGravityResearch.org (M.M.A.); DavidC@QuantumGravityResearch.org (M.M.A.); Klee@QuantumGravityResearch.org (K.I.)

* Correspondence: michel.planat@femto-st.fr

† These authors contributed equally to this work.

Abstract: Revealing the time structure of physical or biological objects is usually performed thanks to the tools of signal processing like the fast Fourier transform, Ramanujan sum signal processing and many other techniques. For space-time topological objects in physics and biology, we propose a type of algebraic processing based on schemes in which the discrimination of singularities within objects is based on the space-time-spin group $SL(2, \mathbb{C})$. Such topological objects possess an homotopy structure encoded in their fundamental group and the related $SL(2, \mathbb{C})$ multivariate polynomial character variety contains a plethora of singularities somehow analogous to the frequency spectrum in time structures. Our approach is applied to a model of quantum computing based on an Akbulut cork in exotic R_4 , to an hyperbolic model of topological quantum computing based on magic states and to microRNAs in genetics. Such diverse topics reveal the manifold of possibilities of using the concept of a scheme spectrum.

Keywords: Finitely generated group; $SL(2, \mathbb{C})$ character variety; algebraic surfaces; schemes; exotic R_4 ; topological quantum computing; microRNAs.

1. Introduction

In signal processing of a time series, the lines of the Fourier spectrum are described by discontinuities. The approach may be generalized with number theory by taking Ramanujan sums as the building blocks of the signal expansion, e.g. see [1].

A more ambitious approach is to use algebraic geometry to reveal the singularities of an object. In recent papers, we explored topics of quantum computing and DNA biology with a common algebraic geometrical tool that we now call $SL(2, \mathbb{C})$ scheme processing. The group $SL(2, \mathbb{C})$ is found in physics in Einstein's field equations [2] and in loop quantum gravity [3]. For us, the necessary ingredient is a finitely generated group π expressing the symmetries of the investigated object. The $SL(2, \mathbb{C})$ character variety [4] associated to π is determined and summarized by its Groebner basis \mathcal{G} [5,6]. The multivariate polynomials in \mathcal{G} may contain isolated (or non-isolated) singularities that characterize the richness of the object. In this essay, the polynomials are reduced to surfaces living in the 3-dimensional projective space $P^3(\mathbb{Q})$ over the rationals.

Isolated singularities of a surface are easy to characterize and classify according to their A-D-E type in the McKay correspondence [7]. For a non zero dimensional singular subset in the surface, we may use tools of schemes for the resolution of singularities [8]. Scheme theory was created by A. Grothendieck to generalize smooth manifolds to algebraic varieties possibly decorated with singularities [9,10]. For our purpose, it is enough to see a scheme as a geometrical object defined by the vanishing of polynomials defined over an affine (or a projective space) like those living in \mathcal{G} .

Previously our approach was restricted to algebraic surfaces containing isolated singularities in applications to topological quantum computing [11] and DNA-RNA transcription

and regulation [12]. From now, $SL(2, \mathbb{C})$ scheme processing is a good option to manage general singular sets. This is needed in our model of quantum computing based on an Akbulut cork in exotic R_4 [13], in our hyperbolic model of topological quantum computing based on magic states [14] and for some transcription factors and microRNAs in genetics [12].

In Section 2, we briefly describe the mathematical formalism used in our paper. It includes the definition of the character variety \mathcal{V} representing the finitely generated group π over the group $SL(2, \mathbb{C})$ and how a Groebner basis \mathcal{G} is obtained from \mathcal{V} in practice. The section mentions the distinction between simple singularities and singularities whose support is not zero dimensional. We then investigate the algebraic geometry of three types of complex objects in physics and biology. The first two objects rely on quantum computing following our previous papers [11,13,14]. Section 3 discusses the symmetries and related representations of the Akbulut cork W , a fundamental object in the theory of exotic 4-manifolds. Section 4 refers to the symmetries and the related representations of an hyperbolic 3-manifold found in the context of magic states in quantum computing. Section 5 investigates a third class of objects, which are a family of biological molecules called microRNAs that regulate the types and amounts of proteins [12]. In Section 6, we provide commentary on our results.

2. Theory

Details about the theory are described below. The corresponding implementation is performed on Sage [15] and Magma [16].

2.1. The $SL(2, \mathbb{C})$ character variety of a finitely generated group and a Groebner basis

Let f_p be a finitely generated group, we describe the representations of f_p in the (double cover alias the group extension of order two of the Lorentz group) $SL(2, \mathbb{C})$, the group of (2×2) matrices with complex entries and determinant 1. The group $SL(2, \mathbb{C})$ may be seen simultaneously as a 'space-time' (a Lorentz group) and a 'quantum' (a spin) group.

Representations of f_p in $SL(2, \mathbb{C})$ are homomorphisms $\rho : f_p \rightarrow SL(2, \mathbb{C})$ with character $\kappa_\rho(g) = \text{tr}(\rho(g))$, $g \in f_p$. The set of characters allows to define an algebraic set by taking the quotient of the set of representations ρ by the group $SL(2, \mathbb{C})$, which acts by conjugation on representations [17].

Such an algebraic set is called the $SL(2, \mathbb{C})$ character variety of f_p . It is made of a sequence of multivariate polynomials called a scheme X . The vanishing of polynomials defines the ideal $\mathcal{I}(X)$ of the scheme X . A Groebner basis $\mathcal{G}(X)$ is a particular set of the polynomial ring $\mathcal{I}(X)$ that has to follow algorithmic rules (similar to the Euclidean division for univariate polynomials).

For the effective calculations of the character variety, we make use of a software on Sage [15]. We also need Magma [16] for the calculation of a Groebner basis, at least for 3- and 4-letter sequences.

2.2. Singularities of an algebraic surface

Simple singularities

The surfaces S of interest in this case are said to be almost not singular in the sense that they have at worst simple singularities. In Magma, it is referred to a simple or A-D-E singularity [7] if it is an isolated singularity on S which is analytically of the type A_n , $n \geq 1, D_n$, $n \geq 4, E_6, E_7$ or E_8 .

The A-D-E type and the number of simple singularities is reflected in our notation. E.g., $S^{(IA_1)}$ means a surface with l singularities of type A_1 , $S^{(A_2)}$ means a surface with a single singularity of type A_2 and $S^{(D_4)}$ means a surface with a single singularity of type D_4 . The Cayley cubic encountered in our previous paper is $\kappa_4^{(4A_1)}(x, y, z) = xyz + x^2 + y^2 + z^2 - 4$ [11]. The Fricke surface of type D_4 is $S^{(D_4)} = xyz + x^2 + y^2 + z^2 - 8(x + y + z) + 28$ [18, Figure 16]. Several other examples can be found in [12].

The relevant Magma command for such simply singular surfaces is *HasOnlySimpleSingularities(S)*.

2.3. Arbitrary singularities

Let $X \in P^3(\mathbb{Q})$ be a singular surface (a surface containing a non zero dimensional singular subset of points). Let $Y \in P^3(\mathbb{Q})$ be a regular surface (devoid of singularities) above X that is, a representation (in a sense to be qualified) $\rho : Y \rightarrow X$. The set of such morphisms ρ belongs to the so-called spectrum $\text{Spec}_X(Y)$ of Y in X .

In our context, the formal desingularization of an (hyper)surface X in $P^3(\mathbb{Q})$ is realized with a proper birational map $Y \rightarrow X$, with Y is regular. The formal desingularization is realized with the introduction of a formal prime divisor $\text{Spec } \widehat{\mathcal{O}_{X,p}} \rightarrow X$, where $p \in X$ is a regular point of codimension 1, $\mathcal{O}_{X,p}$ is the structure sheaf at p of X seen as a scheme and the hat means the completion [19].

Taking the affine surface $S(x, y, z)$, the related homogeneous polynomial defines a hypersurface $X(x, y, z, w) \in P^3(\mathbb{Q})$ so that a formal prime divisor is actually a morphism $\phi : \text{Spec}_\phi[[t]] \rightarrow X$ defined by the \mathbb{Q} -algebra homomorphism $\phi^\# : \mathbb{Q}([x, y, z, w] / \langle X \rangle \rightarrow \mathbb{F}_\phi[[t]]$ where $\mathbb{F}_\phi = \mathbb{Q}(s)[\alpha]$, s a parameter and α is defined by the vanishing of a minimal polynomial.

The relevant Magma command for this case is *FormallyResolveProjectiveHyperSurface(S: AdjComp := true)*. Thanks to the setting *AdjComp := true*, the command returns the number of essential singularities allowing to obtain the formal divisors needed to only compute birational invariants or adjoint spaces [19,20].




Figure 1. The affine singular surface $S_2(x, y, z) = z^4 + 2yz^3 + x^2 - 6yz - 2x - 8$ found in the Groebner basis for the transcription factor Prdm1 [12, Section 3.1].

2.4. Kodaira-Enriques classification

Given an ordinary projective surface S in the projective space P^3 over a number field, if S is birationally equivalent to a rational surface, the software Magma [16] determines the map to such a rational surface and returns its type within five categories. The returned type of S is P^2 for the projective plane, a quadric surface (for a degree 2 surface in P^3), a rational ruled surface, a conic bundle or a degree p Del Pezzo surface where $1 \leq p \leq 9$.

A further classification may be obtained for S in P^3 if S has at most point singularities (unless the singularities may be formally resolved as we described in the previous subsection for the case of characteristic zero). Magma computes the type of S (or rather, the type of the non-singular projective surfaces in its birational equivalence class) according to the classification of Kodaira and Enriques [21]. The first returned value is the Kodaira dimension of S , which is $-\infty$, 0, 1 or 2. The second returned value further specifies the type within the Kodaira dimension $-\infty$ or 0 cases (and is irrelevant in the other two cases).

Kodaira dimension $-\infty$ corresponds to birationally ruled surfaces. The second return in this case is the irregularity $q \geq 0$ of S . So S is birationally equivalent to a ruled surface over a smooth curve of genus q and is a rational surface if and only if q is zero.

Kodaira dimension 0 corresponds to surfaces which are birationally equivalent to a K_3 surface, an Enriques surface, a torus or a bi-elliptic surface.

Every surface of Kodaira dimension 1 is an elliptic surface (or a quasi-elliptic surface in characteristics 2 or 3), but the converse is not true: an elliptic surface can have Kodaira dimension $-\infty$, 0 or 1.

Surfaces of Kodaira dimension 2 are algebraic surfaces of general type.

A singular surface

Let us illustrate our approach with a selected singular surface encountered in the context of the transcription factor Prdm1 in our recent paper [12, Section 3.1]. The affine surface under question is $S_2(x, y, z) = z^4 + 2yz^3 + x^2 - 6yz - 2x - 8$. It contains 9 essential singularities in the desingularization. A 3-dimensional plot of $S_2(x, y, z)$ is in Figure 1. The surface is a conic bundle of K_3 type.

The projective surface is $S_2(x, y, z, w) = z^4 + 2yz^3 + w^2(x^2 - 6yz) - 2xw^3 - 8w^4$. There exists 5 distinct types of minimal polynomial. For one of the divisors, the minimal polynomial is $a^2 + 2s + 1$ and the prime divisor in the desingularization is the morphism $\phi : \text{Spec}_\phi[[t]] \rightarrow X$ defined by

$$x \rightarrow 1, \quad y \rightarrow st, \quad z \rightarrow t, \quad w \rightarrow \alpha t^2 + 0(t^4).$$

The 9 formal morphisms are used to compute the global sections (or adjoints) $\text{Hom}_{i,j}$. For $i, j < 3$, one obtains $\text{Hom}_{1,1} = \text{Hom}_{2,1} = \text{Hom}_{2,2} = \text{Hom}_{2,3} = \text{Hom}_{3,1} = \text{Hom}_{3,2} = \text{Hom}_{3,3} = 0$, $\text{Hom}_{1,2} = [z, w]$, $\text{Hom}_{1,3} = [xz, yz, z^2, xw, yw, zw, w^2]$.

3. $SL(2, \mathbb{C})$ scheme processing in quantum computing based on an Akbulut cork

3.1. A short account of magic states for quantum computing

A quantum state is called magic when it is added to the eigenstates of the d -dimensional Pauli group to obtain universal quantum computation [22,23]. A subset of magic states consists of states associated to minimal informationally complete measurements, also called MIC states, see [14] and references therein. In the context of a 3- or 4-manifold \mathcal{M} , of fundamental group $\pi_1(\mathcal{M})$, a subgroup of index d of $\pi_1(\mathcal{M})$ has cosets organized according to a permutation group \mathcal{P}_d . Then, a permutation is also a permutation matrix and a possible source of MIC states arising from its eigenstates. Finally, for appropriate subgroups of $\pi_1(\mathcal{M})$, a finite geometry \mathcal{F}_d is found from the two-point stabilizer subgroup of \mathcal{P}_d . The coset coordinates of \mathcal{F}_d can be made in one to one correspondence with d -dimensional quantum observables. For example, important quantum geometries such as the Mermin 3×3 grid (at index $d = 9$), the generalized quadrangle of order two $GQ(2 \times 2)$ (at index $d = 15$) are encoding two-qubit contextuality [14].

The Akbulut cork below belongs to the theory of 4-manifolds and features magic states of quantum computing [13]. To our earlier work we add the investigation of $SL(2, \mathbb{C})$ character variety of the relevant fundamental groups. In particular, we put the Akbulut cork in a new perspective, by revealing the singularity spectrum of its $SL(2, \mathbb{C})$ character variety.

3.2. Brief introduction to 4-manifolds

The theory of 4-manifolds is described in books [24–26]. Here we are interested in the decomposition of a 4-manifold into one- and two-dimensional handles as shown in Fig. 2 [24, Fig. 1.1 and Fig. 1.2]. Let B^n and S^n be the n -dimensional ball and the n -dimensional sphere, respectively. An observer is placed at the boundary $\partial B^4 = S^3$ of the 0-handle B^4 and watch the attaching regions of the 1- and 2-handles. The attaching region of 1-handle is a pair of balls B^3 (the yellow balls), and the attaching region of 2-handles is a framed knot (the red knotted circle) or a knot going over the 1-handle (shown in blue). For closed 4-manifolds, there is no need of visualizing a 3-handle since it can be directly attached to the 0-handle. The 1-handle can also be figured out as a dotted circle $S^1 \times B^3$ obtained by

squeezing together the two three-dimensional balls B^3 so that they become flat and close together [25, p. 169] as shown in Fig. 2b.

For the attaching region of a 2- and a 3-handle one needs to enrich our knowledge by introducing the concept of an Akbulut cork to be described in the next paragraph [13, Figure 3].




Figure 2. (a) Handlebody of a 4-manifold with the structure of 1- and 2-handles over the 0-handle B^4 , (b) the structure of a 1-handle as a dotted circle $S^1 \times B^3$, (c) an Akbulut cork $W = 9_{46}(-1, 1)$.

3.3. Akbulut cork

A Mazur manifold is a contractible, compact, smooth 4-manifold (with boundary) not diffeomorphic to the standard 4-ball B^4 [24]. Its boundary is a homology 3-sphere. If we restrict to Mazur manifolds that have a handle decomposition into a single 0-handle, a single 1-handle and a single 2-handle then the manifold has to be of the form of the dotted circle $S^1 \times B^3$ (as in Fig. 2b) (right) union a 2-handle. The simplest object of this type is the Akbulut cork shown in Fig. 2c [27,28].

Given p, q, r (with $p \leq q \leq r$), the Brieskorn 3-manifold $\Sigma(p, q, r)$ is the intersection in the complex 3-space \mathbb{C}^3 of the 5-dimensional sphere S^5 with the surface of equation $z_1^p + z_2^q + z_3^r = 0$. The smallest known Mazur manifold is the Akbulut cork W and its boundary is the Brieskorn homology sphere $\Sigma(2, 5, 7)$. The Akbulut cork has a simple definition in terms of the framings ± 1 of $(-3, 3, -3)$ pretzel knot also called $K = 9_{46}$ [29, Fig. 3]. It has been shown that $\partial W = \Sigma(2, 5, 7) = K(1, 1)$ and $W = K(-1, 1)$.

An exotic R^4 is a differentiable manifold that is homeomorphic but not diffeomorphic to the Euclidean space \mathbb{R}^4 . An exotic R^4 is called small if it can be smoothly embedded as an open subset of the standard \mathbb{R}^4 and is called large otherwise. Here we are concerned with an example of a small exotic R^4 .

According to [29], there exists an involution $f : \partial W \rightarrow \partial W$ that surgers the dotted 1-handle $S^1 \times B^3$ to the 0-framed 2-handle $S^2 \times B^2$ and back, in the interior of W . There exists a smooth contractible 4-manifold V with $\partial V = \partial W$, such that V is homeomorphic but not diffeomorphic to W relative to the boundary [27, Theorem 1]. This leads us to our next paragraph.

3.4. The manifold \bar{W} mediating the Akbulut cobordism between exotic manifolds V and W

A cobordism between two oriented m -manifolds M and N is any oriented $(m + 1)$ -manifold W_0 such that the boundary is $\partial W_0 = \bar{M} \cup N$, where M appears with the reverse orientation. The cobordism $M \times [0, 1]$ is called the trivial cobordism. Next, a cobordism W_0 between M and N is called an h-cobordism if W_0 is homotopically like the trivial cobordism. The h-cobordism due to S. Smale in 1960, states that if M^m and N^m are compact simply-connected oriented M -manifolds that are h-cobordant through the simply-connected $(m + 1)$ -manifold W_0^{m+1} , then M and N are diffeomorphic [26, p. 29].

However this theorem fails in dimension 4. If M and N are cobordant 4-manifolds, then N can be obtained from M by cutting out a compact contractible submanifold W and gluing it back in by using an involution of ∂W .

The h-cobordism under question in our example may be described by attaching an algebraic cancelling pair of 2- and 3-handles to the interior of Akbulut cork W . The 4-manifold \bar{W} mediating V and W resembles the Akbulut cork with the dot replaced by a 0-surgery. The manifold under question is nothing but $L7a6(0,1)(0,1)$ (see [27, p. 355] or [13, Figure 3c]).

3.5. The character variety for an Akbulut cork W

The Sage code we used is as follows

```
from snappy import manifold
M=Manifold('946(-1,1)')
G=M.fundamental_group()
I=G.character_variety_vars_and_polys(as_Ideal=True)
I
```




Figure 3. The surface $S_W(x, y, z)$ found within the Groebner basis of the $SL(2, \mathbb{C})$ character variety for the Akbulut cork W .

The fundamental group ruling the Akbulut cork $W = 9_{46}(-1, 1)$ is the two-generator group

$$\pi_1(W) = \langle a, b | aBAb^2ABabaBABab, a^3BAb^3AB \rangle, \quad A = a^{-1}, B = b^{-1}. \quad (1)$$

With Sage software in Reference [15], or the aforementioned code, we compute the corresponding character variety. Then, from Magma [16], the Groebner basis is found in the form

$$\begin{aligned} \mathcal{G}_W(x, y, z) &= z(z-2)[x+y+f_1(z)][y^2-y+f_2(z)][y+f_3(z)]f_4(z), \\ f_1(z) &= -171/34 z^8 + 913/34 z^7 - 2429/34 z^6 + 4733/34 z^5 - 2443/17 z^4 \\ &\quad - 224/17 z^3 + 1847/17 z^2 - 1103/34 z - 1, \\ f_2(z) &= -95/34 z^8 + 477/34 z^7 - 1221/34 z^6 + 2331/34 z^5 - 1089/17 * z^4 \\ &\quad - 285/17 z^3 + 858/17 z^2 - 337/34 z - 1 \\ f_3(z) &= -71/17 z^7 + 376/17 z^6 - 997/17 z^5 + 1942/17 z^4 - 1993/17 z^3 \\ &\quad - 188/17 z^2 + 1477/17 z - 436/17, \\ f_4(z) &= z^7 - 5 z^6 + 13 z^5 - 25 z^4 + 24 z^3 + 4 z^2 - 18 z + 5. \end{aligned} \quad (2)$$

The factor containing $f_1(z)$ in Equation 3 is the singular surface $S_W(x, y, z)$ shown in Figure 3. It is a rational scroll and a surface of a general type. The factor containing $f_2(z)$ is an hyperelliptic function of genus 7, discriminant ≈ 1.327502101 and points at infinity $(1, 0, 0)$ and $(1, -95/34, 0)$. The factor containing $f_3(z)$ is an ordinary curve. The factor containing $f_4(z)$ is a seventh-order polynomial.

Formal desingularization of the surface $S_W(x, y, z)$

The formal desingularization of a hypersurface X in the three-dimensional projective space $\mathbb{P}_{\mathbb{Q}}^3$ over the rationals \mathbb{Q} is described in [19] and can be explicitly given with Magma [16, Section 122.5.3].

To the surface $X = S_W(x, y, z)$ we associate the degree 8 homogeneous polynomial $S_W(x, y, z, w) = x + y + f_1(z, w) \in \mathbb{Q}(x, y, z, w)$, with $f_1(z, w) = -171/34z^8 + 913/34wz^7 + \dots - 1103/343w^7z - w^8$.

For the projective surface $X = S_W(x, y, z, w)$, the two essential (over the three) singular morphisms are

$$\begin{aligned} x &\rightarrow 1 \\ y &\rightarrow st - 1 \\ z &\rightarrow t \\ w &\rightarrow \alpha t + O(t^8), \end{aligned} \tag{3}$$

$$\begin{aligned} x &\rightarrow 1 \\ y &\rightarrow s \\ z &\rightarrow 34/171(s+1)t^7 + O(t^8) \\ w &\rightarrow O(t^8), \end{aligned} \tag{4}$$

where α has minimal polynomial $-s - \alpha^8 f_1(\alpha^{-1})$.

These formal morphisms are used to compute the global sections (or adjoints) $\text{Hom}_{i,j}$. For the surface $S_W(x, y, z)$ one gets

$$\text{Hom}_{1,1} = \text{Hom}_{2,1} = \text{Hom}_{2,2} = 0, \text{Hom}_{1,2} = [z^6, z^5w, z^4w^2, z^3w^3, z^2w^4, zw^5, w^6] \dots$$

3.6. The character variety for the mediating manifold \bar{W}

The fundamental group $\pi_1(\bar{W})$ of the h-cobordism \bar{W} is as follows [13]

$$\pi_1(\bar{W}) = \langle a, b | a^3 b^2 A B^3 A b^2, (ab)^2 a B^2 A b^2 A B^2 \rangle. \tag{5}$$

The cardinality structure of subgroups of this fundamental group is

$$\eta_d[\pi_1(\bar{W})] = \eta_d[\pi_1(W)] = [1, 0, 0, 0, 1, 1, 2, 0, 0, \mathbf{1}, 0, \mathbf{5}, 4, \mathbf{9}, \mathbf{7}, 1 \dots],$$

where the bold digits refers to our investigation in [13, Table 1] about the existence of a finite geometry obtained with a subgroup of the corresponding index. The smallest case occurs at index 10 and the geometry is the Mermin pentagram, a type of contextual geometry.

As before, with Sage software in Reference [15], we compute the corresponding character variety. Then, from Magma, the Groebner basis is found in the form

$$\begin{aligned}
\mathcal{G}_{\bar{W}} &= (z-2)(z^2-z-1)[x+g_1(z)][y+g_2(z)]g_3(z), \\
g_1(z) &= -z^9+3z^8+5z^7-18z^6-10z^5+43z^4-2z^3-27z^2+3z+4, \\
g_2(z) &= -z^8+2z^7+6z^6-11z^5-15z^4+24z^3+9z^2-10z-2, \\
g_3(z) &= z^7+z^6-5z^5-6z^4+6z^3+5z^2-2z-1.
\end{aligned}
\tag{6}$$

Although the card seq for manifolds W and \bar{W} are the same, we clearly see that the character varieties are distinct. In the later case, it contains two ordinary curves and polynomials of degree 1, 2 and 7.

4. $SL(2, \mathbb{C})$ scheme processing in topological quantum computing

In this section, we are interested in the $SL(2, \mathbb{C})$ character variety of the fundamental group of (hyperbolic) 3-manifold L10n46 (alias otet08₀₀₀₀₂). This manifold describes the 4-fold (irregular) covering of the figure-eight knot $4_1 = K4a1$ [14, Table 2]. It is connected to the magic state describing the contextual geometry of two-qubits (e.g. [14, Figure 1]. or [30, Table 1]). Thanks to the character variety and the related algebraic surfaces, we can add the topological aspect to the previous description.

The Groebner basis of the character variety for the fundamental group of 3-manifold L10n46 may be obtained with Magma. The 3-generator fundamental group is

$$\pi_1(\text{L10n46}) = \langle a, b, c | abAcBac, abbCBccBBBACbcbC \rangle. \tag{7}$$

The character variety $\mathcal{G}_{\text{L10n46}}(e, f, g, h)(x, y, z)$ is now seven-dimensional as in [12] and selected choices of parameters e, f, g and h allow us to determine the algebraic surfaces within the Groebner basis.

For instance, we find

$$\begin{aligned}
\mathcal{G}_{\text{L10n46}}(0, 0, 0, 0) &= yS_1(x, y, z)S_2(x, y, z)S_3(x, y, z) \cdots \\
S_1(x, y, z) &= f_{2, \{ \}}^{(A_1)}(x, y, z) = xyz + xz^2 + x^2 + y^2 + z^2 + yz - x - 6, \\
S_2(x, y, z) &= xz^2 + yz - x - 2, \\
S_3(x, y, z) &= -x^2z^2 + 2x^2 + y^2 + z^2 + 2x - 4.
\end{aligned}
\tag{8}$$




Figure 4. Left: the (degree 3) del Pezzo surface $S_1(x, y, z) = f_{2, \{ \}}^{(A_1)}(x, y, z) = xyz + xz^2 + x^2 + y^2 + z^2 + yz - x - 6$. Middle: the (rational scroll) surface $S_2(x, y, z) = xz^2 + yz - x - 2$. Right: the (del Pezzo degree 4) surface $S_3(x, y, z) = -x^2z^2 + 2x^2 + y^2 + z^2 + 2x - 4$.

The first surface $S_1(x, y, z) = f_{2, \{ \}}^{(A_1)}$ in the product (8) has a single simple singularity of type A_1 whose reduced singular subscheme is of degree 2 with a vanishing support.

The second surface $S_2(x, y, z)$ is a singular rational scroll with a single essential singularity. The corresponding singular morphism $\text{Spec } \mathbb{F}_\phi[[t]] \rightarrow X = S_2(x, y, z, w)$ is

$$\begin{aligned}
x &\rightarrow 1 \\
y &\rightarrow s \\
z &\rightarrow t \\
w &\rightarrow ((-2s^3 - 6s)/(s^2 + 4)\alpha + (-2s^2 - 4)/(s^2 + 4))\alpha t + O(t^3),
\end{aligned}
\tag{9}$$

where α has minimal polynomial $\alpha^2 - s\alpha - 1$.

The third surface $S_3(x, y, z) = -x^2z^2 + 2x^2 + 2x + y^2 + z^2 - 4$ is a singular surface of the degree 4 del Pezzo type. There are 4 essential singularities. We retain one of the two singularities with a nontrivial minimal polynomial. The corresponding singular morphism $\text{Spec } \mathbb{F}_\phi[[t]] \rightarrow X = S_1(x, y, z, w)$ is

$$\begin{aligned}
x &\rightarrow 1 \\
y &\rightarrow s \\
z &\rightarrow t \\
w &\rightarrow \alpha t - 1/(s^4 + 4s^2 + 4)t^2 + (-1/2 s^4 + 9/2)/(s^6 + 6s^4 + 12s^2 + 8)\alpha t^3 + O(t^5)
\end{aligned}
\tag{10}$$

where α has minimal polynomial $\alpha^2 - 1/(s^2 + 2)$.

All three affine surfaces are shown in Figure 4.

5. $SL(2, \mathbb{C})$ scheme processing in microRNAs

In this section, we focus on the $SL(2, \mathbb{C})$ character varieties attached to microRNAs (miRNAs for short). This case was already tackled in our recent paper [12].

The miRNAs are short (approximately 22 nt long) single-stranded RNA molecules playing a fundamental role in the expression and regulation of genes by targeting specific messenger RNAs (mRNAs) for degradation or translational repression. The genes encoding miRNAs are much longer than the processed mature miRNA molecule. There are pre-miRNAs comprising of approximately 70-nucleotides in length which are folded into imperfect stem-loop structures.

Each miRNA is synthesized as an miRNA duplex comprised of two strands (-5p and -3p). However, only one of the two strands becomes active, which is selectively incorporated into the RNA-induced silencing complex in a process known as miRNA strand selection [31,32]. For details about the miRNA sequences, we use the Mir database [33,34].

Disregulation of miRNAs may lead to a disease like cancer. A key microRNA known as an oncomir (involved in immunity and cancer) is mir-155 [12].

Here, we select examples of human miRNAs from the perspective of evolution. The generator of the group π to be considered is a short (about 8-letter long) seed made of two to four distinct bases in the set $\{A, U, G, C\}$. Most of the time, π is close to a free group of rank equal the number of distinct bases in the seed minus one. This point can be checked by the cardinality structure of conjugacy classes subgroups of π (denoted card seq). Exceptions to this rule and the occurrence of singularities (isolated or not) in the corresponding character variety built from π , is a witness of a potential disease. Unlike the case of transcription factors, until now, singularities possibly found with microRNAs are isolated singularities.

According to Reference [35, Table 3], the slowest evolving miRNA gene is hsa-mir-503 (the notation hsa is for the human specie). It is known that mir-503 regulates gene expression from different aspects of pathological processes of diseases, including carcinogenesis, angiogenesis, tissue fibrosis and oxidative stress [36]. The seed region for mir-503-5p is AGCAGCGG and the corresponding Groebner basis for parameters $(e, f, g, h) = (0, 0, 0, 0)$ is very simple: $\mathcal{G}_{mir-503-5p}(0, 0, 0, 0) = \kappa_4^{(4A1)}(x, y, z)$, as shown in Figure 5 (Left).

For $(e, f, g, h) = (1, 1, 0, 0)$, $\mathcal{G}_{mir-503-5p}(1, 1, 0, 0) = -3xyz\kappa_3(x, y, z)$, with $\kappa_3(x, y, z)$ is the Fricke surface found in [11, Section 3.3]. For $(e, f, g, h) = (1, 1, 1, 1)$, there are many more polynomials. One of them defines the Fricke surface $xyz + x^2 + y^2 + z^2 - 2x - y - 2$. The considered seed region for mir-503-3p is GGGUAAU. The surfaces in the Groebner basis are very simple in this case and not even simple singularities lie in them.

One of the fastest evolving microRNA is mir-214 [35, Table 3]. First mir-214 was reported to promote apoptosis in HeLa cells. Presently, mi-214 is implicated in an extensive range of conditions such as cardiovascular diseases, cancers, bone formation and cell differentiation [37]. For mir-214-5p and the seed sequence GCCUGU, one finds the surface $f_{1,\{1:0:0:0\}}^{(A2)}(x, y, z) = xyz + y^2 + z^2 - 4$ within $\mathcal{G}_{mir-214-5p}(0, 0, 0, 0)$, as shown in Figure 5. A surface of the same type $f_{1,\{1:0:0:0\}}^{(A2)}(x, y, z) = xyz + y^2 + z^2y - 2z - 1$ is found in $\mathcal{G}_{mir-214-5p}(1, 1, 1, 1)$. For a longer seed, surfaces are not found to contain singularities.




Figure 5. Left: the Cayley cubic $\kappa_4(x, y, z)$ found in the character variety for the slowest evolving miRNA gene hsa-mir-503. The surface $f_{1,\{1:0:0:0\}}^{(A2)}(x, y, z)$ found in the character variety $\mathcal{G}_{mir-214-5p}$ of the fast evolving gene hsa-mir-214.

6. Conclusion

Over the last few years, the authors of this article have found that some mathematical techniques employed for quantum information processing and quantum computing may also apply to biology at the genome scale. More precisely, group theory and representations of symmetries with characters of finite groups have been used for topological quantum computing (TQC) [14] or elementary particles [38], and the encoding of proteins [39]. Methods for dealing with infinite groups and $SL(2, \mathbb{C})$ representations of such groups in TQC papers [11,40] were similarly employed for transcription factors [12] and miRNAs [12].

Our efforts in this paper are belonging to the field of scheme processing, where the desingularization of discontinuities of algebraic surfaces is a spectrum. The scheme spectrum is a well known concept in commutative algebra. The prime spectrum of a ring is the set of prime ideals of the ring R and denoted by $\text{Spec}(R)$. The sheaf of rings \mathcal{O} is the relevant algebraic geometrical notion introduced by Grothendieck to develop this field [8,41]. We touched this important concept of schemes while investigating non-zero dimensional sets of singularities in surfaces belonging to the $SL(2, \mathbb{C})$ character variety of an infinite group. We are fortunate that the software Magma is designed to implement schemes in a variety of applications (curves, surfaces and more). Another computer algebra system with similar facilities is Singular [42,43].

We have no doubt that scheme theory will play and increasing role in other fields such as cosmology [44] and the field of ‘quantum consciousness’ [45].

Author Contributions: “Conceptualization, M.P.; methodology, M.P and M.A.; software, M.P.; validation, M.A., D.C. and K.I.; formal analysis, D.C., M.P. ; investigation, D.C., M.A., M.P. ; resources, K.I.; data curation, M.P.; writing—original draft preparation, M.P.; writing—review and editing, M.P. and M.A.; visualization, M.A.; supervision, M.P.; project administration, M.P., K.I.; funding acquisition, K.I. All authors have read and agreed to the published version of the manuscript”.

Funding: “This research received no external funding”

Informed Consent Statement: “Not applicable”

Data Availability Statement: “Data are available from the authors after a reasonable demand”

Conflicts of Interest: “The authors declare no conflict of interest”

MDPI Multidisciplinary Digital Publishing Institute

References

- Planat, M.; Minarovjech, M.; Saniga, M. Ramanujan sums analysis of long-period sequences and $1/f$ noise. *EPL* **2009**, *85*, 40005.
- Hehl, F.W.; von der Heyde, P.; Kerlick, G.D.; Nester, J.M., General relativity with spin and torsion: Foundations and prospects. *Rev. Mod. Phys.* **1976**, *48* (3), 393–416.
- Rovelli, C., Vidotto, F.: Covariant Loop Quantum Gravity. Cambridge University Press 1 edition, (2014).
- Character variety. Available online: https://en.wikipedia.org/wiki/Character_variety (accessed on 1 January 2022).
- Bullock, D. Rings of $SL_2(\mathbb{C})$ -characters and the Kauffman bracket skein module. *CM Helvetici* **1997**, *72*, 521–542.
- Gröbner basis. Available online: https://en.wikipedia.org/wiki/Gröbner_basis (accessed on 1 August 2022).
- ADE singularity. Available online: <https://ncatlab.org/nlab/show/ADE+singularity> (accessed on 1 August 2022).
- Reid, M. *Undergraduate commutative algebra*, London Mathematical Society Student Texts, Vol. 29 (Cambridge University Press, Cambridge, USA, 1995).
- Hartshorne, R. *Algebraic geometry*, Graduate texts in mathematics, Vol 52 (Springer, NY, USA, 1977).
- Scheme (mathematics). Available online: [https://en.wikipedia.org/wiki/Scheme_\(mathematics\)](https://en.wikipedia.org/wiki/Scheme_(mathematics)) (accessed on 1 May 2022).
- Planat, M.; Chester, D.; Amaral, M.; Irwin, K. Fricke topological qubits. *Quant. Rep.* **2022**, *4*, 523–532.
- Planat, M.; Amaral, M. M.; Irwin, K. Algebraic morphology of DNA–RNA transcription and regulation. Available at <https://www.preprints.org/manuscript/202212.0256/v1>.
- Planat, M.; Aschheim, R.; Amaral, M.M.; Irwin, Quantum computation and measurements from an exotic space-time R^4 . *Symmetry* **2020**, *12*, 736.
- Planat, M.; Aschheim, R.; Amaral, M.M.; Irwin, Group geometrical axioms for magic states of quantum computing. *Mathematics* **2019**, *7*, 948.
- Python code to compute character varieties. Available online: <http://math.gmu.edu/~slawton3/Main.sagews> (accessed on 1 May 2021).
- Bosma, W.; Cannon, J. J.; Fieker, C. ; Steel, A. (eds). *Handbook of Magma functions*, Edition 2.23; University of Sydney: Sydney, Australia, 2017; 5914pp (accessed on 1 January 2019).
- Goldman, W. M. Trace coordinates on Fricke spaces of some simple hyperbolic surfaces. In *Handbook of Teichmüller theory*, Eur. Math. Soc. , Zürich, **2009**, *13*, 611–684.
- Inaba, M.; Iwasaki, K.; Saito, M. H. Dynamics of the sixth Painlevé equation. *arXiv* **2005**, arXiv:math.AG/0501007
- Beck, T.; Schicho, J. Adjoint computation for hypersurfaces using formal desingularizations. *J. Algebra* **2018**, *320* 3984–3996.
- Schicho, J. Rational parametrization of surfaces. *J. Symbolic Computation* **1998** *26* 1–29.
- Enriques Kodaira classification, available online: https://en.wikipedia.org/wiki/Enriques-Kodaira_classification
- Bravyi, S.; Kitaev, A. Universal quantum computation with ideal Clifford gates and noisy ancillas. *Phys. Rev. A* **2005**, *71*, 022316.
- Veitch, V.; Mousavian, S. A.; Gottesman, D.; Emerson, J. The Resource Theory of Stabilizer Computation. *New J. of Phys.* **2014**, *16*, 013009.
- Akbulut, S. *4-manifolds*, Oxford Graduate Texts in Mathematics, Vol. 25 (Oxford University Press, Oxford, 2016).
- Gompf, R. E.; Stipsicz, A. I. *4-manifolds and Kirby calculus*, Graduate Studies in Mathematics, Vol. 20 (American Mathematical Society, Providence, Rhode Island, 1999).
- Scorpan, A. *The wild world of 4-manifolds* (American Mathematical Society, Providence, Rhode Island, 2011).
- Akbulut, S. A fake compact contractible 4-manifold. *J. Diff. Geom.* **1991**, *33*, 335–356.
- Akbulut, S. An exotic 4-manifold. *J. Diff. Geom.* **1991**, *33*, 357–361.
- Akbulut, S.; Durusoy, S. An involution acting nontrivially on Heegaard-Floer homology. In *Geometry and topology of manifolds*; Fields Inst. Commun., Amer. Math. Soc.: Providence, RI, USA, 2005; Volume 47, pp. 1–9.
- Planat, M. Geometry of contextuality from Grothendieck’s coset space, *Quantum Inf. Process.* **2015**, *14*, 2563–2575.
- Medley, C. M.; Panzade G.; Zinovyeva, A. Y. MicroRNA stran selection: unwinding the rules. *WIREs RNA* **2021**, *12*, e1627.
- Dawson, O.; Piccinini, A. M. miR-155-3p: processing by-product or rising star in immunity and cancer? *Open Biol.* **2022**, *12*, 220070.
- Kozomara, A.; Birgaonu, M.; Griffiths-Jones, S. miRBase: from microRNA sequences to function. *Nucl. Acids Res.* **2019**, *47*, D155–D162.
- miRBase: the microRNA database. Available online: <https://www.mirbase.org/> (accessed on 1 November 2022).
- Fromm, B.; Billipp, T.; Peck, L. E.; Johansen, M.; Tarver, J. E.; King, B. L.; Newcomb, J. M.; Sempere, L. F.; Flatmark, K.; Hovig, E.; Peterson, K. J. A uniform system for the annotation of human microRNA genes and the evolution of the human microRNAome. *Annu. Rev. Genet.* **2015**, *23*, 213–242.

-
36. He, Y.; Cai, Y.; Pail, P. M.; Ren, X.; Xia, Z. The Causes and Consequences of miR-503 Dysregulation and Its Impact on Cardiovascular Disease and Cancer. *Front. Pharmacol.* **2021**, *12*, 629611.
 37. Amin, M. M.J.; Trevelyan, C. J.; Turner, N. A. MicroRNA-214 in health and disease. *Cells* **2021**, *23*, 3274.
 38. Planat, M.; Aschheim, R.; Amaral, M. M. Irwin, K. Informationally complete characters for quark and lepton mixings. *Symmetry* **2020**, *12*, 1000.
 39. Planat, M.; Aschheim, R.; Amaral, M. M.; Fang, F.; Irwin, K. Complete quantum information in the DNA genetic code. *Symmetry* **2020**, *12*, 1993.
 40. Planat, M.; Aschheim, R.; Amaral, M. M.; Fang, F.; Chester, D.; Irwin, K. Character varieties and algebraic surfaces for the topology of quantum computing. *Symmetry* **2022**, *14*, 915.
 41. Catren, G.; Cukierman, F. Grothendieck's theory of schemes and the algebra–geometry duality. *Synthese* **2022**, *200*, 234.
 42. Decker, W.; Greuel, G. M. History of Singular and its relation to Zariski's multiplicity conjecture. *ArXiv* **2012**, arXiv: 2103.00525.
 43. Singular. Available online: <https://www.singular.uni-kl.de> (accessed on 1 January 2023).
 44. Asselmeyer-Maluga, T.; Król, J.; Wilms, A. Big bang and topology. *Symmetry* **2022**, *14*, 1887.
 45. Meijer, D. K.F.; Raggett, S. *Quantum Physics in consciousness studies* Available online: <https://www.deeplook.ir/wp-content/uploads/2016/07/Quantum-Ph-rev-def-2.pdf> (accessed on 1 January 2023).