



HAL
open science

Stroke care pathways according to rural-urban residency

Lola Menant, Olivier Grimaud, Nolwenn Le Meur, Sandrine Kerbrat, Valérie Olié, Serge Timsit

► To cite this version:

Lola Menant, Olivier Grimaud, Nolwenn Le Meur, Sandrine Kerbrat, Valérie Olié, et al.. Stroke care pathways according to rural-urban residency. *Our Food, Our Health, Our Earth: A Sustainable Future for Humanity*, Nov 2023, Dublin (Ireland), Ireland. hal-04285170

HAL Id: hal-04285170

<https://hal.science/hal-04285170v1>

Submitted on 14 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stroke care pathways according to rural-urban residency

L. MENANT (1), O. GRIMAUD (1), N. LE MEUR (1), S. KERBRAT, (2) V. OLIE (3), S. TIMSIT (4)

(1) Ecole des Hautes Etudes en Santé Publique (EHESP), CNRS, INSERM, Arène – UMR 6051 ; (2) DAMAD ; (3) Santé Publique France ; (4) CHU BREST
Contact author: olivier.grimaud@ehesp.fr




Context and Objectives

Stroke is a pathology that traces the performance of the health system, requiring coordination of all of its components (from prevention to rehabilitation). In the French context, a 2019 study showed a higher incidence, but **lower lethality in rural areas** covered by the Brest Stroke Registry [1]. This result is in opposition what was observed in a second study covering the national territory [2].

The **objectives** were **1** - to characterize the care pathways of patients with ischemic stroke and **2** - to study the association with rural-urban residency context of the patient.

Methods

Rural urban indicator



Data

Brest stroke registry - ischemic stroke between 2010 and 2013 (clinical data)

Pairing

French National Health Insurance Data System (SNDS) (healthcare consumption)

Rural urban indicator



Data


SNDS - ischemic stroke in 2018

Typology of pathways

Multichannel Sequence Analysis (MCSA) [3]

+

Hierarchical Ascending Classification (CAH)



Results


BREST AREA

2 499 ischemic stroke
52.9% ♀ 79.5 y
47.1% ♂ 70.8 y

Three typologies of pathways

(Illustrated below according to the antihypertensive consumption indicator)

Prevention of vascular risk (n = 843) | Low consumption of pre-stroke care (n = 1 117) | Post-stroke death (n = 539)




No association between the clustering and rural-urban residency. However:

- In low density area (« rural ») use of care is comparatively higher and case fatality lower compared to strong and intermediate densities,
- In intermediate density area (« peri-urban ») use of care is significantly lower than strong and low densities.

FRANCE

85 250 ischemic stroke
48.8% ♀ 78.4 y
51.2% ♂ 71.6 y



	Severe cases OR [95% CI]	Return home OR [95% CI]	Rehabilitation care OR [95% CI]
Stroke unit: Yes vs No	0.90 [0.86 – 0.94]*	0.94 [0.91 – 0.97]*	1.20 [1.15 – 1.24]*
Intensive care: Yes vs No	6.09 [5.49 – 6.76]*	0.22 [0.19 – 0.25]*	0.77 [0.67 – 0.88]*
Short-Term Hospitalization Units: Yes vs No	0.89 [0.84 – 0.94]*	1.10 [1.05 – 1.15]*	

*adjusted on gender, age, rural-urban residency, charlson score, social protection scheme, clustering of the pre-stroke phase

- Use of care, whether before or after stroke occurrence, did not differ according to rural-urban residency, apart from a higher access to Stroke Unit in large urban areas.

Conclusions

- We did not identify a clear relationship between the typology of stroke care pathways and rural-urban residency, suggesting equity of care,
- The clustering of the acute phase is differentiated by the stroke management services,
- The three period (pre-stroke phase, acute phase and poste-stroke phase) are independent of each other.

[1] Grimaud O, Lachkhem Y, Gao F, et al. Stroke Incidence and Case Fatality According to Rural or Urban Residence: Results From the French Brest Stroke Registry. Stroke. 2019;50(10):2661-2667. doi:10.1161/STROKEAHA.118.024695 ;

[2] Gabet A, Grimaud O, de Peretti C, Béjot Y, Olié V. Determinants of Case Fatality After Hospitalization for Stroke in France 2010 to 2015. Stroke. 2019;50(2):305-312. doi:10.1161/STROKEAHA.118.023495 ;

[3] Gabadinho A, Ritschard G, Müller NS, Studer M. Analyzing and Visualizing State Sequences in R with TraMineR. J Stat Softw. 7 avr 2011;40:1-37