



Laser-induced photodetachment diagnostic for interrogating pulsed ECR-driven plasmas: Application to H- and D- negative ions

M Mitrou, S Béchu, P Svarnas

► To cite this version:

M Mitrou, S Béchu, P Svarnas. Laser-induced photodetachment diagnostic for interrogating pulsed ECR-driven plasmas: Application to H- and D- negative ions. xxxv ICPIG, Jul 2023, Egmond aan Zee, Netherlands. 2023. hal-04285204

HAL Id: hal-04285204

<https://hal.science/hal-04285204>

Submitted on 20 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Laser-induced photodetachment diagnostic for interrogating pulsed ECR-driven plasmas: Application to H⁻ and D⁻ negative ions

ICPIG XXXVth edition, 9 – 14 July 2023
Egmond aan Zee, The Netherlands



M. Mitrou^{1,2}, S. Béchu¹ and P. Svarnas²

¹Université Grenoble Alpes, Grenoble INP (Institute of Engineering), LPSC-IN2P3, 38000 Grenoble, France


²High Voltage Lab., Electrical & Computer Eng. Dept., University of Patras, Rion-Patras 26504, Greece

Discharges sustained in the pulsed mode of operation are of great interest on a fundamental level as they may unveil important information on the plasma kinetics. The present work is devoted to an experimental setup tailored to study **time-resolved processes** relative to negative ion formation in ECR-driven plasmas. It refers to a laser-induced electron photo-detachment system synchronized with **H₂ and D₂ ECR-plasmas, sustained by 2.45 GHz bursts in the kHz range**. The system is combined with electrostatic probe time-resolved measurements and information is obtained both on the formation path and the **yield enhancement of H⁻ and D⁻ negative ions**.

EXPERIMENTAL SETUP


The SCHEME-II+ reactor

- Designed to evaluate plasma-surface interactions relevant to the volume production of H⁻(D⁻) ions via dissociative electron attachment (DEA).
- H₂(D₂) plasmas are produced by a single dipolar ECR module (2.45 GHz, 0-200 W, pulsed with variable frequency and duty cycle).
- Simultaneous, time-resolved, application of the Langmuir-probe and laser-induced photodetachment diagnostics.




METHODOLOGY

Time-resolved measurements require proper synchronization of the laser and Langmuir-probe systems with the driving microwave pulse:




RESULTS

- Probed area within the plasma volume: 75 mm downstream the mid-plane of the magnet of the ECR module
- D₂ (1.6 Pa, 150 W), 100% Quartz coverage of the reactor interior



Meanwhile, in the
CONTINUOUS MODE
of operation:

CONCLUSION



Enhanced negative ion production in the post-discharge

Rapid cooling of the electron population & Extended lifetime of ro-vibrationally excited molecules
Rapid decay of the high-energy electrons & Rapid decay of the atomic species
Time-resolved fourier-transform absorption spectroscopy with synchrotron radiation will be applied in order to gain information on these quantities.

Enhanced negative ion production via DEA:
 $H_2^* + e(<1eV) \rightarrow H^- + H$

Limited losses of negative ions

References:

- [1] S. Béchu *et al.*, J. Quant. Spectr. Radiat. Transfer **257**, 107325 (2020).
- [2] M. Bacal, Rev. Sci. Instrum. **71**, 3981 (2000).
- [3] T. Mosbach *et al.*, Plasma Sources Sci. Technol. **7**, 75-81 (1998).

Acknowledgment:

The present work is partially funded by the "Andreas Mentzelopoulos" PhD scholarship foundation of the University of Patras (for M. Mitrou).

Recent publications:

- [1] M. Mitrou, P. Svarnas, S. Béchu, "H and D production efficiency in a multi-dipole ECR-plasma source as a function of gas pressure", J. Phys.: Conf. Series **2244**, 012007 (2022).
- [2] M. Mitrou, P. Svarnas, S. Béchu, "Experimental study on dynamic effects of H and D negative ions in an ECR-plasma source", J. Phys.: Conf. Series **2244**, 012006 (2022).
- [3] F. Taccogna *et al.*, "Last experimental and theoretical advances in the production of negative ions in Cesium-free plasmas", The European Physical Journal D **75**, 227 (2021).

