



HAL
open science

Biomedical implications of nuclear transport

Nathalie J Arhel, Mark C Field

► **To cite this version:**

Nathalie J Arhel, Mark C Field. Biomedical implications of nuclear transport. FEBS Letters, 2023, 597 (20), pp.2499 - 2500. <10.1002/1873-3468.14752>. <hal-04291349>

HAL Id: hal-04291349

<https://hal.science/hal-04291349v1>

Submitted on 29 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Biomedical implications of nuclear transport

 Nathalie J. Arhel¹ and Mark C. Field^{2,3}

1 Institut de Recherche en Infectiologie de Montpellier (IRIM), CNRS UMR9004, Université de Montpellier, France

2 School of Life Sciences, University of Dundee, UK

3 Institute of Parasitology, Biology Centre, Czech Academy of Sciences, České Budějovice, Czech Republic

A defining feature of eukaryotes is the nucleus, where most of the genetic information resides and is safeguarded within the confines of the nuclear envelope (NE). While this barrier serves to protect the genome and separate transcription from translation, the need to facilitate transport across the NE itself represents a challenge [1]. Nuclear pore complexes (NPCs) support bidirectional transport between the nucleus and cytoplasm and, together with other NE constituents, play critical roles in genome organization, gene expression and nucleocytoplasmic trafficking [2]. As a result, perturbations in the homeostasis of NPC or NE biology are implicated in a range of pathological states, such as neurodegenerative or movement disorders, ageing, cancer, acute necrotizing encephalopathy (ANE) and viral infection [2]. Furthermore, variations in the composition and organization of the NPC are now known to occur between evolutionary lineages with likely significant connections with divergent requirements [3]. Here, in the first of two collections devoted to the

NPC and associated nuclear components, we have assembled a selection of articles with novel perspectives on biomedical implications of NPC biology and nucleocytoplasmic trafficking. Together, these articles explore some of the latest findings and insights into the role of NPCs in essential biological pathways, and the assembly of these fascinating molecular structures.

Firstly, Padilla-Mejia and Field [4] address fundamental concepts of NPC and NE structure and composition throughout eukaryotic evolution. In particular, they discuss trypanosomes, protozoa with novel nuclear components that have evolved to meet the specific demands of trypanosome biology. By examining the nuclear lamina and NPC in trypanosomes, Padilla-Mejia and Field emphasize lineage-specific and conserved aspects of nuclear organization. This exploration is not only scientifically intriguing but also relevant for understanding the mechanisms exploited by these parasitic protozoa in immune evasion and life cycle progression and how they cause disease. Desgraupes *et al.* [5] pursue this by



Nathalie J. Arhel is a CNRS Research Director at the Institut de Recherche en Infectiologie de Montpellier (IRIM), France. After earning BSc and PhD degrees in biochemistry from the University of Bristol, UK, she trained as a post-doctoral researcher in host-virus interactions at the Pasteur Institute in Paris, France, and at the Universitätsklinikum Ulm in Germany, leading to her recruitment as a permanent research fellow by the French CNRS in 2009. She currently leads a research team on the topics of viral infections, nuclear pore biology, and innate immunity.



Mark C. Field is Professor of Cell Biology at the University of Dundee in Scotland and the Institute of Parasitology, České Budějovice, Czech Republic. His research focuses on the African trypanosome, an eukaryotic pathogen, which he uses as a model for evolutionary cell biology and the examination of pathogenic mechanisms and development of therapeutics. He also has interests in the origins of eukaryotes and reconstruction of the last eukaryotic common ancestor.

doi: 10.1002/1873-3468.14752

focusing on a nucleoporin that appeared in metazoans and underwent genomic rearrangements during ape evolution. RANBP2/Nup358 is a cytoplasmic filament nucleoporin and E3 SUMO ligase involved in various cellular processes, which are discussed in depth in the first part of the review. Authors then explore how the dysregulation or mutation of RANBP2 contributes to human pathologies, thus highlighting its importance to human evolution and health.

Next, Kuiper *et al.* [6] delve into NPC assembly, emphasizing the need for careful control over the condensation of phenylalanine-glycine (FG) repeat nucleoporins, which constitute the permeability barrier. Defects in NPC assembly or homeostasis are associated with movement disorders, amyotrophic lateral sclerosis (ALS) and senescence. Recent progress in understanding NPC assembly and its connection to human disorders are discussed, illuminating spatial and temporal control mechanisms required for orderly NPC assembly. In a related area, Cristi *et al.* [7] dissect the critical role of NPCs in maintaining the equilibrium between the nucleus and cytoplasm, elucidating their involvement in neurodegenerative disease pathogenesis. In particular, the importance of the ESCRT-III pathway for NE maintenance and repair, and in preserving functional NPCs, is emphasized. Authors thus highlight how the selective reduction of specific nucleoporins can lead to neurodegenerative pathology. Through a comprehensive description of surveillance pathways that safeguard NPC biogenesis and assembly, we gain a deeper understanding of the pathological cascade leading to diseases such as ALS, frontotemporal dementia (FTD) and Huntington's disease (HD).

Ferreira continues with the topic of NPC assembly and focuses on phase separations during nucleocytoplasmic transport and their impact on proteostasis in the context of neurodegenerative diseases [8]. Specifically, Ferreira addresses how impairments to nucleocytoplasmic transport contribute to dysregulation in cellular partitioning and lead to the formation of protein aggregates—a hallmark of multiple neurodegenerative diseases. A particular focus on RANBP2 is introduced by dissecting its role in the survival of photoreceptor and motor neurons under homeostatic and pathophysiological conditions. Ferreira concludes by highlighting current clinical needs and potential therapeutics targeting neurotoxic aggregates.

Finally, Fichtman and Harel [9] guide us on a journey through the history of microscopy imaging, from the first lenses to modern electron microscopy (EM). By providing a magnified view of individual NPCs, authors showcase recent advances in field emission scanning EM. They then argue that this technique offers a topographic surface perspective, complementing the three-dimensional models

derived from other imaging methods, and which may become an important tool in understanding the manner in which NPC structural variability can lead to pathology, which is absent from other imaging approaches.

We anticipate that this FEBS Letters collection will help readers unravel the complexities of nucleocytoplasmic transport and nuclear pore complex assembly, by offering a comprehensive overview of their roles in health, disease and evolution. The articles here represent a mosaic of research, combining cellular and molecular insights with technological advancements. We invite readers to embark on this journey with us and explore the frontiers of cell biology, where the nucleus and its gatekeepers hold the keys to understanding many aspects of the complexity of life.

References

- 1 Field MC and Rout MP (2022) Coatomer in the universe of cellular complexity. *Mol Biol Cell* **33**, pe8.
- 2 Beck M and Hurt E (2017) The nuclear pore complex: understanding its function through structural insight. *Nat Rev Mol Cell Biol* **18**, 73–89.
- 3 Makarov AA, Padilla-Mejia NE and Field MC (2021) Evolution and diversification of the nuclear pore complex. *Biochem Soc Trans* **49**, 1601–1619.
- 4 Padilla-Mejia NE and Field MC (2023) Evolutionary, structural and functional insights in nuclear organisation and nucleocytoplasmic transport in trypanosomes. *FEBS Lett* **597**, 2501–2518.
- 5 Desgraupes S, Etienne L and Arhel NJ (2023) RANBP2 evolution and human disease. *FEBS Lett* **597**, 2519–2533.
- 6 Kuiper EFE, Prophet SM and Schlieker C (2023) Coordinating nucleoporin condensation and nuclear pore complex assembly. *FEBS Lett* **597**, 2534–2545.
- 7 Cristi AC, Rapuri S and Coyne AN (2023) Nuclear pore complex and nucleocytoplasmic transport disruption in neurodegeneration. *FEBS Lett* **597**, 2546–2566.
- 8 Ferreira PA (2023) Nucleocytoplasmic transport at the crossroads of proteostasis, neurodegeneration and neuroprotection. *FEBS Lett* **597**, 2567–2589.
- 9 Fichtman B and Harel A (2023) The magnified view: from ancient trinkets to single nuclear pore complexes. *FEBS Lett* **597**, 2590–2596.

Correspondence

N. J. Arhel, Institut de Recherche en Infectiologie de Montpellier, CNRS UMR9004, Université de Montpellier, 34090 Montpellier, France

Tel: +33 4 34359453

E-mail: nathalie.arhel@inserm.fr

M. C. Field, School of Life Sciences, University of Dundee, Dundee DD1 5EH, UK

Tel: +44 751 550 7880

E-mail: mfield@mac.com, mfield@dundee.ac.uk