



**HAL**  
open science

## **Assessment of cyanotoxins in water and fish in an African freshwater lagoon (Lagoon Aghien, Ivory Coast) and the application of WHO guidelines**

Eric Kouamé Yao, Mathias Koffi Ahoutou, Mark Olokotum, Sahima Hamlaoui, Emilie Lance, Benjamin Marie, Cécile Bernard, Rosine Yao Djeha, Catherine Quiblier, Jean-François Humbert, et al.

### ► To cite this version:

Eric Kouamé Yao, Mathias Koffi Ahoutou, Mark Olokotum, Sahima Hamlaoui, Emilie Lance, et al.. Assessment of cyanotoxins in water and fish in an African freshwater lagoon (Lagoon Aghien, Ivory Coast) and the application of WHO guidelines. *Environmental Science and Pollution Research*, 2023, 30 (43), pp.97857-97871. 10.1007/s11356-023-29025-3 . hal-04291416

**HAL Id: hal-04291416**

**<https://hal.science/hal-04291416v1>**

Submitted on 20 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Assessment of cyanotoxins in water and fish in an African freshwater lagoon**  
2 **(Lagoon Aghien, Ivory Coast) and the application of WHO guidelines**

3

4 Eric Kouamé Yao<sup>1</sup>, Mathias Koffi Ahoutou<sup>1,2</sup>, Mark Olokotum<sup>3,4</sup>, Sahima Hamlaoui<sup>5</sup>,  
5 Emilie Lance<sup>5,6</sup>, Benjamin Marie<sup>5</sup>, Cécile Bernard<sup>5</sup>, Rosine Yao Djeha<sup>1</sup>, Catherine  
6 Quiblier<sup>5,7</sup>, Jean-François Humbert<sup>\*,8</sup> and Julien Kalpy Coulibaly<sup>1</sup>

7

8 1. Institut Pasteur de Côte d'Ivoire, Abidjan, Côte d'Ivoire

9 2. Université Jean Lorougnon Guédé, Daloa, Côte d'Ivoire

10 3. National Fisheries Resources Research Institute, Jinja, Uganda

11 4. Makerere University, Department of Zoology, Entomology and Fisheries Sciences,  
12 Kampala, Uganda

13 5. Muséum d'Histoire Naturelle de Paris, UMR Molécules de Communication et  
14 Adaptation des Micro-organismes, Paris, France

15 6. Université de Reims Champagne-Ardenne, UMR Stress environnementaux et  
16 biosurveillance des milieux aquatiques, Reims, France

17 7. Université Paris Cité, Paris, France

18 8. Institut National de Recherche pour l'Agriculture l'Alimentation et l'Environnement-  
19 Centre Île-de-France-Versailles-Grignon, Paris, France

20

21 \*Corresponding author :

22 JF Humbert, [jean-francois.humbert@upmc.fr](mailto:jean-francois.humbert@upmc.fr) ; Phone number 33 687803675

23

24 **Abstract**

25 In comparison with northern countries, limited data are available on the occurrence  
26 and potential toxicity of cyanobacterial blooms in lakes and ponds in Sub-Saharan  
27 countries. With the aim of enhancing our knowledge on cyanobacteria and their toxins  
28 in Africa, we performed a 17-month monitoring of a freshwater ecosystem, Lagoon  
29 Aghien (Ivory Coast), which is used for multiple practices by riverine populations and  
30 for drinking water production in Abidjan city. The richness and diversity of the  
31 cyanobacterial community were high and displayed few variations during the entire  
32 survey. The monthly average abundances ranged from  $4.1 \times 10^4$  cell mL<sup>-1</sup> to  $1.8 \times 10^5$   
33 cell mL<sup>-1</sup>, with higher abundances recorded during the dry seasons. Among the five  
34 cyanotoxin families analyzed (anatoxin-a, cylindrospermopsin, homoanatoxin,  
35 microcystins, saxitoxin), only microcystins (MC) were detected with concentrations  
36 ranging from 0 to  $0.364 \mu\text{g L}^{-1}$  in phytoplankton cells, from 32 to  $1,092 \mu\text{g fresh weight}$   
37 (FW) kg<sup>-1</sup> in fish intestines and from 33 to  $383 \mu\text{g FW kg}^{-1}$  in fish livers. Even if the MC  
38 concentrations in water and fish are low, usually below the thresholds defined in WHO  
39 guidelines, these data raise the issue of the relevance of these WHO guidelines for  
40 Sub-Saharan Africa, where local populations are exposed throughout the year to these  
41 toxins in multiple ways.

42

43 **Key words**

44 Africa, Ivory Coast, cyanobacteria, cyanotoxins, freshwater ecosystem, risk  
45 management

## 46 **Introduction**

47 Eutrophication of freshwater ecosystems due to point and diffuse nutrient pollution is  
48 ongoing on the African continent (Jenny et al. 2020). As shown for Lake Victoria (East  
49 Africa), this process is associated with the rapid demographic changes taking place on  
50 this continent (United Nations 2017) and leads to significant land use changes and  
51 increasing pollution pressures on freshwater ecosystems (e.g., Olokotum et al. 2020).  
52 Like anywhere else in the world, one would expect that the ongoing eutrophication of  
53 African lakes would lead to the development of cyanobacterial blooms in these  
54 ecosystems (Nyenje et al. 2010). This has already been well documented in a few  
55 large lakes, such as Lake Victoria (e.g., Krienitz et al. 2002; Olokotum et al. 2021), and  
56 in lakes and reservoirs located in northern and southern Africa (e.g., Oberholster et al.  
57 2010; Matthews and Bernard 2015; Guellati et al. 2017; Hammou et al. 2018).  
58 However, in comparison with northern countries, limited data are available on the  
59 occurrence of cyanobacterial blooms in lakes and ponds located in intertropical areas  
60 (e.g., Merel et al. 2013). In addition, there is also a lack of data on the potential toxicity  
61 of these cyanobacterial blooms, with data available from less than half of African  
62 countries (e.g., Ndlela et al. 2016; Svircev et al. 2019). In contrast, several reports of  
63 freshwater cyanobacterial poisoning in terrestrial wildlife (flamingos and  
64 megaherbivores) have been made, mainly in eastern and southern Africa (Ash and  
65 Patterson 2022).

66 This lack of knowledge on the occurrence and potential toxicity of cyanobacterial  
67 blooms in freshwater ecosystems of numerous sub-Saharan countries is partly  
68 explained by the rarity of institutional monitoring of cyanobacteria in freshwater  
69 ecosystems, and *vice versa* (Codd et al. 2005). In addition, there are also no  
70 management plans, including policy tools, legislation and public communication,

71 dedicated to preventing human exposure to cyanotoxins, despite the potentially high  
72 exposure of human populations to these toxins in developing countries. Indeed, the  
73 majority of populations living around lakes use raw (untreated) water for multiple  
74 purposes, including washing dishes and clothes, bathing, cooking and direct  
75 consumption during fishing expeditions (e.g., Effebi et al. 2017; Roegner et al. 2020  
76 and 2023; Olokotum et al. 2022).

77 In an attempt to improve our knowledge on threat assessment attributed to  
78 cyanobacterial blooms in Sub-Saharan Africa, we performed a 17-month monitoring of  
79 a freshwater lagoon, Lagoon Aghien, located in northern Abidjan city (Ivory Coast).  
80 This lagoon supplies treated water for the Abidjan district and is used for recreational  
81 and multiple domestic activities and for fishing by local populations living in the villages  
82 located along its shores. Finally, this assessment led us to discuss the considerations  
83 for applying the microcystin guideline values provided by the WHO in African countries  
84 and the limits for their use in these countries.

85

## 86 **Material and Methods**


### 87 **Study site**

88 Lagoon Aghien (5°22'N to 5°26'N and 3°49'W and 3°55'W) is a freshwater ecosystem  
89 located in the northern-western part of the Abidjan District (Ivory Coast) (Fig. 1). This  
90 lagoon covers a surface area of 19.5 km<sup>2</sup> with a perimeter of 40.72 km, a volume of 75  
91 km<sup>3</sup> and a maximum depth of 10 m. More than 12,000 inhabitants live on its shores.

92 The lagoon is drained by three rivers (i.e., rivers Bété, Djibi and Mé) (Fig. 1).  
93 The Mé and Bété catchments mainly include natural and agricultural lands, whereas  
94 the Djibi catchment contains mostly urbanized areas. The Djibi and Bété Rivers flow  
95 directly in the western part of the lagoon, while the Mé River flows in the channel

96 between Lagoon Aghien and Lagoon Potou. These three rivers carry many nutrients  
97 (phosphorus and nitrogen), explaining the hypereutrophication of the lagoon (Koffi et  
98 al. 2019; Ahoutou et al. 2021).

99



100

101 **Fig. 1** Locations of the six sampling stations in Lagoon Aghien

102

103 Monthly rainfall values were obtained from the averages of the data collected at  
104 two weather stations (Anyama and Atchokoi).

105

### 106 **Sampling strategy of water**

107 As described in Ahoutou et al. (2021), a monthly monitoring survey was performed  
108 from December 2016 to April 2018 at five sampling stations (St1, St2, St4, St5 and  
109 St6) located on a NW–SE transect and at one additional sampling station (St3) located  
110 close to the shoreline area of Akandje village (Fig. 1). Briefly, water samples were  
111 collected in the first meter of the water column by using an integrated sampler as  
112 described in Laplace-Treyture et al. (2016). The collected water samples were

113 transferred into two clean bottles that were stored in a cool box until they were returned  
114 to the laboratory.

115 For cyanobacterial cell counting and identification, 200 mL of raw water was  
116 fixed in a formaldehyde solution (5% in final concentration). For the extraction of toxins  
117 from cyanobacterial cells, 200 mL of water was filtered through polycarbonate filters  
118 (Whatman® Nuclepore), which were stored at  $-80\text{ }^{\circ}\text{C}$ .

119

## 120 **Dissolved nutrient analyses**

121 All the nutrient analyses were performed following the instructions of the LCK cuvette test  
122 systems provided by of the manufacturer (Hach® Company). The preparation of samples was  
123 performed using LCK cuvette test systems LCK 304, 341, 339 and 349 for ammonium ( $\text{NH}_4$ ),  
124 nitrites ( $\text{NO}_2$ ), nitrates ( $\text{NO}_3$ ) and soluble reactive phosphorus (SRP) respectively, after the  
125 water samples were filtered through nylon membranes (Whatman™, porosity  $0.45\text{ }\mu\text{m}$ ,  
126 diameter 47 mm). Colorimetric determinations of the concentrations were made using a HACH  
127 DR6000 UV VIS spectrophotometer.

128

## 129 **Cyanobacteria identification and counting**

130 The identification and counting of cyanobacteria genera were performed with an  
131 inverted microscope (Nikon Eclipse TS100) according to the Utermöhl method  
132 (Utermöhl 1958) and following the AFNOR 15204 standard. For each sample, a  
133 minimum of 30 fields of view, selected at random and distributed over the entire surface  
134 of the counting chamber, were observed under the microscope such that at least 500  
135 counting units (cells, colonies or filaments/trichomes) were counted. Except for some  
136 cyanobacteria, the cells were directly counted. For *Microcystis*, two categories of  
137 colonies were considered corresponding to their diameters ( $<$  or  $>$   $200\text{ }\mu\text{m}$ ). The mean

138 number of cells per category was estimated under an upright microscope by counting  
139 the number of cells in 30 colonies per category after the colonies were gently spread  
140 between slides and coverslips. For filamentous cyanobacteria (*Aphanizomenon*,  
141 *Cylindrospermopsis/Raphidiopsis*, *Limnothrix*, *Lyngbya*, *Oscillatoria*, *Planktothrix* and  
142 *Pseudanabaena*), filament lengths of 100 µm were considered, and the mean cell  
143 number per 100 µm of filament was estimated on 30 filaments per genus according to  
144 the method described by Catherine et al. (2017). Finally, for all genera, the results of  
145 the cell counts are expressed as numbers of cells per mL.

146 The potential ability of the various genera to produce cyanotoxins was identified  
147 on the basis of a report published by ANSES (2020).

148


## 149 **Fish**

150 The fish were purchased monthly between May 2017 and April 2018 from fishermen  
151 in three villages located around the lagoon (Anyama-Débracadère, Akandjé and  
152 Aghien-Télégraphe). They were transported in a cooler to the laboratory, and then (i)  
153 the length and weight of each individual were measured (Supplementary Fig. S1; Table  
154 1) and (ii) liver, intestinal and muscle tissues were taken from each fish and stored at  
155 -80 °C until analysis.

156 A total of 97 fish belonging to six species were collected: 34 *Chrysichthys*  
157 *nigrodigitatus* (Claroteidae), 33 *Tilapia guineensis* (Cichlidae), 16 *Sarotherodon*  
158 *melanotheon* (Cichlidae), six *Oreochromis niloticus* (Cichlidae), five *Hemichromis*  
159 *fasciatus* (Cichlidae) and three *Hepsetus odoe* (Hepsetidae). These fishes belong to  
160 four trophic groups (i.e., omnivorous for *T. guineensis*, phytoplanktivorous for *S.*  
161 *melanotheon* and *O. niloticus*, zooplanktivorous for *C. nigrodigitatus* and carnivorous



162 for *H. fasciatus* and *H. odoe*) (Blé et al. 2008; Decru et al. 2012; Kidé et al. 2015; Jawad  
 163 et al. 2017)  
 164



165

166 **Supplementary Fig. S1** Average length and weight ( $\pm$  standard deviation) and fish  
 167 number in the six fish species collected in Lagoon Aghien

168 CN = *Chrysichthys nigrodigitatus*; HF = *Hemichromis fasciatus*; HO = *Hepsetus odoe*;  
 169 ON = *Oreochromis niloticus*; SM = *Sarotherodon melanotheron*; TG = *Tilapia*  
 170 *guineensis*  
 171

172 **Table 1** Total lengths (TL), weights and microcystin (MC) -LR and -RR concentrations  
 173 in intestine, liver and muscle of all the fishes collected in Lagoon Aghien

174 Cn = *Chrysichthys nigrodigitatus*; Hf = *Hemichromis fasciatus*; Ho = *Hepsetus odoe*;  
 175 On = *Oreochromis niloticus*; Sm = *Sarotherodon melanotheron*; Tg = *Tilapia*  
 176 *guineensis*  
 177  
 178

Date	Species	TL (mm)	Weight (g)	MC in intestine			MC in liver			MC in muscle
				RR	LR	Total	RR	LR	Total	( $\mu\text{g kg}^{-1}$ FW)
May-17	<i>Tg</i>	187	120	0	0	0	0	0	0	
	<i>Tg</i>	193	125	0	0	0	0	0	0	
	<i>Tg</i>	173	101	0	0	0	74	0	74	
	<i>Sm</i>	125	40	0	0	0	77	0	77	
	<i>Sm</i>	130	46	0	0	0	0	0	0	
	<i>Sm</i>	134	43	180	231	411	68	315	383	
	<i>Cn</i>	188	36	0	0	0	0	0	0	
	<i>Cn</i>	179	26	0	0	0	0	0	0	
	<i>Cn</i>	173	45	0	0	0	69	0	69	
Jun-17	<i>Tg</i>	221	198	0	0	0	0	0	0	
	<i>Tg</i>	202	158	0	0	0	298	0	298	
	<i>Tg</i>	193	151	241	276	517	0	0	0	
	<i>Sm</i>	210	205	46	225	271	189	0	189	
	<i>Sm</i>	208	154	165	0	165	0	0	0	
	<i>Sm</i>	191	143	0	0	0	104	0	104	
	<i>Cn</i>	192	49	0	0	0	0	0	0	
	<i>Cn</i>	184	40	0	0	0	0	0	0	
	<i>Cn</i>	203	82	0	0	0	35	0	35	
Jul-17	<i>Sm</i>	226	235	0	0	0	nd	nd	nd	
	<i>Sm</i>	220	240	0	0	0	nd	nd	nd	
	<i>Sm</i>	200	170	113	0	113	nd	nd	nd	
	<i>Tg</i>	250	210	0	0	0	nd	nd	nd	
	<i>Tg</i>	225	250	0	0	0	nd	nd	nd	
	<i>Cn</i>	170	50	0	0	0	nd	nd	nd	
	<i>Cn</i>	170	50	0	0	0	nd	nd	nd	
Aug-17	<i>Cn</i>	210	80	0	0	0	nd	nd	nd	
	<i>Cn</i>	195	60	0	0	0	nd	nd	nd	
	<i>Cn</i>	220	100	0	0	0	nd	nd	nd	
	<i>Tg</i>	148	76	0	0	0	nd	nd	nd	
	<i>Tg</i>	169	112	368	422	790	nd	nd	nd	
	<i>Tg</i>	180	125	396	696	1092	nd	nd	nd	
	<i>Sm</i>	242	290	0	0	0	nd	nd	nd	
	<i>Sm</i>	243	295	34	0	34	nd	nd	nd	
Sep-17	<i>Cn</i>	160	40	158	422	581	nd	nd	nd	
	<i>Tg</i>	125	32	0	0	0	nd	nd	nd	
	<i>Tg</i>	116	24	0	0	0	nd	nd	nd	
	<i>Cn</i>	165	45	0	0	0	nd	nd	nd	
	<i>Cn</i>	162	36	0	0	0	nd	nd	nd	
Oct-17	<i>Cn</i>	190	47	0	0	0	77	0	77	
	<i>Cn</i>	193	73	0	0	0	0	0	0	
	<i>Cn</i>	237	104	0	0	0	0	0	0	
	<i>Tg</i>	200	166	32	0	32	0	0	0	
	<i>Tg</i>	240	250	0	0	0	0	0	0	

	<i>Tg</i>	170	103	0	0	0	0	0	0	0
<b>Nov-17</b>	<i>Cn</i>	190	34	0	0	0	0	0	0	0
	<i>Cn</i>	210	49	0	0	0	0	0	0	0
	<i>Cn</i>	230	89	0	0	0	0	0	0	0
	<i>Tg</i>	197	151	67	0	67	0	0	0	0
	<i>Tg</i>	194	131	81	0	81	0	0	0	0
	<i>Tg</i>	190	145	0	0	0	41	0	41	0
	<i>Sm</i>	185	136	118	0	118	0	0	0	0
	<i>Sm</i>	195	162	0	0	0	84	0	84	0
	<i>Sm</i>	167	92	76	0	76	39	0	39	0
	<i>Hf</i>	150	81	0	0	0	0	0	0	0
	<i>Hf</i>	180	136	0	0	0	0	0	0	0
	<i>On</i>	167	98	0	0	0	0	0	0	0
	<i>On</i>	175	99	0	0	0	0	0	0	0
	<b>Dec-17</b>	<i>Cn</i>	195	39	0	0	0	nd	nd	nd
<i>Cn</i>		210	66	113	0	113	nd	nd	nd	0
<i>Tg</i>		150	77	0	0	0	nd	nd	nd	0
<i>Tg</i>		140	61	0	0	0	nd	nd	nd	0
<b>Jan-18</b>	<i>Cn</i>	212	117	0	0	0	nd	nd	nd	0
	<i>Cn</i>	160	46	0	0	0	nd	nd	nd	0
	<i>Cn</i>	210	69	0	0	0	nd	nd	nd	0
	<i>Tg</i>	135	49	249	269	518	nd	nd	nd	0
	<i>Tg</i>	165	86	0	0	0	nd	nd	nd	0
	<i>Tg</i>	152	77	0	0	0	nd	nd	nd	0
<b>Feb-18</b>	<i>Ho</i>	250	152	0	0	0	0	0	0	0
	<i>Ho</i>	265	170	227	0	227	0	0	0	0
	<i>Ho</i>	212	74	238	3	241	0	0	0	0
	<i>Cn</i>	190	81	0	0	0	0	0	0	0
	<i>Cn</i>	210	84	0	0	0	33	0	33	0
	<i>Cn</i>	179	54	0	0	0	0	0	0	0
	<i>Tg</i>	146	67	139	0	139	0	0	0	0
	<i>Tg</i>	153	79	166	303	469	0	0	0	0
	<i>Tg</i>	131	56	194	161	355	0	0	0	0
	<i>Hf</i>	126	40	0	0	0	0	0	0	0
	<i>Hf</i>	123	36	0	0	0	0	0	0	0
	<i>Hf</i>	124	35	80	114	194	0	0	0	0
	<i>Sm</i>	153	77	221	112	333	0	0	0	0
	<i>Sm</i>	128	43	78	133	211	50	0	50	0
<b>Mar-18</b>	<i>Cn</i>	230	110	0	0	0	0	0	0	0
	<i>Cn</i>	220	82	155	116	272	0	0	0	0
	<i>Cn</i>	163	46	0	0	0	108	0	108	0
	<i>Tg</i>	150	67	0	0	0	132	0	132	0
	<i>Tg</i>	134	49	188	0	188	0	0	0	0
	<i>Tg</i>	135	53	0	0	0	50	158	209	0
	<i>On</i>	125	43	241	130	371	0	0	0	0
	<i>On</i>	137	46	56	0	56	42	0	42	0

	<i>On</i>	133	50	71	0	71	0	0	0	0
	<i>On</i>	125	46	75	0	75	0	0	0	0
<b>Apr-18</b>	<i>Tg</i>	179	112	0	0	0	nd	nd	nd	0
	<i>Tg</i>	166	87	61	0	61	nd	nd	nd	0
	<i>Tg</i>	193	137	154	160	314	nd	nd	nd	0
	<i>Cn</i>	200	95	0	0	0	nd	nd	nd	0
	<i>Cn</i>	195	100	0	0	0	nd	nd	nd	0
	<i>Cn</i>	165	46	0	0	0	nd	nd	nd	0

181  
182

### 183 **Cyanotoxin identification and quantification from phytoplankton cells and fish**

184 The same protocols as those described in Ahoutou et al. (2022) were used for all toxin  
185 analyses. Briefly, for intracellular cyanotoxins (in the phytoplanktonic biomass), the  
186 extraction was performed from biomass on polycarbonate filters (Whatman®  
187 Nuclepore) stored at -80 °C. After thawing, the filters were placed in sterile glass  
188 containing 4 mL of 75% methanol and then sonicated on ice with an ultrasound probe  
189 (Sonics Vibra Cell). The extracts were centrifuged, and the supernatants were  
190 collected and stored at 4 °C. The resulting pellets underwent a second extraction using  
191 the same protocol and then were centrifuged again to ensure that all toxins were  
192 extracted. These second supernatants were added to the first and centrifuged again.  
193 They were further frozen at -80 °C until cyanotoxin analysis.

194 For cyanotoxin extractions from the fish, the method of Manubolu et al. (2018)  
195 was used with minor adjustments. For each fish, 100 mg of fresh liver, muscle, and  
196 intestinal tissues were used for cyanotoxin extraction. The 100 mg muscle extracts  
197 were further previously freeze-dried and underwent ball milling. All samples were  
198 extracted twice with BuOH:MeOH:H<sub>2</sub>O (1:4:15), submitted to an ultrasonic probe  
199 (Sonics Vibra Cell, 130 Watt, 20 kHz, 60%, 30 sec, three cycles) and centrifuged  
200 (15 300 G, 10 min, 4 °C). The collected supernatants underwent final centrifugation  
201 (15 300 G, 10 min, 4 °C) and were stored at -80 °C until analysis.

202 The identification and quantification of the cyanotoxins in the water and fish  
203 samples were performed by ultrahigh-performance liquid chromatography in tandem  
204 (Elute UHPLC—Bruker) coupled with high-resolution mass spectrometry (Compact  
205 QTOF—Bruker) (UHPLC–MS/MS). All samples were separated on a C18 stationary  
206 phase column (Acclaim RSLC Polar Advance 2, 2.2  $\mu\text{m}$ , Thermo Fisher® 2.1 Å~ 100  
207 mm) along a linear gradient of acidified acetonitrile and acidified ultrapure water and  
208 then analyzed between 50 and 1500 m/z in positive BbCID (MS-MS/MS) mode,  
209 alternating at 2 Hz between low and high-energy MS and MS2 modes, respectively,  
210 with a mass accuracy of less than 0.5 ppm. The injection volume and data acquisition  
211 time were 4  $\mu\text{L}$  and 21 min, respectively. The elution rate was 0.5  $\mu\text{L min}^{-1}$ . The  
212 analytical standards of the cyanotoxins investigated were saxitoxin (STX) (CAS 35554-  
213 086), anatoxin-a (ATX) (CAS 64285-06-9), homoanatoxin (HTX) (CAS 142926-86-1),  
214 nodularin (NOD) (CAS 118399-22-7), cylindrospermopsin (CYL) (CAS 143545-90-8),  
215 and seven variants of microcystins (MC) (variant MC-LR, -LA, -LF, -LW, -LY, -RR and  
216 -YR corresponding to CAS 101043-37-2; CAS 96180-79-9; CAS 154037-70-4; CAS  
217 157622-02-1; CAS 123304-10-9; CAS 11755-37-4; and CAS 10164-48-6,  
218 respectively), all certified one year, purity >99% (Novakit®). Data were integrated using  
219 TASQ 1.1 software (Bruker®). Cyanotoxins were identified according to (i) retention  
220 time, (ii) molecular mass, (iii) isotopic pattern, and (iv) diagnostic ions. They were then  
221 quantified according to the area under the peak signal determined for analytical  
222 standards. *Phormidium favosum* (PMC 240.05), *Raphidiopsis raciborskii* (PMC 99.03),  
223 *Aphanizomenon gracile* (PMC 638.10), and *Microcystis aeruginosa* (PMC 728.11)  
224 strains were used as positive controls for ATX, CYL, STX and MC, respectively. Except  
225 for MC-RR, MC-LR, and MC-YR, other MC structural variants were quantified

226 according to MC-LR equivalence, calculated from the regression curves of the MC-LR  
227 analytical standard.

228 For phytoplankton, the LOD and LOQ were equal to 0.0003 and 0.001  $\mu\text{g L}^{-1}$  of  
229 raw water for MC-RR and 0.001 and 0.004  $\mu\text{g L}^{-1}$  for MC-LR, respectively. For fish  
230 tissues, the LOD and LOQ were equal to 1.3 and 12.8  $\mu\text{g kg}^{-1}$  fresh weight (FW) for  
231 MC-RR and 0.5 and 4.8  $\mu\text{g kg}^{-1}$  for MC-LR, respectively.

232

### 233 **Statistical analyses**

234 Wilcoxon pairwise tests, sign tests, ordinary least squares regression and multiple  
235 linear regression (MLR) were performed by using PAST software v4.11.

236

## 237 **Results**

### 238 **Spatial and temporal variations in cyanobacteria abundances and biovolumes**

239 A total of 15 cyanobacterial genera were found during the monitoring of the lagoon:  
240 *Aphanizomenon*, *Aphanocapsa*, *Aphanothece*, *Chroococcus*, *Dolichospermum*,  
241 *Gomphosphaeria*, *Leptolyngbya*, *Limnothrix*, *Lyngbya*, *Merismopedia*, *Microcystis*,  
242 *Oscillatoria*, *Planktothrix*, *Pseudanabaena* and *Raphidiopsis* (previously  
243 *Cylindrospermopsis*) (Supplementary Table S1).

244

245 **Supplementary Table S1** Cyanobacterial abundances ( $\text{cell mL}^{-1}$ ) and potential toxicity  
246 of the 15 cyanobacterial genera identified during the 17-month monitoring performed  
247 at six sampling stations in Lagoon Aghien  
248



Table with columns: Potential toxicity, ATX-CVI-STX, M\*, ATX-CVI-MC-STX, M\*, M\*, CVI-STX, M\*, M\*, ATX-MC, M\*, ATX-MC, CVI-STX. The table contains numerical data for various sampling dates from May to April. A large watermark 'Capture d'écran' is visible across the middle of the table.

249

250

251


252

253

Few variations were found in the richness and diversity (Shannon and Simpson indices) of the cyanobacterial community at the genus level during the 17-month monitoring of Lagoon Aghien (Fig. 2), except for the decrease in the two diversity

254 indices in January 2018. These low values in richness and diversity were associated  
 255 with a high dominance of the genus *Limnothrix* in the cyanobacterial community (Fig.  
 256 2 and 3).

257



258

259 **Fig. 2** Monthly variations in the richness and diversity (evenness, Shannon and  
 260 Simpson indices calculated on the average cyanobacterial abundances at the six  
 261 sampling stations) of the cyanobacterial community estimated at the genus taxonomic  
 262 level in Lagoon Aghien. Monthly cumulative rainfall values (cm) have been added to  
 263 the figure.

264

265


266

The total cyanobacterial abundances (sum of all genera) estimated during the  
 267 monitoring survey ranged from  $1.6 \times 10^4$  to  $3.1 \times 10^5$  cell mL<sup>-1</sup> (Supplementary Table S1),  
 268 and monthly average abundances ranged from  $4.1 \times 10^4$  cell mL<sup>-1</sup> in July 2017 to  
 269  $1.8 \times 10^5$  cell mL<sup>-1</sup> in March 2017. As shown in Figure 3, two main peaks of total  
 270 cyanobacteria abundances were observed in February-March 2017 and February  
 271 2018. The four dominant cyanobacterial genera in terms of cellular abundance were  
 272 *Raphidiopsis*, *Limnothrix*, *Microcystis* and *Oscillatoria*. The main peak abundances of



273 these four genera were found from January to April 2017 and from December 2017 to  
 274 February 2018 when rainfall was low (Fig. 3).

275




276

277 **Fig. 3** Variations in the monthly average abundances of the total cyanobacteria  
 278 abundances (black curve) and of each cyanobacterial genus abundance (colored  
 279 curves) (mean values from the six sampling stations) and variations in the cumulative  
 280 monthly rainfall (gray bar chart). Standard deviations have only been added for the  
 281 monthly averages of total cyanobacteria abundances. Total cya = Total cyanobacteria  
 282 abundance.

283

284 A negative correlation was found between monthly variations occurring in the  
 285 monthly cumulative rainfall and those occurring in the total cyanobacterial abundance  
 286 averages ( $r^2=0.24$ ; ordinary least squares regression,  $p<0.05$ ) (Supplementary Fig.  
 287 S2).



288

289 **Supplementary Fig. S2** Correlation between Log<sub>10</sub> monthly cyanobacterial  
 290 abundance averages and Log<sub>10</sub> monthly cumulative rainfall values

291


292

293 Finally, among the 15 cyanobacteria genera observed in the lagoon, twelve are  
 294 known to produce cyanotoxins. More specifically, four of them are known to produce  
 295 ATX or HTX, five are known to produce CYL, nine are known to produce MC and four  
 296 are known to produce STX (Supplementary Table S1).

297 As observed for the variations in the total cyanobacterial abundances, the  
 298 variations in the monthly averages of the potentially MC-producing cyanobacteria  
 299 abundances (*Aphanocapsa*, *Dolichospermum*, *Leptolyngbya*, *Limnothrix*,  
 300 *Merismopedia*, *Microcystis*, *Oscillatoria*, *Planktothrix* and *Pseudanabaena*) were  
 301 characterized by a first peak of abundance at the beginning of 2017 followed by a  
 302 strong decrease in June-July 2017 and a progressive increase in average abundance  
 303 values until the end of the monitoring (Supplementary Fig. S3).

304

305



306  
307  
308  
309  
310  
311  
312  
313

**Supplementary Fig. S3** Variations in the monthly average abundances of all cyanobacteria and of cyanobacteria potentially able to produce microcystins (mean values from the six sampling stations) during the monitoring of Lagoon Aghien Total = monthly average abundances of all cyanobacteria; Pot. MC+ = monthly average abundances of cyanobacteria potentially able to produce microcystins

314 Finally, as shown in Fig. 4A and by a Wilcoxon pairwise test (Supplementary  
315 Table S2), Station 6 and, to a lesser extent, Station 3 were characterized by lower  
316 monthly average abundances of cyanobacteria potentially able to produce MC.


317

318 **Supplementary Table S2** p values of the Wilcoxon paired test performed on the  
319 monthly average abundances of cyanobacteria potentially able to produce  
320 microcystins at the six sampling stations during the monitoring of Lagoon Aghien (bold  
321 characters = significant p values)  
322

Stations	St1	St2	St3	St4	St5	St6
St1	-	0,43	<b>0,05</b>	0,61	0,21	<b>0,01</b>
St2		-	0,26	0,46	0,96	<b>0,04</b>
St3			-	<b>0,02</b>	0,21	0,52
St4				-	0,40	<b>0,01</b>
St5					-	0,16
St6						-

323

324  
325  
326



327  
328  
329  
330  
331  
332

**Fig. 4A & B** Box plot of the variations in the monthly average abundances of potentially MC-producing genera (A) and in MC-RR concentrations (B) recorded at the six sampling stations in Lagoon Aghien

333 **Spatial and temporal variations in intracellular cyanotoxin concentrations**

334 During the survey, intracellular STX, ATX, HTX, NOD and CYL were not detected. Two  
335 MC congeners, MC-RR and MC-LR, were detected throughout the year, but MC-LR  
336 concentrations were always under the limits of quantification.

337 MC-RR was detected in cyanobacterial cells during all monitoring (Figure 5).  
338 The MC-RR concentrations ranged from 0 to 0.364  $\mu\text{g L}^{-1}$  (mean value during the  
339 monitoring = 0.08  $\mu\text{g L}^{-1}$ ) and were not correlated with the total abundances of  
340 potentially MC toxigenic cyanobacteria ( $R^2=0.01$ ). There was no correlation between  
341 MC-RR concentrations and the cell abundances of each of the nine genera potentially  
342 able to produce MC. Finally, the MLR analysis performed by using MC concentration  
343 as the dependent variable and cell abundance of each potentially MC-toxigenic  
344 cyanobacteria as the independent variables, showed that MC-RR concentrations were  
345 positively affected by *Microcystis* cell abundance (cell  $\text{mL}^{-1}$ ) and negatively affected by  
346 *Oscillatoria* cell abundance (cell  $\text{mL}^{-1}$ ) ( $p<0.05$ ). The regression model obtained by  
347 MLR was:  $\text{MC-RR} = 0.072 + 1.22 \cdot 10^{-6} (\text{Microcystis}) - 2.99 \cdot 10^{-6} (\text{Oscillatoria})$ .

348 The comparison of these MC-RR concentrations at the six sampling sites shows  
349 that these concentrations were significantly lower at station 6 (median concentration =  
350 0.000  $\mu\text{g L}^{-1}$ , respectively) than at the other stations (median concentrations between  
351 0.055 and 0.079  $\mu\text{g L}^{-1}$ ) (Fig. 4B and Supplementary Table S3).

352  
353 **Supplementary Table S3** p values of the paired sign tests performed on the  
354 intracellular MC-RR concentrations recorded at the six sampling stations (S1 to S6)  
355 during the monitoring of Lagoon Aghien (bold characters = significant p values)  
356


Stations	St1	St2	St3	St4	St5	St6
St1	-	0.80	0.80	0.80	0.80	<b>0.00</b>
St2		-	0.46	0.80	0.80	<b>0.02</b>
St3			-	0.12	0.80	<b>0.00</b>
St4				-	0.80	<b>0.00</b>
St5					-	<b>0.00</b>
St6						-

357

358

359 The monthly variations in MC-RR concentrations were broadly similar at all  
 360 sampling stations, with a first peak in April-May 2017, followed by a sharp decrease in  
 361 June, and then a second peak in September-October 2017, followed by a sharp  
 362 decrease in November 2017 (Fig. 5). These two peaks in MC-RR concentration  
 363 followed an increase in nitrate concentrations in the lagoon (Supplementary Fig. S4).


364



365

366 **Fig. 5.** Variations in the intracellular microcystin-RR concentrations at the six sampling  
 367 stations and of rainfall values during the monitoring of Lagoon Aghien (no value in  
 368 February 2018). MC St1-6= Microcystin concentrations at sampling stations 1 to 6.  
 369 Average MC = Average MC concentrations at the six sampling stations.

370



371  
372  
373  
374  
375  
376  
377

**Supplementary Figure S4.** Variations in the intracellular microcystin-RR and dissolved nutrient average concentrations during the monitoring of Lagoon Aghien (no value in February 2018)

### 378 Temporal variations in microcystin concentrations in fish tissues

379 No STX, ATX, HTX, NOD or CYL were detected in any of the fish tissues analyzed,  
380 while MC-RR and MC-LR were frequently detected in the intestines and livers of all  
381 fish species but never in muscle tissues (Table 1). MC-RR and/or MC-LR were  
382 quantified in 32 of the 97 intestines analyzed (33%) and in 18 of the 61 livers analyzed  
383 (30%) (Table 1).

384 Among the 61 fish for which MC concentrations were quantified in liver and/or  
385 in intestine, MC-RR/LR were quantifiable (i) both in intestine and liver in 8% of the fish  
386 for which intestines and liver were analyzed, (ii) only in liver for 28% and (iii) only in  
387 intestine for 23% of them (Table 1). Total MC concentrations (MC-RR + MC-LR)

388 ranged between 32 and 1,092  $\mu\text{g kg}^{-1}$  FW in the intestines and between 33 and 383  
389  $\mu\text{g kg}^{-1}$  FW in the livers (Table 1).

390 As shown in Table 2, MC was found in a higher proportion of the intestines in  
391 phytoplanktivorous fish (56 and 67% for *O. niloticus* and *S. melanotheron*, respectively)  
392 than in other dietary regimes (from 9 to 39% for all the other fish species). Moreover,  
393 there was a significant difference (Sign test,  $p < 0.05$ ) in the monthly averages of  
394 intestine MC concentrations of phytoplanktivorous-omnivorous fish and  
395 zooplanktivorous-carnivorous fish, with these concentrations being higher in the  
396 intestines of phytoplanktivorous-omnivorous fish. There were no significant differences  
397 (Kruskal–Wallis test) in the MC concentrations estimated in the livers of *T. guineensis*  
398 (omnivorous), *S. melanotheron* (phytoplanktivorous) and *C. nigrodigitatus*  
399 (zooplanktivorous).

400

401 **Table 2** The proportions of the intestines and livers containing microcystins in the six  
402 fish species sampled in Lagoon Aghien

403 C = carnivorous fish; Z = zooplanktivorous fish; P = phytoplanktivorous fish; O = omnivorous  
404 fish

405

	Intestine	Liver
<i>Chrysichthys nigrodigitatus</i> (Z)	3/34 (9%)	5/18 (28%)
<i>Hemichromis fasciatus</i> (C)	1/5 (20%)	0/5 (0%)
<i>Hepsetus odoe</i> (C)	1/3 (33%)	0/3 (0%)
<i>Tilapia guineensis</i> (O)	13/33 (39%)	5/18 (28%)
<i>Sarotherodon melanotheron</i> (P)	9/16 (56%)	7/11 (64%)
<i>Oreochromis niloticus</i> (P)	4/6 (67%)	1/6 (17%)

406

407



408 **Discussion**

409 *Cyanobacterial community composition and dynamics*

410 Fifteen cyanobacterial genera were observed during the survey of Lagoon Aghien,  
411 which is within the same range as data obtained in other tropical freshwater  
412 ecosystems, such as those reported by Olokotum et al. (2022), who identified 16  
413 cyanobacterial genera in two embayments of Lake Victoria (Uganda), and by Dalu and  
414 Wasserman (2018), who identified 13 cyanobacterial genera in a reservoir in  
415 Zimbabwe. In contrast with temperate areas, few variations in the richness of the  
416 cyanobacterial community were observed during the Lagoon Aghien survey, probably  
417 because of the temporal stability of the water temperature of the lagoon (between 26.3  
418 °C in July 2017 and 31.6 °C in March 2017; Ahoutou et al. 2021). Studies performed  
419 in temperate areas have shown that the decrease in water temperature is one of the  
420 main factors associated with a decrease in richness in phytoplankton communities  
421 (e.g., Catherine et al. 2016; Maberly et al. 2022).

422         Among these 15 genera, four were dominant: *Raphidiopsis* (formerly  
423 *Cylindrospermopsis*), *Microcystis*, *Oscillatoria* and *Limnothrix*. The first three genera  
424 are the most frequently reported in studies performed in Sub-Saharan Africa (Svircev  
425 et al.2019), while *Limnothrix* is mainly found in shallow and eutrophic lakes of  
426 temperate areas (e.g., Rucker et al. 1997; Wiedner et al. 2002), even if it is also able  
427 to proliferate in tropical ecosystems, as shown in Brazil by Soares et al. (2009). The  
428 highest cyanobacterial abundances were generally associated with the genus  
429 *Raphidiopsis*, which is known to have high fitness at tropical latitudes and to be able  
430 to outcompete other cyanobacterial genera under these environmental conditions  
431 (e.g., Burford et al. 2016; Jia et al. 2020). Finally, we found that the variations in total  
432 cyanobacterial abundances were negatively correlated with rainfall, as previously

433 observed in other tropical freshwater ecosystems, this negative relationship being  
434 probably explained by the water dilution of cyanobacteria combined with high flushing  
435 rates (e.g., Barros et al. 2017; Giani et al. 2020). As shown by a canonical correspondence  
436 analysis performed by Ahoutou et al. (2021), only 22% of the variation in biovolume values of  
437 the dominant phytoplankton genera (including *Raphidiopsis* and *Limnothrix*) in the Lagoon  
438 Aghien phytoplankton community could be explained by variations in environmental variables.

439

#### 440 *Toxin production*

441 In Lagoon Aghien, 12 of the 15 cyanobacteria genera observed can potentially  
442 produce ATX, HTX, CYL, MC and/or STX (ANSES 2020), but only MC was found in  
443 the phytoplankton biomass. These data are in agreement with those reported in the  
444 reviews of Mowe et al. (2015) and Svircev et al. (2019), showing that MC are the most  
445 frequent cyanotoxins on the African continent during cyanobacterial blooms. Despite  
446 the high abundances of the genus *Raphidiopsis*, known to be a potential producer of  
447 CYL and/or STX (Vico et al. 2020), these two cyanotoxins were not found in Lagoon  
448 Aghien water. In two papers dealing with the production of CYL in Sub-Saharan Africa,  
449 all the *Raphidiopsis* strains isolated in Senegal and Uganda were shown to lack the  
450 genes involved in the biosynthesis of this toxin (Berger et al. 2006; Haande et al. 2008).  
451 These data should also be considered in light of the recent study of Vico et al. (2020),  
452 which suggested by genomic and phylogenetic analyses that central Africa was the  
453 original dispersion center of *Raphidiopsis* (*Cylindrospermopsis*) *raciborskii* and that  
454 these African populations were not able to produce CYL.

455 The intracellular MC concentrations in the water were quite low, always  $<0.4 \mu\text{g}$   
456  $\text{equi. MC-LR eq L}^{-1}$  and were not correlated with the total abundances of  
457 cyanobacterial genera known to be able to produce MC or with the abundance of each  
458 of these genera. On the other hand, our MLR analysis showed that MC-RR

459 concentrations were negatively affected by *Oscillatoria* cell abundances, but positively  
460 affected by *Microcystis* cell abundances. This finding is consistent with the fact that  
461 this genus is considered to be the major MC-producing genus in Africa (see reviews of  
462 Mowe et al. 2015; Chia et al. 2022). It is known that the variations in MC concentration  
463 are partly explained by the variations in the proportions of MC-producing (MC+) and  
464 non-MC-producing (MC-) strains in *Microcystis* populations (e.g., Briand et al. 2009;  
465 Suominen et al. 2017). Their respective proportions were not estimated in our study,  
466 and few data are available in Africa on this topic. Haande et al. (2007) found that only  
467 four of 24 *M. aeruginosa* isolates were able to produce MC, and Ballot et al. (2014)  
468 found that only one of 16 *M. aeruginosa* isolates was able to produce MC. These first  
469 data are interesting to consider, but knowing that there is very high variability in  
470 Europe/North America in the variations of MC+/MC- proportions during a bloom event  
471 or from one bloom to another (e.g., Briand et al. 2009; Sabart et al. 2009; Bozarth et  
472 al. 2010), much more data are necessary to get a real idea of the variability of these  
473 proportions in cyanobacterial populations in Sub-Saharan Africa.

474         The average abundances of potentially MC-producing cyanobacteria and of MC  
475 concentrations in water were lower at sampling Station 6. As shown by Ahoutou et al.  
476 (2022), the values of some physico-chemical parameters, such as turbidity and  
477 phytoplankton community composition and biomass, were significantly different at  
478 Station 6, probably because this station is placed under the direct influence of the Mé  
479 river and Potou channel (Koffi et al. 2019).

480         Another point to consider for explaining these low MC concentrations in Lagoon  
481 Aghien concerns nitrogen availability. In their review, Mowe et al. (2015) reported a  
482 positive correlation between N:P ratio values and MC concentrations from data  
483 collected in several tropical lakes. It is also well known that an increase in N supply

484 leads to an increase in MC cell quotas and concentrations (e.g., Chaffin et al. 2018;  
485 Horst et al. 2014; Wagner et al. 2021). During this monitoring of Lagoon Aghien, the  
486 N:P ratio values ranged between 7 and 14 (Ahoutou et al. 2020), which corresponds  
487 to the low N:P ratio and low microcystin concentrations reported by Mowe et al. (2015).  
488 It is interesting to note that in the present study the two MC peaks followed an increase  
489 in nitrate concentration, suggesting a possible link between these events. Moreover,  
490 during mesocosm experiments performed in Lagoon Aghien (Ahoutou et al. 2022), we  
491 found that the addition of nitrogen, alone or combined with phosphorus in the  
492 mesocosms filled with the water of the lagoon, stimulated MC production by  
493 cyanobacteria. Taken together, these results are interesting to consider in light of the  
494 data provided by Chianu et al. (2012) showing that nitrogen fertilizer consumption is  
495 low in African farming systems compared to northern countries. This suggests that an  
496 increase in the use of nitrogen fertilizers on this continent might lead to increasing MC  
497 concentrations in African freshwater ecosystems in the future.

498         Few data are available on fish contamination by MC in Africa (Abdallah et al.  
499 2021, Roegner et al. 2023), but our data are interesting to compare with those of  
500 Semyalo et al. (2010), who dealt with the MC contamination of Nile tilapia (*O. niloticus*)  
501 in two Ugandan lakes (Lake Mburo and Lake Victoria). The MC concentrations in the  
502 cyanobacterial biomass in Lake Mburo ranged between 0.006 and 0.26  $\mu\text{g eq. MC-LR}$   
503  $\text{L}^{-1}$  and were close to those recorded in Lagoon Aghien (between 0.02 and 0.25  $\mu\text{g MC}$   
504  $\text{L}^{-1}$ ). These MC concentrations were higher in Murchison Bay (Lake Victoria), with  
505 values ranging between 0.2 and 0.7  $\mu\text{g eq. MC-LR L}^{-1}$ , depending on the season. In  
506 the fish intestines, the highest MC concentrations recorded in the *O. niloticus* sampled  
507 in the two Ugandan lakes were  $>300$  and  $>390 \mu\text{g.kg}^{-1}$  FW in Lake Mburo and  
508 Murchison Bay, respectively, whereas the highest MC concentration recorded in

509 Lagoon Aghien was 1,092  $\mu\text{g}\cdot\text{kg}^{-1}$  FW when considering all the fish and 371  $\mu\text{g}\cdot\text{kg}^{-1}$   
510 FW when considering only *O. niloticus*. MC was most frequently found in the intestine  
511 and liver of phytoplanktivorous fish, consistent with the consumption and presence of  
512 cyanobacterial cells in the fish intestine and subsequent accumulation in the liver, the  
513 target organ of MCs. An interesting point was that in carnivorous fish (*H. fasciatus*, *H.*  
514 *odoe*), MC was only quantified in the liver and not in the intestine. This is probably  
515 related to their mode of intoxication via contaminated prey (no direct ingestion of  
516 cyanobacterial cells), the short length of their intestines and their rapid digestion  
517 time, allowing MC to be rapidly assimilated after prey digestion and directed to the liver  
518 for accumulation and metabolism. Such differences in MC accumulation in the gut and  
519 liver of phytoplanktivorous versus carnivorous fish have been previously reported by  
520 Ibeling et al. (2005), Zhang et al. (2009) and Nyakairu et al. (2010).

521 Although similar MC concentrations were found in the fish intestines in Uganda  
522 and Ivory Coast, no MC was found in the fish muscle tissues in Lagoon Aghien, while  
523 low concentrations ( $<6 \mu\text{g}\cdot\text{kg}^{-1}$  FW) were found in these tissues in fish from Mburo and  
524 Victoria Lakes. In the same way, MC concentrations ranging between 0.5 and 1,917  
525  $\mu\text{g}\cdot\text{kg}^{-1}$  FW were found in the muscle tissue of various fish species (including *O.*  
526 *niloticus*) collected in several Ugandan lakes by Poste et al. (2011), and some of these  
527 MC concentrations exceeded the tolerable daily intake guideline defined by the WHO  
528 for chronic exposure to MC. More recently, Roegner et al. (2023) found in the Winam  
529 Gulf (Lake Victoria, Kenya) that the hazard quotients for fish consumed by young  
530 children were 5 to 10 times higher than the permissible levels.

531 This absence of MC in muscle tissues of fish collected in Lagoon Aghien is  
532 intriguing knowing that similar MC concentrations in water and in fish intestine were  
533 found in this lagoon and in two Ugandan lakes (Semyalo et al. 2010), where MC was

534 found in muscle tissues. High MC concentrations in fish muscle have been frequently  
535 estimated by ELISA and this method may overestimate these MC concentrations (see  
536 Ibelings et al. 2021). This could partly explain the lack of MC detection in the muscle  
537 tissue of fish from Lagoon Aghien, but the same analytical approach (LC–MS/MS) as  
538 ours was used in the study of Semyalo et al. (2010), where MC was found in fish  
539 tissues. However, in our study, our LOQ values were quite high (12.8  $\mu\text{g kg}^{-1}$  FW for  
540 MC-RR and 4.8  $\mu\text{g kg}^{-1}$  FW for MC-LR), while the LOQ was not provided by Semyalo  
541 et al. (2010). Consequently, the existence of moderate MC contamination in the muscle  
542 of fish in Lagoon Aghien cannot be ruled out. Furthermore, only free MC was quantified  
543 in our study, whereas much of the accumulated MC may be covalently bound in animal  
544 tissues requiring specific extraction protocols (Neffling et al. 2010, Bouteiller et al.  
545 2021). Finally, as shown in the review paper of Martins and Vasconcelos (2009), fish  
546 are efficient in detoxifying MC, depending on the species and on the environment. In  
547 contrast with fish living in freshwater ecosystems from temperate areas, which are  
548 exposed to MC for only a few months, we showed that the fish community of Lagoon  
549 Aghien is contaminated throughout the year by MC, which could have led this  
550 community to develop active processes for the detoxification of MC.

551 All these data on toxin production by cyanobacteria in Lagoon Aghien raise the  
552 issue of health hazards for populations living around the lagoon. The WHO has  
553 provided guideline values for lifetime and short-term exposure to MC-LR by drinking  
554 water and for exposure by recreational activities (1, 12 and 24  $\mu\text{g L}^{-1}$ , respectively)  
555 (Chorus and Welker 2021). In France, more restrictive guideline values (0.1  $\mu\text{g L}^{-1}$  for  
556 drinking water and 0.3  $\mu\text{g L}^{-1}$  for recreational activities) have been adopted because  
557 the French Agency for Food, Environmental and Occupational Health and Safety

558 (ANSES) has taken into account the findings provided by Chen et al. (2001) showing  
559 reprotoxic effects of MC-LR during chronic low-dose exposure of mice.

560 The WHO TDI for microcystin yields a threshold concentration of microcystin in  
561 fish of 24  $\mu\text{g kg}^{-1}$  FW for a consumer weighing 60 kg and consuming 100 g of fish daily  
562 (Poste et al. 2011). Knowing that no MC was found in the muscle of the six fish species  
563 studied and even if we cannot exclude the presence of a low MC content in fish  
564 muscles, it is likely that the exposure risk by fish consumption of human populations  
565 living around Lagoon Aghien is probably limited. However, local populations consume  
566 some small fish species with viscera (i.e., *Pellonulla leonensis*, which was not sampled  
567 in this study), which could potentially contribute to their exposure to MC, since MC  
568 were found in the intestines of all the species targeted in this study.

569 Taking into account all these guidelines, human populations living around the  
570 lagoon seem to be moderately exposed to adverse health effects of MC because MC  
571 concentrations in water and fish are generally below the WHO thresholds. However, in  
572 agreement with Roegner et al. (2020), we must question the relevancy of these  
573 guideline values, which have been established in developed countries where water  
574 uses are very different and have consequences on the cyanotoxin exposure level. For  
575 example, around Lagoon Aghien, children play for several hours every day of the first  
576 twelve years of their life in water continuously containing approximately 0.2  $\mu\text{g L}^{-1}$  of  
577 MC, as we found, for example, between April 2017 and 2018. Nontreated lagoon water  
578 is also used for meal preparation, to wash children since the first weeks of their life, to  
579 wash dishes and clothes, etc. In addition, these populations eat fish from the lagoon  
580 almost every day, with fish being the main source of protein for them (Ahi 2021; Effebi  
581 et al. 2017). Daily exposure to low MC concentrations by different exposure routes

582 over more than 10 years is not taken into account in the current guidelines, making  
583 children in particular a potentially very vulnerable population.

584 Even if guidelines adapted to developing countries were provided by the WHO,  
585 they would be far from being used in Sub-Saharan African countries, where there is  
586 usually no regular monitoring of cyanobacteria and their toxins in freshwater  
587 ecosystems. As observed for Lagoon Aghien and Lake Guiers (Senegal), which are  
588 used for the production of drinking water for the cities of Abidjan and Dakar,  
589 respectively, and face problems with cyanobacteria, the two main difficulties for the  
590 implementation of long-term monitoring of these resources are (i) the competition  
591 between national water institutions for the leadership of such a program and (ii) the  
592 sustainable funding of this monitoring, none of the ministries concerned agreeing to  
593 ensure it (Mitroi et al. 2022). Beyond the resolution of these governance difficulties,  
594 another key point in Africa will also be to inform the local populations and make them  
595 aware of the health hazards associated with cyanobacterial blooms. Finally, even if  
596 guidelines adapted to the local water practices are provided and monitoring and  
597 awareness of local populations are performed, the issue of the provision of alternative  
598 resources (in water, proteins, etc.) will be the last key step needed to reduce the  
599 exposure of these populations to MC.

600

601



602 **Acknowledgments**

603 We are extremely grateful to the inhabitants and chiefdoms of the villages located  
604 around Lagoon Aghien for their agreement to perform this study on the lagoon. This  
605 study was supported by the French Facility for Global Environment (Fonds Français  
606 pour l'Environnement Mondial, FFEM) (research program WaSAf).

607

608

609 **References**

- 610 Abdallah MF, Van Hassel WHR, Andjelkovic M, Wilmotte A, Rajkovic A (2021)  
611 Cyanotoxins and food contamination in developing countries: Review of their types,  
612 toxicity, analysis occurrence and mitigation strategies. *Toxins* 13 :786.  
613 <https://doi.org/10.3390/toxins13110786>
- 614 Ahi KC (2021) Pratiques locales, représentations sociales et inégalités socio-  
615 environnementales autour des enjeux de gestion intégrée de la lagune Aghien en  
616 Côte d'Ivoire. Dissertation, Université Félix Houphouët-Boigny d'Abidjan
- 617 Ahoutou MK, Djeha RY, Yao EK, Kone M, Quiblier C, Niamen-Ebrottié J et al (2021)  
618 Assessment of some key indicators of the ecological status of an African freshwater  
619 lagoon (Lagoon Aghien, Ivory Coast). *PLoS ONE* 16:e0251065.  
620 <https://doi.org/10.1371/journal.pone.0251065>
- 621 Ahoutou MK, Yao EK, Djeha RY, Kone M, Tambosco K, Duval C et al (2022) Impacts  
622 of nutrient loading and fish grazing on the phytoplankton community and cyanotoxin  
623 production in a shallow tropical lake: Results from mesocosm experiments.  
624 *MicrobiologyOpen* 11:e1278. <https://doi.org/10.1002/mbo3.1278>
- 625 ANSES (2020). Actualisation de l'évaluation des risques liés à la présence de  
626 cyanobactéries et leurs toxines dans les eaux destinées à l'alimentation, les eaux  
627 de loisirs et les eaux destinées aux activités de pêche professionnelle et de loisir :  
628 Avis de l'Anses Rapport d'expertise collective, Anses, 495 p.
- 629 Ash AK, Patterson S (2022) Reporting of freshwater cyanobacterial poisoning in  
630 terrestrial wildlife: A systematic map. *Animals* 12:2423.  
631 <https://doi.org/10.3390/ani12182423>
- 632 Ballot A, Sandvik M, Rundberget T, Botha CJ, Miles CO (2014) Diversity of  
633 cyanobacteria and cyanotoxins in Hartbeespoort Dam, South Africa. *Mar Freshw*  
634 *Res* 65:175-189.
- 635 Barros MUG, Lopes IKC, Carvalho SMD, Neto JC (2017) Impact of filamentous  
636 cyanobacteria on the water quality of two tropical reservoirs. *Rev Braz de Recur*  
637 *Hidr* 22:e6. <https://doi.org/10.1590/2318-0331.011716072>
- 638 Berger C, Ba N, Gugger M, Bouvy M, Rusconi F, Couté A et al (2006) Seasonal  
639 dynamics and toxicity of *Cylindropspermopsis raciborskii* in Lake Guiers (Senegal,  
640 West Africa). *FEMS Microbiol Ecol* 57:355-366. [https://doi.org/10.1111/j.1574-](https://doi.org/10.1111/j.1574-6941.2006.00141.x)  
641 [6941.2006.00141.x](https://doi.org/10.1111/j.1574-6941.2006.00141.x)

642 Blé MC, Alla YL, Kervarec F (2008) Régimes alimentaires de trois principales espèces  
643 piscicoles élevées en Côte d'Ivoire. Fiches Techniques et Documents de  
644 Vulgarisation 10-14.

645 Bouteiller P, Lance E, Guérin T, Biré R (2022) Analysis of total-forms of cyanotoxins  
646 microcystins in biological matrices : A critical review. *Toxins* 14:14080550.  
647 <https://doi.org/10.3390/toxins14080550>

648 Bozarth CS, Schwartz AD, Shepardson JW, Colwell FS, Dreher TW (2010) Population  
649 turnover in a *Microcystis* bloom results in predominantly nontoxic variants late  
650 in the season. *Appl Environ Microbiol* 15:5207-5213.  
651 <https://doi.org/10.1128/AEM.00001-10>

652 Briand E, Escoffier N, Straub C, Sabart M, Quiblier C, Humbert JF (2009)  
653 Spatiotemporal changes in the genetic diversity of a bloom-forming *Microcystis*  
654 *aeruginosa* (cyanobacteria) population. *ISME J* 3:419-429.  
655 <https://doi.org/10.1038/ismej.2008.121>

656 Burford MA, Beardall J, Willis A, Orr PT, Magalhaes LM, Rangel LM et al (2016)  
657 Understanding the winning strategies used by the bloom-forming cyanobacterium  
658 *Cylindrospermopsis raciborskii*. *Harmful Algae* 54:44-53.  
659 <https://doi.org/10.1016/j.hal.2015.10.012>

660 Catherine A, Maloufi S, Mouillot D, Troussellier M, Bernard C (2016) Patterns and  
661 multi-scale drivers of phytoplankton species richness in temperate peri-urban lakes.  
662 *Sci Tot Environ* 559:74-83. <https://doi.org/10.1016/j.scitotenv.2016.03.179>

663 Catherine A, Maloufi S, Congestri R, Viaggiu E, Pilkaityte R (2017) SOP1:  
664 Cyanobacterial samples: Preservation, Enumeration, and biovolume measurement.  
665 In: *Handbook of Cyanobacterial Monitoring and Cyanotoxin Analysis*, Edition: 1st.  
666 Publisher: Wiley, Editors: Jussi Meriluoto, Lisa Spoof, Geoffrey Codd, pp 331–334.  
667 DOI:10.1002/9781119068761

668 Chaffin JD, Davis TW, Smith DJ, Baer MM, Dick GJ (2018) Interactions between  
669 nitrogen form, loading rate, and light intensity on *Microcystis* and *Planktothrix* growth  
670 and microcystin production. *Harmful Algae* 73:84–97.  
671 <https://doi.org/10.1016/j.hal.2018.02.001>

672 Chen Y, Xu J, Li Y, Han X 2011. Decline of sperm quality and testicular function in  
673 male mice during chronic low-dose exposure to microcystin-LR. *Reprod Toxicol*  
674 31:551-557. DOI:[10.1016/j.reprotox.2011.02.006](https://doi.org/10.1016/j.reprotox.2011.02.006)

675 Chia MA, Ameh I, George KC, Balogun EO, Akinyemi SA, Lorenzi AS (2022) Genetic  
676 diversity of microcystin producers (Cyanobacteria) and microcystin congeners in  
677 aquatic resources across Africa: A review paper. *Toxics* 10:772.  
678 <https://doi.org/10.3390/toxics10120772>

679 Chianu JN, Chianu JN, Mairura F (2012) Mineral fertilizers in the farming systems of  
680 sub-Saharan Africa. A review. *Agron Sustain Dev* 32:545-566.  
681 DOI:[10.1007/s13593-011-0050-0](https://doi.org/10.1007/s13593-011-0050-0)

682 Chorus I, Welker M (2021) *Toxic Cyanobacteria in Water*. Second edition. CRC Press,  
683 Boca Raton (FL), on behalf of the World Health Organization, Geneva, 859 p.  
684 <https://doi.org/10.1201/9781003081449>

685 Codd GA, Azevedo SMFO, Bagchi SN, et al (2005) Cyanonet : A global network for  
686 cyanobacterial bloom and toxin risk management. Technical document in  
687 *Hydrology*, n° 76, UNESCO Paris

688 Dalu T, Wasserman RJ (2018) Cyanobacteria dynamics in a small tropical reservoir:  
689 Understanding spatio-temporal variability and influence of environmental variables.  
690 *Sci Tot Environ* 643:835-841. <https://doi.org/10.1016/j.scitotenv.2018.06.256>

691 Decru E, Vreven E, Snoeks J (2012) A revision of the West African *Hepsetus*  
692 (Characiformes: Hepsetidae) with a description of *Hepsetus akawo* sp. nov. and a  
693 redescription of *Hepsetus odoe* (Bloch, 1794). *J Nat Hist* 46:1-23.  
694 <https://doi.org/10.1080/00222933.2011.622055>

695 Effebe KR, N'tain JY, Dabissi D., Diallo S, Tra ABZ et al (2017) Activities and uses of  
696 Aghien lagoon (South-East of Côte d'Ivoire). *J Water Res Prot* 9:11–19.  
697 <http://dx.doi.org/10.4236/jwarp.2017.91002>

698 Giani A, Taranu ZE, von Rückert G, Gregory-Eaves I (2020) Comparing key drivers of  
699 cyanobacteria biomass in temperate and tropical systems. *Harmful Algae*  
700 97:101859. <https://doi.org/10.1016/j.hal.2020.101859>

701 Guellati FZ, Touati H, Tambosco K, Quiblier C, Jumbert JF, Bensouilah M (2017)  
702 Unusual cohabitation and competition between *Planktothrix rubescens* and  
703 *Microcystis* sp. (cyanobacteria) in a subtropical reservoir (Hamman Debagh) located  
704 in Algeria. *Plos One* 12:e0183540. <https://doi.org/10.1371/journal.pone.0183540>

705 Haande S, Ballot A, Rohrlack T, Fastner J, Wiedner C, Edvardsen B (2007) Diversity  
706 of *Microcystis aeruginosa* (Chroococcales, Cyanobacteria) from East-African water  
707 bodies. *Arch Microbiol* 188:15-25. <https://doi.org/10.1007/s00203-007-0219-8>

708 Haande S, Rohrlack T, Ballot A, Roberg K, Skulberg R, Beck M, Wiedner C (2008)  
709 Genetic characterization of *Cylindrospermopsis raciborskii* (Nostocales,  
710 Cyanobacteria) isolates from Africa and Europe. Harmful Algae 7:692-701.  
711 <https://doi.org/10.1016/j.hal.2008.02.010>

712 Hammou HA, Latour D, Samoudi S et al (2018) Occurrence of the first toxic *Microcystis*  
713 bloom in a recent Moroccan reservoir. Wat resour 45:409-417.  
714 <https://doi.org/10.1134/S0097807818030028>

715 Horst GP, Sarnelle O, White JD, Hamilton SK, Kaul RB, Bressie JD (2014) Nitrogen  
716 availability increases the toxin quota of a harmful cyanobacterium, *Microcystis*  
717 *aeruginosa*. Wat Res 54:188–198. <https://doi.org/10.1016/j.watres.2014.01.063>

718 Ibelings BW, Bruning K, de Jonge J, Wolfstein K, Dionisio Pires LM, Postma J, Burger  
719 T (2005) Distribution of microcystins in a lake foodweb: No evidence for  
720 biomagnification. Microb Ecol 49:487-500. [https://doi.org/10.1007/s00248-004-](https://doi.org/10.1007/s00248-004-0014-x)  
721 [0014-x](https://doi.org/10.1007/s00248-004-0014-x)

722 Ibelings BW, Foss A, Chorus I (2021) Food. *In: Toxic Cyanobacteria in Water*. Chorus  
723 & Welker eds. Second edition. CRC Press, Boca Raton (FL), on behalf of the World  
724 Health Organization, Geneva, 368-388. <https://doi.org/10.1201/9781003081449>

725 Jawad LA, Gnohossou P, Toussou AG, Ligas A (2017) Morphometric relationships of  
726 *Coptodon guineensis* and *Sarotherodon melanotheron* (Perciformes, Cichlidae) in  
727 two Lakes of Benin (Western Africa). Turk J Fish Aqu Sci 17:217-221. DOI:  
728 [10.4194/1303-2712-v17\\_1\\_24](https://doi.org/10.4194/1303-2712-v17_1_24)

729 Jenny JP, Anneville O, Arnaud F, Baulaz Y, Bouffard D, Domaizon I al. (2020)  
730 Scientists' warning to humanity: Rapid degradation of the world's large lakes. J Gt  
731 Lakes Res 46:686-702. <https://doi.org/10.1016/j.jglr.2020.05.006>

732 Jia N, Yang Y, Yu G., Wang Y, Qiu P, Li H, Li R (2020) Interspecific competition  
733 reveals *Raphidiopsis raciborskii* as a more successful invader than *Microcystis*  
734 *aeruginosa*. Harmful Algae 97:101858. <https://doi.org/10.1016/j.hal.2020.101858>

735 Kide NG, Dia M, Yarba L, Kone Y, Khalil F, Salhi G et al (2015) Écologie trophique de  
736 *Sarotherodon melanotheron heudelotii* et de *Tilapia guineensis* (Perciformes :  
737 Cichlidae) du Parc National du banc d'Arguin, Mauritanie. Rev Ivoir Sci Technol  
738 25 :188-203.

739 Koffi SE, Koffi TK, Perrin JL, Seguis L, Guilliod M, Gone DL et al (2019) Hydrological  
740 and water quality assessment of the Aghien Lagoon hydrosystem (Abidjan, Côte

741 d'Ivoire). Hydrol Sci J 64:1893-1908. [https://doi.org/10.1080/02626667.2019.](https://doi.org/10.1080/02626667.2019.1672875)  
742 [1672875](https://doi.org/10.1080/02626667.2019.1672875)

743 Krienitz L, Ballot A, Wiegand C, Kotut K, Codd GA, Pflugmacher S (2002) Cyanotoxin-  
744 producing bloom of *Anabaena flos-aquae*, *Anabaena discoidea* and *Microcystis*  
745 *aeruginosa* (Cyanobacteria) in Nyanza Gulf of Lake Victoria, Kenya. J Appl Bot Food  
746 Qual 76:179-183.

747 Laplace-Treyture C, Feret T (2016) Performance of the Phytoplankton Index for Lakes  
748 (IPLAC): A multimetric phytoplankton index to assess the ecological status of water  
749 bodies in France. Ecol Indic 69:686–698. [https://doi.org/10.1016/j.ecolind.2016.](https://doi.org/10.1016/j.ecolind.2016.05.025)  
750 [05.025](https://doi.org/10.1016/j.ecolind.2016.05.025)

751 Maberly C, Chao A, Finlay BJ (2022) Seasonal patterns of phytoplankton taxon  
752 richness in lakes: Effects of temperature, turnover and abundance. Protist  
753 173:125925. <https://doi.org/10.1016/j.protis.2022.125925>

754 Manubolu M, Lee J, Riedl KM, Kua ZX, Collart LP, Ludsin SA (2018) Optimization of  
755 extraction methods for quantification of microcystin-LR and microcystin-RR in fish,  
756 vegetable, and soil matrices using UPLC–MS/MS. Harmful Algae 76 :47-57.  
757 <https://doi.org/10.1016/j.hal.2018.04.00>

758 Martins JC, Vasconcelos VM (2009) Microcystin dynamics in aquatic organisms. J  
759 Toxicol Environ Health B 12:65-82. <https://doi.org/10.1080/10937400802545151>

760 Matthews MW, Bernard S (2015) Eutrophication and cyanobacteria in South Africa's  
761 standing water bodies: A view from space. S Afr J Sci 111:77-84.  
762 <http://dx.doi.org/10.17159/sajs.2015/20140193>

763 Merel S, Villarin MC, Chung K, Snyder S (2013) Spatial and thematic distribution of  
764 research on cyanotoxins. Toxicon 76:118-131. [https://doi.org/10.1016/j.toxicon.](https://doi.org/10.1016/j.toxicon.2013.09.008)  
765 [2013.09.008](https://doi.org/10.1016/j.toxicon.2013.09.008)

766 Mitroi V, Deroubaix JF, Tall Y, Ahi KC, Humbert JF (2022) Rendre compte de la  
767 dégradation des milieux aquatiques. Le rôle des savoirs dans la mise en place des  
768 politiques de protection des ressources en eau en Afrique subsaharienne.  
769 Géocarrefour 96 :1. <https://doi.org/10.4000/geocarrefour.19353>

770 Mowe MAD, Mitrovic SM, Lim RP, Furey A, Yeo DCJ (2015) Tropical cyanobacterial  
771 blooms: a review of prevalence, problem taxa, toxins and influencing environmental  
772 factors. J Limnol 74:205-224. <https://doi.org/10.4081/jlimnol.2014.1005>

773 Ndlela LL, Oberholster PJ, Van Wyk JH, Cheng PH (2016) An overview of  
774 cyanobacterial bloom occurrences and research in Africa over the last decade.  
775 Harmful Algae 60:11-26. <https://doi.org/10.1016/j.hal.2016.10.001>

776 Neffling MR, Lance E, Meriluoto J (2010) Detection of free and covalently bound  
777 microcystins in animal tissues by liquid chromatography–tandem mass  
778 spectrometry. Environ Pollut 158:948-952.  
779 <https://doi.org/10.1016/j.envpol.2009.10.023>

780 [Nyakairu GWA, Nagawa CB, Mbabazy J \(2010\)](#) Assessment of cyanobacteria toxins  
781 in freshwater fish: A case study of Murchison Bay (Lake Victoria) and Lake Mburo,  
782 Uganda. [Toxicon 55:939-946](#). <https://doi.org/10.1016/j.toxicon.2009.07.024>

783 Nyenje PM, Foppen JW, Uhlenbrook S, Kulabako R, Muwanga A (2010)  
784 Eutrophication and nutrient release in urban areas of sub-Saharan Africa- a review.  
785 Sci Total Environ 408:447-455. <https://doi.org/10.1016/j.scitotenv.2009.10.020>

786 Oberholster PJ, Myburgh JG, Ashton PJ, Botha AM (2010) Responses of  
787 phytoplankton upon exposure to a mixture of acid mine drainage and high levels of  
788 nutrient pollution in Lake Loskop, South Africa. Ecotoxicol Environ Saf 73:326-335.  
789 <https://doi.org/10.1016/j.ecoenv.2009.08.011>

790 Olokotum M, Mitroi V, Troussellier M, Semyalo R, Bernard C, Montuelle B et al (2020)  
791 A review of the socioecological causes and consequences of cyanobacterial blooms  
792 in Lake Victoria. Harmful Algae 96:101829. [https://doi.org/10.1016/j.hal.2020.](https://doi.org/10.1016/j.hal.2020.101829)  
793 [101829](#)

794 Olokotum M, Troussellier M, Escalas A, Hamlaoui S, Okello W, Semyalo R et al (2021)  
795 High nutrient loading and climatic parameters influence the dominance and  
796 dissimilarity of toxigenic cyanobacteria in northern bays of Lake Victoria. J Gt Lakes  
797 Res 47:985–996. <https://doi.org/10.1016/j.jglr.2021.04.021>

798 Olokotum M, Humbert JF, Quiblier C, Okello W, Semyalo R, Troussellier M et al (2022)  
799 Toxins in Lake Victoria embayments and during water treatment. Toxins 14:664.  
800 <https://doi.org/10.3390/toxins14100664>

801 Poste AE, Hecky RE, Guildford SJ (2011) Evaluating microcystin exposure risk through  
802 fish consumption. Environ Sci Technol 45:5806-5811.  
803 <https://doi.org/10.1021/es200285c>

804 Roegner A, Sitoki LM, Weirich C, Corman J, Owage D, Umami M, et al (2020) Harmful  
805 algal blooms threaten the health of Peri-urban fisher communities: A case study in



806 Kisumu Bay, Lake Victoria, Kenya. *Expos Health* 12:835-848. doi: 10.1007/s12403-  
807 019-00342-8

808 Roegner A, Corman J, Sitoki LM, Kwena ZA, Ogari Z, Miruka JB, et al (2023) Impacts  
809 of algal blooms and microcystins in fish on small-scale fishers in Winam Gulf, Lake  
810 Victoria: implications for health and livelihood. *Ecol Soc* 28:49.  
811 <https://doi.org/10.5751/ES-13860-280149>

812 Rucker J, Wiedner C, Zippel P (1997) Factors controlling the dominance of *Planktothrix*  
813 *agardhii* and *Limnothrix redekei* in eutrophic shallow lakes. *Hydrobiologia* 342:107-  
814 155.

815 Sabart M, Pobel D, Latour D, Robin J, Salençon MJ, Humbert JF (2009)  
816 Spatiotemporal changes in the genetic diversity in French bloom-forming  
817 populations of the toxic cyanobacteria, *Microcystis aeruginosa*. *Environ Microbiol*  
818 *Rep* 1:263-272. <https://doi.org/10.1111/j.1758-2229.2009.00042.x>

819 Semyalo R, Rohrlack T, Naggawa C, Nyakairu GW (2010) Microcystin concentrations  
820 in Nile Tilapia (*Oreochromis niloticus*) caught from Murchison Bay, Lake Victoria  
821 and lake Mburo: Uganda. *Hydrobiologia* 638:235-244.  
822 <https://doi.org/10.1007/s10750-009-0047-8>

823 Soares MCS, Vidal LO, Roland F, Huszar VLM (2009) Cyanobacterial equilibrium  
824 phases in a small tropical impoundment. *J Plankt Res* 31:1331-1338.  
825 <https://doi.org/10.1093/plankt/fbp079>

826 Suominen S, Brauer VS, Rantala-Ylinen A, Sivonen K, Hiltunen T (2017) Competition  
827 between a toxic and a non-toxic *Microcystis* strain under constant and pulsed  
828 nitrogen and phosphorus supply. *Aquatic Ecol* 51:117-130. DOI:[10.1007/s10452-](https://doi.org/10.1007/s10452-016-9603-2)  
829 [016-9603-2](https://doi.org/10.1007/s10452-016-9603-2)

830 Svircev Z, Lalic D, Savic GB, Tokodi N, Backovic DD, Chen L et al (2019) Global  
831 geographical and historical overview of cyanotoxin distribution and cyanobacterial  
832 poisonings. *Arch Toxicol* 93:2429–2481. DOI:[10.1007/s00204-019-02524-4](https://doi.org/10.1007/s00204-019-02524-4).

833 United Nations (2017) World Population Prospects: The 2017 Revision, Key Findings  
834 and Advance Tables. Working Paper No. ESA/P/WP/248.

835 Utermöhl H (1958) Zur vervollkommnung der quantitativen phytoplankton methodik.  
836 *Mitt Int Theor Angew Limnol* 9:1–38.

837 Vico P, Bonilla S, Cremella B, Aubriot L, Iriarte A, Piccini C (2020) Biogeography of  
838 the cyanobacterium *Raphidiopsis (Cylindrospermopsis) raciborskii*: Integrating



839 genomics, phylogenetic and toxicity data. Mol Phylogenet Evol 148:106824.  
840 <https://doi.org/10.1016/j.ympev.2020.106824>

841 Wagner ND, Quach E, Buscho S, Ricciardelli A, Kannan A, Win Naung S et al (2021)  
842 Nitrogen form, concentration, and micronutrient availability affect microcystin  
843 production in cyanobacterial blooms. Harmful Algae 103:102002.  
844 <https://doi.org/10.1016/j.hal.2021.102002>

845 Wiedner C, Nixdorf B, Heinz R, Wirding B, Neumann U, Weckesser J (2002)  
846 Regulation of cyanobacteria and microcystin dynamics in polymictic shallow lakes.  
847 Arch Hydrobiol 155:383-400. DOI: [10.1127/archiv-hydrobiol/155/2002/383](https://doi.org/10.1127/archiv-hydrobiol/155/2002/383)

848 [Zhang D, Xie P, Liu Y, Qiu T \(2009\)](#) Transfer, distribution and bioaccumulation of  
849 microcystins in the aquatic food web in Lake Taihu, China, with potential risks to  
850 human health. Sci Tot Environ 407:2191-2199.  
851 <https://doi.org/10.1016/j.scitotenv.2008.12.039>

852  
853

#### 854 **Statements and Declarations**

855 **Ethical Approval:** Not applicable.

856 **Consent to Participate:** All the authors listed consent to participate.

857 **Consent to Publish:** The paper is submitted with the consent of all listed authors.

858

859 **Funding:** This work was supported by the Fonds Français pour l'Environnement  
860 Mondial (Program WaSAf). Authors M.K. Ahoutou and M. Olokotum have received a  
861 thesis grant in the framework of this research program.

862 **Competing interests:** The authors have no relevant financial or non-financial interests  
863 to disclose.

864 **Author Contributions:** EKY, MKA., CB., RYD, CQ, JFH and JKC contributed to the  
865 study conception and design. Material preparation, data collection and analysis were  
866 performed by EKY, MKA., MO, SH, EL, BM, CB, RYD, CQr, JFH and JKC. The first  
867 draft of the manuscript was written by EKY, JFH and JKC and all authors commented

868 on previous versions of the manuscript. All authors read and approved the final  
869 manuscript.

870

871 **Data availability.** All the data generated during this study are provided in  
872 supplementary tables

873