

Cooperative Excitation and Many-Body Interactions in a Cold Rydberg Gas

Matthieu Viteau, Paul Huillery, Mark Bason, Nicola Malossi, Donatella Ciampini, Oliver Morsch, Ennio Arimondo, Daniel Comparat, Pierre Pillet

► To cite this version:

Matthieu Viteau, Paul Huillery, Mark Bason, Nicola Malossi, Donatella Ciampini, et al.. Cooperative Excitation and Many-Body Interactions in a Cold Rydberg Gas. Physical Review Letters, 2012, 109 (5), pp.053002. 10.1103/PhysRevLett.109.053002. hal-04291815

HAL Id: hal-04291815 https://hal.science/hal-04291815v1

Submitted on 17 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cooperative Excitation and Many-Body Interactions in a Cold Rydberg Gas

Matthieu Viteau,¹ Paul Huillery,^{1,2,3} Mark G. Bason,¹ Nicola Malossi,^{1,4} Donatella Ciampini,^{1,3,4} Oliver Morsch,¹

Ennio Arimondo,^{1,3,4} Daniel Comparat,² and Pierre Pillet²

¹INO-CNR, Largo Pontecorvo 3, 56127 Pisa, Italy

²Laboratoire Aimé Cotton, Université Paris-Sud, ENS Cachan, Bâtiment 505, 91405 Orsay, France

³Dipartimento di Fisica E. Fermi, Università di Pisa, Largo Pontecorvo 3, 56127 Pisa, Italy

⁴CNISM UdR, Dipartimento di Fisica E. Fermi, Università di Pisa, Largo Pontecorvo 3, 56127 Pisa, Italy

(Received 3 April 2012; published 31 July 2012)

The dipole blockade of Rydberg excitations is a hallmark of the strong interactions between atoms in these high-lying quantum states [M. Saffman, T. G. Walker, and K. Mølmer, Rev. Mod. Phys. **82**, 2313 (2010); D. Comparat and P. Pillet, J. Opt. Soc. Am. B **27**, A208 (2010)]. One of the consequences of the dipole blockade is the suppression of fluctuations in the counting statistics of Rydberg excitations, of which some evidence has been found in previous experiments. Here we present experimental results on the dynamics and the counting statistics of Rydberg excitations of ultracold rubidium atoms both on and off resonance, which exhibit sub- and super-Poissonian counting statistics, respectively. We compare our results with numerical simulations using a novel theoretical model based on Dicke states of Rydberg atoms including dipole-dipole interactions, finding good agreement between experiment and theory.

DOI: 10.1103/PhysRevLett.109.053002

PACS numbers: 32.80.Ee, 03.67.Lx, 42.50.Ct

Atoms excited to high-lying quantum states, so-called Rydberg atoms, are highly polarizable and, therefore, can interact strongly with each other at large distances [1,2]. The study of ultracold gases has opened up new avenues for the investigation of strong atomic interactions. A wealth of potential applications is associated with the manipulation of ultracold atoms excited to Rydberg states interacting through strong van der Waals (vdW) or long-range dipolar interactions, ranging from studies of strongly correlated quantum systems to quantum information [3]. A key signature of interactions between Rydberg atoms is the suppression of fluctuations in the number of excitations due to the dipole blockade. While evidence of the dipole blockade has been observed in several experiments [4–7], prior studies were unable to demonstrate sub-Poissonian behavior with a statistically significant confidence level. Also, those studies were carried out for resonant excitation only, leaving open the question of how Rydberg excitations in the strongly interacting regime evolve for finite detuning from resonance.

The dipole blockade is a hallmark of the strong interactions between Rydberg atoms in an ultracold atomic gas. When an ensemble of atoms is irradiated by laser light resonant with a Rydberg excitation (either using a single laser or a multistep excitation scheme), due to the interaction between Rydberg atoms, the excitation of a particular atom can be suppressed by a neighboring one that is already in a Rydberg state. The radius of influence of an atom in this sense is called the blockade radius, which depends on the interaction strength and the linewidth of the Rydberg excitation. As a result of the dipole blockade the excitation dynamics of the atoms are strongly correlated, leading to a cutoff in the excitation when all the available blockade volumes in the sample have been exhausted [8]. Another way of describing the phenomenon is to view the atoms within a blockade volume as a "superatom," meaning that rather than individual atoms, collective states of several atoms are excited [9]. In the regime where the size of the sample is larger than the blockade radius, quantum correlations are expected to interconnect the whole sample. The description of this fully correlated regime is complex and the role of the atomic correlations on the atomic observables has recently received much attention [2]. In a simple mean field-type model for the interacting Rydberg atoms [10], each Rydberg atom experiences only an average interaction energy, which corresponds to neglecting all quantum correlations in the system. In order to reproduce the observed fluctuations and spatial correlations of the Rydberg excitations, however, more sophisticated approaches are needed [11–15].

While the full dynamics of Rydberg excitations in a cold gas of thousands of atoms is difficult to access experimentally, a number of experiments have observed signatures of the dipole blockade [9], such as the suppression of the excitation of a single atom in the presence of a nearby Rydberg atom [6,7], the spatial correlations between Rydberg excitations [16], and indications of a suppression of fluctuations in the counting statistics near a Förster resonance [5]. In this Letter, we present a comprehensive experimental and theoretical study of the counting statistics of Rydberg excitations of ultracold rubidium atoms in a magneto-optical trap (MOT).

As a full quantum mechanical treatment containing the states of all the atoms in the MOT is not feasible, the results of our experiments are analyzed using an original theoretical model based on the well-established Dicke model of quantum optics. The Dicke model, originally introduced for describing cooperative spontaneous emission or superradiance [17] and subradiance [18], is here modified by including the vdW interactions between the Dicke collective states (DCSs). Our approach leads to a manageable size of the basis set for the simulations, which is only on the order of the number of Rydberg excitations expected, that is, around 10-20 for our parameters, to be compared to our experimental numbers of ultracold atoms between tens and hundreds of thousands. The collective Dicke states contain the full statistical information about the collective Rydberg excitation, not only the average quantities, and therefore allow calculation of all the moments of the excitation statistics, in particular the measured variance of the Rydberg excitations. Furthermore, including a coupling between Dicke states of different symmetry leads to a partially non-Markovian dynamics, which results in better agreement with experimental findings.

In our experiments, rubidium atoms in a MOT are excited to Rydberg states with principal quantum number nbetween 50 and 80 using a two-photon scheme and detected, after field ionization, by a channeltron charge multiplier [19] (for details, see Supplemental Material [20]). The detuning of the resulting coherent excitation from the ground state to a highly excited Rydberg state can be varied by changing the frequency of the second step laser at around 1015 nm (the first step laser at 421 nm is detuned by around a gigahertz from an intermediate resonance in order to avoid population of that state). In the limit of vanishing interactions between the Rydberg atoms each excitation to a Rydberg state is independent of all the others, leading to a Poissonian excitation process. When the interactions are strong, however, the excitation processes are highly correlated and one expects sub-Poissonian counting statistics. These two regimes can be quantified through the Mandel Qparameter [21] defined as

$$Q = \frac{\langle (\Delta N)^2 \rangle}{\langle N \rangle} - 1, \tag{1}$$

where $\langle (\Delta N)^2 \rangle$ and $\langle N \rangle$ are the variance and mean, respectively, for the number N of Rydberg excitations. Experimentally, we determine Q by repeating an excitation-detection cycle for a fixed set of parameters between 50 and 100 times and extract the mean and the variance from that counting distribution. Clearly, for Poissonian processes Q = 0, while sub-Poissonian processes are characterized by a negative Mandel parameter Q < 0 (as, for example, in the case of the photon arrival statistics of squeezed light), with Q = -1 in the limit of complete suppression of fluctuations. When Q > 0 one speaks of a super-Poissonian counting statistics, which can be the result of technical noise or of enhanced fluctuations intrinsic to the dynamics of the system (see discussion below). For detection efficiencies $\eta < 100\%$, the observed value of the Mandel parameter is $Q_D = \eta Q$ [4,5]. This detected value is used in all experimental figures in this Letter. The Supplemental Material [20] reports our experimental verification that a Poissonian distribution $Q_D = 0$ is measured in the direct photoionization process, in which case no suppression of fluctuations is expected.

In Fig. 1, an example of the excitation dynamics to the $71D_{5/2}$ state for atoms on resonance is shown. Initially, a rapid increase in the average detected ion number is concomitant with a decrease in Q_D . After 0.5 μ s the growth in the detected ion number slows down while Q_D fluctuates around -0.4. Taking into account our detector efficiency of $\approx 40\%$ [19], the measurements indicate that on resonance a value close to Q = -1 is obtained. In the long time limit the effect of a varying Rabi frequency across the sample, due to both the distribution of atoms and the spatial



FIG. 1 (color online). Rydberg excitation dynamics in an ultracold gas. The mean number of detected ions $\langle N_D \rangle$ and Q_D are shown, in (a) and (b), respectively, as a function of pulse duration for the resonant excitation of the $71D_{5/2}$ state. The blue laser detuning from the intermediate $6P_{3/2} F = 3$ state is 1 GHz. The number of ground state atoms is 8×10^3 , the average density is 1.2×10^{10} cm⁻³, and the two-photon Rabi frequency is 40 kHz. The error bars are derived from 50 experimental realizations. The continuous lines are the prediction of our complete model, while the dashed line in (a) is the theoretical prediction of a model based on the Dicke symmetric states only. All quantities are rescaled by the detection efficiency.

laser profile and linewidth, cause a dephasing of the excitations, leading to a steady growth in the number of Rydberg atoms excited. In order to obtain reliable statistics with a reasonable number of experimental realizations, we performed numerical simulations to verify how quickly the mean and standard deviation converge for Poissonian and sub-Poissonian distributions. For example, to measure the Mandel parameter of a sub-Poissonian process with $Q \approx -0.5$ to within ± 0.1 requires around 50 realizations.

The resonant excitation dynamics can be easily visualized as a saturation of the maximum number of superatoms that fit into the volume of the MOT, leading to a limiting value for Q around -0.7 in a "close-packing" geometry of blockade spheres. Away from resonance, however, the picture is more complicated. Figure 2 shows the mean of the detected ion number and of Q_D as a function of the detuning from resonance for a fixed time. As expected, for zero detuning Q_D is highly negative (i.e., Q is close to -1). Away from resonance, however, the counting statistics quickly becomes super-Poissonian with Q_D as large as 2-3. One reason for this may be the effect of laser frequency fluctuations on the time scale of the data acquisition periods (around 30 s). On either slope a small shift in frequency corresponds to a large difference in the number of detected ions, leading to a large variance. Our theoretical model discussed below, however, indicates that the measured positive value of the Mandel parameter off resonance is also intrinsic to the excitation dynamics of the system.

For a better understanding of the physics underlying our experimental observations we need a theoretical treatment of our system that gives access to the full counting statistics, and hence to the Q parameter, both on and off resonance. Although recent theoretical work has addressed the expected counting statistics for on-resonant excitation of collective Rydberg states [11,12], so far there have been no predictions for the off-resonant case. We describe our system as an ensemble of N_0 indistinguishable atoms, for which the laser excitation results in a single wave function superposition of a two level system composed of the ground state $|g\rangle$ and the Rydberg state $|R\rangle$, driven by a laser with Rabi frequency Ω and detuning δ . Our approach is based on the DCS [17,22], characterized by the cooperative number r defining the symmetry and the number Nof Rydberg excited atoms, $|r, N\rangle$. As the laser excitation preserves the DCS symmetry, only states with the same symmetry number r are excited when starting from an ultracold atomic sample described by a totally symmetric DCS. When vdW interactions are taken into account, states with the same symmetry and those with different symmetries can be coupled.

Assuming for the initial atomic wave function the fully symmetrical nondegenerate DCS $|r = N_0/2, N\rangle = |N\rangle_s$ and neglecting the vdW coupling between symmetrical and antisymmetrical DCS, we write the atomic wave



FIG. 2 (color online). Detuning dependence of collective Rydberg excitations. The mean number of detected ions $\langle N_D \rangle$ and Q_D are shown for a range of frequencies around the twophoton excitation of state $71D_{1/2}$ in (a) and (b), respectively. The $\langle N_D \rangle$ data are well fitted by a top truncated Lorentzian line shape, except in a frequency range around the maximum. The continuous and dashed lines are the predictions of our complete model, with the calculated average coupling strength and 100 times that coupling strength, respectively. The black dot-dashed line simulates the effect of the laser jitter (assumed to be 0.5 MHz) on Q_D , assuming that Q = 0 independently of the detuning, based on the data of (a). The blue laser detuning from the intermediate $6P_{3/2} F = 3$ state is 1 GHz. The number of ground state atoms is 5×10^4 , the average density is 4×10^{10} cm⁻³, the two-photon Rabi frequency is 40 kHz, and the excitation pulse duration is 0.3 µs.

function as $|\Psi(t)\rangle = \sum_{N=0}^{N_0} a_N(t) |N\rangle_s$. The time evolution of the amplitudes a_N is given by the following coupled equations [23] (with $\hbar = 1$):

$$\dot{a}_{N} = iN\delta a_{N} - i\frac{\Omega}{2} \bigg[\sqrt{(N_{0} - N)(N+1)} a_{N+1} + \sqrt{N(N_{0} - N+1)} a_{N-1} \bigg].$$
(2)

The vdW interaction $W_{ss}^N = W_0 N(N-1)/[2(N_0-1)]$ introduces an additional term

$$-iW_{ss}^{N}a_{N} \tag{3}$$

to the equation of motion for a_N , where W_0 is the sum over all the interaction energies between pairs of Rydberg atoms [10]. The number of excitations in the Dicke ladder is equal to the number of Rydberg excitations, that is, a few tens in our experiment. The solution of the above equations allows one to calculate the cooperative mean and variance values, and all higher moments of the number of Rydberg excitations. It leads to the Rydberg excitation depicted by the dashed line of Fig. 1, in reasonable agreement with the experimental observations for short times, but not in the long time limit. In addition, it predicts asymmetric profiles for the laser detuning dependence in the blockade regime, not matching previous observations [24] and our data on the detected ions vs the laser detuning shown in Fig. 2(a).

The problem is that the vdW interaction couples the fully symmetrical DCSs to the remaining DCSs within a fixed N subspace, which has a large degeneracy given by the binomial coefficient $C_{N_0}^N$. The most appropriate orthonormal basis within that subspace is composed of the symmetrical DCSs $|N\rangle_s$ and the superposition of nonsymmetrical states $\{|N, q\rangle_{ns}\}$, with $1 \le q \le C_{N_0}^N - 1$, for which the vdW interaction matrix is diagonal. This original basis construction, detailed in the Supplemental Material [20], is far more general than the treatment of the vdW interactions provided we know the basis diagonalizing the coupling. Within this basis the vdW matrix contains the terms $W_{ss}^N, W_{qq}^N, W_{sq}^N$ (see Supplemental Material [20]).

By writing the enlarged atomic wave function as a superposition of symmetric and nonsymmetric states, we obtain that the a_N time evolution of Eq. (3) is completed by terms describing the vdW coupling to the amplitude b_N^q of the nonsymmetrical state. As a first approximation, we neglect the laser excitation of the amplitudes b_N^q because of their large degeneracy and the resulting weak occupation for each of them. After the elimination of the equations for b_N^q , the temporal evolution of the symmetric states determined by the coupling with a bath, made up of the remaining collective states, therefore contains both Markovian and non-Markovian memory function contributions. For our experimental conditions, this quite general result can be simplified when calculating the memory functions (see Supplemental Material [20]). In the end, for each number N of Rydberg excitations, the evolution is described by a system of two equations, one for a_N and one for a single level c_N defined as the superposition of all asymmetric DCSs of the fixed N subspace. Thus the bath is modeled by a number of states equal to that of the symmetric DCSs and the total required basis set required for the simulations is approximately twice the number of Rydberg excitations. The bath is Markovian and produces a shift of the symmetric DCS energy precisely compensating the vdW eigenenergy of Eq. (3). This compensation leads to symmetric excitation profiles as in Fig. 2. The vdW

interaction for the symmetric DCS finally replaces Eq. (3) by the following one:

$$-iW_{ss}^{N}c_{N}.$$
 (4)

For the nonsymmetrical DCS

$$\dot{c}_N = i\delta N c_N - iW^N_{ss} a_N. \tag{5}$$

Notice that within the above equations the vdW interaction acts as a coherent coupling between symmetrical and antisymmetrical states, leading to a variety of coherent evolution features, for instance the revivals appearing as narrow peaks in the Q dependence on parameters as the laser detuning (see Fig. 2) or the laser pulse duration. However, for each N manifold the revival conditions are slightly different, and for a large number of Rydberg excitations, the revivals are washed out. The last step of this theoretical analysis is the introduction of the laser excitation into evolution of c_N . We treat it perturbatively, with a limitation on the temporal evolution imposed by the spectral linewidth $\Delta \omega$ of the excitation laser. The final equations are reported within the Supplemental Material [20].

Typical results of numerical simulations using the Dicke model are shown in Figs. 1 and 2 together with our experimental data. Clearly, the salient features of the experiment are reproduced well by our model, both in the dynamics and in the dependence on the detuning. In particular, the off-resonant behavior of the Q factor, which in the experimental data features two symmetric peaks, is also visible in the simulation. The variation in the interatomic interaction curves (see Supplemental Material [20]) is reflected in the uncertainty in our determination of the coupling strength for the theoretical curves in Fig. 2. Perfect quantitative agreement between experiment and theory is not expected as the Dicke model does not, at this time, include the spatially varying atomic density of the MOT, Doppler broadening, the spatial inhomogeneity of the excitation lasers, the full manifold of sublevels of the Rydberg states, and the effects of magnetic and possible residual electric fields, all of which might contribute to the larger linewidth observed experimentally. In particular, the dip in the Rydberg number around zero detuning seen in Fig. 2(a) could be smoothed out when taking into account the density variation over the cloud. In fact, Fig. 3 shows that the variation of Q with the detuning depends strongly on the density (or, equivalently, on the C_6 coefficient that describes the van der Waals interaction between Rydberg atoms). As the density is increased, the value of O on resonance goes from zero to a limiting value of around -0.7, whereas the position and height of the off-resonant peaks symmetric around zero detuning increase steadily. In order to get better agreement, one could either perform experiments with close to uniform atomic densities (e.g., using flattop excitation beams much smaller than the atomic sample) or develop a theoretical description that allows for the nonuniform density to be included.



FIG. 3 (color online). Predictions for the detuning dependence of collective Rydberg excitations. (a) Theoretical dependence of the Q parameter on the laser detuning for a range of atomic densities (in cm⁻³): 1×10^{10} (thin blue line), 4×10^{10} (dashed green line), 10×10^{10} (dot-dashed orange line), and 40×10^{10} (thick red line), respectively. (b) Dependence on the density of the on-resonant Q (blue line) and the value of Q (dot-dashed red line) and position (black dashed line, right-hand scale) of the first off-resonant peaks.

In summary, we have shown clear evidence of sub-Poissonian counting statistics of Rydberg excitations in an ultracold atomic sample, which are a clear signature of many-body quantum correlations in such a system. We have characterized both the dynamics and the dependence on the detuning of the Q parameter and reproduced the main features of both using a novel model. In future studies, it will be interesting to look directly at the spatial correlations between Rydberg excitations [16], at the counting statistics for the antiblockade regime of [25], and at other, more indirect, signatures of Rydberg-Rydberg interactions, for example in electromagnetically induced transparency experiments [26].

This work was supported by the EU through Grants No. 225187-NAMEQUAM and No. 265031-ITN-COHERENCE and the collaboration between University of Pisa and University of Paris-Sud 11. The authors thank M. Allegrini, R. Côté, P. Grangier, T. Pohl, M. Saffman, J. Stanojevic, and M. Weidemüller for useful discussions and J. Radogostowicz for assistance.

- M. Saffman, T.G. Walker, and K. Mølmer, Rev. Mod. Phys. 82, 2313 (2010).
- [2] D. Comparat and P. Pillet, J. Opt. Soc. Am. B **27**, A208 (2010).
- [3] I. Lesanovsky, Physics 4, 71 (2011).
- [4] T. Cubel Liebisch, A. Reinhard, P.R. Berman, and G. Raithel, Phys. Rev. Lett. 95, 253002 (2005); 98, 109903 (E) (2007).
- [5] A. Reinhard, K. C. Younge, and G. Raithel, Phys. Rev. A 78, 060702(R) (2008).
- [6] A. Gaëtan, Y. Miroshnychenko, T. Wilk, A. Chotia, M. Viteau, D. Comparat, P. Pillet, A. Browaeys, and P. Grangier, Nature Phys. 5, 115 (2009).
- [7] E. Urban, T. A. Johnson, T. Henage, L. Isenhower, D. D. Yavuz, T. G.Walker, and M. Saffman, Nature Phys. 5, 110 (2009).
- [8] J. V. Hernández and F. Robicheaux, J. Phys. B 41, 195301 (2008).
- [9] R. Heidemann, U. Raitzsch, V. Bendkowsky, B. Butscher, R. Löw, L. Santos, and T. Pfau, Phys. Rev. Lett. 99, 163601 (2007).
- [10] D. Tong, S. M. Farooqi, J. Stanojevic, S. Krishnan, Y. P. Zhang, R. Côté, E. E. Eyler, and P. L. Gould, Phys. Rev. Lett. 93, 063001 (2004).
- [11] C. Ates, T. Pohl, T. Pattard, and J. M. Rost, J. Phys. B 39, L233 (2006).
- [12] S. Wüster, C. Ates, T. Pohl, P. Deuar, J. F. Corney, and J. M. Rost, Phys. Rev. A 81, 023406 (2010).
- [13] J. Stanojevic and R. Côté, Phys. Rev. A 81, 053406 (2010).
- [14] I. Lesanovsky, Phys. Rev. Lett. 106, 025301 (2011).
- [15] J. V. Hernandez and F. Robicheaux, J. Phys. B **39**, 4883 (2006).
- [16] A. Schwarzkopf, R. E. Sapiro, and G. Raithel, Phys. Rev. Lett. 107, 103001 (2011).
- [17] R. H. Dicke, Phys. Rev. 93, 99 (1954).
- [18] D. Pavolini, A. Crubellier, P. Pillet, L. Cabaret, and S. Liberman, Phys. Rev. Lett. 54, 1917 (1985).
- [19] M. Viteau, J. Radogostowicz, A. Chotia, M. G. Bason, N. Malossi, F. Fuso, D. Ciampini, O. Morsch, I. I. Ryabtsev, and E. Arimondo, J. Phys. B 43, 155301 (2010); 44, 079802(E) (2011).
- [20] See Supplemental Material at http://link.aps.org/ supplemental/10.1103/PhysRevLett.109.053002 for details of the theoretical model.
- [21] L. Mandel, Phys. Rev. Lett. 49, 136 (1982).
- [22] L. Mandel and E. Wolf, *Optical Coherence and Quantum Optics* (Cambridge University Press, Cambridge, England, 1995).
- [23] S. Haroche and J. M. Raimond, *Exploring the Quantum: Atoms, Cavities and Photons* (Oxford University, New York, 2006), p. 220.
- [24] T. Amthor, M. Reetz-Lamour, S. Westermann, J. Denskat, and M. Weidemüller, Phys. Rev. Lett. 98, 023004 (2007).
- [25] T. Amthor, C. Giese, C. S. Hofmann, and M. Weidemüller, Phys. Rev. Lett. **104**, 013001 (2010).
- [26] J. D. Pritchard, D. Maxwell, A. Gauguet, K. J. Weatherill, M. P. A. Jones, and C. S. Adams, Phys. Rev. Lett. 105, 193603 (2010).