



# A synthesis of SNAPO-CO<sub>2</sub> ocean total alkalinity and total dissolved inorganic carbon measurements from 1993 to 2022

Nicolas Metzl, Jonathan Fin, Claire Lo Monaco, Guillaume Bourdin, Samir Alliouane, Jacqueline Boutin, Claude Mignon, Yann Bozec, David Antoine, Pascal Conan, et al.

## ► To cite this version:

Nicolas Metzl, Jonathan Fin, Claire Lo Monaco, Guillaume Bourdin, Samir Alliouane, et al.. A synthesis of SNAPO-CO<sub>2</sub> ocean total alkalinity and total dissolved inorganic carbon measurements from 1993 to 2022. *Earth System Science Data*, In press, 10.5194/essd-2023-308 . hal-04292340v2

HAL Id: hal-04292340

<https://hal.science/hal-04292340v2>

Submitted on 17 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# A synthesis of ocean total alkalinity and dissolved inorganic carbon measurements from 1993 to 2022: the SNAPO-CO<sub>2</sub>-v1 dataset

Nicolas Metzl, Jonathan Fin, Claire Lo Monaco, Claude Mignon, Samir Alliouane, David Antoine, Guillaume Bourdin, Jacqueline Boutin, Yann Bozec, Pascal Conan, et al.

## ► To cite this version:

Nicolas Metzl, Jonathan Fin, Claire Lo Monaco, Claude Mignon, Samir Alliouane, et al.. A synthesis of ocean total alkalinity and dissolved inorganic carbon measurements from 1993 to 2022: the SNAPO-CO<sub>2</sub>-v1 dataset. *Earth System Science Data*, In press, 10.5194/essd-2023-308 . hal-04292340

HAL Id: hal-04292340

<https://hal.science/hal-04292340>

Submitted on 17 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 A synthesis of SNAPO-CO<sub>2</sub> ocean total alkalinity and total dissolved 2 inorganic carbon measurements from 1993 to 2022

3 Nicolas Metzl<sup>1</sup>, Jonathan Fin<sup>1,2</sup>, Claire Lo Monaco<sup>1</sup>, Claude Mignon<sup>1</sup>, Samir Alliouane<sup>3</sup>,  
4 David Antoine<sup>3,4</sup>, Guillaume Bourdin<sup>5</sup>, Jacqueline Boutin<sup>1</sup>, Yann Bozec<sup>6</sup>, Pascal Conan<sup>7,8</sup>,  
5 Laurent Coppola<sup>3,8</sup>, Frédéric Diaz<sup>\*</sup>, Eric Douville<sup>9</sup>, Xavier Durrieu de Madron<sup>10</sup>, Jean-Pierre  
6 Gattuso<sup>3,11</sup>, Frédéric Gazeau<sup>3</sup>, Melek Golbol<sup>8,12</sup>, Bruno Lansard<sup>9</sup>, Dominique Lefèvre<sup>13</sup>,  
7 Nathalie Lefèvre<sup>1</sup>, Fabien Lombard<sup>3,14</sup>, Férial Louanchi<sup>15</sup>, Liliane Merlivat<sup>1</sup>, Léa Olivier<sup>1,16</sup>,  
8 Anne Petrenko<sup>13</sup>, Sébastien Petton<sup>17</sup>, Mireille Pujo-Pay<sup>7</sup>, Christophe Rabouille<sup>9</sup>, Gilles  
9 Reverdin<sup>1</sup>, Céline Ridame<sup>1</sup>, Aline Tribollet<sup>1</sup>, Vincenzo Vellucci<sup>8,12</sup>, Thibaut Wagener<sup>13</sup>, Cathy  
10 Wimart-Rousseau<sup>13,18</sup>

11  
12 <sup>1</sup> Laboratoire LOCEAN/IPSL, Sorbonne Université-CNRS-IRD-MNHN, Paris, 75005, France

13 <sup>2</sup> OSU Ecce Terra, Sorbonne Université-CNRS, Paris, 75005, France

14 <sup>3</sup> Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefranche, LOV, F-06230 Villefranche-sur-  
15 Mer, France

16 <sup>4</sup> Remote Sensing and Satellite Research Group, School of Earth and Planetary Sciences, Curtin University,  
17 Perth WA 6845, Australia

18 <sup>5</sup> School of Marine Sciences, University of Maine, Orono, USA

19 <sup>6</sup> Station Biologique de Roscoff, UMR 7144 – EDYCO-CHIMAR, Roscoff, France

20 <sup>7</sup> Sorbonne Université, CNRS, Laboratoire d'Océanographie Microbienne, LOMIC, F-66650 Banyuls-sur-Mer,  
21 France

22 <sup>8</sup> Sorbonne Université, CNRS, OSU Station Marines, STAMAR, Paris, F-75006, France

23 <sup>9</sup> Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, UMR 8212 CEA- CNRS-UVSQ,  
24 Université Paris-Saclay, 91191 Gif-sur-Yvette, France

25 <sup>10</sup> CEFREM, CNRS-Université de Perpignan Via Domitia, 52 Avenue Paul Alduy, 66860 Perpignan, France

26 <sup>11</sup> Institute for Sustainable Development and International Relations, Sciences Po, 27 rue Saint Guillaume, F-  
27 75007 Paris, France

28 <sup>12</sup> Sorbonne Université, CNRS, Institut de la Mer de Villefranche, IMEV, Villefranche-sur-Mer, F-06230, France

29 <sup>13</sup> Aix Marseille Univ, Université de Toulon, CNRS, IRD, MIO, Marseille, France

30 <sup>14</sup> Research Federation for the study of Global Ocean Systems Ecology and Evolution, FR2022/Tara GOSEE,  
31 75000, Paris, France.

32 <sup>15</sup> CVRM: Laboratoire de Conservation et de Valorisation des Ressources Marines, Ecole Nationale Supérieure  
33 des Sciences de la Mer et de l'Aménagement du Littoral (ENSSMAL), Station de recherche de Sidi Fredj,  
34 Algeria

35 <sup>16</sup> Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

36 <sup>17</sup> Ifremer, Univ Brest, CNRS, IRD, LEMAR, F-29840 Argenton, France

37 <sup>18</sup> GEOMAR Helmholtz Centre for Ocean Research Kiel, 24105 Kiel, Germany

38 \* Passed away 14/3/2021

39 Correspondence to: Nicolas Metzl (nicolas.metzl@locean.ipsl.fr)

40 **Abstract.** Total alkalinity ( $A_T$ ) and dissolved inorganic carbon ( $C_T$ ) in the oceans are important properties  
to  
41 understand the ocean carbon cycle and its link with global change (ocean carbon sinks and sources, ocean  
42 acidification) and ultimately find carbon based solutions or mitigation procedures (marine carbon removal). We  
43 present a database of more than 44 400  $A_T$  and  $C_T$  observations along with basic ancillary data (time and  
space

44 location, depth, temperature and salinity) in various ocean regions obtained since 1993 mainly in the frame of  
45 French projects. This includes both surface and water columns data acquired in open oceans, coastal zones and in

46 the Mediterranean Sea and either from time-series or punctual cruises. Most  $A_T$  and  $C_T$  data in this  
synthesis

47 were measured from discrete samples using the same closed-cell potentiometric titration calibrated with Certified

48 Reference Material, with an overall accuracy of  $\pm 4 \mu\text{mol kg}^{-1}$  for both  $A_T$  and  $C_T$ . The data are provided in two  
49 separate datasets for the global ocean, and for the Mediterranean Sea (<https://doi.org/10.17882/95414>, Metzl et  
50 al., 2023) that offers a direct use for regional or global purposes, e.g.  $A_T$ /Salinity relationships, long-term  $C_T$   
51 estimates, constraint and validation of diagnostics  $C_T$  and  $A_T$  reconstructed fields or ocean carbon and coupled  
52 climate/carbon models simulations, as well as data derived from Biogeochemical-Argo (BGC-Argo) floats.  
53 When associated with other properties, these data can also be used to calculate pH, fugacity of  $\text{CO}_2$  ( $f\text{CO}_2$ ) and  
54 other carbon system properties to derive ocean acidification rates or air-sea  $\text{CO}_2$  fluxes.

55

## 56 **1 Introduction**

57

58 Since 1750, humans activities have added 700 ( $\pm 75$ ) PgC anthropogenic carbon dioxide to the  
59 atmosphere by burning fossil fuels, producing cement and changing land use (Friedlingstein et al., 2022) driving  
60 up the atmospheric carbon dioxide ( $\text{CO}_2$ ) level and leading to unequivocal global change. The ocean plays a  
61 major role in reducing the impact of climate change by absorbing more than 90% of the excess heat in the  
62 climate system (Cheng et al., 2020; von Schuckmann et al., 2020, 2023; IPCC, 2022) and about 25% of human  
63 released  $\text{CO}_2$  (Friedlingstein et al., 2022). However, the oceanic  $\text{CO}_2$  uptake changes the chemistry of seawater  
64 reducing its buffering capacity (Revelle and Suess, 1957; Jiang et al., 2023a) and leading to a process known as  
65 ocean acidification with potential impacts on marine organisms (Fabry et al., 2008; Doney et al., 2009, 2020;  
66 Gattuso et al., 2015). With atmospheric  $\text{CO}_2$  concentrations, surface ocean temperature and ocean heat content,  
67 sea-level, sea-ice and glaciers, the ocean acidification (decrease of pH) is now recognized by the World  
68 Meteorological Organization as one of the 7 key properties for global climate indicators (WMO, 2018). In the  
69 frame of the 2030 Agenda, the United Nations established a set of Sustainable Development Goals (SDG; United  
70 Nations, 2020), including a goal dedicated to the ocean (SDG 14, "Life below water") which calls to "conserve  
71 and sustainably use the oceans, seas and marine resources for sustainable development". Ocean acidification is  
72 specifically referred in the SDG indicator 14.3.1 coordinated at the Intergovernmental Oceanographic  
73 Commission (IOC) of UNESCO. Observing the carbonate system in the oceans and marginal seas and  
74 understanding how this system changes over time is thus highly relevant not only to quantify the global ocean  
75 carbon budget, the anthropogenic  $\text{CO}_2$  inventories or ocean acidification rates, but also to understand and  
76 simulate the processes that govern the complex  $\text{CO}_2$  cycle in the ocean and to better predict the future evolution  
77 of climate and global changes (Eyring et al., 2016; Kwiatkowski et al., 2020; Jiang et al., 2023a).

78 The number and quality of ocean  $f\text{CO}_2$ ,  $A_T$ ,  $C_T$  and pH measurements have increased substantially over  
79 the past few decades. Quality-controlled observations are now regularly assembled in global data syntheses such  
80 as SOCAT (Surface Ocean  $\text{CO}_2$  Atlas, Pfeil et al., 2013; Bakker et al., 2014, 2016) and GLODAP (Global Ocean  
81 Data Analysis Project, Key et al., 2004; Olsen et al., 2016, 2019, 2020; Lauvset et al., 2021, 2022). These  
82 datasets allow evaluation of properties trends in the global ocean, including the change of the ocean  $\text{CO}_2$  sink  
83 (e.g. Wanninkhof et al., 2013; Friedlingstein et al., 2022; Watson et al., 2020), anthropogenic  $\text{CO}_2$  inventories  
84 (e.g. Sabine et al., 2004; Khatiwala et al., 2013; Gruber et al., 2019) and ocean acidification (Lauvset et al.,  
85 2015, 2020; Jiang et al., 2019; Feely et al., 2023; Ma et al., 2023). Thanks to publicly available consistent and  
86 quality controlled databases new methods were recently developed (Carter et al., 2016; Sauzède et al., 2017;  
87 Bittig et al., 2018) to reproduce  $A_T$  and  $C_T$  distributions from other properties like temperature, salinity and  
88 oxygen more often observed in the water column especially from autonomous floats (Claustre et al., 2020;  
89 Mignot et al., 2023). These methods (named CANYON-B and CONTENT, Bittig et al., 2018) are now also used

90 to help decisions on GLODAP data quality control or to fill in observational gaps (Olsen et al., 2019, 2020;  
91 Tanhua et al., 2019, 2021). The GLODAP data products were also successfully used to construct new global  
92 ocean  $A_T$  and  $C_T$  climatological monthly fields in surface and water column using neural network method (e.g.  
93 Broullón et al., 2019, 2020).

94 Following pioneer works that produced various global-ocean climatologies of the sea-surface carbonate  
95 system (Millero et al., 1998; Lee et al., 2000, 2006; Takahashi et al., 2002, 2009, 2014; Sasse et al., 2013; Jiang  
96 et al., 2019), the coupling of  $fCO_2$  data (from SOCAT) and  $A_T$  data (from GLODAP) now enables reconstruction  
97 of the full carbonate system in the surface ocean at monthly scale to investigate temporal trends at decadal scale  
98 (e.g. Gregor and Gruber, 2021; Keppler et al., 2023).

99 International projects such as SOCAT and GLODAP offer important way to synthetize ocean carbon  
100 data. In these projects, each observation is quality controlled offering to users high quality observations for  
101 regional or global analysis, either for processes analysis or to constraint or validate of ocean and coupled  
102 climate/carbon models (CMIP6, e.g. Lerner et al., 2021). SOCAT is a publicly available synthesis product  
103 initiated in 2007 (Metzl et al., 2007) for quality-controlled, surface ocean  $fCO_2$  (fugacity of carbon dioxide)  
104 observations made by the international marine carbon research community (Bakker et al., 2016). The first  
105 SOCAT version was released in 2011 (Pfeil et al., 2013; Sabine et al., 2013), followed by 6 SOCAT versions  
106 (Bakker et al., 2014, 2016). The last version in 2023 includes more than 40 million  $fCO_2$  data with accuracy  
107 better than 5  $\mu\text{atm}$  (Bakker et al., 2023). One important product from SOCAT is the use of data to estimate  
108 global air-sea  $CO_2$  fluxes based on reconstructed  $pCO_2$  fields (e.g. Surface Ocean  $pCO_2$  Mapping  
109 Intercomparison, SOCOM, Rödenbeck et al., 2015). Since 2015, these results are included each year for the  
110 global carbon budget (Le Quere et al., 2015; Friedlingstein et al., 2022).

111 On the other hand, following WOCE/JGOFS era in the 90s when almost all observations were started to  
112 be synthetized in a specific recommended format (Joyce and Corry, 1994), GLODAP focusses on water-column  
113 carbon observations (and other properties). Following the original GLODAP data product (Key et al., 2004), the  
114 project accumulated many new quality controlled observations. One important achievement from GLODAP is  
115 the use of data to estimate the anthropogenic  $CO_2$  inventory or its change over decades (Sabine et al., 2004;  
116 Gruber et al., 2019). Both products, SOCAT and GLODAP, are relevant tools to detect oceanic acidification  
117 rates (Lauvset et al., 2015; Jiang et al., 2019; Feely et al., 2023; Ma et al., 2023).

118 Although these projects include many international ocean observations there are ocean  $CO_2$  related  
119 observations all around the world (published or not published), such as total alkalinity and dissolved inorganic  
120 carbon, that are not included in SOCAT or GLODAP. This is because SOCAT accepts and controls only  $fCO_2$   
121 data, whereas GLODAP includes and controls water-columns data mainly from WOCE/GO-SHIP/CLIVAR  
122 cruises. It should be noticed that many ocean carbon observations in various formats can be also found in  
123 dedicated database such as NCEI/OCADS (former CDIAC-Ocean, Jiang et al., 2023b,  
124 <https://www.ncei.noaa.gov/products/ocean-carbon-acidification-data-system>), PANGAEA  
125 (<https://www.pangaea.de/>) or Seanoe (<https://www.seanoe.org/>). In this context it is recommended to progress in  
126 data synthesis of the ocean carbon observations that would offer new high quality products for the community  
127 (e.g. for GOA-ON, [www.goa-on.org](http://www.goa-on.org), IOC/SDG 14.1.3, <https://oa.iode.org/>, Tilbrook et al., 2019).

128 In this work, we present a synthesis of more than 44 400  $A_T$  and  $C_T$  observations obtained over the  
129 1993-2022 period during various cruises or at time-series stations mainly supported by French projects. This  
130 dataset merges observations measured with the same instruments thus being analytically coherent. Most of the  
131 data have accuracy better than  $\pm 4 \mu\text{mol kg}^{-1}$ , i.e. between the climate ( $\pm 2 \mu\text{mol kg}^{-1}$ ) and weather ( $\pm 10 \mu\text{mol kg}^{-1}$ )

132 <sup>1)</sup> goals (Newton et al, 2015; Bockmon and Dickson, 2015). Hereafter this dataset will be cited as SNAPO-CO2-  
133 V1. We describe the data assemblage and associated quality control and discuss some potential uses of this  
134 dataset.

135

136 **2 Data collections**

137


138 The time series projects and research cruises from which data were collated are listed in Table 1 with  
139 references in the Supplementary file (Table S1) and the sampling locations displayed in Figure 1. Sampling was  
140 performed either from CTD-Rosette casts (Niskin bottles) or from the ship's seawater supply (intake at about 5m  
141 depth depending the ship and swell). Samples collected in 500 mL borosilicate glass bottles were poisoned with  
142 100 to 300  $\mu\text{L}$  of  $\text{HgCl}_2$  depending on the cruises, closed with greased stoppers (Apiezon®) and held tight using  
143 elastic band following the SOP protocol (Dickson et al., 2007). Some samples were also collected in 500 mL  
144 bottles closed with screw caps. After completion of each cruise, discrete samples were returned back to the  
145 LOCEAN laboratory (Paris, France) and stored in a dark room at 4 °C before analysis generally within 2-3  
146 months after sampling (sometimes within a week). Some samples were also measured for specific processes  
147 studies on benthic corals (e.g. Maier et al., 2012; McCulloch et al., 2012) or for mesocosm and culture  
148 experiments but the data are not included in this synthesis as they do not represent natural ocean state (e.g.  
149 addition of Sahara dust during the DUNE project, Ridame et al., 2014).

150 As opposed to  $p\text{CO}_2$ , surface  $A_{\text{T}}$  or  $C_{\text{T}}$  observations are generally obtained from discrete sampling  
151 (measured onboard or onshore). Few cruises offer sea-surface semi-continuous  $A_{\text{T}}$  or  $C_{\text{T}}$  observations (e.g. Metzl  
152 et al., 2006) but new instrumental developments (Seelmann et al., 2020) now enable  $A_{\text{T}}$  measurements on Ship of  
153 Opportunity Program lines (SOOP). In addition to discrete samples analyzed for various projects conducted  
154 mainly in the North Atlantic, Tropical Atlantic, Tropical Pacific, Mediterranean sea and coastal regions (Table  
155 1), we complemented this synthesis with  $A_{\text{T}}$  and  $C_{\text{T}}$  surface observations obtained in the Indian and Southern  
156 oceans during the OISO cruises in 1998-2018 (Metzl et al., 2006; Leseurre et al., 2022; data also available at  
157 NCEI/OCADS: [www.nodc.noaa.gov/ocads/oceans/VOS\\_Program/OISO.html](http://www.nodc.noaa.gov/ocads/oceans/VOS_Program/OISO.html)) and the recent CLIM-EPARSES  
158 cruise conducted in the Mozambique Channel in April 2019 (Lo Monaco et al., 2020, 2021). For OISO cruises  
159 the water-column observations are part of the CARINA (CARbon IN the Atlantic) and GLODAP synthesis  
160 products (Lo Monaco et al., 2010; Olsen et al., 2016, 2019, 2020) and not included here. Excepted when  
161 specified, all data in this synthesis were obtained using the same technique used either in laboratory or at sea (for  
162 OISO 1998-2018 and CLIM-EPARSES 2019 cruises).

163

164 **Table 1:** List of cruises in the SNAPO-CO2-v1 dataset. This is organized by region: Global Ocean and coastal  
 165 zones, and Mediterranean Sea (MedSea). See Tables S1, S2, S4 and S4 in the Supplementary Material for a list  
 166 of laboratories, of CRMs used, for references and for DOI of cruises. Nb = the number of data for each cruise or  
 167 time-series. \* indicates the measurements at sea (surface underway)

170 Cruise/Project	Start	End	Region	Sampling	Nb
173 AWIPEV	2015	2021	Arctic	Surface and sub-surface	195
174 SURATLANT+RREX	1993	2017	North Atlantic	Surface	2832
175 OVIDE	2006	2018	North Atlantic	Surface, Water Column	397
176 STRASSE	2012		North Atlantic	Water Column	205
177 EUREC4A-OA	2020		North Atlantic	Surface, Water Column	135
178 PROTEUS	2010		North Atlantic	Water Column	27
179 CHANNEL	2012	2015	English Channel	Surface	696
180 SOMLIT-Brest	2008	2019	Coastal North Atl	Surface	1174
181 SOMLIT-Roscoff	2009	2019	Coastal North Atl	Surface and 60m	801
182 ECOSCOPA	2017	2019	Coastal North Atl	Surface	67
183 PENZE	2011	2020	River Brittany	Surface and sub-surface	148
184 AULNE	2009	2010	River Brittany	Surface	27
185 ELORN	2009	2009	River Brittany	Surface	28
186 BIOZAIRES	2003	2004	Trop Atlantic	Water Column	87
187 EGEE	2005	2007	Trop Atlantic	Surface	199
188 PIRATA-FR	2009	2017	Trop Atlantic	Surface, Water Column	513
189 PLUMAND	2007		Trop Atlantic	Surface	38
190 OUTPACE	2015		Trop Pacific	Water Column	240
191 PANDORA	2012		Solomon Sea	Water Column	178
192 TARA-Pacific	2016	2018	Trop Pac North Atl	Surface and sub-surface	325
193 TARA-Ocean	2009	2012	Global Ocean	Surface + 400m	123
194 TARA-Microbiome	2021	2022	Atlantic	Surface, Water Column	216
195 ACE	2016	2017	Southern Ocean	Surface, Water Column	135
196 MOBYDICK	2019		Southern Ocean	Water Column	64
197 CLIM-EPARSES *	2019		Indian	Surface	790
198 OISO *	1998	2018	South Indian	Surface	24950
199					
200 DYFAMED	1998	2017	MedSea	Water Column	2118
201 BOUSSOLE	2014	2019	MedSea	Surface + 10m	172
202 SOMLIT-PointB	2007	2015	MedSea Coastal	Surface + 50m	2397
203 ANTARES	2010	2016	MedSea	Water Column	502
204 MOLA	2010	2013	MedSea Coastal	Water Column	66
205 SOLEMIO	2016	2018	MedSea Coastal	Water Column	212
206 MOOSE-GE	2010	2019	MedSea	Water Column	1847
207 LATEX	2010		MedSea	Water Column	51
208 CARBORHONE	2011	2012	MedSea	Water Column	706
209 CASCADE	2011		MedSea	Water Column	218
210 DEWEX	2013		MedSea	Water Column	367
211 SOMBA	2014	2014	MedSea	Water Column	203
212 AMOR-BFLUX	2015		MedSea Coastal	Water Column	6
213 PEACETIME	2017	2017	MedSea	Water Column	233
214 PERLE	2018	2021	MedSea	Water Column	805



**Figure 1:** Locations of  $A_T$  and  $C_T$  data (1993-2022) in the Global Ocean and the Western Mediterranean Sea (white box, insert) in the SNAPO-CO<sub>2</sub>-v1 dataset. Figure produced with ODV (Schlitzer, 2018).

### 3 Method, accuracy, repeatability, inter-comparison and quality control


#### 3.1 Method and accuracy

Since 2003, the discrete samples returned back at SNAPO-CO<sub>2</sub> Service facilities (LOCEAN, Paris), were analyzed simultaneously for  $A_T$  and  $C_T$  by potentiometric titration using a closed cell (Edmond, 1970; Goyet et al., 1991). The same technique was used at sea for surface water underway measurements during OISO and CLIM-EPARSES cruises (indicated by \* in Table 1). For two time series, the dataset also includes measurements obtained before 2000 using other techniques: the DYFAMED time-series observations measured between in 1998 and 2000 in the Mediterranean Sea (Copin-Montégut and Bégoovic, 2002; Coppola et al., 2020; Lange et al., 2023) and the SURATLANT time-series values acquired from 1993 to 1997 in the North Atlantic subpolar gyre (Reverdin et al., 2018). We also include  $A_T$  data in the river Penzé (Brittany) in 2019-2020 (Yann Bozec, SBR/Roscoff, pers. comm.).

In the late 1980s the so-called “JGOFS-IOC Advisory Panel on Ocean CO<sub>2</sub>” recommended the need for standard analysis protocols and for developing Certified Reference Materials (CRMs) for inorganic carbon measurements (Poisson et al., 1990; UNESCO, 1990, 1991). The CRMs were provided to international laboratories by Pr. A. Dickson (Scripps Institution of Oceanography, San Diego, USA), starting in 1990 for  $C_T$  and 1996 for  $A_T$ , respectively. These CRMs were thus always available to us and used to calibrate the measurements (CRM Batch numbers used for each cruise are listed in Supplementary file (Table S2). The concentrations of CRMs we used vary between 2193 and 2426  $\mu\text{mol kg}^{-1}$  for  $A_T$  and between 1968 and 2115  $\mu\text{mol kg}^{-1}$  for  $C_T$  corresponding to the range of concentrations observed in open ocean water. The CRMs accuracy, as indicated in the certificate for each Batch, is around  $\pm 0.5 \mu\text{mol kg}^{-1}$  for both  $A_T$  and  $C_T$  ([www.nodc.noaa.gov/ocads/oceans/Dickson\\_CRM/batches.html](http://www.nodc.noaa.gov/ocads/oceans/Dickson_CRM/batches.html)).

Results of analyses performed on 965 CRM bottles (different Batches) in 2013-2023 are presented in Figure 2. The standard-deviations of the differences of measurements were on average around  $\pm 3.5 \mu\text{mol kg}^{-1}$  for

both  $A_T$  and  $C_T$ . For unknown reasons, the differences were occasionally up to  $10\text{--}15 \mu\text{mol kg}^{-1}$  ( $0.8\%$  of the data, Figure S2). These few CRM measurements were discarded for the data processing. On average, and excluding some outliers, standard-deviations of the differences for 1090 CRM analyses were  $\pm 2.71 \mu\text{mol kg}^{-1}$  for  $A_T$  and  $\pm 2.86 \mu\text{mol kg}^{-1}$  for  $C_T$ , respectively. We did not detect any specific signal for CRM analyses (e.g., larger uncertainty depending on the Batch number or temporal drifts during analyses, Figure 2) but for some cruises the accuracy based on CRMs could be slightly better than  $3 \mu\text{mol kg}^{-1}$  (e.g. Marrec et al., 2014; Touratier et al., 2016; Ganachaud et al., 2017; Wimart-Rousseau et al., 2020a).




**Figure 2:**  $A_T$  (a) and  $C_T$  (b) analyses for different CRM Batches measured in 2013–2023. For these 965 analyses the mean and standard-deviations of the differences with the CRM reference were  $-0.1 (\pm 3.4) \mu\text{mol kg}^{-1}$  for  $A_T$  and  $0.1 (\pm 3.7) \mu\text{mol kg}^{-1}$  for  $C_T$ .

### 3.2 Repeatability

For some projects, duplicates have been regularly sampled (SOMLIT-Point-B, SOMLIT-BREST, BOUSSOLE/DYFAMED) or replicate bottles sampled at selected depths at fixed stations during the cruises (e.g. OUTPACE-2015, SOMBA-2014). Results of  $A_T$  and  $C_T$  repeatability are synthetized in Table 2 and Figure 3 shows example of regular duplicates from the times-series SOMLIT-Point-B in the coastal Mediterranean Sea, SOMLIT-Brest in the Bay of Brest, coastal Iroise Sea and BOUSSOLE/DYFAMED in the Ligurian Sea. For the 26 OISO cruises conducted between 1998 and 2018 and the CLIM-EPARSES cruise in April 2019, the repeatability was evaluated from duplicate analyses (within 20 minutes time) of continuous sea surface underway sampling at the same location (when the ship was stopped). Similarly to what was found for the CRM measurements (Figure S2), differences in duplicates are occasionally higher than  $10\text{--}15 \mu\text{mol kg}^{-1}$  (Figure 3) but most of the duplicates for all projects are within  $0$  to  $3 \mu\text{mol kg}^{-1}$ . Based on the CRM analyses and replicates for

305 different projects, different regions and different periods, we estimated  $A_T$  and  $C_T$  data to be consistent to better  
306 than  $4 \mu\text{mol kg}^{-1}$ .



347 **Figure 3:** Results of duplicate  $A_T$  and  $C_T$  analyses from the time-series (a) SOMLIT-Point-B in the coastal  
348 Mediterranean Sea (Kapsenberg et al., 2017), (b) SOMLIT-BREST in the Bay of Brest, coastal Iroise Sea (Salt  
349 et al., 2016 and unpublished) and (c) BOUSSOLE/DYFAMED in the Ligurian Sea (Merlivat et al., 2018; Golbol  
350 et al., 2020). The plots show differences in duplicates for both  $A_T$  (filled circles) and  $C_T$  (open circles). Standard-  
351 deviations of these duplicates are listed in Table 2.

353  
 354 **Table 2:** Repeatability of  $A_T$  and  $C_T$  analyses for cruises with duplicate analysis. The results are expressed as the  
 355 standard-deviations (Std) of the analysis of replicated samples. Nb = the number of replicates for each Time-  
 356 series or Cruise (a). For the 26 OISO cruises (1998-2018) and for simplicity we list the mean repeatability  
 357 obtained for all cruises.

---

360 Cruise	Nb	Std $A_T$ $\mu\text{mol kg}^{-1}$	Std $C_T$ $\mu\text{mol kg}^{-1}$	Reference
361 OUTPACE	12	3.64	3.68	Wagener et al. (2018)
362 SOMBA	13	2.00	3.30	Keraghel et al. (2020)
363 SOMLIT-Point-B	786	2.63	3.10	Kapsenberg et al. (2017)
364 SOMLIT-Brest	446	3.34	3.67	Salt et al. (2016) + unpub
365 BOUSSOLE	48	3.47	4.02	Merlivat et al. (2018); Golbol et al. (2020)
366 CLIM-EPARSES	122	2.20	2.30	Lo Monaco et al. (2020, 2021)
367 OISO 1998-2018	1162	2.06	2.28	Metzl et al. (2006) and (b)

---

372 (a) See Figure 3 for the results of regular duplicates for 3 time-series (SOMLIT-Point-B, SOMLIT-BREST, and  
 373 BOUSSOLE).

374 (b) Metadata and data available at [www.nodc.noaa.gov/ocads/oceans/VOS\\_Program/OISO.html](http://www.nodc.noaa.gov/ocads/oceans/VOS_Program/OISO.html)

### 377 3.3 Inter-comparisons

378       Inter-comparisons of measurements performed with different technics help to evaluate the quality of the  
 379 data and detect potential biases when merging the data in the same region obtained by different laboratories at  
 380 different periods. This is especially important to interpret long-term trends of  $A_T$  and  $C_T$  as well as for  $p\text{CO}_2$  and  
 381 pH calculated with  $A_T/C_T$  pairs. For ocean acidification studies, this also refers to the “climate goal” for which an  
 382 accuracy for  $A_T$  and  $C_T$  better than  $\pm 2 \mu\text{mol kg}^{-1}$  is needed (Newton et al., 2015; Tilbrook et al., 2019). For the  
 383 projects in this data synthesis, inter-laboratory comparisons were performed occasionally and summarized  
 384 below.

385

#### 386 3.3.1 CHANNEL project

388       As part of the time-series CHANNEL (2012-2015) in the Western English Channel, Marrec et al.  
 389 (2014) analyzed surface samples collected bi-monthly in 2011-2013.  $A_T$  analyses were performed with a TA-  
 390 ALK-2 system (Appolo SciTech.) while  $C_T$  measurements were acquired with an AIRICA system (Marianda  
 391 Inc.). Based on CRM analyses (Batch #92) the accuracy was estimated  $\pm 3 \mu\text{mol kg}^{-1}$  for  $A_T$  and  $\pm 1.5 \mu\text{mol kg}^{-1}$   
 392 for  $C_T$  (Marrec et al., 2014). When comparing with the samples measured at LOCEAN/Paris for the year 2012,  
 393 Marrec et al. (2014) concluded that between the two methods the concentrations were within  $\pm 2 \mu\text{mol kg}^{-1}$  and  
 394  $\pm 3 \mu\text{mol kg}^{-1}$  for  $A_T$  and  $C_T$  respectively. This is close to the “climate goal” offering confident results for long-  
 395 term trend analysis of the carbonate system in this region.

396

#### 397 3.3.2 SURATLANT project

400       In the frame of the SURATLANT project in the subpolar North Atlantic gyre, some samples collected  
 401 at the same time (in 2005, 2006, 2010, 2015, and 2016) were also analyzed onshore for  $A_T$  and/or  $C_T$  by other

402 laboratories using different technics (e.g. coulometric method) and the results summarized by Reverdin et al.  
403 (2018). For  $C_T$ , the mean (and STD) differences between LOCEAN values and from 4 other laboratories range  
404 between -0.7 ( $\pm 4.6$ ) and -6.5 ( $\pm 3.4$ )  $\mu\text{mol kg}^{-1}$  depending on the cruise. For  $A_T$  the mean differences with 2 other  
405 laboratories range from -0.6 ( $\pm 4.1$ )  $\mu\text{mol kg}^{-1}$  to +2.3 ( $\pm 4.8$ )  $\mu\text{mol kg}^{-1}$ .


406

### 407 3.3.3 OVIDE project

408

409 During OVIDE cruises conducted since 2002 in the North Atlantic along a section from Greenland to  
410 Portugal (Lherminier et al., 2007; Mercier et al., 2015) samples have been taken (since 2006) to complement, for  
411 summer, the SURATLANT time-series in the North Atlantic subpolar gyre (NASPG). The OVIDE samples at  
412 the surface and along the water-column at a few stations were measured back at LOCEAN for  $A_T$  and  $C_T$  (Metzl  
413 et al., 2018). This enables us to compare our data with the measurements performed on-board by the IIM group  
414 in Vigo/Spain (e.g. Pérez et al., 2010, 2013, 2018; Vazquez-Rodriguez et al., 2012). The OVIDE data have been  
415 regularly quality controlled in CARINA and GLODAP data products (Velo et al., 2009; Key et al., 2010; Olsen  
416 et al., 2016, 2019, 2020). The results of inter-comparisons are gathered in Table 3. For OVIDE in 2006 we  
417 identified (for unknown reason) a large difference between our original  $A_T$  values compared to the  $A_T$  data  
418 qualified in GLODAP and we thus corrected our  $A_T$  data by +7.2  $\mu\text{mol kg}^{-1}$ . However, no correction was applied  
419 for  $C_T$ . For other OVIDE cruises, differences for  $A_T$  range between -4.5 ( $\pm 4.11$ )  $\mu\text{mol kg}^{-1}$  and -0.05 ( $\pm 3.43$ )  
420  $\mu\text{mol kg}^{-1}$  depending on the cruise (i.e.  $A_T$  measured at LOCEAN was always slightly lower than onboard  
421 measurements). For  $C_T$ , we compared our measurements onshore with  $C_T$  values calculated with  $A_T$  and pH  
422 measured onboard. Most of the mean  $C_T$  differences are slightly positive (i.e.  $C_T$  measured at LOCEAN was  
423 always higher, except for 2010). Taking into account all errors associated with the sampling, the transport of  
424 samples, the instrumentations, the data processing, or the calculations for  $C_T$  using  $A_T/\text{pH}$  pairs (around 8.8  $\mu\text{mol}$   
425  $\text{kg}^{-1}$ , Orr et al, 2018), the comparisons between LOCEAN and IIM data for OVIDE cruises are deemed  
426 acceptable and large differences for both  $A_T$  and  $C_T$  ( $> 4 \mu\text{mol kg}^{-1}$ ) are far from being systematic (Table 3). The  
427 data from SURATLANT and OVIDE can then be merged to complete the time-series in the NASPG in summer  
428 and to better describe the seasonality of the oceanic carbonate system. For example, in 2010, when the North  
429 Atlantic Oscillation (NAO) was strongly negative, the SURATLANT data showed a rapid decrease of  $C_T$   
430 concentrations in the NASPG between early June and August (Figure 4), with  $C_T$  concentrations in August much  
431 lower than other years (Racapé et al., 2014). This leads to a rapid drop in  $f\text{CO}_2$  in 2009-2010, such that the  
432 NASPG was a strong  $\text{CO}_2$  sink (Leseurre et al., 2020). The winter-to-summer seasonal decrease of  $C_T$  in 2010 in  
433 the north NASPG was on average -77  $\mu\text{mol kg}^{-1}$  (Figure 4) much larger than in the climatology (range -50 to -55  
434  $\mu\text{mol kg}^{-1}$ , Takahashi et al., 2014; Reverdin et al., 2018). The OVIDE data in late June 2010 and SURATLANT  
435 in August 2010 confirmed this signal that was linked to a pronounced primary productivity in that period (Figure  
436 4, Henson et al., 2013; Racapé et al., 2014; Mc Kinley et al., 2018). Notice that for this period no  $f\text{CO}_2$   
437 observations were available in July-September 2010 in SOCAT data product and the  $A_T/C_T$  data presented here  
438 could be used to calculate  $f\text{CO}_2$  to complement the  $f\text{CO}_2$  dataset in this region like was done for other periods  
439 (Mc Kinley et al., 2011).

440



**Figure 4:** (a) Time-series of  $C_T$  concentrations ( $\mu\text{mol} \cdot \text{kg}^{-1}$ ) for 2009-2011 in surface waters in the North Atlantic subpolar gyre (zone  $59^{\circ}\text{N}-33^{\circ}\text{W}$ ) based on SURATLANT (open triangles) and OVIDE-2010 (grey circles) data. In 2009, SURATLANT data were available in February, June, September and December, while in 2010 data available in March, June, August and December and in 2011 data only available for March and December. The OVIDE data in late June 2010 completed the temporal cycle and confirmed the strong seasonal signal and low  $C_T$  concentrations in summer 2010 not seen in 2009 (or in 2011 as there is no data in summer). The mean observations for each period describe the  $C_T$  seasonal cycles in 2009 and 2010 (Black squares, dashed line). The monthly surface chlorophyll-a concentrations (Chl-a,  $\text{mg} \cdot \text{m}^{-3}$ ) averaged in the same region based on MODIS are also shown (Green dots and line) highlighting the high productivity during the summer 2010. Chl-a monthly data extracted from MODIS (Giovanni/NASA, last access 3/5/19). (b): Time-series of  $f\text{CO}_2$  ( $\mu\text{atm}$ ) for the same cruises (same symbols) calculated with  $A_T/C_T$  and using the K1, K2 constants from Lueker et al (2000). Mean SST ( $^{\circ}\text{C}$ ) indicated (red triangles). In June 2010 oceanic  $f\text{CO}_2$  decreased by 53  $\mu\text{atm}$  in 2 weeks.

**Table 3:** Comparisons of  $A_T$  and  $C_T$  samples measured back at LOCEAN with measurements onboard by IIM Laboratory (F. Pérez, Vigo, Spain) for OVIDE cruises in the North Atlantic. Nb= Number of samples. ND= No Data. The results listed indicate the mean and standard-deviations of the differences (LOCEAN-IIM). For  $A_T$ , IIM values were measured on-board. For  $C_T$ , IIM values were calculated from  $A_T$  and pH both measured onboard. The IIM data were quality controlled and here taken from the GLODAP data products (Olsen et al, 2016, 2019).

Cruise Year	Nb $A_T$	$A_T$ (LOCEAN) – $A_T$ (IIM) $\mu\text{mol kg}^{-1}$	Nb $C_T$	$C_T$ (LOCEAN) – $C_T$ (IIM) $\mu\text{mol kg}^{-1}$
OVIDE-2006	14	-2.0 ( $\pm 5.9$ ) (*)	14	1.1 ( $\pm 2.5$ )
OVIDE-2008	29	-4.5 ( $\pm 4.1$ )	29	3.8 ( $\pm 3.1$ )
OVIDE-2010	41	-2.0 ( $\pm 2.3$ )	41	-2.4 ( $\pm 3.3$ )
OVIDE-2012	37	-0.1 ( $\pm 8.8$ )	ND	ND
GEOVIDE-2014	57	-0.1 ( $\pm 3.4$ )	54	2.4 ( $\pm 7.9$ )

(\*) for the OVIDE 2006 cruise original difference for  $A_T$  was -9.0 ( $\pm 5.8$ )  $\mu\text{mol kg}^{-1}$  and LOCEAN  $A_T$  data were corrected by +7.2  $\mu\text{mol kg}^{-1}$  based on the mean concentrations in deep layers. No corrections were applied for  $A_T$  and  $C_T$  for other cruises.

497 **3.3.4 PENZE river**

498

499 The comparisons described above concern the open ocean region with  $A_T$  and  $C_T$  concentrations in a  
 500 range of concentrations close to the CRM references (used by the different laboratories). Another example of  
 501 comparison is presented here for samples obtained along a river and thus for waters with low salinity and  $A_T$   
 502 concentrations (river Penzé in North Brittany). In 2019-2020,  $A_T$  was measured at SBR laboratory (Station  
 503 Biologique de Roscoff) by a potentiometric method (using a Titrino-847 plus Metrohm) calibrated with CRM  
 504 (Batch #131) for a final accuracy of  $\pm 2.1 \mu\text{mol kg}^{-1}$  (Gac et al., 2020). Although the samples were measured  
 505 with different technics the  $A_T$ /Salinity relationships are very coherent for both datasets (Figure 5). The  
 506 regressions for each period are for  $A_T$  (in  $\mu\text{mol kg}^{-1}$ ):

507


508 In 2011 (78 samples):  $A_T = 51.525 (\pm 0.944) S + 583.95 (\pm 19.94) (r^2 = 0.975)$

509 In 2019-2020 (70 samples):  $A_T = 54.022 (\pm 1.018) S + 450.23 (\pm 31.53) (r^2 = 0.976)$

510

511 Therefore we added the  $A_T$  data measured in 2019-2020 to complete the synthesis for this location (river Penzé).

512



527 **Figure 5:** Total alkalinity ( $A_T$ ) versus salinity for samples measured in 2011 and 2019 in the river Penzé, North  
 528 Brittany (Gac et al., 2020).  $A_T$  samples were measured at LOCEAN in 2011 (open circles, dashed-line) and at  
 529 SBR laboratory (Roscoff) in 2019 (filled circles, black line).

530

531 **3.4 Quality control assigned flags**

532

533 Identifying each data with an appropriate flag is very convenient for selecting the data (good,  
 534 questionable or bad). Here we used 4 flags for each property (flags 2 = good, 3= questionable, 4=bad, and 9= no  
 535 data) following the WOCE program and used in other data products such as SOCAT (Bakker et al., 2016) or  
 536 GLODAP (Olsen et al., 2016, 2019, 2020; Lauvset et al., 2021). During the data-processing, we first assigned a  
 537 flag for each  $A_T$  and  $C_T$  data based on the standard error in the calculation of  $A_T$  and  $C_T$  concentrations (non-  
 538 linear regression, Dickson et al. 2007). By default, if the standard-deviation on the regression is  $> 1 \mu\text{mol kg}^{-1}$ ,  
 539 we assigned a flag 3 (questionable) although the data could be acceptable and then used for interpretations. Flag  
 540 3 was also assigned when salinity was doubtful or when differences of duplicates were large (e.g.  $\pm 20 \mu\text{mol kg}^{-1}$ ). Flags 4 (bad or certainly bad) were assigned when clear anomalies were detected for unknown reason (e.g. a  
 541

sample probably not fixed with  $\text{HgCl}_2$ ). A secondary quality control was performed by the PIs of each project based on data inspection, duplicates,  $A_T$ /Salinity relationship, or the mean observations in deep layers where large variability in  $A_T$  and  $C_T$  is unlikely to occur from year to year. An example presents all data from the MOOSE-GE cruises conducted in 2010-2019 in the Mediterranean Sea (Coppola et al., 2020; Testor et al., 2010) where clear outliers have been identified (Figure S3). For the 10 MOOSE-GE cruises and a total of 1847  $A_T$  and  $C_T$  analyses, 26 were identified flagged as bad (flag 4), 139 for  $A_T$  and 141 for  $C_T$  listed as questionable (flag 3) and 1682 for  $A_T$  and 1680 for  $C_T$  considered as good data (flag 2, i.e. more than 90%). Similar control was performed for each project.

The synthesis of various cruises in the same region and period also offers verification and secondary control of the data. For example, several cruises were conducted in the Mediterranean Sea in 2014 (MOOSE-GE, SOMBA, ANTARES and DYFAMED). The mean values of  $C_T$  and  $A_T$  in the deep layers ( $> 1800\text{m}$ ) for each cruise confirmed the coherence of the data (Table 4). This enabled to merge the different datasets for interpretations of the temporal trends and processes driving the  $\text{CO}_2$  cycle (Coppola et al., 2019, 2020; Ulles et al., 2023) or to train and validate a regional neural network to reconstruct the carbonate system (e.g. CANYON-MED, Fourrier et al., 2020, 2022).

**Table 4:** Mean observations in the deep layers ( $> 1800\text{m}$ ) of the Western Mediterranean Sea for different cruises conducted in 2014. Results in deep layers ( $> 1800\text{m}$ ) for the DEWEX cruise in 2013 and the PEACETIME cruise in 2017 in the same region are also listed. N- $A_T$  and N- $C_T$  are  $A_T$  and  $C_T$  normalized at salinity = 38. Nb = number of data (with flag 2). Standard-deviations are in brackets. References for these cruises are listed in Supplementary Material.

Cruise	Period	Nb	Pot. Temp (°C)	Salinity (PSU)	N- $A_T$ ( $\mu\text{mol kg}^{-1}$ )	N- $C_T$ ( $\mu\text{mol kg}^{-1}$ )
All cruises	Feb/Dec-2014	76	12.905 (0.007)	38.486 (0.005)	2562.9 (5.3)	2303.7 (4.7)
ANTARES	Feb/Nov-2014	14	12.913 (0.004)	38.488 (0.006)	2564.0 (3.8)	2301.9 (3.5)
DYFAMED	Mar/Dec-2014	9	12.905 (0.0016)	38.487 (0.004)	2560.1 (5.0)	2304.3 (6.8)
MOOSE-GE	Jul-2014	21	12.909 (0.004)	38.487 (0.005)	2565.6 (4.6)	2303.5 (4.1)
SOMBA	Aug/Sep-2014	32	12.899 (0.005)	38.483 (0.005)	2561.5 (5.6)	2304.6 (4.8)
DEWEX	Feb/Apr-2013	44	12.903 (0.010)	38.588 (0.006)	2556.0 (4.3)	2294.0 (5.7)
PEACETIME	May/Jun-2017	7	12.904 (0.002)	38.486 (0.003)	2567.2 (10.6)	2308.1 (8.9)

The total number of data for the Global Ocean and the Mediterranean Sea are gathered in Table 5 with corresponding flags for each property. Overall, the synthesis includes more than 94% of good data for both  $A_T$


595 and  $C_T$ . About 5% are questionable and 2% are likely bad. Overall, we believe that all data (with flag 2) in this  
 596 synthesis have an accuracy better than 4  $\mu\text{mol kg}^{-1}$  for both  $A_T$  and  $C_T$ , the same as for quality-controlled data in  
 597 GLODAP (Olsen et al., 2020; Lauvset et al., 2021). The uncertainty ranges between the “Climate goal” (2  $\mu\text{mol}$   
 598  $\text{kg}^{-1}$ ) and the “Weather Goal” (10  $\mu\text{mol kg}^{-1}$ ) for ocean acidification studies (Newton et al., 2015; Tilbrook et al.,  
 599 2019). This accuracy is also relevant to validate or constraint data-based methods that reconstruct  $A_T$  and  $C_T$   
 600 fields with an error of around 10-15  $\mu\text{mol kg}^{-1}$  for both properties (Bittig et al., 2018; Broullón et al., 2019, 2020;  
 601 Fourrier et al., 2020; Chau et al., 2023).

602  
 603 **Table 5:** Number of Temperature, Salinity,  $A_T$  and  $C_T$  data in the synthesis identified for flags 2, 3, 4, 9. The  
 604 data are given for the full data-set Global Ocean and for the Mediterranean Sea. Last column is the percentage of  
 605 flag 2 (Good).

	Flag 2	Flag 3	Flag 4	Flag 9	% Flag 2
<hr/>					
Global Ocean					
Temperature	43538	410	0	478	99.07
Salinity	44033	319	2	71	99.28
$A_T$	39331	2144	1165	1787	92.24
$C_T$	39921	2091	1148	1279	92.50
<hr/>					
Mediterranean Sea					
Temperature	9843	1	0	65	99.99
Salinity	9879	8	2	20	99.99
$A_T$	8853	425	411	220	91.37
$C_T$	8854	451	389	211	91.33

### 628 3.5 Using $A_T$ and $C_T$ to calculate $f\text{CO}_2$ and pH and compare with $f\text{CO}_2$ and pH measurements

629  
 630 For some projects, the  $A_T$  and  $C_T$  data presented in this synthesis were used to calibrate or validate in  
 631 situ  $f\text{CO}_2$  sensors (Bozec et al., 2011; Marrec et al., 2014; Merlivat et al., 2018). The  $A_T$  and  $C_T$  data were also  
 632 used to calculate  $f\text{CO}_2$  and to derive associated air-sea  $\text{CO}_2$  fluxes, especially during periods when no direct  $f\text{CO}_2$   
 633 measurements were available (e.g. in the North Atlantic, Figure 4, Watson et al., 2009; Mc Kinley et al., 2011).  
 634 For example, Marrec et al. (2014) successfully used the calculated  $p\text{CO}_2$  (with  $A_T/C_T$  pairs) to adjust the drift of  
 635 the  $p\text{CO}_2$  data recorded with a Contros-HydroC/ $\text{CO}_2$  FT sensor mounted on a FerryBox for regularly sampling  
 636 the Western English Channel (CHANNEL project). Here we show the results for the period 2012-2014 (Figure  
 637 6). In this region the total alkalinity is relatively constant over time; the average of  $A_T$  for 528 samples at  
 638 different seasons and years is  $2334.4 (\pm 7.2) \mu\text{mol kg}^{-1}$ . On the opposite, the  $C_T$  concentrations show distinctive  
 639 seasonality, with higher concentrations in winter and lower in summer when biological activity is pronounced  
 640 (Marrec et al., 2013, 2014; Kitidis et al., 2019). This controls the seasonal  $p\text{CO}_2$  distribution revealed each year  
 641 in both measured and calculated  $p\text{CO}_2$  (Figure 6). For 528 co-located samples the mean difference between  
 642 calculated and measured  $p\text{CO}_2$  is  $-1.9 (\pm 11.9) \mu\text{atm}$  with no distinct differences depending on the season and  
 643 year.  
 644



**Figure 6:** (a): Time-series of  $A_T$  (brown triangles, right Y-axis),  $C_T$  (blue triangles, right Y-axis),  $p\text{CO}_2$  calculated (open triangles, left Y-axis) and  $p\text{CO}_2$  measured (filled circles) in the Western English Channel in 2012-2014 (Marrec et al., 2014). (b): Measured  $p\text{CO}_2$  versus calculated  $p\text{CO}_2$  for the same samples. The mean difference ( $p\text{CO}_{2\text{cal}}-p\text{CO}_{2\text{mes}}$ ) for 528 samples is  $-1.9 \mu\text{atm} (\pm 11.9) \mu\text{atm}$ . Data from Marrec and Bozec (2016 a,b; 2017). Localization of the samples is shown in the inserted map.

In the Ligurian Sea, following the first high frequency in situ  $f\text{CO}_2$  measurements in 1995-1997 at the DYFAMED time-series station (Hood and Merlivat, 2001), a new CARIOCA  $f\text{CO}_2$  sensor was deployed at that location in 2013 (BOUSSOLE project, Merlivat et al., 2018). The CARIOCA sensor was calibrated with regular  $A_T/C_T$  analyses performed at LOCEAN. Based on these data, the mean difference between CARIOCA- $f\text{CO}_2$  measurements and calculated- $f\text{CO}_2$  data was estimated to be around  $\pm 4.4 \mu\text{atm}$  for 2013-2015, i.e. the same order than the precision of the CARIOCA sensor ( $\pm 5 \mu\text{atm}$ , Merlivat et al., 2018). Here we extend the results for the period 2013-2018 (Golbol et al., 2020; data also in SOCAT version v2021, Bakker et al., 2016) and compared the CARIOCA  $f\text{CO}_2$  time-series with  $A_T$  and  $C_T$  data from different cruises (BOUSSOLE, DYFAMED and MOOSE-GE) selected in the layer 0-20m at that location (Figure S4). For 67 co-located samples at different seasons and years, the mean difference between calculated and measured  $f\text{CO}_2$  ( $f\text{CO}_{2\text{cal}}-f\text{CO}_{2\text{mes}}$ ) was  $-3.7 \mu\text{atm}$  ( $\pm 10.8$ )  $\mu\text{atm}$  where  $f\text{CO}_2$  was calculated from  $A_T/C_T$  pairs using the constant from Lueker et al (2000). At that location, the alkalinity is relatively constant over 2013-2018 with an average concentration of  $2569.8 (\pm 13.2) \mu\text{mol kg}^{-1}$ .  $C_T$  concentrations show a clear seasonality, decreasing by around  $50 \mu\text{mol kg}^{-1}$  from winter to late summer driving the large seasonal cycle of  $f\text{CO}_2$  (range  $80 \mu\text{atm}$ ) revealed in both measured and calculated values (here  $f\text{CO}_2$  is normalized at  $13^\circ\text{C}$ , Figure S4). In addition to calibration purposes, a regional  $A_T/\text{Salinity}$  relationship was derived from the  $A_T$  data measured at that location and successfully used to construct time-

series of  $C_T$  and pH calculated from the high-frequency CARIOCA  $f\text{CO}_2$  data to investigate and interpret the long-term change of  $f\text{CO}_2$  and acidification in the Ligurian Sea (Merlivat et al., 2018; Coppola et al., 2020).

A CARIOCA sensor was also deployed in 2003 near the SOMLIT-Brest time-series site in the Bay of Brest (Bozec et al., 2011; Salt et al., 2016). As for BOUSSOLE in the Ligurian Sea, samples collected for  $A_T$  and  $C_T$  were used for validation of the  $p\text{CO}_2$  recorded by the CARIOCA sensor and the comparison with calculated  $p\text{CO}_2$  showed a good agreement, i.e.  $p\text{CO}_{2\text{cal}} = 0.98 * p\text{CO}_{2\text{mes}} + 7 \mu\text{atm}$  (Bozec et al., 2011). CARIOCA sensors were also deployed on moorings in the Tropical Atlantic (PIRATA project, e.g. Lefèvre et al., 2008, 2016; Parard et al., 2010). With the discrete  $A_T$  and  $C_T$  data included in this synthesis (EGEE and PIRATA-FR cruises), the  $f\text{CO}_2$  data from CARIOCA sensor associated with an adapted  $A_T$ /Salinity relationship were used to derive pH (Lefèvre et al., 2016) or  $C_T$  time-series to evaluate net community production in the Eastern tropical Atlantic (Parard et al., 2010; Lefèvre and Merlivat 2012).

Although this is not a direct instrumental inter-comparison, differences between  $p\text{CO}_2$  (or  $f\text{CO}_2$ ) calculated using  $A_T/C_T$  pairs with direct  $p\text{CO}_2$  measurements give a glimpse of the quality of  $A_T$  and  $C_T$  data in this synthesis given the uncertainty attached to the  $p\text{CO}_2$  or pH calculations (Orr et al., 2015). For example, in the frame of the SURATLANT project in the North Atlantic, calculated  $f\text{CO}_2$  data were compared with co-located  $f\text{CO}_2$  measurements for different seasons and years (Figure S5). The mean differences ( $f\text{CO}_{2\text{cal}} - f\text{CO}_{2\text{mes}}$ ) ranged between  $-4.3 \mu\text{atm} (\pm 12.9) \mu\text{atm}$  (2004-2007, 74 co-located samples) and  $-3.0 (\pm 12.1) \mu\text{atm}$  (2014-2015, 98 co-located samples). The differences are almost the same for different years (and seasons) and are thus attributed to method uncertainties (including sampling time, measurement errors, and data processing). Based on these comparisons and the consistency between data we are confident that the  $A_T$  and  $C_T$  data presented in this synthesis could be used to calculate  $f\text{CO}_2$  (and pH) and interpret temporal changes and drivers of these parameters as well as to estimate air-sea  $\text{CO}_2$  fluxes in the North Atlantic (e.g. Corbière et al., 2007; Schuster et al., 2009, 2013; Watson et al., 2009; Metzl et al., 2010; Mc Kinley et al., 2011; Reverdin et al., 2018; Kitidis et al., 2019; Leseurre et al., 2020).

The  $A_T$  and  $C_T$  data in this synthesis have been also successfully used for  $f\text{CO}_2$  and air-sea  $\text{CO}_2$  fluxes calculations in other regions: the tropical Atlantic (Koffi et al., 2010), the tropical Pacific (Moutin et al., 2018; Wagener et al., 2018), the Solomon sea (Ganachaud et al., 2017) or the Mediterranean sea and coastal zones (De Carlo et al., 2013; Marrec et al., 2015; Kapsenberg et al., 2017; Coppola et al., 2020; Keraghel et al., 2020; Wimart-Rousseau et al., 2020a; Gattuso et al., 2023).

In addition,  $A_T$  and  $C_T$  data in the surface and the water-column are also relevant to calculate pH and evaluate its rate of change for addressing ocean acidification topic in different regions (Kapsenberg et al., 2017; Ganachaud et al., 2017; Wagener et al., 2018; Coppola et al., 2020; Leseurre et al., 2020; Lo Monaco et al., 2021). At the time-series station ECOSCOPA in the Bay of Brest (Fleury et al., 2023; Petton et al., 2023), pH calculated with  $A_T/C_T$  data were compared with direct pH measurements (Figure S6). In 2017-2019, pH (at standard temperature  $25^\circ\text{C}$ , pH-25C) was always lower than 8 and presented a large seasonal signal of 0.3 (high pH values in spring, low in winter). The mean difference between calculated and measured pH-25C for 46 samples was equal to  $+0.013 (\pm 0.010)$  which is in the range of the pH uncertainty evaluated by error propagation when calculated from  $A_T/C_T$  pairs ( $A_T$  and  $C_T$  error of  $\pm 3 \mu\text{mol kg}^{-1}$  leads to pH error of  $\pm 0.0144$ ). Part of these  $A_T$  and  $C_T$  data used to calculate pH also helped for interpreting the response of marine species to acidification, e.g. pteropodes or coccolithophores (*Emiliana huxleyi*) in the Mediterranean Sea (Howes et al., 2015, 2017; Meier et al., 2014) or in the Southern Ocean (Beaufort et al., 2011). The  $A_T$  and  $C_T$  data were also

734 supporting environmental analysis in coral reef ecosystems in the tropical Pacific (TARA Expedition, Douville  
735 et al., 2022; Lombard et al., 2023; Canesi et al., 2023).

736


#### 737 **4 Global distribution and relationships from $A_T$ and $C_T$ based on the SNAPO-CO<sub>2</sub> dataset**

738


739 The surface distribution in the global ocean based on the SNAPO-CO<sub>2</sub> dataset is presented in Figure 7  
740 for  $A_T$  and  $C_T$ . In the open ocean, high  $A_T$  concentrations are identified in the subtropics in all basins (Jiang et al.,  
741 2014; Takahashi et al., 2014) with highest concentrations up to 2484  $\mu\text{mol kg}^{-1}$  in the central North Atlantic  
742 (STRASSE cruise in August 2012, 26°N/36°W). In surface and at depth, the  $A_T/\text{Salinity}$  and  $A_T/C_T$  relationships  
743 are clearly identified and structured at regional scale (Figure 8).

744

745  **$A_T [\mu\text{mol/kg}] @ Depth [m]=\text{first}$**




757  **$C_T [\mu\text{mol/kg}] @ Depth [m]=\text{first}$**



773 **Figure 7:** Distribution of  $A_T$  (top) and  $C_T$  (bottom) concentrations ( $\mu\text{mol}\cdot\text{kg}^{-1}$ ) in surface waters (0-10m). Only  
774 data with flag 2 are presented in these figures. Figures produced with ODV (Schlitzer, 2018).

775

776




**Figure 8:** Relationships between  $A_T$  and Salinity (upper panel) and  $A_T$  versus  $C_T$  (lower panel) for samples in surface waters (0-10m and SSS > 31) (left) and in the water column below 100m (right). Only data with flag 2 are presented. The color scales correspond to the temperature (left) or the latitude (right). Some location of data are identified: Mediterranean Sea (Med Sea), Red Sea, Tropical Pacific (PAC), North Atlantic (NATL) and Southern Ocean (SO). Figures produced with ODV (Schlitzer, 2018).

In the Eastern tropical Atlantic (ETA) where the Congo River impacts the salinity field (Vangriesheim et al., 2009),  $A_T$  concentrations range between 2100 and 2400  $\mu\text{mol kg}^{-1}$ . The regional  $A_T$ /Salinity relationship in the ETA based on data from the EGEE cruises in 2005-2007 (Koffi et al., 2010) is robust and validated with more recent measurements from PIRATA-FR cruises in 2010-2019 (Lefevre et al., 2021). The strong  $A_T$ /Salinity relationship in the ETA was also recognized using data from the TARA-MICROBIOME cruise in May-July 2022 (Figure S7). Low salinity (< 30) and low  $A_T$  (1700-2200  $\mu\text{mol kg}^{-1}$ ) are also observed in the Western tropical Atlantic near the Amazon River plume. The  $A_T$ /Salinity relationships in both river plume regions are very similar (Figure S7).

For  $C_T$ , the lowest concentrations were observed in the coastal regions of the Tropical Atlantic, on the eastern side in the Gulf of Guinea (BIOZAIRe cruise in 2003, 6°S/11°E,  $C_T=1390 \mu\text{mol kg}^{-1}$ , Vangriesheim et al., 2009) and on the Western side in coastal zone off French Guyana (PLUMAND cruise in 2007, 5°N/51°W,  $C_T=1512 \mu\text{mol kg}^{-1}$ , Lefèvre et al., 2010). Such low  $C_T$  concentrations were also observed around 5°N/51°W in the Amazon River plume during the recent EUREC4A-OA cruise in 2020 and the TARA-MICROBIOME cruise in 2021 ( $C_T=1451 \mu\text{mol kg}^{-1}$ ) leading to low oceanic  $f\text{CO}_2$  (< 350  $\mu\text{atm}$ ) and a  $\text{CO}_2$  sink in this region (Olivier et al., 2022).


832        The high  $C_T$  concentrations were mainly observed in the Southern Ocean (OISO and ACE cruises)  
 833        south of the Polar Front around 50°S linked to the upwelling of  $C_T$ -rich deep water (Figure 7, Metzl et al., 2006;  
 834        Wu et al., 2019; Chen et al., 2022). This leads to a high  $C_T/A_T$  ratio and a high Revelle factor in the Southern  
 835        Ocean (Figure 9, Fassbender et al., 2017). The high  $C_T$  content and low temperature in the Southern Ocean also  
 836        lead to low calcite and aragonite saturation state ( $\Omega$ ) (Takahashi et al., 2014; Jiang et al., 2015). We calculate  $\Omega$   
 837        from  $A_T$  and  $C_T$  data at insitu temperature, salinity and pressure. At present the surface ocean is not under-  
 838        saturated with regard to aragonite (Figure 10); however, under-saturation levels ( $\Omega-Ar<1$ ) were found around  
 839        500 m in the Southern Ocean (ACE cruise in 2017, MODYDICK cruise in 2018), 1000 m in the Tropical Pacific  
 840        (PANDORA 2012 and OUTPACE 2015 cruises) and 2200 m in the North Atlantic (OVIDE 2012 and 2014  
 841        cruises, see also Turk et al., 2017) (Figure 10). Samples at 400 m from the TARA-Oceans cruise in 2009-2012  
 842        also indicated aragonite under-saturation in the Equatorial Atlantic, Equatorial Pacific, as well as off South  
 843        America (73°W-34°S, Chile) associated to equatorial or eastern boundary upwelling systems (Feely et al., 2012;  
 844        Lauvset et al., 2020).

845        In surface,  $\Omega-Ar>3$  is found in the latitudinal band 45°S-54°N and  $\Omega-Ar<3$ , below the critical  
 846        threshold of  $\Omega-Ar=3.25$  that represents a limit for distribution of tropical coral reefs (Hoegh-Guldberg et al.,  
 847        2007) is observed at very few locations in the tropics.




869        **Figure 9:** Distribution of the  $C_T/A_T$  ratio (top) and the Revelle factor (bottom) in surface waters (0-10m). Only  
 870        data with flag 2 were used. Figures produced with ODV (Schlitzer, 2018).

871  
 872  
 873  
 874  
 875  
 876  
 877  
 878  
 879  
 880  
 881  
 882  
 883  
 884  
 885  
 886  
 887  
 888  
 889  
 890  
 891  
 892  
 893  
 894  
 895  
 896  
 897  
 898  
 899  
 900  
 901  
 902  
 903  
 904  
 905  
 906  
 907  
 908  
 909  
 910  
 911  
 912  
 913  
 914  
 915  
 916  
 917  
 918  
 919



**Figure 10:** (a): Distribution of the aragonite saturation state ( $\Omega$ -Ar) in surface waters (0-10m). Only data with flag 2 were used. (b): Depth profiles (100-3000m) of  $\Omega$ -Ar at few locations in the Tropical Pacific, Atlantic and Southern Oceans. Stations where under-saturation is detected ( $\Omega$ -Ar<1) at depth are identified in the inserted map. Figures produced with ODV (Schlitzer, 2018).

Compared to the open ocean,  $A_T$  concentrations are much higher in the Mediterranean Sea (Copin-Montégut, 1993; Schneider et al., 2007; Álvarez et al., 2023) with values up to  $2600 \mu\text{mol kg}^{-1}$  (Figure 8). The  $A_T$  and  $C_T$  data obtained in 1998-2019 show on average a clear contrast between the northern and southern regions of the Western Mediterranean sea (Figure 11 a, b) with higher concentration in the Ligurian Sea and the Gulf of Lion (Gemayel et al., 2015). However, the basin scale average distribution view smoothed the meso-scale signals recognized in the Mediterranean Sea (e.g. Bosse et al., 2017; Petrenko et al., 2017). In the Gulf of Lion the synthesis of 11 cruises conducted from May 2010 to June 2011 (CARBORHONE, CASCADE, LATEX, MOLA, MOOSE-GE) highlights the contrasting distributions of  $A_T$  and  $C_T$  in the coastal zones and off shore (Figure 11 c, d). The averaging of all data in 1998-2019 also smoothed the seasonal signal and the inter-annual variability described below.




**Figure 11:** Distribution of  $A_T$  (a) and  $C_T$  (b) in  $\mu\text{mol kg}^{-1}$  in surface waters of the Western Mediterranean Sea (0-10m) from all data for 1998-2019. Detailed distribution of  $A_T$  (c) and  $C_T$  (d) in  $\mu\text{mol kg}^{-1}$  in surface waters of the Gulf of Lion for the period 2010-2011 only (cruises CARBORHONE, CASCADE, LATEX, MOLA, MOOSE-GE). Figures produced with ODV (Schlitzer, 2018).

## 5 Temporal variations of $A_T$ and $C_T$ : examples from the SNAPO-CO<sub>2</sub> dataset

Time-series stations such as BATS, ESTOC, HOT in the subtropics and stations in the Irminger Sea or in the Iceland Sea are the only way to detect the long-term change in the ocean carbonate system in the surface and the water column (Bates et al., 2014). These important time-series help to understand driving processes (e.g. Hagens and Middelburg, 2016) and are often used to validate the  $p\text{CO}_2$ ,  $A_T$ ,  $C_T$ , or pH reconstructed fields (e.g. Rödenbeck et al., 2013; Broullón et al., 2019, 2020; Keppler et al., 2020; Gregor and Gruber, 2021; Chau et al., 2023; Ma et al., 2023).

Here we show examples of the temporal surface variations at locations where data were obtained for more than 10 years (Figure 12). We thus selected the following contrasting regions: the North Atlantic subpolar gyre (NASPG around 60°N/30°W, period 1993-2018), the Equatorial Atlantic (at 2°N-2°S/12°W-8°W, period 2005-2017), the Indian Ocean subtropical sector (26-35°S/50-56°E, period 1998-2018), the Indian Ocean high latitude (54-60°S/60-70°E, period 1998-2018), the Ligurian Sea (around DYFAMED station, 43.5-42.5°N/5.5-9°E, period 1998-2019) and times-series stations in the coastal zones off Britanny (period 2008-2019).



**Figure 12:** Time-series of (a) sea surface temperature (°C), (b)  $A_T$  ( $\mu\text{mol kg}^{-1}$ ) and (c)  $C_T$  ( $\mu\text{mol kg}^{-1}$ ) in 6 regions: the North Atlantic subpolar gyre (NASPG 1993-2018, green diamond), the Equatorial Atlantic (ATL-EQ, 2005-2017, green triangle), the Indian subtropical sector (IO-SBT, red circle) and high latitude (IO-POOZ, blue circle) (1998-2018), the Ligurian Sea (MED, 1998-2019, orange square) and times-series stations in the coastal zones off Brittany (BRIT, period 2008-2019, black dots). Trends (dashed lines and values) are shown when relevant for the discussion ( $C_T$  trends listed in Table 6).

**Table 6:** Trend of  $C_T$  ( $\mu\text{mol kg}^{-1} \text{ yr}^{-1}$ ) and corresponding standard error in 5 selected regions where data were available for more than 10 years (data are shown in Figure 12). The projects/cruises for selection of the data in each domain are indicated.

Region (acronym)	Period	$C_T$ trend ( $\mu\text{mol kg}^{-1} \text{ yr}^{-1}$ )	Season	Projects/Cruises
North Atlantic (NASPG)	1994-2014	+0.72 (0.17)	Jan-Feb	SURATLANT
Indian Subtropic (IO-SBT)	1998-2018	+0.65 (0.12)	Jan-Feb	OISO
Indian South. (IO-POOZ)	1998-2018	+0.67 (0.04)	Jan-Feb	OISO
Ligurian Sea (MED)	1998-2019	+0.68 (0.18)	All seasons	DYFAMED, BOUSSOLE, MOOSE-GE
Coast Brittany (BRIT)	2008-2019	+1.72 (0.28)	All seasons	Brest, Roscoff, ECOSCOPIA, PENZE

In the 6 regions, there was a progressive warming most clearly detected in the Mediterranean Sea (e.g. Nykjaer, 2009). From 1998 to 2019 the warming in the Ligurian Sea was  $+0.1208^{\circ}\text{C yr}^{-1}$  ( $\pm 0.0227$ ) (Figure 12). In the equatorial Atlantic, the apparent rapid increase of temperature of  $+0.48^{\circ}\text{C yr}^{-1}$  ( $\pm 0.04$ ) in 2005-2017 from the selected data indicated a change in water masses and circulation. The colder sea surface in 2005 was associated with the so-called Atlantic Cold Tongue (ACT) which was one of the most intense ATC since 1982 (Caniaux et al., 2011). The ACT also leads to significant changes in oceanic  $f\text{CO}_2$  and air-sea  $\text{CO}_2$  fluxes (Parard et al., 2010; Koseki et al., 2023) and explained the high  $C_T$  concentrations observed in 2005 in this region (Figure 12, Koffi et al., 2010).

Total alkalinity presents rather homogenous concentrations in the NASGP and the south Indian Ocean. Inter-annual variability of  $A_T$  is pronounced in the equatorial Atlantic ranging between 2245 and 2378  $\mu\text{mol kg}^{-1}$ . This is mainly related to salinity as normalized  $A_T$  values ( $N-A_T$ , for salinity= 35) do not show such inter-annual variability (Mean  $N-A_T = 2295.7 \pm 4.6 \mu\text{mol kg}^{-1}$ ,  $n= 67$  for 2005-2017, not shown). In the coastal zones off Brittany, the  $A_T$  is also highly variable (Salt et al., 2016; Gac et al., 2021) ranging between 2150 and 2386  $\mu\text{mol kg}^{-1}$  (Figure 12).

An interesting signal is the progressive increase of  $A_T$  in the Mediterranean Sea. The positive  $A_T$  trend of  $+0.53 (\pm 0.11) \mu\text{mol kg}^{-1} \text{yr}^{-1}$  ( $n=538$ ) in 1998-2019 in the region offshore was also observed at the coastal station SOMLIT-Point-B in 2007-2015 but with a faster increase of  $+2.08 (\pm 0.19) \mu\text{mol kg}^{-1} \text{yr}^{-1}$  (Kapsenberg et al., 2017). Close to the DYFAMED site, at station SOMLIT-Point-B, the  $A_T$  trend was not linked to salinity temporal changes as a positive  $N-A_T$  trend was also reported,  $+0.52 (\pm 0.07) \mu\text{mol kg}^{-1} \text{yr}^{-1}$  (not shown). Based on data from the PERLE cruises in 2018-2021 a significant increase in  $A_T$  was also identified in the Eastern Mediterranean Sea (Wimart-Rousseau et al., 2021). Along with the increase of  $C_T$  and the warming, the  $A_T$  increase would impact on the  $f\text{CO}_2$ , air-sea  $\text{CO}_2$  fluxes and pH temporal changes (Merlivat et al., 2018). Processes explaining the  $A_T$  increase in the Mediterranean Sea are still unexplained and deserve further investigations (Coppola et al., 2019).

As expected, because of the anthropogenic  $\text{CO}_2$  uptake the  $C_T$  concentrations increased in most regions (Figure 12, Table 6). This is identified in the Indian Ocean (in the subtropics and the high latitude), in the Mediterranean Sea, and in coastal waters off Brittany. However, the signal is more complex in the NASPG. As previously shown the  $C_T$  trend in the NASPG depends on seasons and decades (Metzl et al., 2010; Reverdin et al., 2018; Fröb et al., 2019; Leseurre et al., 2020). Here we selected only the data in January-February from the SURATLANT cruises leading a  $C_T$  trend of  $+0.72 (\pm 0.17) \mu\text{mol kg}^{-1} \text{yr}^{-1}$ . Compared to the regions further north the  $C_T$  trend in the NASPG is about half the  $C_T$  trends of  $+1.44 (\pm 0.23) \mu\text{mol kg}^{-1} \text{yr}^{-1}$  observed in the Iceland Sea (Olafsson et al., 2009) or  $+1.48 (\pm 0.22) \mu\text{mol kg}^{-1} \text{yr}^{-1}$  at station M in the Norwegian Sea (Skjelvan et al., 2022).


In the coastal zones off Brittany, although there are large seasonal and inter-annual variabilities (Gac et al., 2021), an annual  $C_T$  trend of  $+1.72 (\pm 0.28) \mu\text{mol kg}^{-1} \text{yr}^{-1}$  is detected over 10 years (2009 to 2019). The same is observed in the Mediterranean Sea where the  $C_T$  offshore trend of  $+0.69 (\pm 0.18) \mu\text{mol kg}^{-1} \text{yr}^{-1}$  is low compared to what was observed in the coastal zone (SOMLIT-Point-B,  $+2.97 (\pm 0.20) \mu\text{mol kg}^{-1} \text{yr}^{-1}$ , Kapsenberg et al., 2017).

In the southern Indian Ocean,  $C_T$  concentrations also increased in both subtropical and high latitudes, two regions where the primary productivity is relatively low (oligotrophic regime in the subtropics and High Nutrient Low Chlorophyll regime, HNLC, south of the Polar Front). With the data selected for austral summer

(January-February) the  $C_T$  trends appeared almost similar in these two regions, around  $+0.65 \mu\text{mol kg}^{-1} \text{yr}^{-1}$  (Table 6).

Finally, in the Equatorial Atlantic the selected data around  $0^{\circ}$ - $10^{\circ}\text{W}$  highlighted the large variability linked to the oceanic circulation. Detecting a  $C_T$  trend as well as a possible link with anthropogenic carbon uptake, at least with the data available in 2005-2017, appears to be intricate as it has been previously discussed for the period 2006-2013 (Lefèvre et al., 2016). However, the signal of the  $C_T$  increase is better identified north or south of the Equator in the Eastern tropical Atlantic sector (Lefèvre et al., 2021).

In the water column  $A_T$  and  $C_T$  data from dedicated cruises were used to evaluate the anthropogenic  $\text{CO}_2$  ( $C_{\text{ant}}$ ) distribution and pH change since pre-industrial era (e.g. PANDORA cruise, Ganachaud et al., 2017; OUTPACE cruise, Wagener et al., 2018; SOMBA cruise, Keraghel et al., 2020). Time-series at DYFAMED station also enabled to investigate the temporal variability of  $C_T$ ,  $A_T$  and  $C_{\text{ant}}$  in the water column (Touratier and Goyet, 2009; Coppola et al., 2020; Fourrier et al., 2022). As an example of the observed temporal variations at depth we selected the data in the layer 950-1050m in the Ligurian Sea from different cruises (Figure 13). At that depth both  $A_T$  and  $C_T$  present some large anomalies especially noticed in 2013 (lower  $A_T$  and  $C_T$  in February 2013, DEWEX cruise) and in 2018 (higher  $A_T$  and  $C_T$  in May 2018, MOOSE-GE cruise) the later probably linked to episodic convective process that occurred in winter 2018 (Fourrier et al., 2022; Coppola et al., 2023). During the strong convection event in 2013 the positive anomalies of  $A_T$  and  $C_T$  were mostly identified in the upper layers (Figure 12c, Ulles et al, 2023).



**Figure 13:** Time-series of  $A_T$  ( $\mu\text{mol kg}^{-1}$ ) and  $C_T$  ( $\mu\text{mol kg}^{-1}$ ) in the Ligurian Sea (1998-2019) in the layer 950-1050m. Annual mean (grey triangles) was calculated from all data each year (black dots). The trends (dashed line) based on annual mean are  $+0.82 (\pm 0.15) \mu\text{mol kg}^{-1} \text{yr}^{-1}$  for  $A_T$  and  $+1.20 (\pm 0.12) \mu\text{mol kg}^{-1} \text{yr}^{-1}$  for  $C_T$ . In this layer data selected are from cruises ANTARES, CASCADE, DEWEX, DYFAMED, MOOSE-GE and PEACETIME (location of stations shown in the inserted map).

In this region the long-term increase of  $A_T$  indicates that in addition to the anthropogenic  $\text{CO}_2$  signal other processes are at play to explain the rapid  $C_T$  trend of  $+1.20 (\pm 0.12) \mu\text{mol kg}^{-1} \text{yr}^{-1}$  at depth compared to that observed in surface (Figure 12). The signal at depth is probably linked to the variations of the deep convection and mixing with Levantine intermediate water (LIW, Margerier et al., 2020) with higher  $A_T$  and  $C_T$

concentrations. The long-term increase of  $A_T$  and  $C_T$  at depth (here at 1000m, Figure 13) was also observed below 2000m (Coppola et al., 2020) a signal that has to be investigated in dedicated analysis using other properties ( $O_2$ , nutrients, following Fourrier et al. (2022) for the period 2012-2020) and a larger dataset in the Mediterranean Sea (e.g. GLODAP).

## 6 Using $A_T$ and $C_T$ data to validate observations from autonomous instruments


The dataset presented in this synthesis would also offer interesting observations to validate properties ( $A_T$  and  $C_T$ ) derived from BGC-Argo floats equipped with pH sensors (e.g. Bushinsky et al., 2019; Mazloff et al., 2023; Mignot et al., 2023). The water column in situ  $A_T$  and  $C_T$  data obtained during the Antarctic Circumpolar Expedition (ACE) in 2016-2017 were collected at location where SOCCOM floats were launched (Walton and Thomas, 2018). A SOCCOM float (WMO ID 5905069) was launched on January 11<sup>th</sup> 2017 at 55°S-96°E south of the Polar Front in the southern Indian Ocean. The pH, temperature and salinity data from the float were then used to derive  $A_T$  and  $C_T$  profiles (here using a multiple linear regression (MLR) algorithm, Williams et al., 2016, 2017). In the top layers the discrete ACE data (Figure 14) present large variability of  $A_T$  and  $C_T$  concentrations not captured in the records derived from the float (MLR method somehow smooth the profiles). However, given the uncertainty in reconstructed  $A_T$  from float data (5.6  $\mu\text{mol kg}^{-1}$ ) the average values in the first 100m were almost identical ( $A_{T-\text{ACE}} = 2285.1 (\pm 4.4) \mu\text{mol kg}^{-1}$  and  $A_{T-\text{float}} = 2278.3 (\pm 0.7) \mu\text{mol kg}^{-1}$ ;  $C_{T-\text{ACE}} = 2139.7 (\pm 9.2) \mu\text{mol kg}^{-1}$  and  $C_{T-\text{float}} = 2141.1 (\pm 3.2) \mu\text{mol kg}^{-1}$ ). Moreover below 200m, profiles from the float are coherent compared to the  $A_T$  and  $C_T$  measurements (Figure 14). This is encouraging for using float data to explore the seasonal variability of  $A_T$  and  $C_T$  in the Southern Ocean (e.g. Williams et al., 2018; Johnson et al., 2022) and the estimation of anthropogenic  $\text{CO}_2$  in the water column in this sector (Figure 14). Here the  $C_{\text{ant}}$  concentrations were calculated below 200m (corresponding to the temperature minimum of the winter winter in the SO and using the TrOCA method, Touratier et al., 2007). The float data suggest that  $C_{\text{ant}}$  concentrations are positive down to about 1000m, with maximum values in subsurface. In 2017 the mean  $C_{\text{ant}}$  concentration at 200m was  $49.1 (\pm 9.0) \mu\text{mol kg}^{-1}$ . Below that depth,  $C_{\text{ant}}$  decreased and reduced to  $+29.8 (\pm 8.5) \mu\text{mol kg}^{-1}$  in the layer 300-400m. To complement the  $C_{\text{ant}}$  inventories based on GLODAP data product (e.g. Gruber et al., 2019)  $C_{\text{ant}}$  estimates derived from BGC-Argo floats as evaluated here in the Southern Ocean could be applied in other locations as was previously tested in the North Pacific (Li et al., 2019).

In surface water as the  $A_T$  derived from the float data are deduced using MLR or LIAR methods (Williams et al., 2017; Carter et al., 2016), the  $A_T$  data in the SNAPO-CO<sub>2</sub> synthesis could also be used to identify  $A_T$  anomalies not always captured from floats. This is particularly relevant in coccolithophores blooms areas when low  $A_T$  concentrations and high  $p\text{CO}_2$  are observed (e.g. Balch et al., 2016 in the Southern Ocean; Robertson et al., 1994 in the North Atlantic).

## 7 Summary and suggestions

The ocean data synthesized in this product are based on measurements of  $A_T$  and  $C_T$  performed between the period 1993 and 2022 with an accuracy of  $\pm 4 \mu\text{mol kg}^{-1}$ . It offers a large data set of  $A_T$  and  $C_T$  for the global ocean and regional biogeochemical studies. It includes more than 44 400 surface and water column observations in all oceanic basins, in the Mediterranean Sea, in the coastal zones, near coral reefs, and in rivers. For the open ocean this complements the SOCAT and GLODAP data products (Bakker et al., 2016; Lauvset et al., 2022). For

the coastal sites this also complements the synthesis of coastal time-series only done around North America (Fassbender et al., 2018; Jiang et al., 2021; OCADS, 2023).



**Figure 14:** Profiles of (a)  $A_T$  ( $\mu\text{mol kg}^{-1}$ ) and (b)  $C_T$  ( $\mu\text{mol kg}^{-1}$ ) observed at station ACE-20 (55°S-95°E, 11/1/17, black triangles) compared with the profiles deduced from the SOCCOM float (WMO code 5905069) launched at that location (first data on January 12<sup>th</sup> 2017, grey circles). The location/drift of the float in 2017-2018 is shown on the inserted map. (c) Hovmöller section (Pressure/time) of anthropogenic CO<sub>2</sub> concentrations ( $C_{ant}$  in  $\mu\text{mol kg}^{-1}$ ) estimated from the float data ( $A_T$ ,  $C_T$ , O<sub>2</sub>, T) below 200m (period January 2017–February 2018). Section produced with ODV (Schlitzer, 2018).

The SNAPO-CO<sub>2</sub> dataset enables to investigate seasonal variations to decadal trends of  $A_T$  and  $C_T$  in various oceanic provinces. In regions where data are available for more than 2 decades in surface water (North Atlantic, Ligurian Sea, Southern Indian Ocean, and coastal regions), all time-series show an increase in  $C_T$ . Excepted in the Mediterranean Sea,  $A_T$  appears relatively constant over time, although the  $A_T$  content present significant inter-annual variability such as in the NASPG or in the coastal zones including near the Congo and Amazon Rivers plumes.

This dataset represents independent data for validation of reconstructed  $A_T$  or  $C_T$  fields using various methods (e.g. Rödenbeck et al., 2013, 2015; Sauzède et al., 2017; Turk et al., 2017; Bittig et al., 2018; Broullón et al., 2019, 2020; Land et al., 2019; Keppler et al., 2020; Fourier et al., 2020; Gregor and Gruber, 2021; Sims et al., 2023; Chau et al., 2023). It is also useful to validate Earth System Models (ESM) that currently present bias to reproduce the seasonal cycle of  $C_T$  and  $A_T$  due to inadequate representation of biogeochemical cycles, including the coupling of biological and physical processes (e.g. Pilcher et al., 2015; Mongwe et al., 2018; Lerner et al., 2021). This should be resolved for confident in future projections of the productivity, ocean acidification, and the responses of the marine ecosystems (e.g. Kwiatowkki et al., 2020). Recall that OBGM or ESM models calculate pCO<sub>2</sub> from  $A_T/C_T$  pairs and the simulated annual CO<sub>2</sub> flux might be correct when

1186 compared to observations but for wrong reasons (e.g. Goris et al., 2018, Lerner et al., 2021). For example, it has  
1187 been shown that biases in  $A_T$  in ESM models led to an overestimation of the oceanic  $f\text{CO}_2$  trend and thus  
1188 uncertainty when predicting the oceanic anthropogenic  $\text{CO}_2$  uptake (Lebehot et al., 2019). The simulated  
1189 seasonal cycle of  $f\text{CO}_2$  is also uncertain in ESM models especially in high latitudes (e.g. Joos et al., 2023). It is  
1190 thus important to attempt validating ESM models with  $A_T$  and  $C_T$  data such as presented in this synthesis.

1191 This dataset would also serve for validating autonomous platforms capable of measuring pH and  $f\text{CO}_2$   
1192 variables and, along with SOCAT and GLODAP datasets, provides an additional reference dataset for the  
1193 development and validation of regional biogeochemical models for simulating air-sea  $\text{CO}_2$  fluxes. It is also  
1194 essential for training and validating neural networks capable of predicting variables in the carbonate system,  
1195 thereby enhancing observations of marine  $\text{CO}_2$  at different spatial and temporal scales.

1196 The data presented here are available online on the Seanoe servor (Metzl et al., 2023,  
1197 <https://doi.org/10.17882/95414>) and is divided in two files: one for the Global Ocean, and one for the  
1198 Mediterranean Sea. The sources of the original datasets (doi) with the associated references are listed in the  
1199 Supplementary Material (Table S3, S4). We invite the users to comment on any anomaly that would have not  
1200 been detected or to suggest potential misqualification of data in the present product (e.g. data probably good  
1201 although assigned with flag 3, probably wrong). The SNAPO-CO<sub>2</sub> dataset will be regularly updated on Seanoe  
1202 data servor with new observations controlled and archived.

## 1203 1204 8 Data availability

1205 Data presented in this study are available at Seanoe: <https://www.seanoe.org>, <https://doi.org/10.17882/95414>  
1206 (Metzl et al., 2023).

1207 *Author contributions.* NM prepared the data synthesis, the figures and wrote the draft of the manuscript with  
1208 contributions from all authors. JF measured the discrete samples since 2014, with the help from CM and CLM,  
1209 and prepared the individual reports for each project. NM and JF pre-qualified the discrete  $A_T/C_T$  data. CLM and  
1210 NM are co-Is of the ongoing OISO project and qualified the underway  $A_T/C_T$  data from OISO and CLIM-  
1211 EPARSES cruises. All authors have contributed either to organizing cruises, sample collection and/or data  
1212 qualification, and reviewed the manuscript.

1214 *Competing interest.* The authors declare that they have no conflict of interest.

1216 *Acknowledgments.* The  $A_T$  and  $C_T$  data presented in this study were measured at the SNAPO-CO<sub>2</sub> facility  
1217 (Service National d'Analyse des Paramètres Océaniques du CO<sub>2</sub>) housed by the LOCEAN laboratory and part of  
1218 the OSU ECCE Terra at Sorbonne University and INSU/CNRS analytical services. Support by INSU/CNRS, by  
1219 OSU ECCE Terra and by LOCEAN, is gratefully acknowledged as well as support by different French “Services  
1220 nationaux d’Observations”, such as OISO/CARAUS, SOMLIT, PIRATA, SSS and MOOSE. We thank the  
1221 research infrastructure ICOS (Integrated Carbon Observation System) France for funding a large part of the  
1222 analyses. We thank the French oceanographic fleet (“Flotte océanographique française”) for financial and  
1223 logistic support for most cruises listed in this synthesis and for the OISO program  
1224 (<https://campagnes.flotteoceanographique.fr/series/228/>). We acknowledge the MOOSE program (Mediterranean  
1225 Ocean Observing System for the Environment, <https://campagnes.flotteoceanographique.fr/series/235/fr/>)  
1226 coordinated by CNRS-INSU and the Research Infrastructure IILICO (CNRS-IFREMER). AWIPEV-CO<sub>2</sub> was  
1227 coordinated by AWI and the Research Infrastructure IILICO (CNRS-IFREMER).

1228 supported by the Coastal Observing System for Northern and Arctic Seas (COSYNA), the two Helmholtz large-  
1229 scale infrastructure projects ACROSS and MOSES, the French Polar Institute (IPEV) as well as the European  
1230 Union's Horizon 2020 research and innovation projects Jericho-Next (No 871153 and 951799), INTAROS (No  
1231 727890) and FACE-IT (No 869154). The BOUSSOLE time series project was funded by the Centre National  
1232 d'Etudes Spatiales (CNES) and the European Space Agency (ESA ESRIN contract 4000119096/17/I-BG), and  
1233 the first three years of the CO<sub>2</sub> time series at that site were funded by the French Agence Nationale de la  
1234 Recherche (ANR). The EURECA4-OA cruise was also supported by the EURECA4-OA JPI Ocean and Climate  
1235 program. We thank Tara Ocean Foundation and many institutes and funding agencies for supporting TARA  
1236 cruises since 2009. The OISO program was supported by the French institutes INSU (Institut National des  
1237 Sciences de l'Univers) and IPEV (Institut Polaire Paul-Emile Victor), OSU Ecce-Terra (at Sorbonne Université),  
1238 and the French program SOERE/Great-Gases. The CLIM-EPARSES cruise was supported by TAAF (Terres  
1239 Australes et ANtarctiques Françaises), Fondation du Prince Albert II de Monaco, IRD, OSU Ecce-Terra, CNRS,  
1240 MNHN, LOCEAN and LSCE laboratory. Data from the float launched during the ACE cruise were made freely  
1241 available by the Southern Ocean Carbon and Climate Observations and Modeling (SOCCOM) Project funded by  
1242 the National Science Foundation, Division of Polar Programs (NSF PLR -1425989), supplemented by NASA,  
1243 and by the International Argo Program and the NOAA programs that contribute to it. The Argo Program is part  
1244 of the Global Ocean Observing System (<http://doi.org/10.17882/42182>, <http://argo.jcompos.org>). We thank  
1245 Frédéric Merceur (IFREMER) for preparing the page and data availability on Seanoe. We thank Patrick  
1246 Raimbault (retired, former at MIO, Marseille) for managing the MOOSE project until 2019. We thank all  
1247 colleagues and students who participated to the cruises and have carefully collected the precious seawater  
1248 samples. We warmly acknowledge our colleague Christian Brunet (retired) for his supportive help for the  
1249 analysis since the start of the Service facility SNAPO-CO<sub>2</sub>. We would like to pay tribute to our late colleague  
1250 Frédéric Diaz who contributed to the LATEX cruise in 2010. We thank the associated editor Xingchen Wang to  
1251 manage this manuscript, Marta Álvarez and a (more or less) anonymous reviewer for their suggestions that  
1252 helped to improve this article.

1253

## 1254 **References**

1255

1256 Álvarez, M., Catalá, T. S., Civitarese, G., Coppola, L., Hassoun, A. E.R., Ibello, V., Lazzari, P., Lefevre, D.,  
1257 Macías, D., Santinelli, C. and Ulses, C.: Chapter 11 - Mediterranean Sea general biogeochemistry, Editor(s):  
1258 Katrin Schroeder, Jacopo Chiggiato, Oceanography of the Mediterranean Sea, Elsevier, Pages 387-451,  
1259 <https://doi.org/10.1016/B978-0-12-823692-5.00004-2>., 2023.

1260

1261 Bakker, D. C. E., Pfeil, B., Smith, K., Hankin, S., Olsen, A., Alin, S. R., Cosca, C., Harasawa, S., Kozyr, A.,  
1262 Nojiri, Y., O'Brien, K. M., Schuster, U., Telszewski, M., Tilbrook, B., Wada, C., Akl, J., Barbero, L., Bates, N.  
1263 R., Boutin, J., Bozec, Y., Cai, W.-J., Castle, R. D., Chavez, F. P., Chen, L., Chierici, M., Currie, K., De Baar, H.  
1264 J. W., Evans, W., Feely, R. A., Fransson, A., Gao, Z., Hales, B., Hardman-Mountford, N. J., Hoppema, M.,  
1265 Huang, W.-J., Hunt, C. W., Huss, B., Ichikawa, T., Johannessen, T., Jones, E. M., Jones, S., Jutterstrøm, S.,  
1266 Kitidis, V., Kötzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Manke, A. B., Mathis, J. T., Merlivat,  
1267 L., Metzl, N., Murata, A., Newberger, T., Omar, A. M., Ono, T., Park, G.-H., Paterson, K., Pierrot, D., Ríos, A.  
1268 F., Sabine, C. L., Saito, S., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Sieger, R., Skjelvan, I., Steinhoff, T.,  
1269 Sullivan, K. F., Sun, H., Sutton, A. J., Suzuki, T., Sweeney, C., Takahashi, T., Tjiputra, J., Tsurushima, N., Van  
1270 Heuven, S. M. A. C., Vandemark, D., Vlahos, P., Wallace, D. W. R., Wanninkhof, R. and Watson, A. J.: An  
1271 update to the Surface Ocean CO<sub>2</sub> Atlas (SOCAT version 2). Earth System Science Data, 6, 69-90.  
1272 doi:10.5194/essd-6-69-2014. 2014

1273

- 1274 Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa,  
 1275 S., Jones, S. D., Nakaoka, S.-I., Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B.,  
 1276 Wada, C., Wanninkhof, R., Alin, S. R., Balestrini, C. F., Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F.,  
 1277 Boutin, J., Bozec, Y., Burger, E. F., Cai, W.-J., Castle, R. D., Chen, L., Chierici, M., Currie, K., Evans, W.,  
 1278 Featherstone, C., Feely, R. A., Fransson, A., Goyet, C., Greenwood, N., Gregor, L., Hankin, S., Hardman-  
 1279 Mountford, N. J., Harley, J., Hauck, J., Hoppema, M., Humphreys, M. P., Hunt, C. W., Huss, B., Ibánhez, J. S.  
 1280 P., Johannessen, T., Keeling, R., Kitidis, V., Körtzinger, A., Kozyr, A., Krasakopoulou, E., Kuwata, A.,  
 1281 Landschützer, P., Lauvset, S. K., Lefèvre, N., Lo Monaco, C., Manke, A., Mathis, J. T., Merlivat, L., Millero, F.  
 1282 J., Monteiro, P. M. S., Munro, D. R., Murata, A., Newberger, T., Omar, A. M., Ono, T., Paterson, K., Pearce, D.,  
 1283 Pierrot, D., Robbins, L. L., Saito, S., Salisbury, J., Schlitzer, R., Schneider, B., Schweitzer, R., Sieger, R.,  
 1284 Skjelvan, I., Sullivan, K. F., Sutherland, S. C., Sutton, A. J., Tadokoro, K., Telszewski, M., Tuma, M., Van  
 1285 Heuven, S. M. A. C., Vandemark, D., Ward, B., Watson, A. J., and Xu, S.: A multi-decade record of high-quality  
 1286 fCO<sub>2</sub> data in version 3 of the Surface Ocean CO<sub>2</sub> Atlas (SOCAT), *Earth Syst. Sci. Data*, 8, 383–413,  
 1287 doi:10.5194/essd-8-383-2016. 2016.
- 1288
- 1289 Bakker, D.C.E., Alin, S.R., Bates, N.R., Becker, M., Feely, R. A., Gritzalis, T. , Jones, S. D., Kozyr, A., Lauvset,  
 1290 S. K., Metzl, N., Munro, D.R., Nakaoka, S.-I., Nojiri, Y., O'Brien, K., Olsen, A., Pierrot, D., Rehder, G.,  
 1291 Steinhoff, T., Sutton, A., Sweeney, C., Tilbrook, B., Wada, C., Wanninkhof, R., and all >100 SOCAT  
 1292 contributors: An alarming decline in the ocean CO<sub>2</sub> observing capacity. Available at [www.socat.info](http://www.socat.info). 2023.
- 1293
- 1294 Balch, W.M., Bates, N.R., Lam, P.J., Twining, B.S., Rosengard, S. Z., Bowler, B.C., Drapeau, D.T., Garley, R.,  
 1295 Lubelczyk, L.C., Mitchell, C. and Rauschenberg S.: Factors regulating the Great Calcite Belt in the Southern  
 1296 Ocean and its biogeochemical significance. *Global Biogeochem. Cycles*, 30, doi:10.1002/2016GB005414, 2016
- 1297
- 1298 Beaufort, L., Probert, I., de Garidel-Thoron, T., Bendif, E.M., Ruiz-Pino, D., Metzl, N., Goyet, C., Buchet, N.,  
 1299 Coupel, P., Grelaud, M., Rost, B., Rickaby, R.E.M., and de Vargas C.: Sensitivity of coccolithophores to  
 1300 carbonate chemistry and ocean acidification. *Nature*, doi:10.1038/nature10295. 2011
- 1301
- 1302 Bittig, H.C., Steinhoff, T., Claustre, H., Fiedler, B., Williams, N.L., Sauzède, R., Körtzinger, A. and Gattuso, J.-  
 1303 P.: An Alternative to Static Climatologies: Robust Estimation of Open Ocean CO<sub>2</sub> Variables and Nutrient  
 1304 Concentrations From T, S, and O<sub>2</sub> Data Using Bayesian Neural Networks. *Front. Mar. Sci.* 5:328. doi:  
 1305 10.3389/fmars.2018.00328, 2018
- 1306
- 1307 Bockmon, E. E., and Dickson, A. G.,: An inter-laboratory comparison assessing the quality of seawater carbon  
 1308 dioxide measurements. *Marine Chemistry*, 171, 36-43, doi:10.1016/j.marchem.2015.02.002, 2015.
- 1309
- 1310 Bosse, A., Testor, P., Mayot, N., Prieur, L., D'Ortenzio, F., Mortier, L., Le Goff, H., Gourcuff, C., Coppola, L.,  
 1311 Lavigne, H. and Raimbault, P.: A submesoscale coherent vortex in the Ligurian Sea: From dynamical barriers to  
 1312 biological implications. *J. Geophys. Res. Oceans*, 122, doi:10.1002/2016JC012634., 2017.
- 1313
- 1314 Bozec, Y., Merlivat, L., Baudoux, A.-C., Beaumont, L., Blain, S., Bucciarelli, E., Danguy, T., Grossteffan, E.,  
 1315 Guillot, A., Guillou, J., Répécaud, M., and Tréguer, P.: Diurnal to inter-annual dynamics of pCO<sub>2</sub> recorded by a  
 1316 CARIOCA sensor in a temperate coastal ecosystem (2003–2009). *Marine Chemistry*, 126, 1-4, 13-26,  
 1317 10.1016/j.marchem.2011.03.003. 2011
- 1318
- 1319 Broullón, D., Pérez, F. F., Velo, A., Hoppema, M., Olsen, A., Takahashi, T., Key, R. M., Tanhua, T., González-  
 1320 Dávila, M., Jeansson, E., Kozyr, A., and van Heuven, S. M. A. C.: A global monthly climatology of total  
 1321 alkalinity: a neural network approach, *Earth Syst. Sci. Data*, 11, 1109–1127, <https://doi.org/10.5194/essd-11-1109-2019>. 2019
- 1322
- 1323 Broullón, D., Pérez, F. F., Velo, A., Hoppema, M., Olsen, A., Takahashi, T., Key, R. M., Tanhua, T., Santana-  
 1324 Casiano, J. M., and Kozyr, A.: A global monthly climatology of oceanic total dissolved inorganic carbon: a  
 1325 neural network approach, *Earth Syst. Sci. Data*, 12, 1725–1743, <https://doi.org/10.5194/essd-12-1725-2020>.  
 1326 2020
- 1327

- 1328  
1329 Bushinsky, S. M., Landschützer, P., Rödenbeck, C., Gray, A. R., Baker, D., Mazloff, M. R., Resplandy L.,  
1330 Johnson K. S., and Sarmiento, J. L.: Reassessing Southern Ocean air-sea CO<sub>2</sub> flux estimates with the addition of  
1331 biogeochemical float observations. *Global Biogeochemical Cycles*, 33. doi: 10.1029/2019GB006176, 2019.  
1332  
1333 Canesi, M., Douville, E., Montagna, P., Taviani, M., Stolarski, J., Bordier, L., Dapoigny, A., Coulibaly, G. E. H.,  
1334 Simon, A.-C., Agelou, M., Fin, J., Metzl, N., Iwankow, G., Allemand, D., Planes, S., Moulin, C., Lombard, F.,  
1335 Bourdin, G., Troublé, R., Agostini, S., Banaigs, B., Boissin, E., Boss, E., Bowler, C., de Vargas, C., Flores, M.,  
1336 Forcioli, D., Furla, P., Gilson, E., Galand, P. E., Pesant, S., Sunagawa, S., Thomas, O., Thurber, R. V., Voolstra,  
1337 C. R., Wincker, P., Zoccola, D., and Reynaud, S.: Differences in carbonate chemistry up-regulation of long-lived  
1338 reef-building corals. *Sci. Rep.* 13, 11589, Doi: 10.1038/s41598-023-37598-9, 2023.  
1339  
1340 Caniaux, G., Giordani, H., Redelsperger, J-L., Guichard, F., Key, E. and Wade, M.: Coupling between the  
1341 Atlantic cold tongue and the West African monsoon in boreal spring and summer. *J. Geophys. Res.*, 119,  
1342 C04003, doi:10.1029/2010JC006570., 2011.  
1343  
1344 Cariou, T., and Bozec, Y.: COMOR-CARBORHONE 1 cruise, RV L'Europe,  
1345 https://doi.org/10.17600/11060060, 2011a.  
1346  
1347 Cariou, T., and Bozec, Y.: COMOR-CARBORHONE 2 cruise, RV Téthys II,  
1348 https://doi.org/10.17600/11450150, 2011b.  
1349  
1350 Cariou, T., and Bozec, Y.: CARBORHONE 3 cruise, RV Téthys II, https://doi.org/10.17600/12450020, 2012a.  
1351  
1352 Cariou, T., and Bozec Y.: CARBORHONE 4 cruise, RV Téthys II, https://doi.org/10.17600/12450140, 2012b.  
1353  
1354 Carter, B. R., Williams, N. L., Gray, A. R., and Feely, R. A.: Locally interpolated alkalinity regression for global  
1355 alkalinity estimation, *Limnol. Oceanogr. Methods*, 14(4), 268–277, doi:10.1002/lom3.10087, 2016.  
1356  
1357 Chau, T.-T.-T., Gehlen, M., Metzl, N., and Chevallier, F.: CMEMS-LSCE: A global 0.25-degree, monthly  
1358 reconstruction of the surface ocean carbonate system, *Earth Syst. Sci. Data Discuss.* [preprint],  
1359 https://doi.org/10.5194/essd-2023-146, in review, 2023.  
1360  
1361 Chen, H., Haumann, F. A., Talley, L. D., Johnson, K. S., and Sarmiento, J. L.: The deep ocean's carbon exhaust.  
1362 *Global Biogeochemical Cycles*. doi: <https://doi.org/10.1002/essoar.10507757.1>, 2022  
1363  
1364 Cheng, L. J., Abraham, J., Zhu, J., Trenberth, K. E., Fasullo, J., Boyer, T., Locarnini, R., Zhang, B., Yu, F. J.,  
1365 Wan, L. Y., Chen, X. R., Song, X. Z., Liu, Y. L., and Mann, M. E.: Record-setting ocean warmth continued in  
1366 2019, *Adv. Atmos. Sci.* 37, 137-142. <https://doi.org/10.1007/s00376-020-9283-7>, 2020  
1367  
1368 Claustre, H., Johnson, K. S., and Takeshita, Y.: Observing the Global Ocean with Biogeochemical-Argo. *Annual  
1369 Review of Marine Science*, 12: 23-48 | DOI: [10.1146/annurev-marine-010419-010956](https://doi.org/10.1146/annurev-marine-010419-010956), 2020.  
1370  
1371 Conan, P., Guieux, A., and Vuillemin, R.: MOOSE (MOLA), https://doi.org/10.18142/234, 2020.  
1372  
1373 Copin-Montégut, C.: Alkalinity and carbon budgets in the Mediterranean Sea, *Global Biogeochemical Cycles*,  
1374 vol. 7, pp. 915–925, 1993.  
1375  
1376 Copin-Montégut, C., and Bégovic, M.: Distributions of carbonate properties and oxygen along the water column  
1377 (0–2000 m) in the central part of the NW Mediterranean Sea (Dyfamed site): influence of the winter vertical  
1378 mixing on air-sea CO<sub>2</sub> and O<sub>2</sub> exchanges. *Deep-Sea Research II* 49, 2049–2066, [https://doi.org/10.1016/S0967-0645\(02\)00027-9](https://doi.org/10.1016/S0967-0645(02)00027-9), 2002.  
1379  
1380 Coppola, L., and Diamond-Riquier, E.: MOOSE (DYFAMED), <https://doi.org/10.18142/131>, 2008.

- 1382  
1383 Coppola, L., Raimbault, P., Mortier, L., and Testor, P.: Monitoring the environment in the northwestern  
1384 Mediterranean Sea, Eos, 100, <https://doi.org/10.1029/2019EO125951>. Published on 25 July 2019.  
1385  
1386 Coppola, L., Boutin, J., Gattuso, J.-P., Lefèvre, D., and Metzl, N.: The Carbonate System in the Ligurian Sea. In  
1387 *The Mediterranean Sea in the Era of Global Change: Evidence from 30 years of multidisciplinary study of the*  
1388 *Ligurian Sea*, C. Migon, P. Nival, A. Sciandra, Eds. (ISTE Science Publishing LTD, London, UK, 2020), vol. 1,  
1389 chap. 4, pp. 79-104. ISBN: 9781786304285. <https://doi.org/10.1002/9781119706960.ch4>, 2020.  
1390  
1391 Coppola, L., Fourrier, M., Pasqueron de Fommervault, O., Poteau, A., Riquier, E. D. and Béguery, L.:  
1392 Highresolution study of the air-sea CO<sub>2</sub> flux and net community oxygen production in the Ligurian Sea by a  
1393 fleet of gliders. *Front. Mar. Sci.* 10:1233845. doi: 10.3389/fmars.2023.1233845, 2023  
1394  
1395 Corbière, A., Metzl, N., Reverdin, G., Brunet, C., and Takahashi, T.: Interannual and decadal variability of the  
1396 oceanic carbon sink in the North Atlantic subpolar gyre. *Tellus B*, Vol. 59, issue 2, 168-179, DOI:  
1397 10.1111/j.1600-0889.2006.00232, 2007.  
1398  
1399 D'Ortenzio, F. and Taillandier, V: BIO-ARGO-MED-2018 cruise, RV Téthys II,  
1400 <https://doi.org/10.17600/18000550>, 2018.  
1401  
1402 De Carlo, E. H., Mousseau, L., Passafiume, O., Drupp, P. and Gattuso J.-P.: Carbonate chemistry and air-sea  
1403 CO<sub>2</sub> flux in a NW Mediterranean bay over a four-year period: 2007-2011. *Aquatic Geochemistry*  
1404 doi:10.1007/s10498-013-9217-4, 2013.  
1405  
1406 Dickson, A. G., Sabine, C. L., and Christian, J. R.: Guide to best practices for ocean CO<sub>2</sub> measurements, North  
1407 Pacific Marine Science Organization, Sidney, British Columbia, 191, <https://doi.org/10.25607/OPB-1342>, 2007.  
1408  
1409 Division Plans de DMI – SHOM : PROTEUS2010\_LEG1 cruise, RV Pourquoi pas ?,  
1410 <https://doi.org/10.17600/10030040>, 2010.  
1411  
1412 Division Plans de DMI – SHOM: PROTEVSMED\_PERLE\_2018 cruise, RV L'Atalante,  
1413 <https://campagnes.flotteoceanographique.fr/campaign>, 2018.  
1414  
1415 Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A., Ocean acidification: The other CO<sub>2</sub> problem. *Annual*  
1416 *Review of Marine Science*, 1(1), 169–192. 10.1146/annurev.marine.010908.163834, 2009  
1417  
1418 Doney, S. C., Busch, D. S., Cooley, S. R., and Kroeker, K. J.: The Impacts of Ocean Acidification on Marine  
1419 Ecosystems and Reliant Human Communities. *Annual Review of Environment and Resources* 45:1,  
1420 <https://doi.org/10.1146/annurev-environ-012320-083019>. 2020  
1421  
1422 Douville, E., Bourdin, G., Lombard, F., Gorsky, G., Fin, J., Metzl, N., Pesant, S., and Tara Pacific Consortium:  
1423 Seawater carbonate chemistry dataset collected during the Tara Pacific Expedition 2016-2018. PANGAEA,  
1424 <https://doi.org/10.1594/PANGAEA.944420>, 2022.  
1425  
1426 Durrieu de Madron, X.: CASCADE cruise, RV L'Atalante, <https://doi.org/10.17600/11010020>, 2011.  
1427  
1428 Durrieu de Madron, X., and Conan, P.: PERLE2 cruise, RV Pourquoi pas ?, <https://doi.org/10.17600/18000865>,  
1429 2018  
1430  
1431 Edmond, J. M.: High precision determination of titration alkalinity and total carbon dioxide content of sea water  
1432 by potentiometric titration, *Deep-Sea Res.*, 17, 737–750, [https://doi.org/10.1016/0011-7471\(70\)90038-0](https://doi.org/10.1016/0011-7471(70)90038-0), 1970.  
1433  
1434 Eldin, G. : PANDORA cruise, RV L'Atalante, <https://doi.org/10.17600/12010050>, 2012.  
1435

- 1436 Eyring, V., Righi, M., Lauer, A., Evaldsson, M., Wenzel, S., Jones, C., Anav, A., Andrews, O., Cionni, I., Davin,  
 1437 E. L., Deser, C., Ehbrecht, C., Friedlingstein, P., Gleckler, P., Gottschaldt, K.-D., Hagemann, S., Juckes, M.,  
 1438 Kindermann, S., Krasting, J., Kunert, D., Levine, R., Loew, A., Mäkelä, J., Martin, G., Mason, E., Phillips, A. S.,  
 1439 Read, S., Rio, C., Roehrig, R., Senftleben, D., Sterl, A., van Ulft, L. H., Walton, J., Wang, S., and Williams, K.  
 1440 D.: ESMValTool (v1.0) – a community diagnostic and performance metrics tool for routine evaluation of Earth  
 1441 system models in CMIP, Geosci. Model Dev., 9, 1747–1802, doi:10.5194/gmd-9-1747-2016, 2016.
- 1442
- 1443 Fabry, V. J., Seibel, B. A., Feely, R. A. and Orr, J. C.: Impacts of ocean acidification on marine fauna and  
 1444 ecosystem processes. *ICES J. Mar. Sci.* 65, 414–432. <https://doi.org/10.1093/icesjms/fsn048>, 2008.
- 1445
- 1446 Fassbender, A. J., Sabine, C. L., and Palevsky, H. I.: Nonuniform ocean acidification and attenuation of the  
 1447 ocean carbon sink, *Geophys. Res. Lett.*, 44, 8404–8413, doi:10.1002/2017GL074389., 2017.
- 1448
- 1449 Fassbender, A. J., Alin, S. R., Feely, R. A., Sutton, A. J., Newton, J. A., Krembs, C., Bos, J., Keyzers, M., Devol,  
 1450 A., Ruef, W., and Pelletier, G.: Seasonal carbonate chemistry variability in marine surface waters of the US  
 1451 Pacific Northwest, *Earth Syst. Sci. Data*, 10, 1367–1401, <https://doi.org/10.5194/essd-10-1367-2018>, 2018.
- 1452
- 1453 Feely, R. A., Sabine, C. L., Byrne, R. H., Millero, F. J., Dickson, A. G., Wanninkhof, R., et al.: Decadal changes  
 1454 in the aragonite and calcite saturation state of the Pacific Ocean. *Global Biogeochemical Cycles*, 26, GB3001.  
 1455 <https://doi.org/10.1029/2011GB004157>, 2012.
- 1456
- 1457 Feely, R. A., Jiang, L.-Q., Wanninkhof, R., Carter, B. R., Alin, S. R., Bednaršek, N., and Cosca, C. E.: Acidification of the global surface ocean: What we have learned from observations. *Oceanography*,  
 1458 <https://doi.org/10.5670/oceanog.2023.222>, 2023
- 1460
- 1461 Fleury, E., Petton, S., Benabdelmouna, A., and Pouvreau, S., (coord.): Observatoire national du cycle de vie de  
 1462 l'huître creuse en France. Rapport annuel ECOSCPA 2022. R.INT.BREST RBE/PFOM/PI 2023-1, 2023.
- 1463
- 1464 Fourrier, M., Coppola, L., Claustre, H., D'Ortenzio, F., Sauzède, R. and Gattuso, J.-P.: A regional neural  
 1465 network approach to estimate water-column nutrient concentrations and carbonate system variables in the  
 1466 Mediterranean Sea: CANYON-MED. *Frontiers in Marine Science*, 7:620,  
 1467 <https://www.frontiersin.org/articles/10.3389/fmars.2020.00620>, 2020.
- 1468
- 1469 Fourrier, M., Coppola, L., D'Ortenzio, F., Migon, C., and Gattuso, J.-P.: Impact of intermittent convection in the  
 1470 northwestern Mediterranean Sea on oxygen content, nutrients, and the carbonate system. *Journal of Geophysical  
 1471 Research: Oceans*, 127, e2022JC018615. <https://doi.org/10.1029/2022JC018615>, 2022
- 1472
- 1473 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C., Luijkx, I.,  
 1474 T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P.,  
 1475 Jackson, R. B., Alin, S. R., Alkama, R., Arneth, A., Arora, V. K., Bates, N. R., Becker, M., Bellouin, N., Bittig,  
 1476 H. C., Bopp, L., Chevallier, F., Chini, L. P., Cronin, M., Evans, W., Falk, S., Feely, R. A., Gasser, T., Gehlen,  
 1477 M., Gkrizalis, T., Gloege, L., Grassi, G., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Houghton, R. A., Hurt, G. C., Iida, Y., Ilyina, T., Jain, A. K., Jersild, A., Kadono, K., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lindsay, K., Liu, J., Liu, Z., Marland, G., Mayot, N., McGrath, M. J., Metzl, N., Monacci, N. M., Munro, D. R., Nakao, S.-I., Niwa, Y., O'Brien, K., Ono, T., Palmer, P. I., Pan, N., Pierrot, D., Pocock, K., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Rodriguez, C., Rosan, T. M., Schwinger, J., Séférian, R., Shutler, J. D., Skjelvan, I., Steinhoff, T., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tanhua, T., Tans, P. P., Tian, X., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., Walker, A. P., Wanninkhof, R., Whitehead, C., Willstrand Wranne, A., Wright, R., Yuan, W., Yue, C., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2022, *Earth Syst. Sci. Data*, 14, 4811–4900, <https://doi.org/10.5194/essd-14-4811-2022>, 2022.
- 1477
- 1478 Fröb, F., Olsen, A., Becker, M., Chafik, L., Johannessen, T., Reverdin, G., and Omar, A.: Wintertime fCO<sub>2</sub>  
 1479 variability in the subpolar North Atlantic since 2004. *Geophysical Research Letters*, 46,  
 1490 <https://doi.org/10.1029/2018GL080554>, 2019.

- 1491  
1492 Gac, J-P., Marrec, P., Cariou, T., Guillerm, C., Macé, E., Vernet, M., and Bozec, Y.: Cardinal buoys: An  
1493 opportunity for the study of air-sea CO<sub>2</sub> fluxes in coastal ecosystems. *Front. Mar. Sci.* doi:  
1494 10.3389/fmars.2020.00712. 2020.  
1495  
1496 Gac, J.-P., Marrec, P., Cariou, T., Grossstefan, E., Macé, E., Rimmelin-Maury, P., Vernet, M., and Bozec, Y.:  
1497 Decadal Dynamics of the CO<sub>2</sub> System and Associated Ocean Acidification in Coastal Ecosystems of the North  
1498 East Atlantic Ocean. *Front. Mar. Sci.* 8:688008. doi:10.3389/fmars.2021.688008, 2021.  
1499  
1500 Ganachaud, A., Cravatte, S., Sprintall, J., Germinoaud, C., Alberty, M., Jeandel, C., Eldin, G., Metzl, N., Bonnet,  
1501 S., Benavides, M., Heimbürger, L.-E., Lefèvre, J., Michael, S., Resing, J., Queroué, F., Sarthou, G., Rodier, M.,  
1502 Berthelot, H., Baurand, F., Grelet, J., Hasegawa, T., Kessler, W., Kilepak, M., Lacan, F., Privat, E., Send, U.,  
1503 Van Beek, P., Souhaut, M. and Sonke, J. E.: The Solomon Sea: its circulation, chemistry, geochemistry and  
1504 biology explored during two oceanographic cruises. *Elem Sci Anth*, 5: 33, DOI:  
1505 <https://doi.org/10.1525/elementa.221>, 2017.  
1506  
1507 Gattuso, J.-P., Magnan, A., Billé, R., Cheung, W. W. L., Howes, E. L., Joos, F., Allemand, D., Bopp, L., Cooley,  
1508 S., Eakin, M., Hoegh-Guldberg, O., Kelly, R. P., Pörtner, H.-O., Rogers, A. D., Baxter, J. M., Laffoley, D.,  
1509 Osborn, D., Rankovic, A., Rochette, J., Sumaila, U. R., Treyer, S., and Turley, C.: Contrasting futures for ocean  
1510 and society from different anthropogenic CO<sub>2</sub> emissions scenarios. *Science* 349:aac4722.doi:  
1511 10.1126/science.aac4722, 2015.  
1512  
1513 Gattuso, J.-P., Alliouane, S., and Mousseau, L.: Seawater carbonate chemistry in the Bay of Villefranche, Point  
1514 B (France), January 2007 - September 2020. PANGAEA, <https://doi.org/10.1594/PANGAEA.727120>, 2021.  
1515  
1516 Gattuso, J.-P., Alliouane, S., and Fischer, P.: High-frequency, year-round time series of the carbonate chemistry  
1517 in a high-Arctic fjord (Svalbard), *Earth Syst. Sci. Data*, 15, 2809–2825, <https://doi.org/10.5194/essd-15-2809-2023>, 2023.  
1518  
1519  
1520 Gattuso, J.-P., Alliouane, S., and Fischer, P.: High-frequency, year-round time series of the carbonate chemistry  
1521 in a high-Arctic fjord (Svalbard) v2. PANGAEA, <https://doi.org/10.1594/PANGAEA.960131>, 2023.  
1522  
1523 Gemayel, E., Hassoun, A. E. R., Benallal, M. A., Goyet, C., Rivaro, P., Abboud-Abi Saab, M. , Krasakopoulou,  
1524 E., Touratier, F. , and Ziveri, P.: Climatological variations of total alkalinity and total inorganic carbon in the  
1525 Mediterranean Sea surface waters. *Earth Syst. Dynam.*, 6, 789-800, doi:10.5194/esd-6-789-2015. 2015.  
1526  
1527 Golbol, M., Vellucci, V., and Antoine, D.: BOUSSOLE, <https://doi.org/10.18142/1>, 2000.  
1528  
1529 Golbol M., Boutin J., Merlivat L., Vellucci, V., and Antoine, D.: Dissolved Inorganic Carbon and Total  
1530 Alkalinity sampled at Boussole site in the Mediterranean Sea. SEANOE. <https://doi.org/10.17882/71911>, 2020.  
1531  
1532 Goris, N., Tjiputra, J. F., Olsen, A., Schwinger, J., Lauvset, S. K. and Jeansson, E.: Constraining projection-  
1533 based estimates of the future North Atlantic carbon uptake, *J. Climate*, 31, 3959–3978,  
1534 <https://doi.org/10.1175/JCLI-D-17-0564.1>,2018.  
1535  
1536 Goyet, C., Beauverger, C., Brunet, C., and Poisson, A.: Distribution of carbon dioxide partial pressure in surface  
1537 waters of the Southwest Indian Ocean, *Tellus B: Chemical and Physical Meteorology*, 43:1, 1-11, DOI:  
1538 [10.3402/tellusb.v43i1.15242](https://doi.org/10.3402/tellusb.v43i1.15242), 1991.  
1539  
1540 Gregor, L. and Gruber, N.: OceanSODA-ETHZ: a global gridded data set of the surface ocean carbonate system  
1541 for seasonal to decadal studies of ocean acidification, *Earth Syst. Sci. Data*, 13, 777–808,  
1542 <https://doi.org/10.5194/essd-13-777-2021>, 2021.  
1543

- 1544 Gruber, N., Clement, D. , Carter, B. R., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Key, R. M.,  
1545 Kozyr, A., Lauvset, S. K., Lo Monaco, C. , Mathis, J. T., Murata, A., Olsen, A., Perez, F. F., Sabine, C. L.,  
1546 Tanhua, T., and Wanninkhof, R.: The oceanic sink for anthropogenic CO<sub>2</sub> from 1994 to 2007, Science vol. 363  
1547 (issue 6432), pp. 1193-1199. DOI: 10.1126/science.aau5153, 2019.
- 1548
- 1549 Guieu, C., Desboeufs, K., Albani, S., et al.: BIOGEOCHEMICAL dataset collected during the PEACETIME  
1550 cruise. SEANOE. <https://doi.org/10.17882/75747>, 2020.
- 1551
- 1552 Hagens, M., and Middelburg, J. J.: Attributing seasonal pH variability in surface ocean waters to governing  
1553 factors, Geophys. Res. Lett., 43, doi:10.1002/2016GL071719. 2016.
- 1554
- 1555 Henson, S. A., Painter, S. C., Holliday, N. P., Stinchcombe, M. C., and Giering, S. L. C.: Unusual subpolar  
1556 North Atlantic phytoplankton bloom in 2010: Volcanic fertilization or North Atlantic Oscillation?, J. Geophys.  
1557 Res. Oceans, 118, 4771–4780, doi:10.1002/jgrc.20363, 2013.
- 1558
- 1559 Hoegh-Guldberg, O., Mumby, P.J., Hooten, A.J., Steneck, R.S., Greenfield, P., Gomez, E., Harvell, C.D., Sale,  
1560 P.F., Edwards, A.J., Caldeira, K., Knowlton, N., Eakin, C.M., Iglesias-Prieto, R., Muthiga, N., Bradbury, R.H.,  
1561 Dubi, A., and Hatziolos, M.E.: Coral reefs under rapid climate change and ocean acidification. Science 14,  
1562 1737–1742. <https://doi.org/10.1126/science.1152509>, 2007.
- 1563
- 1564 Hood, E.M., and Merlivat, L.: Annual to interannual variations of fCO(2) in the northwestern Mediterranean  
1565 Sea: Results from hourly measurements made by CARIOCA buoys, 1995-1997, Journal Of Marine Research,  
1566 59, 113-131, doi: 10.1357/002224001321237399. 2001
- 1567
- 1568 Howes, E., Stemmann, L., Assailly, C., Irisson, J.-O., Dima, M., Bijma, J., Gattuso, J.-P.: Pteropod time series  
1569 from the North Western Mediterranean (1967-2003): impacts of pH and climate variability. Mar Ecol Prog Ser  
1570 531: 193-206, doi: 10.3354/meps11322. 2015.
- 1571
- 1572 Howes, E. L., Eagle, R., Gattuso, J.-P., and Bijma, J.: Comparison of Mediterranean pteropod shell biometrics  
1573 and ultrastructure from historical (1910 and 1921) and present day (2012) samples provides baseline for  
1574 monitoring effects of global change. *PLoS ONE* 12:e0167891. 2017.
- 1575
- 1576 IPCC. Changing Ocean, Marine Ecosystems, and Dependent Communities. in The Ocean and Cryosphere in a  
1577 Changing Climate 447–588 (Cambridge University Press, 2022). doi:10.1017/9781009157964.007. 2022
- 1578
- 1579 Jiang, Z.-P., Tyrrell, T., Hydes, D. J., Dai, M., and Hartman, S. E.: Variability of alkalinity and the alkalinity-  
1580 salinity relationship in the tropical and subtropical surface ocean, Global Biogeochem. Cycles, 28, 729–742,  
1581 doi:10.1002/2013GB004678, 2014.
- 1582
- 1583 Jiang, L.-Q., Feely, R. A., Carter, B. R., Greeley, D. J., Gledhill, D. K., and Arzayus K. M.: Climatological  
1584 distribution of aragonite saturation state in the global oceans, Global Biogeochem. Cycles, 29, 1656–1673,  
1585 doi:10.1002/2015GB005198, 2015.
- 1586
- 1587 Jiang, L.-Q., Carter, B. R., Feely, R. A., Lauvset, S. K. and Olsen, A.: Surface ocean pH and buffer capacity:  
1588 past, present and future, *Sci Rep*, 9(1), 1–11, doi:10.1038/s41598-019-55039-4. 2019.
- 1589
- 1590 Jiang, L.-Q., Feely, R. A., Wanninkhof, R., Greeley, D., Barbero, L., Alin, S., Carter, B. R., Pierrot, D.,  
1591 Featherstone, C., Hooper, J., Melrose, C., Monacci, N., Sharp, J. D., Shellito, S., Xu, Y.-Y., Kozyr, A., Byrne, R.  
1592 H., Cai, W.-J., Cross, J., Johnson, G. C., Hales, B., Langdon, C., Mathis, J., Salisbury, J., and Townsend, D. W.:  
1593 Coastal Ocean Data Analysis Product in North America (CODAP-NA) – an internally consistent data product for  
1594 discrete inorganic carbon, oxygen, and nutrients on the North American ocean margins. Earth System Science  
1595 Data, 13(6), 2777–2799. <https://doi.org/10.5194/essd-13-2777-2021>, 2021
- 1596

- 1597 Jiang, L.-Q., Dunne, J., Carter, B. R., Tjiputra, J. F., Terhaar, J., Sharp, J. D., et al.: Global surface ocean  
1598 acidification indicators from 1750 to 2100. *Journal of Advances in Modeling Earth Systems*, 15,  
1599 e2022MS003563. <https://doi.org/10.1029/2022MS003563>, 2023a
- 1600
- 1601 Jiang, L.Q., Kozyr, A., Relph, J.M. *et al.* The Ocean Carbon and Acidification Data System. *Sci Data* 10, 136.  
1602 <https://doi.org/10.1038/s41597-023-02042-0>, 2023b
- 1603
- 1604 Johnson, K. S., Mazloff, M. R., Bif, M. B., Takeshita, Y., Jannasch, H. W., Maurer, T. L., et al.: Carbon to  
1605 nitrogen uptake ratios observed across the Southern Ocean by the SOCCOM profiling float array. *Journal of*  
1606 *Geophysical Research: Oceans*, 127, e2022JC018859. <https://doi.org/10.1029/2022JC018859>, 2022.
- 1607
- 1608 Joos, F., Hameau, A., Frölicher, T. L., and Stephenson, D. B.: Anthropogenic attribution of the  
1609 increasing seasonal amplitude in surface ocean pCO<sub>2</sub>. *Geophysical Research Letters*, 50, e2023GL102857.  
1610 <https://doi.org/10.1029/2023GL102857>, 2023.
- 1611
- 1612 Joyce, T. and Corry, C., eds: Requirements for WOCE Hydrographic Programme Data Reporting. WHPO  
1613 Publication 90-1 Revision 2, WOCE Report 67/91, Woods Hole, Mass., USA, May 1994.
- 1614
- 1615 Kapsenberg, L., Alliouane, S., Gazeau, F., Mousseau, L., and Gattuso, J.-P.: Coastal ocean acidification and  
1616 increasing total alkalinity in the northwestern Mediterranean Sea, *Ocean Sci.*, 13, 411-426, doi:10.5194/os-13-  
1617 411-2017, 2017.
- 1618
- 1619 Keppler, L., Landschützer, P., Gruber, N., Lauvset, S. K., and Stemmler, I.: Seasonal carbon dynamics in the  
1620 near-global ocean. *Global Biogeochemical Cycles*, 34, e2020GB006571. doi:10.1029/2020GB006571, 2020.
- 1621
- 1622 Keppler, L., Landschützer, P., Lauvset, S.K., and Gruber, N.: Recent trends and variability in the oceanic storage  
1623 of dissolved inorganic carbon. *Global Biogeochemical Cycles*, 37, e2022GB007677. Doi:  
1624 10.1029/2022GB007677, 2023.
- 1625
- 1626 Keraghel, M. A., Louanchi, F., Zerrouki, M., Kaci, M. A., Aït-Ameur, N., Labaste, M., Legoff, H., Taillandier,  
1627 V., Harid, R., and Mortier, L.: Carbonate system properties and anthropogenic carbon inventory in the Algerian  
1628 Basin during SOMBA cruise (2014): Acidification estimate, *Marine Chemistry*,  
1629 <https://doi.org/10.1016/j.marchem.2020.103783>. 2020.
- 1630
- 1631 Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J.,  
1632 Mordy, C., and Peng, T. H.: A global ocean carbon climatology: Results from Global Data Analysis Project  
1633 (GLODAP), *Global Biogeochemical Cycles*, 18, GB4031, <https://doi.org/10.1029/2004GB002247>, 2004.
- 1634
- 1635 Key, R. M., Tanhua, T., Olsen, A., Hoppema, M., Jutterström, S., Schirnick, C., van Heuven, S., Kozyr, A., Lin,  
1636 X., Velo, A., Wallace, D. W. R., and Mintrop, L.: The CARINA data synthesis project: introduction and  
1637 overview, *Earth Syst. Sci. Data*, 2, 105–121, <https://doi.org/10.5194/essd-2-105-2010>, 2010.
- 1638
- 1639 Khatiwala, S., Tanhua, T., Mikaloff Fletcher, S., Gerber, M., Doney, S. C., Graven, H. D., Gruber, N.,  
1640 McKinley, G. A., Murata, A., Ríos, A. F., and Sabine, C. L.: Global ocean storage of anthropogenic carbon,  
1641 *Biogeosciences*, 10, 2169-2191, <https://doi.org/10.5194/bg-10-2169-2013>, 2013.
- 1642
- 1643 Kitidis, V., Shutler, J. D., Ashton, I., Warren, M., Brown, I., Findlay, H., Hartman, S. E., Sanders, R.,  
1644 Humphreys, M., Kivimäe, C., Greenwood, N., Hull, T., Pearce, D., McGrath, T., Stewart, B. M., Walsham, P.,  
1645 McGovern, E., Bozec, Y., Gac, J.-P., van Heuven, S., Hoppema, M., Schuster, U., Johannessen, T., Omar, A.,  
1646 Lauvset, S. K., Skjelvan, I., Olsen, A., Steinhoff, T., Kötzinger, A., Becker, M., Lefèvre, N., Diverrès, D.,  
1647 Gkrizalis, T., Cattrijssse, A., Petersen, W., Voynova, Y., Chapron, B., Grouazel, A., Land, P. E., Sharples, J., and  
1648 Nightingale, P. D.: Winter weather controls net influx of atmospheric CO<sub>2</sub> on the north-west European shelf. *Sci*  
1649 *Rep* 9, 20153, doi:10.1038/s41598-019-56363-5. 2019.
- 1650

- 1651 Koffi U., Lefevre, N., Kouadio, G., and Boutin, J.: Surface CO<sub>2</sub> parameters and air-sea CO<sub>2</sub> fluxes distribution  
1652 in the eastern equatorial Altantic Ocean. *J. Marine Systems*, doi:10.1016/j.jmarsys/2010.04.010. 2010.
- 1653
- 1654 Koffi, U., Georges, K., and Boutin, J.: Partial pressure (or fugacity) of carbon dioxide, dissolved inorganic  
1655 carbon, alkalinity, temperature, salinity and other variables collected from Surface underway, discrete sample  
1656 and profile observations using CTD, Carbon dioxide (CO<sub>2</sub>) gas analyzer and other instruments from ANTEA  
1657 and L'ATALANTE in the Gulf of Guinea, North Atlantic Ocean and South Atlantic Ocean from 2005-06-09 to  
1658 2007-09-30 (NCEI Accession 0108086). [indicate subset used]. NOAA National Centers for Environmental  
1659 Information. Dataset. [https://doi.org/10.3334/cdiac/otg.egee1\\_6](https://doi.org/10.3334/cdiac/otg.egee1_6). Accessed [date]., 2013
- 1660
- 1661 Koseki, S., Tjiputra, J., Fransner, F. et al.: Disentangling the impact of Atlantic Niño on sea-air CO<sub>2</sub> flux. *Nat*  
1662 *Commun* 14, 3649. <https://doi.org/10.1038/s41467-023-38718-9>, 2023.
- 1663
- 1664 Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J. R., Dunne, J. P., Gehlen, M.,  
1665 Ilyina, T., John, J. G., Lenton, A., Li, H., Lovenduski, N. S., Orr, J. C., Palmieri, J., Santana-Falcón, Y.,  
1666 Swinger, J., Séférian, R., Stock, C. A., Tagliabue, A., Takano, Y., Tjiputra, J., Toyama, K., Tsujino, H.,  
1667 Watanabe, M., Yamamoto, A., Yool, A., and Ziehn, T.: Twenty-first century ocean warming, acidification,  
1668 deoxygenation, and upper-ocean nutrient and primary production decline from CMIP6 model projections,  
1669 *Biogeosciences*, 17, 3439–3470, <https://doi.org/10.5194/bg-17-3439-2020>, 2020.
- 1670
- 1671 Land, P. E., Findlay, H. S., Shutler, J. D., Ashton, I. G., Holding, T., Grouazel, A., Girard-Ardhuin, F., Reul, N.,  
1672 Piolle, J. F., Chapron, B., and Quilfen, Y.: Optimum satellite remote sensing of the marine carbonate system  
1673 using empirical algorithms in the global ocean, the Greater Caribbean, the Amazon Plume and the Bay of  
1674 Bengal. *Remote Sensing of Environment*, 235, p.111469, doi: 10.1016/j.rse.2019.111469, 2019.
- 1675
- 1676 Lange, N., Fiedler, B., Álvarez, M., Benoit-Cattin, A., Benway, H., Buttigieg, P. L., Coppola, L., Currie, K.,  
1677 Flecha, S., Honda, M., Huertas, I. E., Lauvset, S. K., Muller-Karger, F., Körtzinger, A., O'Brien, K. M.,  
1678 Ólafsdóttir, S. R., Pacheco, F. C., Rueda-Roa, D., Skjelvan, I., Wakita, M., White, A., and Tanhua, T.: Synthesis  
1679 Product for Ocean Time-Series (SPOTS) – A ship-based biogeochemical pilot, *Earth Syst. Sci. Data Discuss.*  
1680 [preprint], <https://doi.org/10.5194/essd-2023-238>, in review, 2023.
- 1681
- 1682 Lauvset, S. K., Gruber, N., Landschützer, P., Olsen, A., and Tjiputra, J.: Trends and drivers in global surface  
1683 ocean pH over the past 3 decades. *Biogeosciences*, 12, 1285-1298, doi:10.5194/bg-12-1285-2015, 2015
- 1684
- 1685 Lauvset, S. K., Carter, B. R., Perez, F. F., Jiang, L.-Q., Feely, R. A., Velo, A., and Olsen, A.: Processes Driving  
1686 Global Interior Ocean pH Distribution, *Global Biogeochem. Cycles*, 34, e2019GB006 229,  
1687 <https://doi.org/10.1029/2019GB006229>, 2020.
- 1688
- 1689 Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Álvarez, M., Becker, S., Brown, P. J.,  
1690 Carter, B. R., Cotrim da Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Jeansson, E.,  
1691 Jutterström, S., Jones, S. D., Karlsen, M. K., Lo Monaco, C., Michaelis, P., Murata, A., Pérez, F. F., Pfeil, B.,  
1692 Schirnick, C., Steinfeldt, R., Suzuki, T., Tilbrook, B., Velo, A., Wanninkhof, R., Woosley, R. J., and Key, R. M.:  
1693 An updated version of the global interior ocean biogeochemical data product, GLODAPv2.2021, *Earth Syst. Sci.*  
1694 *Data*, 13, 5565–5589, <https://doi.org/10.5194/essd-13-5565-2021>, 2021.
- 1695
- 1696 Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Alin, S., Álvarez, M., Azetsu-Scott,  
1697 K., Barbero, L., Becker, S., Brown, P. J., Carter, B. R., da Cunha, L. C., Feely, R. A., Hoppema, M., Humphreys,  
1698 M. P., Ishii, M., Jeansson, E., Jiang, L.-Q., Jones, S. D., Lo Monaco, C., Murata, A., Müller, J. D., Pérez, F. F.,  
1699 Pfeil, B., Schirnick, C., Steinfeldt, R., Suzuki, T., Tilbrook, B., Ulfsbo, A., Velo, A., Woosley, R. J., and Key, R.  
1700 M.: GLODAPv2.2022: the latest version of the global interior ocean biogeochemical data product, *Earth Syst.*  
1701 *Sci. Data*, 14, 5543–5572, <https://doi.org/10.5194/essd-14-5543-2022>, 2022.
- 1702
- 1703 Lebehot, A. D., Halloran, P. R., Watson, A. J., McNeill, D., Ford, D. A., Landschützer, P., Lauvset, S. K.,  
1704 and Schuster, U.: Reconciling Observation and Model Trends in North Atlantic Surface CO<sub>2</sub>, *Global*  
1705 *Biogeochem. Cy.*, 33, 1204–1222, <https://doi.org/10.1029/2019GB006186>, 2019.

- 1706  
1707 Lee, K., Wanninkhof, R., Feely, R. A., Millero, F. J., and Peng, T.-H.: Global relationships of total inorganic  
1708 carbon with temperature and nitrate in surface seawater, *Global Biogeochem. Cy.*, 14, 979–994,  
1709 https://doi.org/10.1029/1998GB001087, 2000.  
1710  
1711 Lee, K., Tong, L.T., Millero, F.J., Sabine, C.L., Dickson, A.G., Goyet, C., Park, G.H., Wanninkhof, R., Feely,  
1712 R.A., and Key, R.M.: Global relationships of total alkalinity with salinity and temperature in surface waters of  
1713 the world's oceans. *Geophys. Res. Lett.* 33, L19605. doi10.1029/2006GL027207. 2006.  
1714  
1715 Lefèvre, D.: MOOSE (ANTARES), <https://doi.org/10.18142/233>, 2010.  
1716  
1717 Lefèvre, N., Guillot, A., Beaumont, L, and Danguy, T.: Variability of fCO<sub>2</sub> in the Eastern Tropical Atlantic  
1718 from a moored buoy. *J. of Geophysical Research-Oceans*, Volume: 113 Issue: C1, DOI:  
1719 10.1029/2007JC004146. 2008.  
1720  
1721 Lefèvre N., and Merlivat, L.: Carbon and oxygen net community production in the eastern tropical Atlantic  
1722 estimated from a moored buoy. *Global biogeochemical cycles*, 26(1), 1-14.  
1723 https://doi.org/10.1029/2010GB004018. 2012.  
1724  
1725 Lefèvre, N., Diverres, D., and Gallois, F.: Origin of CO<sub>2</sub> undersaturation in the western tropical Atlantic: Tellus  
1726 Series B Chemical and Physical Meteorology. Volume: 62 Issue: 5 Special Issue: SI Pages: 595-607 DOI:  
1727 10.1111/j.1600-0889.2010.00475.x, 2010.  
1728  
1729 Lefèvre N.: Carbon parameters along a zonal transect. SEANOE. <https://doi.org/10.17882/58575>, 2010.  
1730  
1731 Lefèvre, N., Veleda, D., Araujo, M., Caniaux, G.: Variability and trends of carbon parameters at a time series in  
1732 the eastern tropical Atlantic. *Tellus B*, Co-Action Publishing, 68, pp.30305. 10.3402/tellusb.v68.30305. 2016.  
1733  
1734 Lefèvre, N., Mejia, C., Khvorostyanov, D., Beaumont, L., and Koffi, U.: Ocean Circulation Drives the  
1735 Variability of the Carbon System in the Eastern Tropical Atlantic. *Oceans*, 2021, 2, 126–148.  
1736 <https://doi.org/10.3390/oceans2010008>, 2021.  
1737  
1738 Lefèvre, N.: Discrete profile measurements of dissolved inorganic carbon, total alkalinity, temperature and  
1739 salinity collected from R/V Antea French PIRATA cruise in Eastern Tropical Atlantic Ocean from 2009-07-10  
1740 to 2010-10-01 (NCEI Accession 0171193), 2018a.  
1741  
1742 Lefèvre, N.: Discrete surface measurements of dissolved inorganic carbon, total alkalinity, temperature and  
1743 salinity collected from R/V Le Suroit French PIRATA cruise in Eastern Tropical Atlantic Ocean from 2011-05-  
1744 03 to 2011-05-25 (NCEI Accession 0171197), 2018b.  
1745  
1746 Lefèvre, N.: Discrete profile measurements of dissolved inorganic carbon, total alkalinity, temperature and  
1747 salinity collected from R/V Le Suroit French PIRATA cruise in Eastern Tropical Atlantic Ocean from 2012-03-  
1748 21 to 2012-04-30 (NCEI Accession 0171195), 2018c.  
1749  
1750 Lefèvre, N.: Discrete surface measurements of dissolved inorganic carbon, total alkalinity, temperature and  
1751 salinity collected from R/V Le Suroit French PIRATA cruise in Eastern Tropical Atlantic Ocean from 2013-05-  
1752 11 to 2013-06-18 (NCEI Accession 0171189), 2018d.  
1753  
1754 Lefèvre, N.: Discrete profile measurements of dissolved inorganic carbon, total alkalinity, temperature and  
1755 salinity collected from R/V Le Suroit French PIRATA cruise in Eastern Tropical Atlantic Ocean from 2014-04-  
1756 10 to 2014-05-19 (NCEI Accession 0171194), 2018e.  
1757  
1758 Lefèvre, N.: Discrete surface measurements of dissolved inorganic carbon, total alkalinity, temperature and  
1759 salinity collected from R/V Thalassa French PIRATA cruise in Eastern Tropical Atlantic Ocean from 2015-03-  
1760 18 to 2015-04-15 (NCEI Accession 0171196), 2018f.  
1761

- 1762 Lefèvre, N.: Discrete surface measurements of dissolved inorganic carbon, total alkalinity, temperature and  
1763 salinity collected from R/V Thalassa French PIRATA cruise in Eastern Tropical Atlantic Ocean from 2016-03-  
1764 08 to 2016-04-11 (NCEI Accession 0171190), 2018g.
- 1765 Lefèvre, N.: Discrete surface measurements of dissolved inorganic carbon, total alkalinity, temperature and  
1766 salinity collected from R/V Thalassa French PIRATA cruise in Eastern Tropical Atlantic Ocean from 2017-02-  
1767 26 to 2017-03-30 (NCEI Accession 0171191), 2018h.
- 1769 Le Quéré, C., Moriarty, R., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Friedlingstein, P., Peters,  
1770 G. P., Andres, R. J., Boden, T. A., Houghton, R. A., House, J. I., Keeling, R. F., Tans, P., Arneth, A., Bakker, D.  
1771 C. E., Barbero, L., Bopp, L., Chang, J., Chevallier, F., Chini, L. P., Ciais, P., Fader, M., Feely, R. A., Gkritzalis,  
1772 T., Harris, I., Hauck, J., Ilyina, T., Jain, A. K., Kato, E., Kitidis, V., Klein Goldewijk, K., Koven, C.,  
1773 Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lima, I. D., Metzl, N., Millero, F., Munro, D. R.,  
1774 Murata, A., Nabel, J. E. M. S., Nakaoka, S., Nojiri, Y., O'Brien, K., Olsen, A., Ono, T., Pérez, F. F., Pfeil, B.,  
1775 Pierrot, D., Poulter, B., Rehder, G., Rödenbeck, C., Saito, S., Schuster, U., Schwinger, J., Séférian, R., Steinhoff,  
1776 T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van der Laan-Luijkx, I. T., van der Werf, G. R., van  
1777 Heuven, S., Vandemark, D., Viovy, N., Wiltshire, A., Zaehle, S., and Zeng, N.: Global Carbon Budget 2015,  
1778 Earth Syst. Sci. Data, 7, 349–396, <https://doi.org/10.5194/essd-7-349-2015>, 2015.
- 1780 Lerner, P., Romanou, A., Kelley, M., Romanski, J., Ruedy, R., and Russell, G.: Drivers of Air-Sea CO<sub>2</sub> Flux  
1781 Seasonality and its Long-Term Changes in the NASA-GISS model CMIP6 submission. Journal of Advances in  
1782 Modeling Earth Systems, 13, e2019MS002028. [Doi:10.1029/2019MS002028](https://doi.org/10.1029/2019MS002028), 2021.
- 1784 Leseurre, C., Lo Monaco, C., Reverdin, G., Metzl, N., Fin, J., Olafsdottir, S., and Racapé, V.: Ocean carbonate  
1785 system variability in the North Atlantic Subpolar surface water (1993–2017), Biogeosciences, 17, 2553–2577,  
1786 <https://doi.org/10.5194/bg-17-2553-2020>, 2020
- 1788 Leseurre, C., Lo Monaco, C., Reverdin, G., Metzl, N., Fin, J., Mignon, C., and Benito, L.: Summer trends and  
1789 drivers of sea surface fCO<sub>2</sub> and pH changes observed in the southern Indian Ocean over the last two decades  
1790 (1998–2019), Biogeosciences, 19, 2599–2625, <https://doi.org/10.5194/bg-19-2599-2022>, 2022.
- 1792 Lherminier, P., Mercier, H., Gourcuff, C., Alvarez, M., Bacon, S., and Kermabon, C.: Transports across the 2002  
1793 Greenland-Portugal OVIDE section and comparison with 1997. J. Geophys. Res., 112(C7), C07003,  
1794 [doi:10.1029/2006JC003716](https://doi.org/10.1029/2006JC003716). 2007
- 1796 Li, B. F., Watanabe, Y. W., Hosoda, S., Sato, K., and Nakano, Y.: Quasireal-time and high-resolution  
1797 spatiotemporal distribution of ocean anthropogenic CO<sub>2</sub>. Geophysical Research Letters, 46, 4836–4843.  
1798 <https://doi.org/10.1029/2018GL081639>, 2019.
- 1800 Lo Monaco, C., Álvarez, M., Key, R. M., Lin, X., Tanhua, T., Tilbrook, B., Bakker, D. C. E., van Heuven, S.,  
1801 Hoppema, M., Metzl, N., Ríos, A. F., Sabine, C. L., and Velo, A.: Assessing the internal consistency of the  
1802 CARINA database in the Indian sector of the Southern Ocean, Earth Syst. Sci. Data, 2, 51–70,  
1803 <https://doi.org/10.5194/essd-2-51-2010>, 2010.
- 1805 Lo Monaco, C., Metzl, N., Fin, J., and Tribollet, A.: Sea surface measurements of dissolved inorganic carbon  
1806 (DIC) and total alkalinity (TALK), temperature and salinity during the R/V Marion-Dufresne cruise CLIM-  
1807 EPARSES (EXPOCODE 35MV20190405) in the Indian Ocean and Mozambique Channel from 2019-04-04 to  
1808 2019-04-30. (NCEI Accession 0212218). [indicate subset used]. NOAA National Centers for Environmental  
1809 Information. Dataset. <https://doi.org/10.25921/26rw-w185>. Accessed [date]. 2020.
- 1811 Lo Monaco, C., Metzl, N., Fin, J., Mignon, C., Cuet, P., Douville, E., Gehlen, M., Trang Chau, T.T., and  
1812 Tribollet, A.: Distribution and long-term change of the sea surface carbonate system in the Mozambique Channel  
1813 (1963-2019), Deep-Sea Research Part II, <https://doi.org/10.1016/j.dsr2.2021.104936>, 2021
- 1815

- 1816 Lombard, F., Bourdin, G. , Pesant, S., Agostini, S., Baudena, A., Boissin, E., Cassar, N., Clampitt, M., Conan,  
1817 P., Da Silva, O., Dimier, C., Douville, E., Elineau, A., Fin, J., Flores, J.-M., Ghiglione, J.-F., Hume, B. C. C.,  
1818 Jalabert, L., John, S. G., Kelly, R. L., Koren, I., Lin, Y., Marie, D., McMinds, R., Mériguet, Z., Metzl, N., Paz-  
1819 García, D. A., Luiza Pedrotti, M., Poulain, J., Pujo-Pay, M., Ras, J., Reverdin, G., Romac, S., Röttinger, E.,  
1820 Vardi, A., Voolstra, C. R., Moulin, C., Iwankow, G., Banaigs, B., Bowler, C., de Vargas, C., Forcioli, D., Furla,  
1821 P., Galand, P. E., Gilson, E., Reynaud, S., Sunagawa, S., Thomas, O., Troublé, R., Vega Thurber, R., Wincker,  
1822 P., Zoccola, D., Allemand, D., Planes, S., Boss, E., and Gorsky, G.: Open science resources from the Tara  
1823 Pacific expedition across the surface ocean and coral reef ecosystems. *Sci Data* 10, 324 (2023).  
1824 <https://doi.org/10.1038/s41597-022-01757-w>, 2022.
- 1825
- 1826 Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean pCO<sub>2</sub> calculated from dissolved inorganic carbon,  
1827 alkalinity, and equations for K-1 and K-2: validation based on laboratory measurements of CO<sub>2</sub> in gas and  
1828 seawater at equilibrium. *Marine Chemistry* 70, 105-119. [https://doi.org/10.1016/S0304-4203\(00\)00022-0](https://doi.org/10.1016/S0304-4203(00)00022-0), 2000.
- 1829
- 1830 Ma, D., Gregor, L., and Gruber, N: Four decades of trends and drivers of global surface ocean acidification.  
1831 *Global Biogeochemical Cycles*, 37, e2023GB007765. [10.1029/2023GB007765](https://doi.org/10.1029/2023GB007765), 2023.
- 1832
- 1833 Maier, C., Watremez, P., Taviani, M., Weinbauer, M. G, and Gattuso, J-P.: Calcification rates and the effect of  
1834 ocean acidification on Mediterranean cold-water corals. *Proceedings of the Royal Society B-Biological Sciences*,  
1835 279(1734), 1716-1723, doi:10.1098/rspb.2011.1763. 2012.
- 1836
- 1837 Margirier, F., Testor, P., Heslop, E. et al. : Abrupt warming and salinification of intermediate waters interplays  
1838 with decline of deep convection in the Northwestern Mediterranean Sea. *Sci Rep* 10, 20923. [10.1038/s41598-020-77859-5](https://doi.org/10.1038/s41598-020-77859-5), 2020.
- 1840
- 1841 Marrec, P., Cariou, T., Collin, E., Durand, A., Latimier, M., Macé, E., Morin, P., Raimund, S., Vernet, M., and  
1842 Bozec, Y.: Seasonal and latitudinal variability of the CO<sub>2</sub> system in the western English Channel based on  
1843 Voluntary Observing Ship (VOS) measurements. *Marine Chemistry*, 155 (2013): 29–41. 2013.
- 1844
- 1845 Marrec, P., Cariou, T., Latimier, M., Macé, E., Morin, P., Vernet, M., and Bozec, Y.: Spatio-temporal dynamics  
1846 of biogeochemical processes and air-sea CO<sub>2</sub> fluxes in the Western English Channel based on two years of  
1847 FerryBox deployment. *Journal of Marine Systems*, Special Issue: 5<sup>th</sup> FerryBox Workshop.  
1848 [10.1016/j.jmarsys.2014.05.010](https://doi.org/10.1016/j.jmarsys.2014.05.010). 2014.
- 1849
- 1850 Marrec, P., Cariou, T., Macé, E., Morin, P., Salt, L. A., Vernet, M., Taylor, B., Paxman, K., and Y. Bozec:  
1851 Dynamics of air-sea CO<sub>2</sub> fluxes in the northwestern European shelf based on voluntary observing ship and  
1852 satellite observations, *Biogeosciences*, 12, 5371-5391, doi:10.5194/bg-12-5371-2015. 2015
- 1853
- 1854 Marrec, P., and Bozec, Y.: Partial pressure (or fugacity) of carbon dioxide, dissolved inorganic carbon, alkalinity  
1855 and salinity collected from Surface underway observations using Carbon dioxide (CO<sub>2</sub>) gas analyzer and other  
1856 instruments from ARMORIQUE in the English Channel from 2012-04-25 to 2013-01-03 (NCEI Accession  
1857 0157472). Version 1.1. NOAA National Centers for Environmental Information. Dataset.  
1858 doi:10.3334/CDIAC/OTG.COAST\_FERRYBOX\_ROSCOFF\_PLYMOUTH\_2012 [access date]. 2016a.
- 1859
- 1860 Marrec, P., and Bozec, Y.: Partial pressure (or fugacity) of carbon dioxide, dissolved inorganic carbon, alkalinity  
1861 and salinity collected from Surface underway observations using Carbon dioxide (CO<sub>2</sub>) gas analyzer and other  
1862 instruments from ARMORIQUE in the English Channel from 2013-03-15 to 2013-12-22 (NCEI Accession  
1863 0157444). Version 1.1. NOAA National Centers for Environmental Information. Dataset.  
1864 doi:10.3334/CDIAC/OTG.COAST\_FERRYBOX\_ROSCOFF\_PLYMOUTH\_2013 [access date]. 2016b.
- 1865
- 1866 Marrec, P., and Bozec, Y.: Partial pressure (or fugacity) of carbon dioxide, dissolved inorganic carbon, alkalinity  
1867 and salinity collected from surface underway observations using Carbon dioxide (CO<sub>2</sub>) gas analyzer and other  
1868 instruments from ARMORIQUE in the English Channel from 2014-03-18 to 2014-10-09 (NCEI Accession  
1869 0163193). Version 1.1. NOAA National Centers for Environmental Information. Dataset.  
1870 doi:10.3334/CDIAC/OTG.COAST\_FERRYBOX\_ROSCOFF\_PLYMOUTH\_2014 [access date]. 2017.

- 1871  
1872 Mazloff, M. R., Verdy, A., Gille, S. T., Johnson, K. S., Cornuelle, B. D., and Sarmiento, J.: Southern Ocean  
1873 acidification revealed by biogeochemical-Argo floats. *Journal of Geophysical Research: Oceans*, 128,  
1874 e2022JC019530. <https://doi.org/10.1029/2022JC019530>, 2023.
- 1875  
1876 McCulloch, M., Trotter, J., Montagna, P., Falter, J., Dunbar, R., Freiwald, A., Försterra, G., López Correa, M.,  
1877 Maier, C., Rüggeberg, A., and Taviani, M.: Resilience of cold-water scleractinian corals to ocean acidification:  
1878 Boron isotopic systematics of pH and saturation state up-regulation. *Geochimica et Cosmochimica Acta*, Volume  
1879 87, 21-34. <http://dx.doi.org/10.1016/j.gca.2012.03.027>. 2012
- 1880  
1881 McKinley, G. A., Fay, A. R., Takahashi, T., and Metzl, N.: Convergence of atmospheric and North Atlantic  
1882 carbon dioxide trends on multidecadal timescales. *Nature Geoscience*. doi:10.1038/NGEO1193. 2011.
- 1883  
1884 McKinley, G. A., Ritzer, A. L. and Lovenduski, N. S.: Mechanisms of northern North Atlantic biomass  
1885 variability, *Biogeosciences*, 15(20), 6049–6066, doi:<https://doi.org/10.5194/bg-15-6049-2018>, 2018.
- 1886  
1887 Meier, K. J. S., Beaufort, L., Heussner, S., and Ziveri, P.: The role of ocean acidification in *Emiliania huxleyi*  
1888 coccolith thinning in the Mediterranean Sea, *Biogeosciences*, 11, 2857–2869, <https://doi.org/10.5194/bg-11-2857-2014>, 2014.
- 1889  
1890 Mercier, H., Lherminier, P., Sarafanov, A., Gaillard, F., Daniault, N., Desbruyères, D., Falina, A., Ferron, B.,  
1891 Huck, T., and Thierry, V.: Variability of the meridional overturning circulation at the Greenland-Portugal Ovide  
1892 section from 1993 to 2010. *Progress in Oceanography*, 132, 250-261, doi:[10.1016/j.pocean.2013.11.001](https://doi.org/10.1016/j.pocean.2013.11.001). 2015
- 1893  
1894 Merlivat, L., Boutin, J., Antoine, D., Beaumont, L., Golbol, M., and Vellucci, V.: Increase of dissolved inorganic  
1895 carbon and decrease in pH in near-surface waters in the Mediterranean Sea during the past two decades,  
1896 *Biogeosciences*, 15, 5653-5662, <https://doi.org/10.5194/bg-15-5653-2018>, 2018.
- 1897  
1898 Metzl, N., Brunet, C., Jabaud-Jan, A., Poisson, A., and Schauer, B.: Summer and winter air-sea CO<sub>2</sub> fluxes in  
1899 the Southern Ocean *Deep Sea Res I*, 53, 1548-1563, doi:10.1016/j.dsr.2006.07.006. 2006.
- 1900  
1901 Metzl, N., Tilbrook, B., Bakker, D., Le Quéré, C., Doney, S., Feely, R., Hood M., and Dargaville, R.: Global  
1902 Changes in Ocean Carbon: Variability and Vulnerability. *Eos, Transactions of the American Geophysical Union*  
1903 88 (28): 286-287. doi: 10.1029/2007EO280005, 2007.
- 1904  
1905 Metzl, N., Corbière, A., Reverdin, G., Lenton, A., Takahashi, T., Olsen, A., Johannessen, T., Pierrot, D.,  
1906 Wanninkhof, R., Ólafsdóttir, S. R., Olafsson, J., and Ramonet, M.: Recent acceleration of the sea surface fCO<sub>2</sub>  
1907 growth rate in the North Atlantic subpolar gyre (1993-2008) revealed by winter observations, *Global  
1908 Biogeochem. Cycles*, 24, GB4004, doi:10.1029/2009GB003658, 2010.
- 1909  
1910 Metzl, N., and Lo Monaco, C.: OISO - OCÉAN INDIEN SERVICE D'OBSERVATION,  
1911 <https://doi.org/10.18142/228>, 1998.
- 1912  
1913 Metzl, N., Pierre, C., and Vangriesheim, A.: Hydrographic and Chemical measurements during the R/V  
1914 L'Atalante BIOZaire III Cruise in the Atlantic Ocean (14 December, 2003 - 7 January 2004).  
1915 <http://cdiac.esd.ornl.gov/ftp/oceans/BIOZAIRED3>. Carbon Dioxide Information Analysis Center, Oak Ridge  
1916 National Laboratory, US Department of Energy, Oak Ridge, Tennessee. doi:  
1917 10.3334/CDIAC/OTG.BIOZAIRED3, 2016.
- 1918  
1919 Metzl, N., Ferron, B. Lherminier, P. Sarthou, G. Thierry, V.: Discrete profile measurements of dissolved  
1920 inorganic carbon (DIC), total alkalinity (TALK), temperature and salinity during the multiple ships Observatoire  
1921 de la variabilité interannuelle et décennale en Atlantique Nord (OVIDE) project, OVIDE-2006, OVIDE-2008,  
1922 OVIDE-2010, OVIDE-2012, OVIDE-2014 cruises in the North Atlantic Ocean from 2006-05-23 to 2014-06-30
- 1923

- 1924 (NCEI Accession 0177219). Version 1.1. NOAA National Centers for Environmental Information. Dataset.  
1925 doi:10.25921/v0qt-ms48 [access date], 2018.
- 1926
- 1927 Metzl, N., Fin, J., Lo Monaco, C., et al.: A synthesis of total alkalinity and dissolved inorganic carbon  
1928 measurements in the global ocean (1993-2022) SNAPO-CO2-V1 dataset. SEANOE.  
1929 <https://doi.org/10.17882/95414>, 2023.
- 1930 Mignot, A., Claustre, H., Cossarini, G., D'Ortenzio, F., Gutknecht, E., Lamouroux, J., Lazzari, P., Perruche, C.,  
1931 Salon, S., Sauzède, R., Taillardier, V., and Teruzzi, A.: Using machine learning and Biogeochemical-Argo  
1932 (BGC-Argo) floats to assess biogeochemical models and optimize observing system design, *Biogeosciences*, 20,  
1933 1405–1422, <https://doi.org/10.5194/bg-20-1405-2023>, 2023.
- 1934
- 1935 Millero, F. J., Lee, K. and Roche, M.: Distribution of alkalinity in the surface waters of the major oceans. *Mar.*  
1936 *Chem.* **60**, 111–130. [https://doi.org/10.1016/S0304-4203\(97\)00084-4](https://doi.org/10.1016/S0304-4203(97)00084-4). 1998.
- 1937
- 1938 Mongwe, N. P., Vichi, M., and Monteiro, P. M. S: The seasonal cycle of *p*CO<sub>2</sub> and CO<sub>2</sub> fluxes in the Southern  
1939 Ocean: Diagnosing anomalies in CMIP5 earth system models. *Biogeosciences*, 15(9), 2851–2872.  
1940 <https://doi.org/10.5194/bg-15-2851-2018>, 2018.
- 1941
- 1942 Mortier, L., Ait Ameur, N., and Taillardier, V. : SOMBA-GE-2014 cruise, RV Téthys II,  
1943 <https://doi.org/10.17600/14007500>, 2014.
- 1944
- 1945 Moutin, T., and Bonnet, S. : OUTPACE cruise, RV L'Atalante, <https://doi.org/10.17600/15000900>, 2015.
- 1946
- 1947 Moutin, T., Wagener, T., Caffin, M., Fumenia, A., Gimenez, A., Baklouti, M., Bouruet-Aubertot, P., Pujo-Pay,  
1948 M., Leblanc, K., Lefevre, D., Helias Nunige, S., Leblond, N., Grossi, O., and de Verneil, A.: Nutrient  
1949 availability and the ultimate control of the biological carbon pump in the western tropical South Pacific Ocean,  
1950 *Biogeosciences*, 15, 2961–2989, <https://doi.org/10.5194/bg-15-2961-2018>, 2018.
- 1951
- 1952 Newton, J.A., Feely, R. A., Jewett, E. B., Williamson, P. and Mathis, J.: Global Ocean Acidification Observing  
1953 Network: Requirements and Governance Plan. Second Edition, GOA-ON,  
1954 <https://www.iaea.org/sites/default/files/18/06/goa-on-second-edition-2015.pdf>, 2015.
- 1955
- 1956 Nykjaer, L.: Mediterranean Sea surface warming 1985-2006. *Clim. Res.* 39, 11–17. doi: 10.3354/cr00794, 2009.
- 1957
- 1958 Obernosterer, I.: MOBYDICK-THEMISTO cruise, RV Marion-Dufresne, <https://doi.org/10.17600/18000403>,  
1959 2018.
- 1960
- 1961 OCADS: Coastal Carbon Data, [https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/coastal\\_carbon\\_data.html](https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/coastal_carbon_data.html), 2023
- 1963
- 1964 Olafsson, J., Olafsdottir, S.R., Benoit-Cattin, A., Danielsen, M., Arnarson, T.S., and Takahashi, T.: Rate of  
1965 Iceland Sea acidification from time series measurements. *Biogeosciences* 6, 2661–2668.  
1966 <https://doi.org/10.5194/bg-6-2661-2009>, 2009.
- 1967
- 1968 Olivier, L., Boutin, J., Reverdin, G., Lefèvre, N., Landschützer, P., Speich, S., Karstensen, J., Labaste, M.,  
1969 Noisel, C., Ritschel, M., Steinhoff, T., and Wanninkhof, R.: Wintertime process study of the North Brazil  
1970 Current rings reveals the region as a larger sink for CO<sub>2</sub> than expected, *Biogeosciences*, 19, 2969–2988,  
1971 <https://doi.org/10.5194/bg-19-2969-2022>, 2022.
- 1972
- 1973 Olsen, A., Key, R. M., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T.,  
1974 Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Pérez, F. F., and Suzuki, T.: The Global  
1975 Ocean Data Analysis Project version 2 (GLODAPv2) – an internally consistent data product for the world ocean,  
1976 *Earth Syst. Sci. Data*, 8, 297–323, <https://doi.org/10.5194/essd-8-297-2016>, 2016.

- 1977
- 1978 Olsen, A., Lange, N., Key, R. M., Tanhua, T., Álvarez, M., Becker, S., Bittig, H. C., Carter, B. R., Cotrim da  
 1979 Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Jeansson, E., Jones, S. D., Jutterström, S.,  
 1980 Karlsen, M. K., Kozyr, A., Lauvset, S. K., Lo Monaco, C., Murata, A., Pérez, F. F., Pfeil, B., Schirnick, C.,  
 1981 Steinfeldt, R., Suzuki, T., Telszewski, M., Tilbrook, B., Velo, A., and Wanninkhof, R.: GLODAPv2.2019 – an  
 1982 update of GLODAPv2, *Earth Syst. Sci. Data*, 11, 1437–1461, <https://doi.org/10.5194/essd-11-1437-2019>, 2019.  
 1983
- 1984 Olsen, A., Lange, N., Key, R. M., Tanhua, T., Bittig, H. C., Kozyr, A., Álvarez, M., Azetsu-Scott, K., Becker, S.,  
 1985 Brown, P. J., Carter, B. R., Cotrim da Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M.,  
 1986 Jeansson, E., Jutterström, S., Landa, C. S., Lauvset, S. K., Michaelis, P., Murata, A., Pérez, F. F., Pfeil, B.,  
 1987 Schirnick, C., Steinfeldt, R., Suzuki, T., Tilbrook, B., Velo, A., Wanninkhof, R., and Woosley, R. J.: An updated  
 1988 version of the global interior ocean biogeochemical data product, GLODAPv2.2020, *Earth Syst. Sci. Data*, 12,  
 1989 3653–3678, <https://doi.org/10.5194/essd-12-3653-2020>, 2020.
- 1990
- 1991 Orr, J. C., Epitalon, J.-M., and Gattuso, J.-P.: Comparison of ten packages that compute ocean carbonate  
 1992 chemistry, *Biogeosciences*, 12(5), 1483–1510, doi:10.5194/bg-12-1483-2015, 2015.  
 1993
- 1994 Orr, J. C., Epitalon, J.-M., Dickson, A. G., and Gattuso, J.-P.: Routine uncertainty propagation for the marine  
 1995 carbon dioxide system, *Marine Chemistry*, Vol. 207, 84–107, doi:10.1016/j.marchem.2018.10.006., 2018.  
 1996
- 1997 Parard, G., Lefèvre, N., and Boutin, J.: Sea water fugacity of CO<sub>2</sub> at the PIRATA mooring at 6°S, 10°W. Tellus-  
 1998 B, DOI: 10.1111/j.1600-0889.2010.00503.x. 2010.  
 1999
- 2000 Pérez F. F., Vázquez-Rodríguez, M., Mercier, H., Velo, A., Lherminier, P. and Ríos, A. F.: Trends of  
 2001 anthropogenic CO<sub>2</sub> storage in North Atlantic water masses. *Biogeosciences*, 7, 1789–1807, [doi:10.5194/bg-7-1789-2010](https://doi.org/10.5194/bg-7-1789-2010), 2010.  
 2003
- 2004 Pérez, F. F., Mercier, H., Vazquez-Rodriguez, M., Lherminier, P., Velo, A., Pardo, P., Roson, G., and Rios, A.:  
 2005 Reconciling air-sea CO<sub>2</sub> fluxes and anthropogenic CO<sub>2</sub> budgets in a changing North Atlantic. *Nature  
 2006 Geosciences*, 6, 146–152, [doi:10.1038/ngeo1680](https://doi.org/10.1038/ngeo1680), 2013.  
 2007
- 2008 Pérez, F., Fontela, M., García-Ibáñez, M. et al. : Meridional overturning circulation conveys fast acidification to  
 2009 the deep Atlantic Ocean. *Nature* 554, 515–518. Doi: 10.1038/nature25493, 2018.  
 2010
- 2011 Pesant, S., Not, F., Picheral, M., Kandels-Lewis, S., Le Bescot, N., Gorsky, G., Iudicone, D., Karsenti, E.,  
 2012 Speich, S., Troublé, R., Dimier, C., Searson, S., and Tara Oceans Consortium Coordinators: Open science  
 2013 resources for the discovery and analysis of Tara Oceans data. *Scientific Data* 2:150023. doi:  
 2014 10.1038/sdata.2015.23, 2015.  
 2015
- 2016 Petrenko, A.: LATEX10 cruise, RV Téthys II, <https://doi.org/10.17600/10450150>, 2010.  
 2017
- 2018 Petrenko, A.A., Doglioli, A.M., Nencioli, F., Kersalé, M., Hu, Z., and d'Ovidio, F.: A review of the LATEX  
 2019 project: mesoscale to submesoscale processes in a coastal environment. *Ocean Dynam.*, doi: 10.1007/s10236-  
 2020 017-1040-9, 2017.  
 2021
- 2022 Petton, S., Pouvreau, S., and Fleury, E. : ECOSCOPA network : high frequency environmental database.  
 2023 SEANOE. <https://doi.org/10.17882/86131>, 2023.  
 2024
- 2025 Pfeil, B., Olsen, A., Bakker, D. C. E., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., Metzl, N.,  
 2026 Sabine, C. L., Akl, J., Alin, S. R., Bates, N., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J.,  
 2027 Chavez, F. P., Chen, A., Cosca, C., Fassbender, A. J., Feely, R. A., González-Dávila, M., Goyet, C., Hales,  
 2028 B., Hardman-Mountford, N., Heinze, C., Hood, M., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M.,  
 2029 Johannessen, T., Jones, S. D., Key, R. M., Kötzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N.,  
 2030 Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A.,

- 2031 Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot,  
 2032 D., Poisson, A., Ríos, A. F., Santana-Casiano, J. M., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R.,  
 2033 Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K.,  
 2034 Telszewski, M., Thomas, H., Tilbrook, B., Tjiputra, J., Vandemark, D., Veness, T., Wanninkhof, R., Watson,  
 2035 A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: A uniform, quality controlled Surface Ocean CO<sub>2</sub>  
 2036 Atlas (SOCAT), *Earth Syst. Sci. Data*, 5, 125–143, doi:10.5194/essd-5-125-2013, 2013.  
 2037
- 2038 Picheral, M., Searson, S., Taillandier, V., Bricaud, A., Boss, E., Ras, J., Claustre, H., Ouhssain, M., Morin, P.,  
 2039 Coppola, L., Gattuso, J.-P., Metzl, N., Thuillier, D., Gorsky, G., Tara Oceans Consortium, Coordinators; Tara  
 2040 Oceans Expedition, Participants: Vertical profiles of environmental parameters measured on discrete water  
 2041 samples collected with Niskin bottles during the Tara Oceans expedition 2009–2013.  
 2042 doi:10.1594/PANGAEA.836319, 2014.  
 2043
- 2044 Pilcher, D. J., Brody, S. R., Johnson, L., and Bronselaer, B.: Assessing the abilities of CMIP5 models to  
 2045 represent the seasonal cycle of surface ocean pCO<sub>2</sub>, *J. Geophys. Res. Oceans*, 120, 4625–4637,  
 2046 doi:10.1002/2015JC010759, 2015.  
 2047
- 2048 Poisson, A., Culkin, F., and Ridout, P.: Intercomparison of CO<sub>2</sub> measurements. *Deep Sea Research Part A.*  
 2049 *Oceanographic Research Papers*, 37, 10, 1647–1650, [https://doi.org/10.1016/0198-0149\(90\)90067-6](https://doi.org/10.1016/0198-0149(90)90067-6), 1990.  
 2050
- 2051 Pujo-Pay, M., Durrieu de Madron, X., and Conan, P.: PERLE3 cruise, RV Pourquoi pas ?,  
 2052 <https://doi.org/10.17600/18001342>, 2020.  
 2053
- 2054 Pujo-Pay, M., Durrieu de Madron, X., and Conan, P.: PERLE4 cruise, RV L'Atalante,  
 2055 <https://doi.org/10.17600/18001980>, 2021.  
 2056
- 2057 Rabouille C.: AMOR-BFLUX cruise, RV Téthys II, <https://doi.org/10.17600/15008700>, 2015.  
 2058
- 2059 Racapé, V., Metzl, N., Pierre, C., Reverdin, G., Quay, P.D., and Olafsdottir, S. R.: The seasonal cycle of the  
 2060 d<sub>13</sub>C<sub>DIC</sub> in the North Atlantic Subpolar Gyre. *Biogeosciences*, 11, 6, 1683–1692, doi:10.5194/bg-11-1683-2014,  
 2061 2014.  
 2062
- 2063 Revelle, R., and Suess, H. E.: Carbon dioxide exchange between atmosphere and ocean and the question of an  
 2064 increase of atmospheric CO<sub>2</sub> during the past decades. *Tellus* 9, 18–27. doi:10.1111/j.2153-  
 2065 3490.1957.tb01849.x., 1957.  
 2066
- 2067 Reverdin, G.: STRASSE cruise, RV Thalassa, <https://doi.org/10.17600/12040060>, 2012.  
 2068
- 2069 Reverdin, G., Metzl, N., Olafsdottir, S., Racapé, V., Takahashi, T., Benetti, M., Valdimarsson, H., Benoit-Cattin,  
 2070 A., Danielsen, M., Fin, J., Naamar, A., Pierrot, D., Sullivan, K., Bringas, F., and Goni, G.: SURATLANT: a  
 2071 1993–2017 surface sampling in the central part of the North Atlantic subpolar gyre, *Earth Syst. Sci. Data*, 10,  
 2072 1901–1924, <https://doi.org/10.5194/essd-10-1901-2018>, 2018.  
 2073
- 2074 Reverdin, G., Metzl, N., Olafsdottir, S., Racapé, V., Takahashi, T., Benetti, M., Valdimarsson, H., Quay, P. D.,  
 2075 Benoit-Cattin, A., Danielsen, M., Fin, J., Naamar, A., Pierrot, D., Sullivan, K., Bringas, F., and Goni, G.:  
 2076 SURATLANT: a surface dataset in the central part of the North Atlantic subpolar gyre. SEANOE.  
 2077 <https://doi.org/10.17882/54517>, 2022.  
 2078
- 2079 Ridame, C., Dekaezemacker, J., Guieu, C., Bonnet, S., L'Helguen, S., and Malien, F.: Contrasted Saharan dust  
 2080 events in LNLC environments: impact on nutrient dynamics and primary production, *Biogeosciences*, 11, 4783–  
 2081 4800, <https://doi.org/10.5194/bg-11-4783-2014>, 2014.  
 2082

- 2083 Robertson, J. E., Robinson, C., Turner, D. R., Holligan, P., Watson, A. J., Boyd, P., Fernandez, E., and Finch,  
2084 M.: The impact of a coccolithophore bloom on oceanic carbon uptake in the northeast Atlantic during summer  
2085 1991, Deep Sea Res., Part I, 41(2), 297–314, 1994.
- 2086
- 2087 Rödenbeck, C., Keeling, R. F., Bakker, D. C. E., Metzl, N., Olsen, A., Sabine, C., and Heimann, M.: Global  
2088 surface-ocean pCO<sub>2</sub> and sea-air CO<sub>2</sub> flux variability from an observation-driven ocean mixed-layer scheme,  
2089 Ocean Sci., 9, 193–216, <https://doi.org/10.5194/os-9-193-2013>, 2013.
- 2090
- 2091 Rödenbeck, C., Bakker, D. C. E., Gruber, N., Iida, Y., Jacobson, A.R., Jones, S., Landschützer, P., Metzl, N.,  
2092 Nakaoka, S., Olsen, A., Park, G.-H., Peylin, P., Rodgers, K. B., Sasse, T. P., Schuster, U., Shutler, J. D., Valsala,  
2093 V., Wanninkhof, R., Zeng, J. Data-based estimates of the ocean carbon sink variability – First results of the  
2094 Surface Ocean pCO<sub>2</sub> Mapping intercomparison (SOCOM). Biogeosciences 12: 7251-7278. doi:10.5194/bg-12-  
2095 7251-2015, 2015.
- 2096
- 2097 Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S.,  
2098 Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T.-H., Kozyr, A., Ono, T., and Rios, A. F.: The Oceanic  
2099 Sink for Anthropogenic CO<sub>2</sub>, Science, 305, 367-371, <https://doi.org/10.1126/science.1097403>, 2004.
- 2100
- 2101 Sabine, C. L., Hankin, S., Koyuk, H., Bakker, D. C. E., Pfeil, B., Olsen, A., Metzl, N., Kozyr, A., Fassbender,  
2102 A., Manke, A., Malczyk, J., Akl, J., Alin, S. R., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-  
2103 J., Chavez, F. P., Chen, A., Cosca, C., Feely, R.A., González-Dávila, M., Goyet, C., Hardman-Mountford, N.,  
2104 Heinze, C., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Key, R. M., Kötzinger, A.,  
2105 Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T.,  
2106 Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin,  
2107 X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Salisbury, J., Santana-  
2108 Casiano, J. M., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff,  
2109 T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Vandemark, D., Veness,  
2110 T., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: Surface Ocean CO<sub>2</sub> Atlas (SOCAT)  
2111 gridded data products, Earth Syst. Sci. Data, 5, 145-153, doi:10.5194/essd-5-145-2013, 2013.
- 2112
- 2113 Salt, L. A., Beaumont, L., Blain, S., Buccarelli, E.,Grossteffan, E., Guillot, A., L'Helguen, S., Merlivat, L.,  
2114 Répécaud, M., Quéméner, L.,Rimmelin-Maury, P., Tréguer, P., and Bozec, Y.: The annual and seasonal  
2115 variability of the carbonate system in the Bay of Brest (Northwest Atlantic Shelf, 2008–2014). Marine  
2116 Chemistry, doi:10.1016/j.marchem.2016.09.003. 2016.
- 2117
- 2118 Sasse, T. P., McNeil, B. I., and Abramowitz, G.: A novel method for diagnosing seasonal to inter-annual surface  
2119 ocean carbon dynamics from bottle data using neural networks, Biogeosciences, 10, 4319–4340,  
2120 <https://doi.org/10.5194/bg-10-4319-2013>, 2013.
- 2121
- 2122 Sauzède, R., Claustre, H., Pasqueron de Fommervault, O., Bittig, H., Gattuso, J.-P., Legendre, L. and Johnson,  
2123 K. S.: Estimates of water-column nutrients and carbonate system parameters in the global ocean: A novel  
2124 approach based on neural networks. *Front. Mar. Sci.* 4:128. doi:10.3389/fmars.2017.00128, 2017.
- 2125
- 2126 Seelmann, K., Steinhoff, T., Aßmann, S., and Kötzinger, A.: Enhance Ocean Carbon Observations: Successful  
2127 Implementation of a Novel Autonomous Total Alkalinity Analyzer on a Ship of Opportunity. *Front. Mar. Sci.*  
2128 7:571301.doi: 10.3389/fmars.2020.571301, 2020.
- 2129
- 2130 Schlitzer, R.: Ocean Data View, Ocean Data View, <http://odv.awi.de> (last access: 13 March 2019), 2018.
- 2131
- 2132 Schneider, A., Wallace, D. W. R., and Kötzinger, A.: Alkalinity of the Mediterranean Sea, *Geophys. Res. Lett.*,  
2133 34, L15608, doi:10.1029/2006GL028842, 2007.
- 2134

- 2135 Schuster, U., Watson, A.J., Bates, N., Corbière, A., Gonzalez-Davila, M., Metzl, N., Pierrot, D. and Santana-  
2136 Casiano, M.: Trends in North Atlantic sea surface pCO<sub>2</sub> from 1990 to 2006. Deep-Sea Res II,  
2137 doi:10.1016/j.dsr2.2008.12.011, 2009.
- 2138
- 2139 Schuster, U., McKinley, G. A., Bates, N., Chevallier, F., Doney, S. C., Fay, A. R., González-Dávila, M., Gruber,  
2140 N., Jones, S., Krijnen, J., Landschützer, P., Lefèvre, N., Manizza, M., Mathis, J., Metzl, N., Olsen, A., Rios, A.  
2141 F., Rödenbeck, C., Santana-Casiano, J. M., Takahashi, T., Wanninkhof, R., and Watson, A. J.: An assessment of  
2142 the Atlantic and Arctic sea-air CO<sub>2</sub> fluxes, 1990–2009, Biogeosciences, 10, 607–627,  
2143 https://doi.org/10.5194/bg-10-607-2013, 2013.
- 2144
- 2145 Sims, R. P., Holding, T. M., Land, P. E., Piolle, J.-F., Green, H. L., and Shutler, J. D.: OceanSODA-UNEXE: a  
2146 multi-year gridded Amazon and Congo River outflow surface ocean carbonate system dataset, Earth Syst. Sci.  
2147 Data, 15, 2499–2516, https://doi.org/10.5194/essd-15-2499-2023, 2023.
- 2148
- 2149 Skjelvan, I., Lauvset, S.K., Johannessen, T., et al.: Decadal trends in Ocean Acidification from the Ocean  
2150 Weather Station M in the Norwegian Sea, Journal of Marine Systems, <https://doi.org/10.1016/j.jmarsys.2022.103775>, 2022.
- 2152
- 2153 Speich, S., and The Embarked Science Team: EUREC4A-OA. Cruise Report. 19 January – 19 February 2020.  
2154 Vessel: L'ATALANTE. <https://doi.org/10.13155/80129>, 2021
- 2155
- 2156 Takahashi, T., Sutherland, S. C., Sweeney, C., Poisson, A., Metzl, N., Tilbrook, B., Bates, N., Wanninkhof, R.,  
2157 Feely, R. A., Sabine, C., Olafsson, J., and Nojiri, Y.: Global Sea-Air CO<sub>2</sub> Flux Based on Climatological Surface  
2158 Ocean pCO<sub>2</sub>, and Seasonal Biological and Temperature Effect. *Deep-Sea Res. II*, 49, 9-10, 1601-1622,  
2159 [https://doi.org/10.1016/S0967-0645\(02\)00003-6](https://doi.org/10.1016/S0967-0645(02)00003-6). 2002
- 2160
- 2161 Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B.,  
2162 Friederich, G., Chavez, F., Sabine, C., Watson, A. J., Bakker, D. C., Schuster, U., Metzl, N., Yoshikawa-Inoue,  
2163 H., Ishii, M., Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T.  
2164 S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C., Delille, B., Bates, N., and de Baar, H. J.:  
2165 Climatological mean and decadal change in surface ocean pCO<sub>2</sub>, and net sea air CO<sub>2</sub> flux over the global  
2166 oceans. *Deep-Sea Res. II*, 56 (8-10), 554–577, <http://dx.doi.org/10.1016/j.dsr2.2008.12.009>. 2009.
- 2167
- 2168 Takahashi, T., Sutherland, S. C., Chipman, D. W., Goddard, J. G., Ho, C., Newberger, T., Sweeney, C. and  
2169 Munro, D. R.: Climatological distributions of pH, pCO<sub>2</sub>, total CO<sub>2</sub>, alkalinity, and CaCO<sub>3</sub> saturation in the  
2170 global surface ocean, and temporal changes at selected locations. *Marine Chemistry*, 164, 95–125,  
2171 doi:10.1016/j.marchem.2014.06.004. 2014.
- 2172
- 2173 Tanhua, T., Pouliquen, S., Hausman, J., O'Brien, K., Bricher, P., de Bruin, T., Buck, J. J. H., Burger, E. F.,  
2174 Carval, T., Casey, K. S., Diggs, S., Giorgetti, A., Glaves, H., Harscoat, V., Kinkade, D., Muelbert, J. H.,  
2175 Novellino, A., Pfeil, B., Pulsifer, P. L., Van de Putte, A., Robinson, E., Schaap, D., Smirnov, A., Smith, N.,  
2176 Snowden, D., Spears, T., Stall, S., Tacoma, M., Thijssse, P., Tronstad, S., Vandenberghe, T., Wengren, M.,  
2177 Wyborn, L. and Zhao, Z.: Ocean FAIR Data Services. *Front. Mar. Sci.* 6:440. doi: 10.3389/fmars.2019.00440,  
2178 2019.
- 2179
- 2180 Tanhua, T., Lauvset, S.K., Lange, N. et al.: A vision for FAIR ocean data products. *Commun Earth Environ* 2,  
2181 136. <https://doi.org/10.1038/s43247-021-00209-4>, 2021
- 2182
- 2183 Testor, P., Bosse, A., and Coppola, L.: MOOSE-GE, <https://doi.org/10.18142/235>, 2010.
- 2184
- 2185 Testor, P.: DEWEX-MERMEX 2013 LEG1 cruise, RV Le Suroît, <https://doi.org/10.17600/13020010>, 2013.
- 2186
- 2187 Tilbrook, B., Jewett, E. B., DeGrandpre, M. D., Hernandez-Ayon, J. M., Feely, R. A., Gledhill, D. K., Hansson,  
2188 L., Isensee, K., Kurz, M. L., Newton, J. A., Siedlecki, S. A., Chai, F., Dupont, S., Graco, M., Calvo, E., Greeley,

- 2189 D., Kapsenberg, L., Lebrec, M., Pelejero, C., Schoo, K. L., and Telszewski, M.: An Enhanced Ocean  
2190 Acidification Observing Network: From People to Technology to Data Synthesis and Information Exchange.  
2191 *Frontiers in Marine Science*, 6, 337, DOI:10.3389/fmars.2019.00337, 2019.
- 2192
- 2193 Touratier, F., Azouzi, L. and Goyet, C.: CFC-11,  $\Delta^{14}\text{C}$  and  ${}^3\text{H}$  tracers as a means to assess anthropogenic CO<sub>2</sub>  
2194 concentrations in the ocean. *Tellus B*, 59(2), 318–325, doi:10.1111/j.1600-0889.2006.00247.x, 2007.
- 2195
- 2196 Touratier, F., and Goyet, C.: Decadal evolution of anthropogenic CO<sub>2</sub> in the north western Mediterranean Sea  
2197 from the mid-1990's to the mid-2000's. *Deep Sea Research Part I*.doi:10.1016/j.dsr.2009.05.015, 2009.
- 2198
- 2199 Touratier, F., Goyet, C., Houpert, L., Durrieu de Madron, X., Lefèvre, D., Stabholz, M., and Guglielmi, V.: Role  
2200 of deep convection on anthropogenic CO<sub>2</sub> sequestration in the Gulf of Lions (northwestern Mediterranean Sea).  
2201 *Deep-Sea Research Part I*. doi.org/10.1016/j.dsr.2016.04.003, 2016.
- 2202
- 2203 Turk, D., Dowd, M., Lauvset, S. K., Koelling, J., Alonso-Pérez, F. and Pérez, F. F.: Can Empirical Algorithms  
2204 Successfully Estimate Aragonite Saturation State in the Subpolar North Atlantic? *Front. Mar. Sci.* 4:385. doi:  
2205 10.3389/fmars.2017.00385, 2017.
- 2206
- 2207 Ulles, C., Estournel, C., Marsaleix, P., Soetaert, K., Fourrier, M., Coppola, L., Lefèvre, D., Touratier, F., Goyet,  
2208 C., Guglielmi, V., Kessouri, F., Testor, P., and Durrieu de Madron, X.: Seasonal dynamics and annual budget of  
2209 dissolved inorganic carbon in the northwestern Mediterranean deep convection region, *Biogeosciences Discuss.*  
2210 [preprint], <https://doi.org/10.5194/bg-2022-219>, in review, accepted, 2023.
- 2211
- 2212 UNESCO: Intercomparison of total alkalinity and total inorganic carbon determinations in seawater. *UNESCO*  
2213 *Tech. Pap. Mar. Sci.* 59., 1990
- 2214
- 2215 UNESCO: Reference materials for oceanic carbon dioxide measurements. *UNESCO Tech. Pap. Mar. Sci.* 60.,  
2216 1991
- 2217
- 2218 United Nations. The Sustainable Development Goals 2020, 68pp. <https://unstats.un.org/sdgs/report/2020/>, 2020
- 2219
- 2220 Vangriesheim A., Pierre, C., Aminot, A., Metzl, N., Baurand, F., and Caprais, J.-C.: The influence of Congo  
2221 river discharges in the surface and deep layers of the Gulf of Guinea. *Deep-Sea Res. II*, doi:  
2222 10.1016/j.dsr2.2009.04.002, 2009.
- 2223
- 2224 Vazquez-Rodriguez, M., Perez, F., Velo, A., Rios, A., and Mercier, H.: Observed acidification trends in the  
2225 North Atlantic water masses. *Biogeosciences*, 9, 5217-5230, doi:10.5194/bg-9-5217-2012, 2012.
- 2226
- 2227 Velo, A., Perez, F. F., Brown, P., Tanhua, T., Schuster, U., and Key, R. M.: CARINA alkalinity data in the  
2228 Atlantic Ocean, *Earth Syst. Sci. Data*, 1, 45–61, <https://doi.org/10.5194/essd-1-45-2009>, 2009.
- 2229
- 2230 von Schuckmann, K., Cheng, L., Palmer, M. D., Hansen, J., Tassone, C., Aich, V., Adusumilli, S., Beltrami, H.,  
2231 Boyer, T., Cuesta-Valero, F. J., Desbruyères, D., Domingues, C., García-García, A., Gentine, P., Gilson, J.,  
2232 Gorfer, M., Haimberger, L., Ishii, M., Johnson, G. C., Killick, R., King, B. A., Kirchengast, G., Kolodziejczyk,  
2233 N., Lyman, J., Marzeion, B., Mayer, M., Monier, M., Monselesan, D. P., Purkey, S., Roemmich, D., Schweiger,  
2234 A., Seneviratne, S. I., Shepherd, A., Slater, D. A., Steiner, A. K., Straneo, F., Timmermans, M.-L., and Wijffels,  
2235 S. E.: Heat stored in the Earth system: where does the energy go?, *Earth Syst. Sci. Data*, 12, 2013–2041,  
2236 <https://doi.org/10.5194/essd-12-2013-2020>, 2020.
- 2237
- 2238 von Schuckmann, K., Minière, A., Gues, F., Cuesta-Valero, F. J., Kirchengast, G., Adusumilli, S., Straneo, F.,  
2239 Ablain, M., Allan, R. P., Barker, P. M., Beltrami, H., Blazquez, A., Boyer, T., Cheng, L., Church, J.,  
2240 Desbruyeres, D., Dolman, H., Domingues, C. M., García-García, A., Giglio, D., Gilson, J. E., Gorfer, M.,  
2241 Haimberger, L., Hakuba, M. Z., Hendricks, S., Hosoda, S., Johnson, G. C., Killick, R., King, B., Kolodziejczyk,  
2242 N., Korosov, A., Krinner, G., Kuusela, M., Landerer, F. W., Langer, M., Lavergne, T., Lawrence, I., Li, Y.,

- 2243 Lyman, J., Marti, F., Marzeion, B., Mayer, M., MacDougall, A. H., McDougall, T., Monselesan, D. P., Nitzbon,  
2244 J., Otosaka, I., Peng, J., Purkey, S., Roemmich, D., Sato, K., Sato, K., Savita, A., Schweiger, A., Shepherd, A.,  
2245 Seneviratne, S. I., Simons, L., Slater, D. A., Slater, T., Steiner, A. K., Suga, T., Szekely, T., Thiery, W.,  
2246 Timmermans, M.-L., Vanderkelen, I., Wijffels, S. E., Wu, T., and Zemp, M.: Heat stored in the Earth system  
2247 1960–2020: where does the energy go?, *Earth Syst. Sci. Data*, 15, 1675–1709, <https://doi.org/10.5194/essd-15-1675-2023>, 2023.
- 2249
- 2250 Wagener, T., Metzl, N., Caffin, M., Fin, J., Helias Nunige, S., Lefevre, D., Lo Monaco, C., Rougier, G., and  
2251 Moutin, T.: Carbonate system distribution, anthropogenic carbon and acidification in the western tropical South  
2252 Pacific (OUTPACE 2015 transect), *Biogeosciences*, 15, 5221–5236, <https://doi.org/10.5194/bg-15-5221-2018>,  
2253 2018a.
- 2254
- 2255 Wagener, T., Metzl, N., Caffin, M., Fin, J., Helias Nunige, S., Lefevre, D., Lo Monaco, C., Rougier, G., and  
2256 Moutin, T.: Discrete profile measurements of dissolved inorganic carbon (DIC), total alkalinity (TALK),  
2257 temperature, salinity and other parameters during the R/V L'Atalante "Oligotrophy from Ultra-oligoTrophy  
2258 PACific Experiment" (OUTPACE) cruise (EXPOCODE 35A320150218) in the South Pacific Ocean from 2015–  
2259 02-18 to 2015-04-03 (NCEI Accession 0177706). Version 1.1. NOAA National Centers for Environmental  
2260 Information. Dataset. doi:[10.25921/wbkb-0q19](https://doi.org/10.25921/wbkb-0q19) [access date], 2018b.
- 2261
- 2262 Walton, D. W. H., and Thomas, J.: Cruise Report - Antarctic Circumnavigation Expedition (ACE) 20th  
2263 December 2016 - 19th March 2017 (1.0). Zenodo. <https://doi.org/10.5281/zenodo.1443511>, 2018.
- 2264
- 2265 Wanninkhof, R., Park, G.-H., Takahashi, T., Sweeney, C., Feely, R., Nojiri, Y., Gruber, N., Doney, S. C.,  
2266 McKinley, G. A., Lenton, A., Le Quéré, C., Heinze, C., Schwinger, J., Graven, H., and Khatiwala, S.: Global  
2267 ocean carbon uptake: magnitude, variability and trends, *Biogeosciences*, 10, 1983–2000, doi:[10.5194/bg-10-1983-2013](https://doi.org/10.5194/bg-10-1983-2013), 2013.
- 2268
- 2269
- 2270 Watson, A. J., Schuster, U., Bakker, D .C. E., Bates, N., Corbiere, A., Gonzalez-Davila, M., Freidrich, T.,  
2271 Hauck, J., Heinze, C., Johannessen, T., Koertzinger, A., Metzl, N., Olafsson, J., Olsen, A., Oschlies, A., Padin,  
2272 X., Pfeil, B., Rios, A., Santana-Casiano, M, Steinhoff, T., Telszewski, M., Wallace, D. W. R., and Wanninkhof,  
2273 R.: Tracking the variable North Atlantic sink for atmospheric CO<sub>2</sub> , *Science* ,326, 1391,  
2274 doi:[10.1126/science.1177394](https://doi.org/10.1126/science.1177394). 2009.
- 2275
- 2276 Watson, A. J., Schuster, U., Shutler, J.D. et al.: Revised estimates of ocean-atmosphere CO<sub>2</sub> flux are consistent  
2277 with ocean carbon inventory. *Nat Commun* 11, 4422, <https://doi.org/10.1038/s41467-020-18203-3>, 2020.
- 2278
- 2279 Williams, N. L., Juranek, L. W., Johnson, K. S., Feely, R. A., Riser, S. C., Talley, L. D., et al.: Empirical  
2280 algorithms to estimate water column pH in the Southern Ocean. *Geophysical Research Letters*, 43, 3415–3422.  
2281 <https://doi.org/10.1002/2016GL068539>, 2016.
- 2282
- 2283 Williams, N. L., Juranek, L. W., Feely, R. A., Johnson, K. S., Sarmiento, J. L., Talley, L. D., Dickson,  
2284 A. G., Gray, A. R., Wanninkhof, R., Russell, J. L., Riser, S. C., and Takeshita, Y.: Calculating surface  
2285 ocean pCO<sub>2</sub> from biogeochemical Argo floats equipped with pH: An uncertainty analysis, *Global Biogeochem.*  
2286 *Cycles*, 31, 591–604, doi:[10.1002/2016GB005541](https://doi.org/10.1002/2016GB005541)., 2017.
- 2287
- 2288 Williams, N. L., Juranek, L. W., Feely, R. A., Russell, J. L., Johnson, K. S.,and Hales, B.: Assessment of the  
2289 carbonate chemistry seasonal cycles in the Southern Ocean from persistent observational platforms. *Journal of*  
2290 *Geophysical Research: Oceans*, 123. <https://doi.org/10.1029/2017JC012917>, 2018.
- 2291
- 2292 Wimart-Rousseau, C., Lajaunie-Salla, K., Marrec, P., Wagener, T., Raimbault,P., Lagadec, V., Lafont, M.,  
2293 Garcia, N., Diaz, F., Pinazo, C., Yohia, C., Garcia, F., Xueref-Remy,I., Blanc, P.-E., Armengaud, A., and  
2294 Lefèvre, D.: Temporal variability of the carbonate system and air-sea CO<sub>2</sub> exchanges in a Mediterranean human-  
2295 impacted coastal site. *Estuarine, Coastal and Shelf Science*. <https://doi.org/10.1016/j.ecss.2020.106641>, 2020a.
- 2296

2297 Wimart-Rousseau, C., Wagener, T., Raimbault, P., Lagadec, V., Lafont, M., Garcia, N., and Lefèvre, D.:  
2298 Oceanic carbonate chemistry measurements from discret samples collected at the SOLEMIO station (Bay of  
2299 Marseille - North western Mediterannean Sea) between 2016 and 2019. SEANOE.  
2300 <https://doi.org/10.17882/72356>, 2020b.

2301

2302 Wimart-Rousseau, C., Wagener, T., Álvarez, M., Moutin, T., Fourrier, M., Coppola, L., Niclas-Chirurgien, L.,  
2303 Raimbault, P., D'Ortenzio, F., Durrieu de Madron, X., Taillardier, V., Dumas, F., Conan, P., Pujo-Pay, M. and  
2304 Lefèvre, D.: Seasonal and Interannual Variability of the CO<sub>2</sub> System in the Eastern Mediterranean Sea: A Case  
2305 Study in the North Western Levantine Basin. *Front. Mar. Sci.* 8:649246. doi: 10.3389/fmars.2021.649246, 2021

2306

2307 WMO/GCOS, 2018: <https://gcos.wmo.int/en/global-climate-indicators>, 2018

2308

2309 Wu, Y., Hain, M. P., Humphreys, M. P., Hartman, S., and Tyrrell, T.: What drives the latitudinal gradient in  
2310 open-ocean surface dissolved inorganic carbon concentration?, *Biogeosciences*, 16, 2661-2681,  
2311 <https://doi.org/10.5194/bg-16-2661-2019>, 2019.

2312

2313