



Chemical and olfactory analysis of essential oils of *Hedychium gardnerianum*, *Hedychium flavescens*, *Pittosporum senacia* and *Psidium cattleianum* from Reunion Island

Marine Canton, Marina Thierry, Sylvain Antoniotti

► To cite this version:

Marine Canton, Marina Thierry, Sylvain Antoniotti. Chemical and olfactory analysis of essential oils of *Hedychium gardnerianum*, *Hedychium flavescens*, *Pittosporum senacia* and *Psidium cattleianum* from Reunion Island. *Journal of Essential Oil & Plant Composition*, 2023, 1 (3), pp.298-311. 10.58985/jeopc.2023.v01i03.39 . hal-04293587

HAL Id: hal-04293587

<https://hal.science/hal-04293587>

Submitted on 18 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 **Chemical and olfactory analysis of essential oils of *Hedychium gardnerianum*,**
2 ***Hedychium flavescens*, *Pittosporum senacia* and *Psidium cattleianum* from**
3 **Reunion Island**

4 Marine Canton¹, Marina Thierry¹ and Sylvain Antoniotti^{1,2,*}

5 ¹ *Université Côte d'Azur*

6 *Institut d'Innovation et de Partenariats Arômes Parfums Cosmétiques*

7 *Espace J.-L. Lions, 4 traverse Dupont, 06130 Grasse, France*

8 ² *Université Côte d'Azur, CNRS*

9 *Institut de Chimie de Nice*

10 *Parc Valrose, 06108 Nice cedex 2, France*

11 *Correspondence: sylvain.antoniotti@univ-cotedazur.fr

12

ABSTRACT

Four essential oils (EOs), rarely described in the literature, and never for samples originating from Reunion Island, were studied by gas chromatography with flame ionisation detector (GC-FID) and gas chromatography coupled with mass spectrometry and olfactometry (GC/MS-O). The chemical composition, as well as the main olfactory properties, of the following four EOs were determined: longose (*Hedychium gardnerianum*), yellow ginger (*Hedychium flavescens*), bois de joli coeur (*Pittosporum senacia*), and Chinese guava (*Psidium cattleianum*). The chemical compositions were found to be rich in mono- and sesquiterpene hydrocarbons. The main components of *H. gardnerianum* flowers EO was β -farnesene (12.1%), α -cadinol (9.7%) and α -farnesene (7.1%). The composition of essential oil from *Hedychium flavescens* leaves is reported for the first time with β - and α -pinene (47.8 and 18.7%, respectively) as well as β -caryophyllene (17.5%) as the main compounds. The main components of *P. senacia* leaves EO and *P. cattleianum* floral tops EO was identified as nonane (36.2%) and β -caryophyllene (43.7%), respectively. With such a chemical composition, herbal, citrus, green, pine tree, and in some cases floral and woody odors were determined by GC/MS-O.

Keywords: natural complex substance • chemical composition • natural scent • olfactometry • joli coeur

INTRODUCTION

The genus *Hedychium* is part of the Zingiberaceae family and includes about 80 species distributed mainly in Asia. *H. gardnerianum* is a plant native to the Himalayas whose stem can be 2 meters long and its leaves 30 cm long. Its flowers can be pink or orange yellow [1-3]. It is considered an invasive plant on most of the Azores archipelago [1]. The rhizomes of the *Hedychium* species are known to be fleshy and aromatic. Some species are cultivated only to extract the fragrant essential oil from their rhizomes. Aerial parts of the *Hedychium* species can be used in the paper industry and its flowers as culinary ingredients [4]. The bloom of the *Hedychium* species is very brief and usually occurs during the monsoon. The chemical profile of *Hedychium* sp. EO is reported as complex and comprising monoterpene and sesquiterpene derivatives. The compounds α -pinene, β -pinene, eucalyptol, linalool and nerolidol, although quite ubiquitous, are described as markers of the genus *Hedychium* [4].

The genus *Pittosporum* belongs to the Pittosporaceae family and encompasses ca. 200 species [5]. Among these, *Pittosporum senacia* is a species found on Indian Ocean Islands. The chemical composition of EO from the whole plant from Mauritius was described in 2020 and 2021 by Jugreet et al. [5, 6]. In their most recent article, the distillation yield for *P. senacia* was found to be 0.77%. The chemical composition mainly comprised monoterpene hydrocarbons (up to 71.9%), and particularly β -myrcene (62.2%). Other compounds found were germacrene D (7.8%), limonene (3.4%), and β -

phellandrene (2.9%). In an earlier article from 1998, EO from the leaves of *P. senecioioides*, endemic to Madagascar, was studied [7]. With a distillation yield of 0.67%, 95% of the chemical composition of the EO was determined by GC-FID and GC/MS. With monoterpene hydrocarbons accounting for only 20.4%, including β -myrcene (6.3%), α -terpinene (4.6%) and camphene (3%), the main compounds were found to be sesquiterpene hydrocarbons and derivatives (69.5%) such as α -cadinol (19%), α -muurolol (15.9%), and δ -cadinene (11.3%). In the same study, the antimicrobial activity of this EO against *Staphylococcus aureus* was shown to outperform streptomycin, while it was in the same order of magnitude as streptomycin against *Streptococcus faecalis* [7].

The genus *Psidium* belongs to the Myrtaceae family which contains ca. 5500 species [8, 9]. This genus is known for presenting rich essential oil bearing plants [9]. *Psidium cattleianum* is a species found in Oceania, Brazil, North America and the Caribbean [8]. The shrub is typically between 1 and 4 meters in height. The EO from the leaves of *P. cattleianum* has been described as having antifungal activities against *Candida spp.* and antibacterial activities against several strains such as *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* in qualitative analyses with the antibiotic ampicillin as control [8].

The chemical composition of EO from *P. cattleianum* leaves has been studied with plants coming mainly from Brazil [8-12]. Older studies describe *P. cattleianum* EO harvested in Hawaii [13], Cuba [14], and California [15].

As part of a project dedicated to the chemical and sensory evaluation of plants from Reunion Island for applications in perfumery, the chemical composition and olfactory properties of four essential oils (EOs) for which literature data are scarce or non-existent were studied. The four species, from the Zingiberaceae, Pittosporaceae, and Myrtaceae botanical families illustrated in Figure 1 are listed hereafter:

- Longose (*Hedychium gardnerianum* Sheppard ex Ker-Gawl) – flowers,

- Yellow ginger (*Hedychium flavescens* Carey ex Roscoe) – leaves,

- Bois de joli coeur (*Pittosporum senacia* Putt.) – leaves,

- Chinese guava (*Psidium cattleianum* Afzel. Ex Sabine) – floral tops.

The aim of these studies was to determine the chemical composition of these essential oils and evaluate their olfactory profile for potential applications in perfumery.

MATERIAL AND METHODS

Plant materials

H. gardnerianum, *H. flavescens*, *P. senacia* and *P. cattleianum* were collected in 2020 on Reunion Island, in the wild between 800 and 1300 meters of altitude and their essential oils were obtained from OLICA (19 Rue Fangourin, Saint-Leu 97424, La Réunion, FRANCE).

The four pictures of the species in Figure 1 are licensed under the Creative Commons Attribution-Share Alike 3.0 Unported license without modification. Authors are (1) and (4) B. Navez, (2) Forest and Kim Starr, (3) Change-ecorce.




Figure 1. The four species studied: (1) *H. gardnerianum*, flowers; (2) *H. flavescens*, leaves; (3) *P. senacia*, leaves; (4) *P. cattleianum*, floral tops.

GC/MS-FID analyses

The analyses of the four EOs chemical composition were performed on an Agilent GC 7820 chromatograph coupled to an Agilent 5977B electron ionization mass spectrometer and equipped with a flame ionization detector. The column used was Agilent HP5-MS capillary column (30 m x 0.25 mm i.d., film thickness 0.25 μ m). The sample of EO was diluted to 1% v/v in ethyl acetate. One microliter of the sample was injected with a split ratio of 1/10. Carrier gas was helium with a flow rate of 1.2 mL/min. The temperatures of injector and source was 250 °C and 230 °C, respectively. The oven temperature

was programmed to stay 4 min at 40°C, then to rise from 40 to 200°C with an increase of 2 °C/min and finally from 200 to 270 °C with an increase of 8 °C/min. For the MS, ionization energy was 70 eV and the range was from 35 to 350 m/z. The FID detector was set at 270 °C with an air flow of 400 mL/min and a hydrogen flow of 40 mL/min.

Characterization of the EOs

GC/MS was used for identification and GC-FID for quantification. The identification of EO constituents was carried out by matching the retention index (RI) determined against a series of C7-C40 alkanes as well as by matching the EI-MS mass spectra obtained with various databases (NIST20, Wiley6n and internal database).

GC/MS-O analyses

GC/MS-O analyses were performed on a Perkin Elmer Clarus 690 chromatograph coupled with a Perkin Elmer Clarus SQ8T mass selective detector and equipped with a Perkin Elmer sniffing port. The column was a Perkin Elmer Elite 5-MS (30m x 0.32mm i.d., film thickness 0.25µm). The sample of EO was diluted to 10% v/v in ethyl acetate. One microliter of the sample was injected with a split of 72 mL/min (inlet pressure: 23 psi). The split between the MS and the Olfactometry detector was 1:7 (v/v). The oven temperature was programmed to stay 2 min at 60 °C, to rise from 60 to 200 °C with an increase of 5 °C/min, then from 200 to 280 °C with an increase of 8 °C/min and finally held at 280 °C for 5 min. The solvent delay was 5.5 min. Four panelists carried out

the GC/MS-O analysis in duplicate, on each of the four EOs. Compounds were annotated by matching the EI-MS mass spectra obtained with databases (NIST20).

RESULTS AND DISCUSSION

Hedychium gardnerianum and *Hedychium flavescens*

Essential oils obtained from the rhizomes of the two species *H. gardnerianum* and *H. flavescens* have been described in the literature [2-4, 16-18]. For example, Ray et al. (2018) studied 10 species of *Hedychium*, including *H. gardnerianum* and *H. flavescens*, cultivated under the same conditions in India [4]. After harvesting, the EO of each plant was collected from the rhizomes and analyzed by GC/MS and GC-FID. The distillation yield of *H. gardnerianum* and *H. flavescens* species was about 0.20%, which is lower than other species of the genus such as *H. thyrsiforme* (0.75%) and *H. gracile* (0.65%), and higher than others such as *H. flavum* (0.10%) and *H. greenii* (0.05%). The EO of *H. gardnerianum* was described as dark yellow and that of *H. flavescens* as dark brown. The chemical compositions of the two EOs were determined, accounting for 75.9% and 92.8% of oils of *H. gardnerianum* and *H. flavescens*, respectively. Both were composed mainly of monoterpene derivatives, mostly β -pinene and α -pinene (28.03% and 21.15% for *H. gardnerianum* and 29.76% and 13.17% for *H. flavescens*, respectively). Eucalyptol, one of the main compounds of *H. flavescens* (12.80%), was found in a much smaller fraction in *H. gardnerianum* (0.42%) [4].

The EO of *H. gardnerianum* flowers has been very little studied; to the best of our knowledge, only two papers published in 2002 and 2003 described the EO obtained from this part of the plant [1, 2] and EO extracted from the leaves of *H. flavescens* has never been described. Medeiros et al. (2003) studied the chemical composition of EO from *H. gardnerianum* flowers from three locations on San Miguel Island in the Azores [1]. The composition of these was determined using GC/MS analysis alone, accounting for 87%, 94% and 91% of the total EO from the three locations, respectively. Although they came from the same island, significant differences in the relative percentages of compounds were observed, although GC/MS alone can hardly provide reliable quantitative information. The main compounds of the three EOs were monoterpene hydrocarbons with α -pinene (8.38-18.13%), β -pinene (5.06-11.99%), p-cymene (3.85-8.16%), γ -terpinene (3.15-14.43%) and the sesquiterpene derivatives α -cadinol (6.42-14.59%), β -caryophyllene (7.04-8.89%), δ -cadinene (4.89-8.76%) and τ -muurolol (2.64-5.86%). The chemical composition of EO from *H. gardnerianum* flowers from Fiji was studied by Smith et al. [2]. Up to 75% of its chemical composition was characterized by GC/MS-FID analyses. β -Caryophyllene (17.4%) and β -pinene (17.0%) were found to be the two main compounds.

***Hedychium gardnerianum* flowers EO analysis**

The sample of *H. gardnerianum* flowers EO was found to be chemically very complex (Figure 2). As shown in Table 1, a hundred compounds were detected by monodimensional gas chromatography and those identified

accounted for more than 87% of the EO. The sample was mainly composed of sesquiterpene derivatives including β -farnesene (12.05%), α -cadinol (9.71%), α -farnesene (7.09%), τ -muurolol (5.89%) and δ -cadinene (5.83%). The monoterpene derivatives identified accounted for ca. 20% with α -pinene, β -pinene and humulene as main representatives (4.48%, 4.46% and 3.93%, respectively). The sample also contained diterpene derivatives including kaur-16-ene (2.42%).

Our results are in agreement with the article by Medeiros et al. (2003) concerning the study of the Eos of longose leaves and flowers (*H. gardnerianum*) from the island of San Miguel (Azores) [1]. In their study, the main compounds were α -pinene, β -pinene, p-cymene, γ -terpinene, β -caryophyllene, β -cadinene and α -cadinol. These compounds were all present in the sample studied herein. However, β -farnesene, the main compound in our sample, was identified to a much lesser extent (ca. 3%) in the study by Medeiros et al (2003). This could suggested a particular role of farnesene in plant defense particular to the Reunion Island territory for a compound known to be associated with insect attraction [19].




Figure 2. Chromatogram of *Hedychium gardnerianum* flowers essential oil.

Table 1. Chemical composition of *Hedychium gardnerianum* flowers essential oil.

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
1	17.625	4.48	α-pinene	C ₁₀ H ₁₆	931	937
2	18.563	0.08	camphene	C ₁₀ H ₁₆	945	952
3	20.344	0.15	sabinene	C ₁₀ H ₁₆	971	974
4	20.493	4.46	β-pinene	C ₁₀ H ₁₆	973	979
5	21.682	0.21	β -myrcene	C ₁₀ H ₁₆	991	991
6	22.460	0.39	α -phellandrene	C ₁₀ H ₁₆	1002	1005
7	22.853	0.04	3-carene	C ₁₀ H ₁₆	1008	1011
8	23.339	0.24	α -terpinene	C ₁₀ H ₁₆	1015	1017
9	23.892	1.46	<i>p</i> -cymene	C ₁₀ H ₁₄	1022	1025
10	24.175	0.52	limonene	C ₁₀ H ₁₆	1026	1030
11	24.363	0.08	eucalyptol	C ₁₀ H ₁₆ O	1029	1032
12	25.000	0.23	(<i>Z</i>)- β -ocimene	C ₁₀ H ₁₆	1038	1038
13	25.705	2.97	(<i>E</i>)-β-ocimene	C ₁₀ H ₁₆	1048	1049
14	26.355	2.46	γ-terpinene	C ₁₀ H ₁₆	1057	1060
15	28.431	0.09	terpinolene	C ₁₀ H ₁₆	1086	1088
16	28.888	0.06	methyl benzoate	C ₈ H ₈ O ₂	1093	1094
17	29.371	2.22	linalool	C ₁₀ H ₁₈ O	1100	1099

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
18	30.504	0.12	(<i>E</i>)-4.8-Dimethyl- nona-1,3,7-triene	C ₁₁ H ₁₈	1116	1116
19	31.954	0.01	sabinol	C ₁₀ H ₁₆ O	1137	1143
20	33.804	0.09	endo-borneol	C ₁₀ H ₁₈ O	1164	1167
21	34.654	0.10	terpinen-4-ol	C ₁₀ H ₁₈ O	1176	1177
22	35.662	0.11	methyl salicylate	C ₈ H ₈ O ₃	1191	1192
23	41.904	0.06	bornyl acetate	C ₁₂ H ₂₀ O ₂	1285	1285
24	41.921	0.06	2-undecanone	C ₁₁ H ₂₀ O ₂	1285	1294
25	43.170	0.01	indole	C ₈ H ₇ N	1304	1295
26	45.202	0.05	δ-elemene	C ₁₅ H ₂₄	1336	1338
27	45.945	0.16	α-cubebene	C ₁₅ H ₂₄	1348	1351
28	47.567	0.19	α-copaene	C ₁₅ H ₂₄	1374	1376
29	48.614	0.21	β-elemene	C ₁₅ H ₂₄	1391	1391
30	49.628	0.04	α-gurjunene	C ₁₅ H ₂₄	1407	1409
31	50.219	1.38	β-caryophyllene	C ₁₅ H ₂₄	1417	1419
32	51.67	0.07	guaia-6,9-diene	C ₁₅ H ₂₄	1442	1443
33	52.281	3.93	humulene	C ₁₅ H ₂₄	1452	1454
34	52.640	12.05	(6<i>E</i>)-β-farnesene	C ₁₅ H ₂₄	1458	1457
35	52.848	0.01	allo-aromadendrene	C ₁₅ H ₂₄	1462	1461
36	53.484	0.18	γ-gurjunene	C ₁₅ H ₂₄	1472	1473
37	53.680	0.45	γ-muurolene	C ₁₅ H ₂₄	1476	1477
38	53.914	0.96	germacrene D	C ₁₅ H ₂₄	1480	1481
39	54.061	0.21	α-curcumene	C ₁₅ H ₂₂	1482	1483
40	54.537	0.17	Bicyclosesqui- phellandrene	C ₁₅ H ₂₄	1490	1489
41	54.822	1.67	bicyclogermacrene	C ₁₅ H ₂₄	1495	1495

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
42	55.070	1.05	α -muurolene	C ₁₅ H ₂₄	1499	1499
43	55.461	0.43	β -cadinene	C ₁₅ H ₂₄	1506	1518
44	55.646	7.09	(3E,6E)α-farnesene	C ₁₅ H ₂₄	1509	1508
45	55.848	1.03	γ -cadinene	C ₁₅ H ₂₄	1513	1513
46	56.449	5.83	δ-cadinene	C ₁₅ H ₂₄	1523	1524
47	56.891	0.09	cadina-1,4-diene	C ₁₅ H ₂₄	1531	1532
48	57.186	0.24	α -cadinene	C ₁₅ H ₂₄	1536	1538
49	57.476	0.12	α -calacorene	C ₁₅ H ₂₀	1542	1542
50	57.897	3.00	elemol	C ₁₅ H ₂₆ O	1549	1549
51	58.246	0.18	204 161 121 105 93 69	-	1555	-
52	58.548	0.14	204 161 121 109 93 41	-	1561	-
53	58.767	2.11	(E)-nerolidol	C ₁₅ H ₂₆ O	1564	1564
54	59.140	0.59	202 69 41 79 93 109 55	-	1571	-
55	59.372	4.33	222 161 119 105 91 81	-	1575	-
56	59.444	0.55	spathulenol	C ₁₅ H ₂₄ O	1576	1576
57	59.735	0.75	caryophyllene oxide	C ₁₅ H ₂₄ O	1582	1581
58	60.000	0.19	204 161 119 105 91 81	-	1586	-
59	60.332	0.18	202 159 97 83 69 55	-	1592	-
60	60.622	1.41	220 121 107 93 88 67	-	1597	-
61	60.851	0.26	204 122 107 93 81 69	-	1602	-
62	61.179	1.46	humulene oxide	C ₁₅ H ₂₄ O	1608	1604
63	61.358	0.55	222 179 105 91 69 41	-	1611	-
64	62.25	1.35	220 119 93 81 67 41	-	1628	-
65	62.413	0.97	γ -eudesmol	C ₁₅ H ₂₆ O	1631	1631
66	63.024	5.89	τ-muurolol	C ₁₅ H ₂₆ O	1642	1642
67	63.225	0.84	204 161 119 105 95	-	1646	-

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
68	63.411	0.67	β -eudesmol	C ₁₅ H ₂₆ O	1649	1649
69	63.749	9.71	α-cadinol	C ₁₅ H ₂₆ O	1655	1653
70	64.460	0.16	200 157 142 123 93 69	-	1669	-
71	64.929	0.10	cadalene	C ₁₅ H ₁₈	1677	1674
72	65.504	0.38	204 161 119 105 84 81	-	1688	-
73	66.212	0.12	220 177 159 131 117	-	1701	-
74	67.089	0.06	220 187 159 145 131	-	1719	-
75	68.005	0.21	oplopanone	C ₁₅ H ₂₆ O ₂	1736	1730
76	70.851	0.06	1-octadecene	C ₁₈ H ₃₆	1792	1793
77	71.662	0.07	147 119 91 77 69 55	-	1808	-
78	72.253	0.10	187 159 119 107 93 77	-	1821	-
79	74.520	0.79	benzyl salicylate	C ₁₄ H ₁₂ O ₃	1867	1869
80	80.459	0.06	97 91 83 69 57 55	-	1993	-
81	81.149	0.05	(8 β .13 β)-kaur-16-ene	C ₂₀ H ₃₂	2008	2012
82	82.173	2.42	kaur-16-ene	C ₂₀ H ₃₂	2032	2041
83	85.762	0.11	Coronarin E	C ₂₀ H ₂₈ O	2123	2136
84	87.726	0.03	207 91 83 77 69 57	-	2194	-
85	89.042	0.04	298 146 123 91 77	-	2259	-
86	89.229	0.04	281 109 96 81 67 55	-	2269	-
87	89.325	0.26	9-tricosene	C ₂₃ H ₄₆	2274	2278
88	89.806	0.15	324 99 85 71 57 43	-	2299	-
89	90.032	0.34	298 174 146 131 109	-	2313	-
90	90.719	0.03	207 151 81 69 55 40	-	2358	-
91	90.956	0.52	314 190 162 95 81 55	-	2373	-
92	92.293	0.05	283 109 95 82 67 55	-	2470	-
93	92.346	0.16	281 111 97 83 69 57	-	2475	-

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
94	92.402	0.19	290 108 95 79 67 55	-	2479	-
95	92.658	0.04	327 113 99 85 71 57	-	2498	-
Total identified		87.43 %				
Monoterpenes / Monoterpenoid s		20.28 %				
Sesquiterpenes / Sesquiterpenoid s		63.21 %				

RT: retention time; RI: retention indices; RI litt. from NIST 2020 database. NI:

Not identified (12.55%). Bold: main compounds.

Hedychium flavescens leaves EO analysis

The EO sample of *H. flavescens* leaves was found to be relatively simple and to contain mainly three compounds: β -pinene (47.81%), α -pinene (18.74%) and β -caryophyllene (17.47%), which together accounted for 84.02% of the sample (Figure 3). As shown in Table 2, in total, 96.26% of the EO was identified. In addition of these three compounds, monoterpene and sesquiterpene derivatives were found in smaller proportions such as the monoterpene hydrocarbons D-limonene (0.60%), β -myrcene (0.16%), γ -terpinene (0.05%) and sesquiterpene hydrocarbons bicyclogermacrene (2.65%), germacrene B (1.29%) and β -cadinene (1.14%).




Figure 3. Chromatogram of *Hedychium flavescens* leaves essential oil.

Table 2. Chemical composition of *Hedychium flavescens* leaves essential oil.

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
1	17.625	18.74	α-pinene	C ₁₀ H ₁₆	931	937
2	18.553	0.04	camphene	C ₁₀ H ₁₆	945	952
3	20.553	47.81	β-pinene	C ₁₀ H ₁₆	974	979
4	21.715	0.16	β-myrcene	C ₁₀ H ₁₆	991	991
5	23.562	0.05	α-terpinene	C ₁₀ H ₁₆	1018	1017
6	24.093	0.02	<i>p</i> -cymene	C ₁₀ H ₁₄	1025	1025
7	24.183	0.60	limonene	C ₁₀ H ₁₆	1026	1030
8	24.352	0.42	eucalyptol	C ₁₀ H ₁₈ O	1029	1032
9	25.113	0.05	(<i>Z</i>)-β-ocimene	C ₁₀ H ₁₆	1040	1038
10	25.799	0.05	(<i>E</i>)-β-ocimene	C ₁₀ H ₁₆	1049	1049
11	26.391	0.05	γ-terpinene	C ₁₀ H ₁₆	1058	1060
12	28.483	0.03	α-terpinolene	C ₁₀ H ₁₆	1087	1088
13	29.398	0.02	linalool	C ₁₀ H ₁₈ O	1100	1099
14	31.920	0.01	pinocarveol	C ₁₀ H ₁₆ O	1137	1138
15	34.619	0.02	terpinen-4-ol	C ₁₀ H ₁₈ O	1176	1177
16	35.661	0.03	1-dodecene	C ₁₂ H ₂₄	1191	1190

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
17	45.195	0.90	δ -elemene	C ₁₀ H ₁₈ O	1336	1338
18	47.567	0.02	α -copaene	C ₁₁ H ₁₈	1374	1376
19	48.606	0.24	β -elemene	C ₁₅ H ₂₄	1391	1391
20	49.650	0.06	β -maaliene	C ₁₅ H ₂₄	1408	1398
21	50.269	17.47	β-caryophyllene	C ₁₅ H ₂₄	1418	1419
22	51.215	0.33	γ -elemene	C ₁₅ H ₂₄	1434	1434
23	51.673	0.66	guaia-6,9-diene	C ₁₅ H ₂₄	1442	1443
24	51.996	1.19	204 161 133 119 105	-	1447	-
25	52.257	1.20	humulene	C ₁₅ H ₂₄	1452	1454
26	52.658	0.27	(3 <i>E</i> ,6 <i>E</i>)- α -farnesene	C ₁₅ H ₂₄	1458	1457
27	52.683	0.27	allo-aromadendrene	C ₁₅ H ₂₄	1459	1461
28	53.430	0.12	204 161 121 105 93 55	-	1471	-
29	53.659	0.08	204 161 133 119 105	-	1475	-
30	53.902	0.55	germacrene D	C ₁₅ H ₂₄	1479	1481
31	54.202	0.02	204 189 147 122 108	-	1484	-
32	54.369	0.08	β -selinene	C ₁₅ H ₂₄	1487	1486
33	54.521	0.14	204 189 161 104 91	-	1490	-
34	54.818	2.65	bicyclogermacrene	C ₁₅ H ₂₄	1495	1495
35	55.057	0.02	161 136 105 95 91 53	-	1499	-
36	55.298	0.45	204 147 107 93 81 67	-	1503	-
37	55.454	1.14	β -cadinene	C ₁₅ H ₂₄	1506	1518
38	56.277	0.89	204 161 136 121 105	-	1520	-
39	56.647	0.02	122 119 107 91 77	-	1527	-
40	57.524	0.03	α -calacorene	C ₁₅ H ₂₀	1542	1542
41	58.010	0.02	106 91 79 41	-	1551	-
42	58.235	1.29	germacrene B	C ₁₅ H ₂₄	1555	1557

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
43	59.428	0.07	spathulenol	C ₁₅ H ₂₄ O	1576	1576
44	59.706	0.49	caryophyllene oxyde	C ₁₅ H ₂₄ O	1581	1581
45	59.987	0.44	204 161 119 105 91 81	-	1586	-
46	60.843	0.08	161 105 93 81 77 67	-	1601	-
47	61.360	0.18	204 179 161 119 105	-	1610	-
48	62.099	0.07	204 161 119 105 91	-	1624	-
49	62.309	0.43	isospathulenol	C ₁₅ H ₂₄ O	1627	1638
50	63.668	0.01	204 161 121 91 81	-	1651	-
Total identified		96.26 %				
Monoterpenes / Monoterpenoid s		68.08 %				
Sesquiterpenes / Sequiterpenoid s		28.16 %				

RT: retention time; RI: retention indices; RI litt. from NIST 2020 database. NI:

Not identified (3.72%). Bold: main compounds.

To our knowledge, the chemical composition of the EO obtained solely from the leaves of *H. flavescens* have not been published so far. In the most recent article on the EO of *H. flavescens* rhizomes, 78.05% of a sample obtained from a plant grown in India could be identified.[4] The main compounds described were β -pinene (24.77%) and α -pinene (11.17%), both present in large proportions in our sample (47.81 and 18.74%, respectively), together with β -caryophyllene and bicyclogermacrene.

***Pittosporum senacia* leaves EO analysis**

The relatively simple composition of the EO from *Pittosporum senacia* leaves was determined for more than 99% of the total EO (Figure 4). As presented in Figure 4. Chromatogram of *Pittosporum senacia* leaves essential oil.

Table 3, its chemical profile essentially contained four compounds: the C9 linear alkane hydrocarbon nonane (36.24%) and three monoterpene derivatives, β -pinene (25.11%), β -myrcene (19.19%) and α -pinene (7.36%), which together accounted for 87.9%. Moreover, the chemical profile included monoterpene derivatives such as limonene (1.46%), (Z)- and (E)- β -ocimene (0.01% and 0.09%, respectively), sesquiterpene derivatives like germacrene D (3.68%), humulene (0.40%), as well as other alkanes like decane (0.09%) and undecane (0.56%).




Figure 4. Chromatogram of *Pittosporum senacia* leaves essential oil.

Table 3. Chemical composition of *Pittosporum senacia* leaves essential oil.

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
1	15.492	36.24	nonane	C ₉ H ₂₀	900	900

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
2	17.233	0.03	α -thujene	C ₁₀ H ₁₆	925	929
3	17.578	7.36	α-pinene	C₁₀H₁₆	930	937
4	18.984	0.06	camphene	C ₁₀ H ₁₆	951	952
5	20.464	25.11	β-pinene	C₁₀H₁₆	973	979
6	21.675	19.19	β-myrcene	C₁₀H₁₆	991	991
7	22.628	0.09	decane	C ₁₀ H ₂₂	1004	1000
8	22.843	0.05	α -phellandrene	C ₁₀ H ₁₆	1008	1005
9	23.605	0.07	α -terpinene	C ₁₀ H ₁₆	1018	1017
10	24.131	0.06	<i>p</i> -cymene	C ₁₀ H ₁₄	1026	1025
11	24.183	1.46	limonene	C ₁₀ H ₁₆	1026	1030
12	25.145	0.01	(<i>Z</i>)- β -ocimene	C ₁₀ H ₁₆	1040	1038
13	25.818	0.09	(<i>E</i>)- β -ocimene	C ₁₀ H ₁₆	1050	1049
14	26.414	0.22	γ -terpinene	C ₁₀ H ₁₆	1058	1060
15	28.474	0.08	terpinolene	C ₁₀ H ₁₆	1087	1088
16	29.383	0.56	undecane	C ₁₁ H ₂₄	1100	1100
17	30.518	0.04	(<i>E</i>)-4,8-dimethyl- nona-1,3,7-triene	C ₁₁ H ₁₈	1116	1116
18	35.662	0.05	1-dodecene	C ₁₂ H ₂₄	1191	1190
19	36.632	0.05	decanal	C ₁₀ H ₂₀ O	1205	1206
20	45.180	0.81	δ -elemene	C ₁₅ H ₂₄	1336	1338
21	45.941	0.03	α -cubebene	C ₁₅ H ₂₄	1348	1351
22	47.279	0.05	α -ylangene	C ₁₅ H ₂₄	1370	1372
23	47.557	0.12	α -copaene	C ₁₅ H ₂₄	1374	1376
24	48.602	0.56	β -elemene	C ₁₅ H ₂₄	1391	1391
25	50.222	0.48	β -caryophyllene	C ₁₅ H ₂₄	1417	1419
26	50.795	0.15	β -copaene	C ₁₅ H ₂₄	1427	1432

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
27	51.134	0.15	γ -elemene	C ₁₅ H ₂₄	1433	1434
28	51.405	0.05	α -guaiene	C ₁₅ H ₂₄	1437	1439
29	51.682	0.10	guaia-6,9-diene	C ₁₅ H ₂₄	1442	1443
30	52.006	0.20	161 133 105 91 79 69	-	1447	-
31	52.242	0.40	humulene	C ₁₅ H ₂₄	1451	1454
32	52.842	0.03	<i>cis</i> -muurola- 4(15),5-diene	C ₁₅ H ₂₄	1461	1463
33	53.910	3.68	germacrene D	C ₁₅ H ₂₄	1479	1481
34	54.213	0.03	β -selinene	C ₁₅ H ₂₄	1485	1486
35	54.519	0.06	161 105 91 44	-	1490	-
36	54.720	0.07	161 105 91	-	1493	-
37	55.057	0.06	α -muurolene	C ₁₅ H ₂₄	1499	1499
38	55.304	0.21	α -elemene	C ₁₅ H ₂₄	1503	1462
39	55.834	0.11	γ -cadinene	C ₁₅ H ₂₄	1512	1513
40	56.407	0.39	cadina-1(10),4-diene	C ₁₅ H ₂₄	1523	1524
41	56.650	0.03	122 43	-	1527	-
42	58.244	0.65	germacrene B	C ₁₅ H ₂₄	1555	1557
43	60.289	0.10	1-hexadecene	C ₁₆ H ₃₂	1591	1592
44	61.085	0.04	rosifoliol	C ₁₅ H ₂₆ O	1606	1600
45	63.633	0.09	τ -muurolol	C ₁₅ H ₂₆ O	1653	1642
46	70.847	0.03	1-octadecene	C ₁₈ H ₃₆	1792	1793
Total identified			99.09 %			
Monoterpenes / Monoterpenoids			53.78 %			
Sesquiterpenes / Sesquiterpenoids			8.19 %			

RT: retention time; RI: retention indices; RI litt. from NIST 2020 database.

NI: Not identified (0.41%). Bold: main compounds.

Many of the metabolites present in the study by B. S. Jugreet et al.

published in 2021 on the EO of *P. senacia* (whole plant) from Mauritius [5], were also identified in our sample such as β -myrcene, germacrene D, α -muurolene, limonene, α -pinene, α -thujene, p-cymene, β -ocimene, β -elemene, β -caryophyllene, and humulene. The alkanes nonane, decane, and undecane were not reported in the article by B. S. Jugreet et al. However, some alkanes, including nonane, have already been identified in the EO of the genus *Pittosporum*, such as in an endemic species of the Philippines, *P. resiniferum* [20]. Interestingly, nonane, a compound described as insect attractant,[21] was found in 36.24% in our sample.

***Psidium cattleianum* floral tops EO analysis**

In the most recent study, a distillation yield of 0.83% was reported for EO from *P. cattleianum* leaves [8]. A total of 68 compounds were observed within the oil, 60 of which could be identified. β -Caryophyllene was the main compound found in 14.7%, followed by eucalyptol (11.7%) and γ -muurolene (5.6%). The second most recent paper, published in 2019, describes the EO of the leaves of *P. cattleianum*. The EO was mainly composed of sesquiterpene and monoterpene derivatives (47.8% and 28.7%, respectively). A total of 46 compounds were identified with β -caryophyllene (23.4%), caryophyllene oxide (11.4%) and α -pinene (11.3%) as the main compounds [9]. Our results are in

agreement with this study (Figure 5), whose main compounds described were β -caryophyllene and α -pinene at 23.42% and 11.31%, respectively, caryophyllene oxide being present in 1.67%. Indeed, the chemical profile of the EO sample from *P. cattleianum* floral tops was identified at 88.41%. As shown in Table 4, nearly 70% of the composition was consisting in sesquiterpene derivatives including the main compound, β -caryophyllene, present at 43.68%. In addition, fifteen monoterpene derivatives have been observed and identified. The four main monoterpenes were β -myrcene (4.67%), α -pinene (4.50%), (Z)- β -ocimene (3.23%) and terpinolene (2.98%).




Figure 5. Chromatogram of *Psidium cattleianum* leaves essential oil.

Table 4. Chemical composition of *Psidium cattleianum* leaves essential oil.

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
1	16.610	0.03	5,5-Dimethyl-1-vinylbicyclo[2,1,1]hexane	C ₁₀ H ₁₆	916	921

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
2	17.214	0.15	α -thujene	C ₁₀ H ₁₆	925	929
3	17.611	4.50	α-pinene	C ₁₀ H ₁₆	931	937
4	18.920	0.01	camphene	C ₁₀ H ₁₆	950	952
5	20.475	0.19	β -pinene	C ₁₀ H ₁₆	973	979
6	21.677	4.67	β-myrcene	C ₁₀ H ₁₆	991	991
7	22.467	0.24	α -phellandrene	C ₁₀ H ₁₆	1002	1005
8	22.864	0.14	3-carene	C ₁₀ H ₁₆	1008	1011
9	23.360	0.12	4-carene	C ₁₀ H ₁₆	1015	1009
10	23.901	0.19	<i>p</i> -cymene	C ₁₀ H ₁₄	1022	1025
11	24.181	1.07	limonene	C ₁₀ H ₁₆	1026	1030
12	24.984	3.23	(Z)-β-ocimene	C ₁₀ H ₁₆	1038	1038
13	25.710	0.56	(E)- β -ocimene	C ₁₀ H ₁₆	1048	1049
14	26.346	0.63	γ -terpinene	C ₁₀ H ₁₆	1057	1060
15	28.421	2.98	terpinolene	C ₁₀ H ₁₆	1086	1088
16	40.039	0.23	linalyl acetate	C ₁₂ H ₂₀ O ₂	1257	1257
17	45.984	0.27	161 136 121 105 93 67	-	1349	-
18	47.293	0.44	α -ylangene	C ₁₅ H ₂₄	1370	1372
19	47.581	1.18	α -copaene	C ₁₅ H ₂₄	1374	1376
20	48.127	0.07	α -bourbornene	C ₁₅ H ₂₄	1383	1384
21	48.598	0.12	204 121 108 93 81 55	-	1391	
22	49.495	0.03	tetradecene	C ₁₄ H ₂₈	1405	1392
23	49.641	0.03	204 189 161 119 105	-	1408	-
24	50.374	43.68	β-caryophyllene	C ₁₅ H ₂₄	1420	1419
25	50.818	0.42	γ -elemene	C ₁₅ H ₂₄	1427	1434
26	51.387	0.07	aromadendrene	C ₁₅ H ₂₄	1437	1440
27	51.672	0.10	204 161 133 119 105 91	-	1442	-

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
28	52.009	0.16	204 161 119 105 91 79	-	1447	-
29	52.278	6.49	humulene	C ₁₅ H ₂₄	1452	1454
30	52.709	0.18	9-epi-caryophyllene	C ₁₅ H ₂₄	1459	1466
31	53.640	2.32	γ-murolene	C ₁₅ H ₂₄	1475	1477
32	53.874	0.50	α-amorphene	C ₁₅ H ₂₄	1479	1482
33	54.197	1.27	β-selinene	C ₁₅ H ₂₄	1484	1486
34	54.307	0.29	204 133 119 107 93 79	-	1486	-
35	54.517	0.25	204 189 161 133 91	-	1490	-
36	54.727	1.70	204 161 133 119 105 93	-	1493	-
37	55.059	0.34	α-murolene	C ₁₅ H ₂₄	1499	1499
38	55.454	0.51	204 161 134 119 105	-	1506	-
39	55.575	0.26	122 109 93 79 69	-	1508	-
40	55.837	0.87	β-bisabolene	C ₁₅ H ₂₄	1513	1509
41	56.175	0.88	γ-cadinene	C ₁₅ H ₂₄	1519	1513
42	56.401	2.38	cadina-1(10),4-diene	C ₁₅ H ₂₄	1523	1524
43	57.015	2.35	204 189 161 133 105 91	-	1533	-
44	57.184	0.74	204 189 161 133 105	-	1536	-
45	57.385	2.06	selina-3,7(11)-diene	C ₁₅ H ₂₄	1540	1542
46	57.987	0.11	161 109 95 91 79 69	-	1551	-
47	58.238	1.84	germacrene B	C ₁₅ H ₂₄	1555	1557
48	58.604	0.17	204 189 133 119 105	-	1562	-
49	58.749	0.33	nerolidol	C ₁₅ H ₂₆ O	1564	1564
50	58.978	0.19	204 161 123 111 69 55	-	1568	-
51	59.701	1.67	caryophyllene oxyde	C ₁₅ H ₂₄ O	1581	1581
52	60.284	0.44	204 161 119 105 91 79	-	1591	-
53	60.593	0.07	204 161 119 107 93 79	-	1597	-

Peak #	RT (min)	Area %	Compound	Formula	RI	RI litt.
54	60.746	0.10	161 133 121 93 82 77	-	1600	-
55	60.956	0.09	204 161 133 119 105 95	-	1603	-
56	61.162	0.17	147 138 109 96 93 67	-	1607	-
57	61.623	0.29	202 187 131 123 91	-	1616	-
58	62.084	0.71	204 161 119 105 91	-	1624	-
59	62.221	0.78	204 161 119 105 91	-	1627	-
60	62.424	0.54	204 179 161 119 105	-	1631	-
61	62.642	0.14	159 136 107 91 79 69	-	1635	-
62	62.969	1.09	τ -cadinol	C ₁₅ H ₂₆ O	1641	1640
63	63.198	0.36	204 161 119 105 93	-	1645	-
64	63.489	0.41	202 187 121 105 91	-	1651	-
65	63.630	0.87	τ -muurolol	C ₁₅ H ₂₆ O	1653	1642
66	64.190	0.20	204 189 161 133 107	-	1664	-
67	65.113	0.06	126 119 111 77 55	-	1681	-
68	65.807	0.49	eudesm-7(11)-en-4-ol	C ₁₅ H ₂₆ O	1694	1692
Total identified		88.41 %				
Monoterpenes / Monoterpenoids		18.70 %				
Sesquiterpenes / Sesquiterpenoids		69.99 %				

RT: retention time; RI: retention indices; RI litt. from NIST 2020 database.

NI: Not identified (11.61%). Bold: main compounds.

Olfactory analysis

Olfactory evaluation of the four essential oils was performed by a panel of 4 persons by GC/MS-O analysis. The objective was to obtain an overview of

the main contributors to the overall scent of each EO which was found to be generally woody, a family of olfactory properties highly priced in modern perfumery.

In Table 5, the main aroma-active compounds and their odor properties are presented.

Table 5. Overview of the main contributors of the four EOs.

	RT (min)	RI _{apolar}	Annotated compound	Scent description
<i>H. gardnerianum</i> , flowers EO	9.55	1144	Sabinol*	Floral
	17.23	N.D.	Unknown**	Woody
	22.48	16.94	Unknown	Floral
<i>H. flavescens</i> , leaves, EO	8.55	1093	Terpinolene	Woody
	17.35	1512	<i>trans</i> - α -bisabolene	Woody, rose
<i>P. senacia</i> , leaves, EO	6.61	N.D.	β -pinene	Pine, Green
	8.53	1092	Terpinolene	Woody
<i>P. cattleianum</i> , floral tops, EO	6.42	979	β -pinene	Pine, Green
	8.26	1078	Unknown**	Woody
	11.80	1261	Unknown	Citrus, spicy

*Tentative assignment. **No peak detectable.

Hedychium gardnerianum flowers EO presented three characteristic areas, a woody area identified by 2 panelists out of 4, and two floral area for 2 panelists out of 4.

GC/MS-O analysis of EO from *Hedychium flavescens* leaves revealed a main contributor with a fresh, woody and green scent which was annotated as

γ -elemene. The presence of woody, earthy, and green area was highlighted by 5 panelists out of 5, floral and rose area by 4 panelists out of 5 and mint, herbal, and citrus area by 2 panelists out of 5.

Main aroma-active compounds of *Pittosporum senacia* leaves EO were annotated as β -myrcene with a green, woody and spicy scent and as terpinolene with a moss and woody scent. The GC/MS-O analysis revealed herbal, and pine tree area for 5 panelists out of 5 as well as floral and citrus area for 5 panelists out of 5.

Finally, *Psidium cattleianum* floral tops EO showed three characteristic scents with β -myrcene and two unknown compounds, eluting at 8.9 min and 13.25 min, respectively, both with floral scent. In addition of these markers, pine tree scent was found by 4 panelists out of 4, woody, and earthy area by 4 panelists out of 4, floral area by 2 panelists out of 4, and citrus with mossy/earthy area by 2 panelists out of 4. In addition to these main contributors, others smaller contributors as linalool, with floral and fresh scent, was detected in each of the four EOs GC/MS-O analyses.

CONCLUSION

In general, the chemical composition of an EO varies according to many factors such as the year of harvest, the geographical area, the climate, the storage of the raw material, the duration of storage of the plant before extraction, the extraction process [22]. Two EOs from the same species may

therefore have a different chemical composition. Nevertheless, the chemical profile and chemical markers may be specific to a given plant or chemotype. The analysis and authentication tasks can be sometimes complicated by conformity or adulteration issues [23, 24].

In this context, the main goal of our work was to characterize for the first time the EOs of the four species harvested on Reunion Island and to compare their chemical profiles with those described in the literature with samples collected in other locations. We used two complementary analyses, gas chromatography coupled with mass spectrometry (GC/MS) and with flame ionization detector (GC-FID), to obtain the most complete description. Additionally, an olfactory analysis was performed on these four EOs by gas chromatography coupled with mass spectrometry and olfactometry (GC/MS-O). The chemical composition of each essential oil was identified in more than 87.43% for *H. gardnerianum* flowers, 96.26% for *H. flavescens* leaves, 99.09% for *P. senecio* leaves and 88.41% for *P. cattleianum* floral tops.

The chemical composition of *H. gardnerianum* EO, mostly composed of mono- and sesquiterpene hydrocarbons of low complexity, was found to be consistent with other EOs within natural variability due to differences in their geographical origin (islands from Atlantic, Pacific, and Indian oceans). The sample was mainly composed of sesquiterpene derivatives including β -farnesene (12.05%), α -cadinol (9.71%), α -farnesene (7.09%), τ -muurolol (5.89%)

and δ -cadinene (5.83%). The monoterpene derivatives identified accounted for about 20% (mostly α -pinene, β -pinene and humulene).

For the EO from *H. flavescens* leaves, our study provided the first description of its chemical composition, which appeared to be rather simple and to contain mostly hydrocarbons. The main compounds were found to be β -pinene (47.81%), α -pinene (18.74%) and β -caryophyllene (17.47%), which together accounted for 84.02% of the sample.

For the *P. senacia*, while β -myrcene was found to be an important constituent of the EO, as was the case for material from Madagascar and Mauritius, which are geographically close to French Reunion, the composition was different but consistent, and the striking difference was the presence of nonane. The oil primarily contained nonane (36.24%), β -pinene (25.11%), β -myrcene (19.19%) and α -pinene (7.36%), together accounting for 87.9% of the composition.

Lastly, the composition of the *P. cattleianum* EO was very consistent with other studies, with 70% sesquiterpene derivatives including the main compound, β -caryophyllene, present at 43.68%.

With these compositions mostly based on mono- and sesquiterpene hydrocarbons, it came as no surprise that the panelists described the scent of these EOs as herbal, citrus, green, pine tree, and in some cases floral and woody.

With these results in hand, these endemic essential oils from Reunion could receive in the future further attention for applications in fragrance and cosmetic products. Future work could include the determination of odor impact molecules by aroma extract dilution analysis based on the GC-O studies presented herein.

AUTHOR CONTRIBUTIONS

CONCEPTUALIZATION, M.C AND S.A.; METHODOLOGY, M.C AND S.A.; INVESTIGATION, M.C, M.T. AND S.A.; RESOURCES, S.A.; DATA CURATION, M.C, M.T. AND S.A.; WRITING – ORIGINAL DRAFT PREPARATION, M.C AND S.A.; WRITING – REVIEW & EDITING, M.C, M.T. AND S.A.; SUPERVISION, S.A.; PROJECT ADMINISTRATION, M.C, M.T. AND S.A.

ACKNOWLEDGEMENTS

This work was supported by Université Côte d’Azur, CNRS and Technopole de la Réunion in the framework of the Insola Scent project (Sophie Afchain). We are grateful to Charlotte Richard and Julie Abdalla for contributing to the olfactory evaluation during GC-O analyses and Dr. Catherine Buchanan for proofreading.

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

- [1] Medeiros, J.R., Campos, L.B., Mendonça, S.C., Davin, L.B. and Lewis, N.G. Composition and antimicrobial activity of the essential oils from invasive species of the Azores, *Hedychium gardnerianum* and *Pittosporum undulatum*. *Phytochemistry* 2003, 64, 561-565.
- [2] Ali, S., Sotheeswaran, S., Tuiwawa, M. and Smith, R.M. Comparison of the composition of the essential oils of *Alpinia* and *Hedychium* species-essential oils of Fijian plants, Part 1. *J. Essent. Oil Res.* 2002, 14, 409-411.
- [3] Weyerstahl, P., Marschall, H., Thefeld, K. and Subba, G.C. Constituents of the essential oil from the rhizomes of *Hedychium gardnerianum* Roscoe. *Flavour Fragr. J.* 1998, 13, 377-388.
- [4] Ray, A., Sudipta, J., Basudeba, K., Ambika, S., Pratap, C.P., Sanghamitra, N. and Namita, M. Volatile metabolite profiling of ten *Hedychium* species by gas chromatography mass spectrometry coupled to chemometrics. *Ind. Crops Prod.* 2018, 126, 135-142.
- [5] Jugreet, B.S., Ibrahime, S.K., Zengin, G., Abdallah, H.H. and Mahomoodally, M.F. GC/MS profiling, in vitro and in silico pharmacological screening and principal component analysis of essential oils from three exotic and two endemic plants from Mauritius. *Chem. Biodiv.* 2021, 18, e2000921.
- [6] Jugreet, B.S. and Mahomoodally, M.F. Essential oils from 9 exotic and endemic medicinal plants from Mauritius shows in vitro antibacterial and antibiotic potentiating activities. *S. Afr. J. Bot.* 2020, 132, 355-362.
- [7] Mananjarasoa, E., Mananjarasoa, E., Rakotovao, M., Ramanoelina, A.R.P., Andriantsiferana, M.H. and Ravaonindrina, N. Composition and antimicrobial activity of leaf oil of *Pittosporum senacia* var. *coursii* Cufodontis. *J. Essent. Oil Res.* 1998, 10, 459-462.
- [8] Savoldi, T.L., Glamočlija, J., Soković, M., Gonçalves, J.E., Ruiz, S.P., Linde, G.A., Gazim, Z.C. and Colauto, N.B. Antimicrobial activity of essential oil from *Psidium cattleianum* Afzel. ex Sabine leaves. *Blacpma* 2020, 19, 614-627.
- [9] Vasconcelos, L.C., de Souza Santos, E., de Oliveira Bernardes, C., da Silva Ferreira, M.F., Ferreira, A., Tuler, A.C., Carvalho, J.A.M., Pinheiro, P.F. and Praça-Fontes, M.M. Phytochemical analysis and effect of the essential oil of *Psidium* L. species on the initial development and mitotic activity of plants. *Environ. Sci. Pollut. Res.* 2019, 26, 26216-26228.
- [10] Scur, M.C., Pinto, F.G.S., Pandini, J.A., Costa, W.F., Leite, C.W. and Temponi, L.G. Antimicrobial and antioxidant activity of essential oil and different plant extracts of *Psidium cattleianum* Sabine. *Braz. J. Biol.* 2016, 76, 101-108.
- [11] Castro, M.R., Victoria, F.N., Oliveira, D.H., Jacob, R.G., Savegnago, L. and Alves, D. Essential oil of *Psidium cattleianum* leaves: antioxidant and antifungal activity. *Pharm. Biol.* 2015, 53, 242-250.
- [12] Marques, F.A., Wendler, E.R., Sales Maia, B.H.L.N., Coffani-Nunes, J.V., Campana, J. and Guerrero, P.G. Volatile oil of *Psidium cattleianum* Sabine from the Brazilian atlantic forest. *J. Essent. Oil Res.* 2008, 20, 519-520.
- [13] Chen, J., Lichwa, J. and Ray, C. Essential oils of selected Hawaiian plants and associated litters. *J. Essent. Oil Res.* 2007, 19, 276-278.
- [14] Pino, J.A., Bello, A., Urquiola, A., Marbot, R. and Martí, M. Leaf oils of *Psidium parvifolium* Griseb. and *Psidium cattleianum* Sabine from Cuba. *J. Essent. Oil Res.* 2004, 16, 370-371.

- [15] Tucker, A.O., Maciarello, M.J. and Landrum, L.R. Volatile leaf oils of American Myrtaceae. III. *Psidium cattleianum* Sabine, *P. friedrichsthalianum* (Berg) Niedenzu, *P. guajava* L., *P. guineense* Sw., and *P. sartorianum* (Berg) Niedenzu. J. Essent. Oil Res. 1995, 7, 187-190.
- [16] Suksathan, R., Sookkhee, S., Anuntalabhochai, S. and Chansakaow, S. Chemical composition and antibacterial activity of rhizome oils from five *Hedychium* species. Nat. Prod. Commun. 2013, 8, 519-522.
- [17] Sabulal, B., Varughese, G., Dan, M. and Pradeep, S. Chemical composition and antimicrobial activities of the essential oils from the rhizomes of four *Hedychium* Species from South India. J. Essent. Oil Res. 2007, 19, 93-97.
- [18] Gurib-Fakim, A., Maudarbaccus, N., Leach, D., Doimo, L. and Wohlmuth, H. Essential oil composition of Zingiberaceae species from Mauritius. J. Essent. Oil Res. 2002, 14, 271-273.
- [19] Wang, Q., Liu, H., Zhang, M., Liu, S., Hao, Y. and Zhang, Y. MdMYC2 and MdERF3 positively co-regulate α -farnesene biosynthesis in apple. Front. Plant Sci. 2020, 11,
- [20] Alimboyoguen, A.B., Castro-Cruz, K.D., Altena, I.v. and Ragasa, C. Triterpenes from *Pittosporum resiniferum* Hemsl. Int. J. Toxicol. Pharmacol. Res. 2016, 8, 261-262.
- [21] Mendesil, E., Bruce, T.J.A., Woodcock, C.M., Caulfield, J.C., Seyoum, E. and Pickett, J.A. Semiochemicals used in host location by the coffee berry borer, *Hypothenemus hampei*. Journal of Chemical Ecology 2009, 35, 944-950.
- [22] Arruda, M., Viana, H., Rainha, N., Neng, N.R., Rosa, J.S., Nogueira, J.M.F. and Barreto, M.d.C. Anti-acetylcholinesterase and antioxidant activity of essential oils from *Hedychium gardnerianum* Sheppard ex Ker-Gawl. Molecules 2012, 17, 3082-3092.
- [23] T. K. T. Do, F. Hadji-Minaglou, S. Antoniotti and Fernandez, X. Essential oil authenticity. Trends Anal. Chem. 2015, 66, 146–157.
- [24] Pierson, M., Fernandez, X. and Antoniotti, S. Type and magnitude of non-compliance and adulteration in neroli, mandarin and bergamot essential oils purchased on-line: potential consumer vulnerability. Sci. Rep. 2021, 11, 11096.