



HAL
open science

Towards integrated pest and pollinator management in tropical crops

Isabelle Merle, Juliana Hipólito, Fabrice Requier

► **To cite this version:**

Isabelle Merle, Juliana Hipólito, Fabrice Requier. Towards integrated pest and pollinator management in tropical crops. *Current Opinion in Insect Science*, 2022, 50, pp.100866. 10.1016/j.cois.2021.12.006 . hal-04293793

HAL Id: hal-04293793

<https://hal.science/hal-04293793>

Submitted on 22 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Towards integrated pest and pollinator management in tropical crops**

2

3 Isabelle Merle¹, Juliana Hipólito^{2,3}, Fabrice Requier^{1,*}

4 ¹ Université Paris-Saclay, CNRS, IRD, UMR Évolution, Génomes, Comportement et

5 Écologie, 91198, Gif-sur-Yvette, France

6 ² Instituto de Biologia, Universidade Federal da Bahia, Salvador, BA, Brasil

7 ³ Instituto Nacional de Pesquisas da Amazônia, Manaus, AM, Brasil

8 * Correspondence: Fabrice Requier; fabrice.requier@egce.cnrs-gif.fr

9

10 **ORCID**

11 *Isabelle Merle* <https://orcid.org/0000-0003-3402-2892>

12 *Juliana Hipólito* <https://orcid.org/0000-0002-0721-3143>

13 *Fabrice Requier* <https://orcid.org/0000-0003-1638-3141>

14

15 **Running head:** IPPM in tropical crops

16

17 **Word count:**

18 Abstract = 134 words (120 words suggested)

19 Main text = 2,317 words (2,000 words suggested)

20 References = 156 (including 102 for the systematic review)

21 Artwork = 1 figure + 1 table + 1 box (max. 4)

22

23

24 **Abstract**

25 Biotic pollination and pest control are two critical insect-mediated ecosystem services
26 that support crop production. Although management of both services is usually
27 treated separately, the new paradigm of *Integrated Pest and Pollinator Management*
28 (IPPM) suggests synergetic benefits by considering them together. We reviewed the
29 management practices in two major tropical perennial crops: cocoa and coffee, to
30 assess IPPM applications under the tropics. We found potential synergies and
31 antagonisms among crop pest and pollination management, however, very few
32 studies considered these interactions. Interestingly, we also found management
33 practices focusing mainly on a single service mediated by insects although species
34 can show multiple ecological functions as pests, natural enemies, or pollinators. The
35 tropics represent a promising area for the implementation of IPPM and future
36 research should address this concept to move towards a more sustainable
37 agriculture.

38

39 **Keywords:** Biodiversity; Pest control; Cocoa; Coffee; Ecosystem services; IPPM;
40 Landscape management; Pollination

41 **Introduction**

42 Insect-mediated ecosystem services are critical in agriculture [1**]. Beyond interac-
43 tions between crops and their abiotic environment, crop yield is driven by ecosystem
44 services including pollination and pest control and pollination, which are governed by
45 antagonistic and mutualistic biotic interactions [2]. Animal pollination is critical for
46 yield improvement of three-quarters of the world's major crops [3]. Overall, 5 to 8% of
47 global crop production would be lost without animal pollination [4]. A wide diversity of
48 insects are often assessed as efficient pollinators [5,6*] but pollination limitation is
49 frequently driven by a lack of insect pollinators [7]. Crop pests herbivory is another
50 well-known driver of important crop yield losses due to a decrease in plant survival,
51 growth, and reproduction [8]. Worldwide yield losses are estimated from 7.9% to
52 15.1% [8]. Biological pest control is defined as a process by which predators, parasi-
53 toids or pathogens reduce pest abundance. Apart from alleviating the negative im-
54 pact from crop pest, managing the diversity of natural pest enemies to support crop
55 yield could help reduce both the use of synthetic pesticides in agriculture [9*] and the
56 associated detrimental effects on biodiversity and human well-being [10]. Given the
57 evidence of benefits from pest control and pollination on crop production, manage-
58 ment practices should be developed to foster the ecosystem services and mitigate
59 the ecosystem disservices provided by insects [1**,11].

60 Crop pest control is commonly performed by means of species management
61 approaches such as pesticide use [12–14]. However, many concerns are associated
62 with over-reliance on pesticide applications including not only pest resistance, out-
63 breaks of secondary pests, but also loss of natural enemies [9*]. Alternatively, there
64 is evidence that landscape management can promote the presence of natural ene-
65 mies, and thus enhance biological control which could reduce pesticide application
66 [18**–21**]. Pest control practices are distinguished between human input-based
67 practices and biodiversity based, respectively [18**,19**], along with the concept of
68 Integrated Pest Management (IPM). On the other hand, the management of crop pol-
69 lination was also divided among different strategies from human input-based practic-
70 es (e.g. the implementation of managed beehives [11]) to habitat and resource pro-
71 motion to maintain or attract wild pollinators [20,21]. Nevertheless, the management
72 practices of pest control and pollination services are usually treated independently
73 [18**]. IPM has been used as a response to mitigate pollinator decline, yet IPM itself
74 is not necessarily "pollinator friendly" [18**]. The impact on yield is affected by the

75 success of both pollination and insect pest control services, thus a more holistic ap-
76 proach is essential.

77

78 **New paradigm of *Integrated Pest and Pollinator Management (IPPM)***

79 The concept of Integrated Pest and Pollinators Management (IPPM) emerged in
80 2015 with the main goal of considering adverse effects and synergies of pest control
81 management on pollinators and *vice versa* [18**,19**,22,23]. IPPM has been further
82 defined by categorizing and prioritizing co-management practices to limit adverse
83 environmental impacts. A pyramid has thus been proposed as a decision support tool
84 for choosing preventive and biodiversity-based management practices rather than
85 curative and human-based ones [18**,19**] (**Figure 1**). At the top of the pyramid are
86 the human-based practices, such as pesticide application, which are widely used for
87 their effectiveness on target species but are also highly criticized for their effects on
88 beneficial insects [24]. These practices involve a direct management of targeted spe-
89 cies (e.g. controlling pests or fostering pollinators) and are differentiated in two man-
90 agement strategies according to the use of biotic or abiotic inputs (**Figure 1**). The
91 biodiversity based-practices stand at the base of the pyramid and are differentiated
92 among field and landscape management strategies [19**] (**Figure 1**). Landscape
93 components, namely habitat quantity and density, have a primary role in maintaining
94 insect communities [25**]. Landscape simplification affects pollination and pest con-
95 trol services through negative effects on the abundance, richness and diversity of
96 pollinators and predators [26**]. Consequently, the objective in such practices is no
97 longer to directly manage the arthropod population but to manage their dynamics
98 through the provision of nesting sites and food resources. However, while recent
99 studies applied the concept of IPPM in field-based trials [25**], IPPM remains essen-
100 tially conceptual since its establishment in 2015 [23]. Here, we carried out a system-
101 atic review of the management practices of pest control and pollination services in
102 two major tropical crops: cocoa and coffee, in order to assess for the first time IPPM
103 applications on those crops.

104

105 **IPPM application in tropical crops**

106 Coffee (*Coffea canephora* and *Coffea arabica*) and cocoa (*Theobroma cacao*) are
107 two tropical crops with different levels of animal-mediated pollination required for
108 yield improvement. The species *T. cacao* and *C. canephora* have a high dependence

109 on pollinators for fruit set, *C. arabica* has a moderate dependence [3]. Coffee and
110 cocoa suffer important pressures from crop pests [27,28]. We performed our system-
111 atic review on these two crops by selecting the scientific publications published since
112 the emergence of the IPPM concept in 2015 (**Box 1**). Despite the importance of in-
113 sects to coffee and cocoa –as pollinators, pests and natural enemies– very few stud-
114 ies have applied the IPPM concept. We found no study addressing both pollination
115 and pest control services in cocoa and only three in coffee. Some studies assessed
116 the effects of shade cover in agroforestry systems, ground cover, and distance to
117 forest [29*,30]–or the use of vector-disseminated biopesticides [31]–on both services
118 (**Table 1**). Otherwise, most studies focused on a single service or a single insect
119 species mediating that service. Nevertheless, some studies considered a more holis-
120 tic framework by assessing arthropods’ community without considering their ecologi-
121 cal function [32,33*]. Other studies focused only on pest and predator communities
122 [17,34–47] or pollinators and predators [48*–51*].

123 Overall, we found fewer experiments conducted on the pollination service
124 compared to the pest control service (**Figure 1**). Moreover, in cocoa and coffee
125 crops, pollination management seems to depend to a greater extent on the concept
126 of IPM. Conversely to pest management, the promotion of pollination relies more on
127 biodiversity-based practices at the landscape scale than on human practices. Con-
128 sidering both crops, the impact of abiotic and biotic inputs is studied on the pollination
129 service in only 10% of cases (6 experiments) compared to 90% on the pest control
130 service (53 experiments). Conversely, 69% of the experimental units (35 experi-
131 ments) investigated the effect of landscape-level practices on pollination, compared
132 to 31% (16 experiments) on pest control (**Figure 1**). While IPPM encourages the use
133 of landscape-level practices as a priority, plot-level practices are the most common
134 since they represent 57% of the practices studied on cocoa (11 pollination experi-
135 ments vs. 19 pest control experiments) and 49% on coffee (36 pollination experi-
136 ments vs. 49 pest control experiments). This particularity could be explained by the
137 fact that coffee and cocoa plantations are complex systems where the implementa-
138 tion of agroecological practices within the plot, such as agroforestry [52], is easier
139 than in annual crop plantations. Importantly, we found that landscape level manage-
140 ment strategies are rarely studied in cocoa compared to coffee for the pollination ser-
141 vice. It is likely that this difference is explained by the short distance foraging behav-

142 ior of the main cocoa pollinators (e.g. small flies) [53] compared to those of coffee
143 crop (e.g. bees) [54*].

144

145 **Synergies and antagonisms among crop pest and pollination managements**

146 Complementary and synergistic effects can occur in crop management practices with
147 potential positive interactions between pollinators and natural enemies on crop yield
148 [55]. We examined such synergies but also potential antagonisms among the prac-
149 tices performed for crop pest and pollination management in cocoa and coffee. Sev-
150 eral practices were studied specifically on one of the two services (i.e. pollination or
151 pest control) in both cocoa and coffee, although there were fewer studies on cocoa
152 (**Table 1**). The large majority of the practices involving abiotic and biotic inputs (i.e.
153 pesticides, traps, biopesticides and biocontrol agents) were evaluated for their effec-
154 tiveness on pest control but rarely on pollination service (**Table 1**). However, these
155 practices can have antagonistic effects by negatively affecting the pollination service
156 [24] or the community of beneficial arthropods [56]. Surprisingly, we found an exper-
157 iment showing an increase of pollination service linked with pesticide use. This result
158 can be explained by the family of the pesticide used, a fungicide, and by its effect on
159 the plant physiology by increasing flower retention capability that could attract more
160 insect pollinators [57]. Similarly, several biodiversity-based practices applied at the
161 plot level and landscape level (e.g. mechanical management such as sanitary har-
162 vesting, pruning and exclusion netting, and the practices of biodiversity conservation
163 such as bird-mediated predation) were only considered affecting pest control (**Table**
164 **1**). Nevertheless, these practices may also lead to antagonistic adverse effects on
165 pollination or pest control services if, for example, birds also predate on pollinators
166 and predators [22].

167 On the other hand, practices such as promoting floral resources in the plot and
168 enhancing abundance of semi-natural habitats at the landscape scale are almost ex-
169 clusively studied on the pollination service. However, studies carried out in temperate
170 regions, i.e. where farmers can receive financial incentive to set flower strip, show
171 benefits of such practices on pest control [58], suggesting that floral resources and
172 semi-natural habitats could represent an interesting approach that offers a synergetic
173 practice on both services. For instance, breeding substrates, vegetal ground cover
174 and agroforestry are practices that could validate the IPPM concept through syner-
175 getic effects among pest control and pollination. However, the impact of agroforestry

176 on insect-mediated services is mixed (**Table 1**). This result is likely due to the large
177 diversity of agroforestry systems with variability in tree size and the species of shade
178 trees [59] and the differences in the survey scales chosen by the different studies.
179 Shade tree species may affect their attractiveness and repellency to arthropod spe-
180 cies. The size of the trees influences the microclimate, which in turn affects the de-
181 velopment cycle of arthropods in various ways, depending on the species considered
182 [52,53]. Surprisingly, most of the practices studied at the landscape scale aim to
183 promote forest areas as arthropod-friendly habitats. Indeed, little attention has been
184 paid to the landscape composition in terms of crops [60]. However, these crops can
185 share the same pest or pollinator species assemblage, or can host arthropods per-
186 forming multiple functions in the cropping system.

187

188 **Multiple services mediated by insects**

189 Our systematic review confirms that pest control and pollination management are
190 mainly treated separately in cocoa and coffee (**Figure 1**). Interestingly, we also found
191 that most management practices focus on targeted insect species for a single service
192 (i.e. pollination or pest control) although numerous species have multiple ecological
193 functions as crop pests, natural enemies or pollinators. In other words, very few
194 works consider the multiple ecological functions of insects while the species are usu-
195 ally classified by their major ecological function for example pollinators, pests, or nat-
196 ural enemies. Nevertheless, many insects can provide multiples services improving
197 crop production due to their double ecological functions over their development stag-
198 es (e.g. larvae as herbivore vs. adult as pollinator) or over different crops. For in-
199 stance, *Forcipomyia* spp (Diptera) are pollinators in cocoa [53] and also natural ene-
200 mies of *Dione juno* (Lepidoptera: Nymphalidae) [61] that damages many tropical
201 crops (e.g. passion fruit, tomato, sugarcane, among others) [62,63]. In coffee, *Syr-*
202 *phus* sp. (Diptera: Syrphidae), *Allobaccha brevis* (Diptera: Syrphidae) and *Hemipyrel-*
203 *lia* sp. (Diptera: Calliphoridae) are recorded as pollinators but these species are also
204 reported as insect natural enemies in passion fruit and avocado [62,64].

205 On the other hand, other insects can have adverse effects (e.g. providing a
206 service and a disservice) such as Sciaridae that are pollinators in cocoa [65] but are
207 also considered as pests in watermelon [66]. *Trigona spinipes* Fabr. (Hymenoptera:
208 Apidae) is a pollinator in coffee that can be detrimental to coffee fruit set [67] and is
209 also considered as a pest due to its damages of open galleries on fruits reducing fruit

210 quality [68]. The *Azteca sericeasur* ant is known to be an effective predator of the
211 coffee berry borer [69], a major coffee pest, but its aggressiveness towards pollina-
212 tors visiting the coffee flowers has been studied little [56]. Overall, it is accepted that
213 improving insect diversity in croplands benefits crop yield [70]. However, despite di-
214 versity being important, studies that evaluate whether an insect is an efficient and or
215 effective pollinator, predator or has multiple ecological functions are valuable [54*].
216 For instance, *Trigona jaty* (Hymenoptera: Apidae, meliponinae) is a flower visitor in
217 cocoa but is considered as an inefficient pollinator given its pollen robbing action that
218 weakens flower fitness [71]. Pollination efficiency of many flower visitors in coffee
219 and cocoa crops are not yet tested and although studies suggest their ecological
220 function as pollinators, they can also act as pests or natural enemies in other crops.
221 Neglecting the multiple ecological functions of insects and their efficiency carrying out
222 these ecosystem services and disservices would considerably limit crop yield im-
223 provement.

224

225 **Perspectives**

226 The tropics are home to the highest diversity and abundance of arthropods [72], and
227 as such, represent an excellent area for considering IPPM approaches in agricultural
228 systems. Through the study of two tropical crops, we found that few experiments
229 have investigated the effect of agricultural practices simultaneously on pests, natural
230 enemies and pollinator communities despite IPPM recommendations. Although it is
231 rather recognized that analyzing the influence of a single ecosystem service on crop
232 yield can be complex due to context-dependency, studying combined effects of
233 pollinator and pest managements implies interacting relationships that are current
234 challenges in applied ecology [55]. Integrating pest and pollinator management
235 strategies in agriculture is therefore pioneering and represents future a direction
236 towards sustainable food production. Consideration of the landscape scale also
237 invokes the possibility of multiple ecological functions that insects may have [22] as
238 pests of one crop and pollinators of a neighboring crop. This consideration should be
239 even more important when species richness and abundance are high. In the future,
240 the growing awareness of the landscape scale effect and the multiple ecological
241 functions of arthropods in relation to crops should lead to further consideration of the
242 neighboring agricultural landscape, i.e. the nature of the established crops and their
243 pollinator, pests and predator assemblages.

244

245 **Acknowledgments**

246 This publication was made possible through support provided by the IRD. Special
247 thanks go to Clémence Riva and Adrian David González Chaves for constructive
248 comments on the manuscript. We also thank Elsa Blareau for critical reading of
249 English used in the text.

250

251 **Declarations of interest:** none.

252

253 **References and recommended reading**

254 Papers of particular interest, published within the period of review, have been
255 highlighted of special interest (*) or of outstanding interest (**).

256 1**. Gagic V, Marcora A, Howie L: **Additive and interactive effects of pollination
257 and biological pest control on crop yield.** *J Appl Ecol* 2019, **56**:2528–2535.

258 This study conducted a greenhouse experiment and a field study to disentangle the
259 potential and realized contribution of insect pollination to cotton boll retention and
260 yield under various pest pressures and biocontrol levels. They found additive and
261 interactive effects of insect pollination and biocontrol depending on the level of pest
262 pressure. A very interesting result was that at high pest abundances, boll retention
263 and final yield were higher when pollinators were present.

264 2. Pywell RF, Heard MS, Woodcock BA, Hinsley S, Ridding L, Nowakowski M,
265 Bullock JM: **Wildlife-friendly farming increases crop yield: evidence for
266 ecological intensification.** *Proc R Soc B* 2015, **282**:20151740.

267 3. Klein A-M, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen
268 C, Tscharntke T: **Importance of pollinators in changing landscapes for world
269 crops.** *Proc R Soc B* 2007, **274**:303–313.

270 4. Aizen MA, Garibaldi LA, Cunningham SA, Klein AM: **How much does
271 agriculture depend on pollinators? Lessons from long-term trends in crop
272 production.** *Annals of Botany* 2009, **103**:1579–1588.

273 5. Potts SG, Imperatriz-Fonseca V, Ngo HT, Aizen MA, Biesmeijer JC, Breeze TD,
274 Dicks LV, Garibaldi LA, Hill R, Settele J, et al.: **Safeguarding pollinators and
275 their values to human well-being.** *Nature* 2016, **540**:220–229.

276 6*. Rader R, Cunningham SA, Howlett BG, Inouye DW: **Non-Bee Insects as**
277 **Visitors and Pollinators of Crops: Biology, Ecology, and Management.** *Annu*
278 *Rev Entomol* 2020, **65**:391–407.

279 This review synthesizes the diversity and the importance of non-bee insects as crop
280 pollinators. They recorded the visitors and pollinators of 105 global food crops that
281 are known to benefit from animal pollinators. Non-bee insects are commonly
282 observed visiting crop flowers, including mostly Diptera, Coleoptera, Lepidoptera and
283 Hymenoptera. Interestingly, they also discussed the distinction between flower
284 visitation and pollination efficiency for some species.

285 7. Reilly JR, Artz DR, Biddinger D, Bobiwash K, Boyle NK, Brittain C, Brokaw J,
286 Campbell JW, Daniels J, Elle E, et al.: **Crop production in the USA is**
287 **frequently limited by a lack of pollinators.** *Proc R Soc B* 2020, **287**:20200922.

288 8. Oerke EC: **Crop losses to pests.** *J Agric Sci* 2006, **144**:31–43.

289 9*. Martin EA, Feit B, Requier F, Friberg H, Jonsson M: **Assessing the resilience of**
290 **biodiversity-driven functions in agroecosystems under environmental**
291 **change.** In *Advances in Ecological Research*. . Elsevier; 2019:59–123.

292 In this book chapter, authors reviewed and summarized dimensions of ecosystem
293 service resilience in agroecosystems, and the disturbances they are suffering under
294 environmental change. They concluded that studying the resilience of ecosystem
295 services in agroecosystems is still in its infancy, but novel tools and approaches can
296 catalyze further steps to assess and improve the resilience of biodiversity-driven
297 agroecosystem functions under environmental change.

298 10. Bommarco R, Kleijn D, Potts SG: **Ecological intensification: harnessing**
299 **ecosystem services for food security.** *Trends in Ecology & Evolution* 2013,
300 **28**:230–238.

301 11. Garibaldi LA, Requier F, Rollin O, Andersson GK: **Towards an integrated**
302 **species and habitat management of crop pollination.** *Current Opinion in*
303 *Insect Science* 2017, **21**:105–114.

304 12. Goulson D, Nicholls E, Botias C, Rotheray EL: **Bee declines driven by**
305 **combined stress from parasites, pesticides, and lack of flowers.** *Science*
306 2015, **347**:1255957.

307 13. Bigirimana J, Gerard A, Mota-Sanchez D, Gut LJ: **Options for Managing**
308 ***Antestiopsis thunbergii* (Hemiptera: Pentatomidae) and the Relationship of**

- 309 **Bug Density to the Occurrence of Potato Taste Defect in Coffee.** *Florida*
310 *Entomologist* 2018, **101**:580–586.
- 311 14. Vanhove W, Feng Y, Yu M, Hafiz IO, Vanhoudt N, Van Damme PLJ, Zhang A:
312 **Evaluation of attract-and-kill strategy for management of cocoa pod borer,**
313 **Conopomorpha cramerella, in Malaysia cocoa plantation.** *International*
314 *Journal of Pest Management* 2020, **66**:155–162.
- 315 15. Aristizábal N, Metzger JP: **Landscape structure regulates pest control**
316 **provided by ants in sun coffee farms.** *J Appl Ecol* 2019, **56**:21–30.
- 317 16. Jimenez-Soto E, Morris JR, Letourneau DK, Philpott SM: **Vegetation**
318 **connectivity increases ant activity and potential for ant-provided biocontrol**
319 **services in a tropical agroforest.** *Biotropica* 2019, **51**:50–61.
- 320 17. Medeiros HR, Grandinete YC, Manning P, Harper KA, Cutler GC, Tyedmers P,
321 Righi CA, Ribeiro MC: **Forest cover enhances natural enemy diversity and**
322 **biological control services in Brazilian sun coffee plantations.** *Agron Sustain*
323 *Dev* 2019, **39**:50.
- 324 18**. Egan PA, Dicks LV, Hokkanen HMT, Stenberg JA: **Delivering Integrated**
325 **Pest and Pollinator Management (IPPM).** *Trends in Plant Science* 2020,
326 **25**:577–589.
- 327 This review provides conceptual schemes and decision supporting tools related to
328 the new paradigm of Integrated Pest and Pollinator Management (IPPM). This review
329 standardizes the implementation of actions that promote complementarity between
330 integrated pest management and pollination management.
- 331 19**. Lundin O, Rundlöf M, Jonsson M, Bommarco R, Williams NM: **Integrated pest**
332 **and pollinator management – expanding the concept.** *Front Ecol Environ*
333 2021, **19**:283–291.
- 334 This review expands the concept of Integrated Pest and Pollinator Management
335 (IPPM) by suggesting categories of management strategies and their importance in
336 moving toward this concept. Authors described the actions grouped in these
337 categories and synthesized their effects on pollinator, pest and predator
338 communities.
- 339 20. Hipólito J, Boscolo D, Viana BF: **Landscape and crop management**
340 **strategies to conserve pollination services and increase yields in tropical**
341 **coffee farms.** *Agriculture, Ecosystems & Environment* 2018, **256**:218–225.

342 21**. Toledo-Hernández M, Tscharrntke T, Tjoa A, Anshary A, Cyio B, Wanger TC:
343 **Landscape and farm-level management for conservation of potential**
344 **pollinators in Indonesian cocoa agroforests.** *Biological Conservation* 2021,
345 **257**:109106.

346 The objective of this study was to determine the effect of plot and landscape scale
347 factors on the abundance of arthropods visiting cocoa flowers and thus potential
348 pollinators of this highly insect pollination dependent crop. The results indicate that
349 the presence of semi-natural habitats increases the abundance of ants and Diptera
350 visiting the flowers while the distance to the forest does not affect them. Other factors
351 such as abundance of cocoa flowers, leaf litter or shade cover may also have a
352 beneficial effect on the abundance of some of these potential pollinators.

353 22. Saunders ME, Peisley RK, Rader R, Luck GW: **Pollinators, pests, and**
354 **predators: Recognizing ecological trade-offs in agroecosystems.** *Ambio*
355 2016, **45**:4–14.

356 23. Biddinger DJ, Rajotte EG: **Integrated pest and pollinator management —**
357 **adding a new dimension to an accepted paradigm.** *Current Opinion in Insect*
358 *Science* 2015, **10**:204–209.

359 24. Henry M, Beguin M, Requier F, Rollin O, Odoux J-F, Aupinel P, Aptel J,
360 Tchamitchian S, Decourtye A: **A Common Pesticide Decreases Foraging**
361 **Success and Survival in Honey Bees.** *Science* 2012, **336**:348–350.

362 25**. Martin EA, Dainese M, Clough Y, Báldi A, Bommarco R, Gagic V, Garratt
363 MPD, Holzschuh A, Kleijn D, Kovács - Hostyánszki A, et al.: **The interplay of**
364 **landscape composition and configuration: new pathways to manage**
365 **functional biodiversity and agroecosystem services across Europe.** *Ecol*
366 *Lett* 2019, **22**:1083–1094.

367 This study assessed how spatial arrangement of crop fields and other habitats in
368 landscapes impact arthropods and their functions for pollination and pest control.
369 They found that enhancing edge density in agroecosystems can increase the
370 abundance of pollinator and natural enemy species, leading to an improvement of
371 pollination and pest control services, respectively. These results suggest that
372 enhancing edge density in agroecosystems can promote yield-enhancing ecosystem
373 services.

374 26**. Dainese M, Martin EA, Aizen MA, Albrecht M, Bartomeus I, Bommarco R,
375 Carvalho LG, Chaplin-Kramer R, Gagic V, Garibaldi LA, et al.: **A global**
376 **synthesis reveals biodiversity-mediated benefits for crop production.**
377 *Science Advances* 2019, 5 :eaax0121.

378 This study analyzed whether pest control and crop pollination services can be
379 maintained by a few dominant species or rely on high richness. They found that
380 maintaining high species richness of pollinators and natural enemies was more
381 important to support ecosystem services than providing an abundance of dominant
382 species. Biodiversity conservation in croplands is therefore critical for providing
383 ecosystem services and sustainable food production.

384 27. Adomako B: **Causes and extent of yield losses in cocoa progenies.** *Trop*
385 *Sci* 2007, **47**:22–25.

386 28. Cerda R, Avelino J, Gary C, Tixier P, Lechevallier E, Allinne C: **Primary and**
387 **Secondary Yield Losses Caused by Pests and Diseases: Assessment and**
388 **Modeling in Coffee.** *PLoS ONE* 2017, **12**:e0169133.

389 29*. Iverson AL, Gonthier DJ, Pak D, Ennis KK, Burnham RJ, Perfecto I, Ramos
390 Rodriguez M, Vandermeer JH: **A multifunctional approach for achieving**
391 **simultaneous biodiversity conservation and farmer livelihood in coffee**
392 **agroecosystems.** *Biological Conservation* 2019, **238**:108179.

393 This study presents a multifunctional analysis to investigate the economic and
394 biodiversity conservation impacts of agricultural practices commonly implemented in
395 complex coffee-based agroecosystems. The impact study focuses on practices such
396 as the promotion of ground vegetation heterogeneity in the plot; the conservation of
397 nearby forest areas; the provision of shade by agroforestry systems; and predation
398 by vertebrates. In a second step, economic incentives for farmers are explored.

399 30. Classen A, Peters MK, Fergner SW, Helbig-Bonitz M, Schmack JM, Maassen
400 G, Schleuning M, Kalko EKV, Böhning-Gaese K, Steffan-Dewenter I:
401 **Complementary ecosystem services provided by pest predators and**
402 **pollinators increase quantity and quality of coffee yields.** *Proc R Soc B* 2014,
403 **281**:20133148.

404 31. Flores S, Campos S, Villaseñor A, Valle Á, Enkerlin W, Toledo J, Liedo P,
405 Montoya P: **Sterile males of *Ceratitis capitata* (Diptera: Tephritidae) as**
406 **disseminators of *Beauveria bassiana* conidia for IPM strategies.** *Biocontrol*
407 *Science and Technology* 2013, **23**:1186–1198.

- 408 32. Akesse-Ransford G, Owusu EO, Kyerematen R, Adu-Acheampong S:
409 **Arthropod diversity of cocoa farms under two management systems in the**
410 **Eastern and Central regions of Ghana.** *Agroforest Syst* 2021, **95**:791–803.
- 411 33*. Dewi VS, A Nurariaty A, Sulastria, Tuwo M: **Arthropoda diversity in organic**
412 **cocoa farming in Bantaeng District.** In *IOP Conference Series: Earth and*
413 *Environmental Science.* . 2020:012164.
- 414 This study compares arthropod abundance and diversity between an organically
415 managed and a conventionally managed plantation in Indonesia. Organic
416 management increased arthropod abundance by 50%. By classifying these
417 arthropods according to their function as pest or predator, it is shown that predatory
418 arthropods, and thus potential pest regulation services, are favored.
- 419 34. Khoo KC, Ho CT: **The influence of *Dolichoderus thoracicus***
420 **(Hymenoptera: Formicidae) on losses due to *Helopeltis theivora***
421 **(Heteroptera: Miridae), black pod disease, and mammalian pests in cocoa in**
422 **Malaysia.** *Bull Entomol Res* 1992, **82**:485–491.
- 423 35. Bisseleua DHB, Fotio D, Yede, Missoup AD, Vidal S: **Shade Tree Diversity,**
424 **Cocoa Pest Damage, Yield Compensating Inputs and Farmers' Net Returns**
425 **in West Africa.** *PLoS ONE* 2013, **8**:e56115.
- 426 36. Maas B, Clough Y, Tscharntke T: **Bats and birds increase crop yield in**
427 **tropical agroforestry landscapes.** *Ecol Lett* 2013, **16**:1480–1487.
- 428 37. Gras P, Tscharntke T, Maas B, Tjoa A, Hafsah A, Clough Y: **How ants, birds**
429 **and bats affect crop yield along shade gradients in tropical cacao**
430 **agroforestry.** *J Appl Ecol* 2016, **53**:953–963.
- 431 38. Novais SMA, Macedo-Reis LE, Neves FS: **Predatory beetles in cacao**
432 **agroforestry systems in Brazilian Atlantic forest: a test of the natural enemy**
433 **hypothesis.** *Agroforest Syst* 2017, **91**:201–209.
- 434 39. Novais SMA, Macedo-Reis LE, DaRocha WD, Neves FS: **Effects of habitat**
435 **management on different feeding guilds of herbivorous insects in cacao**
436 **agroforestry systems.** *Rev Biol Trop* 2016, **64**:763–777.
- 437 40. Vandermeer J, Perfecto I, Nuñez GI, Phillpott S, Ballinas AG: **Ants (*Azteca***
438 **sp.) as potential biological control agents in shade coffee production in**
439 **Chiapas, Mexico.** *Agroforestry Systems* 2002, **56**:271–276.

- 440 41. Borkhataria RR, Collazo JA, Groom MJ: **Additive Effects Of Vertebrate**
441 **Predators On Insects In A Puerto Rican Coffee Plantation.** *Ecological*
442 *Applications* 2006, **16**:696–703.
- 443 42. Borkhataria RR, Collazo JA, Groom MJ: **Species abundance and potential**
444 **biological control services in shade vs. sun coffee in Puerto Rico.**
445 *Agriculture, Ecosystems & Environment* 2012, **151**:1–5.
- 446 43. Peters VE, Greenberg R: **Fruit Supplementation Affects Birds but not**
447 **Arthropod Predation by Birds in Costa Rican Agroforestry Systems.**
448 *Biotropica* 2013, **45**:102–110.
- 449 44. Karungi J, Nambi N, Ijala AR, Jonsson M, Kyamanywa S, Ekbohm B: **Relating**
450 **shading levels and distance from natural vegetation with hemipteran pests**
451 **and predators occurrence on coffee.** *J Appl Entomol* 2015, **139**:669–678.
- 452 45*. Bagny Beilhe L, Roudine S, Quintero Perez JA, Allinne C, Daout D, Mauxion
453 R, Carval D: **Pest-regulating networks of the coffee berry borer**
454 **(*Hypothenemus hampei*) in agroforestry systems.** *Crop Protection* 2020,
455 **131**:105036.
- 456 In this study, field data were analyzed using structural equation modelling to better
457 understanding the effect of plant diversity associated with agroforestry systems on
458 the regulation of the coffee berry borer, a major coffee pest. The effects of plant
459 diversity on this regulation are numerous and sometimes indirect in the ecosystem
460 formed by the pest, its predators and their environment. Some effects may even
461 depend on the evaluation mode of the regulation service, since shade cover
462 increases the infestation of this pest but decreases the intensity of the damage
463 caused to the plant.
- 464 46. Amaral DS, Venzon M, Pallini A, Lima PC, DeSouza O: **A diversificação da**
465 **vegetação reduz o ataque do bicho-mineiro-do-cafeeiro *Leucoptera***
466 ***coffeella* (Guérin-mèneville) (Lepidoptera: Lyonetiidae)?** *Neotrop entomol*
467 2010, **39**:543–548.
- 468 47. Mendonça JMA de, Carvalho GA, Guimarães RJ, Reis PR, Rocha LCD:
469 **Produtos naturais e sintéticos no controle de *Leucoptera coffeella* (Guérin-**
470 **Mèneville & Perrottet, 1842) (Lepdoptera: Lyonetiidae) e seus efeitos sobre**
471 **a predação por vespas.** *Ciênc agrotec* 2006, **30**:892–899.

472 48*. Vandromme M, Trekels H, Sepúlveda Ruiz N, Somarriba E, Vanschoenwinkel
473 B: **Exploring the suitability of bromeliads as aquatic breeding habitats for**
474 **cacao pollinators**. *Hydrobiologia* 2019, **828**:327–337.

475 This study focuses on the effect of the morphology and the height of implantation of
476 bromeliads in cocoa plantations on the abundance of larvae belonging to the
477 potential pollinators and pest predators of this crop. Thus, the water contained in
478 these bromeliads shelters these pollinator larvae in 66% of the cases and their
479 abundance is more important when these plants are placed higher and when their
480 water reservoir is bigger.

481 49. Philpott SM, Uno S, Maldonado J: **The Importance of Ants and High-Shade**
482 **Management to Coffee Pollination and Fruit Weight in Chiapas, Mexico**.
483 *Biodivers Conserv* 2006, **15**:487–501.

484 50. Veddeler D, Tylianakis J, Tscharrntke T, Klein A-M: **Natural enemy diversity**
485 **reduces temporal variability in wasp but not bee parasitism**. *Oecologia* 2010,
486 **162**:755–762.

487 51*. Medeiros HR, Martello F, Almeida EAB, Mengual X, Harper KA, Grandinete
488 YC, Metzger JP, Righi CA, Ribeiro MC: **Landscape structure shapes the**
489 **diversity of beneficial insects in coffee producing landscapes**. *Biological*
490 *Conservation* 2019, **238**:108193.

491 This study analyzed whether compositional and configurational landscape
492 heterogeneity influence the diversity of beneficial insects in Brazilian coffee farms.
493 Through field collections on 16 landscapes in south-eastern Brazil, they surveyed
494 insects that revealed some interesting patterns. Forest loss in the surrounding
495 landscape resulted in wasp and bee species loss. Complementary high variations in
496 landscape diversity and edge density between landscapes resulted in species
497 replacement suggesting that pairs of landscapes with high and low diversity of habitat
498 types and edge density harbor different species.

499 52. Jha S, Bacon CM, Philpott SM, Ernesto Méndez V, Läderach P, Rice RA:
500 **Shade Coffee: Update on a Disappearing Refuge for Biodiversity**.
501 *BioScience* 2014, **64**:416–428.

502 53. Zakariyya F, Sulistyowati E, Rahayu DS: **Abundance of Pollinator Insect**
503 **(*Forcipomyia* spp.) of Cocoa Under Some Shade Trees**. *Pelita Perkebunan*
504 2016, **32**:91–100.

- 505 54*. Hipólito J, Nunes DO, Angel-Coca C, Mahlmann T, Galetto L, Viana BF:
506 **Performance, Effectiveness, and Efficiency of Honeybees as Pollinators of**
507 **Coffea arabica (Gentianales, Rubiaceae).** *Neotrop Entomol* 2020, **49**:501–510.
508 This study evaluated the influence of nectar traits on visitor’s performance throughout
509 flower lifetime and honeybees are effective and efficient for coffee pollination in Bahia
510 Brazil. They found that honeybees spend enough time to pollinate coffee flowers
511 even after one visit. However, coffee can be limited by the quality of pollen grains
512 moved by pollinators instead of quantity and for that reason pollinator diversity and
513 pollinator-friendly management is essential to guarantee crop production.
- 514 55. Garibaldi LA, Andersson GKS, Requier F, Fijen TPM, Hipólito J, Kleijn D,
515 Pérez-Méndez N, Rollin O: **Complementarity and synergisms among**
516 **ecosystem services supporting crop yield.** *Global Food Security* 2018, **17**:38–
517 47.
- 518 56. Vannette RL, Bichier P, Philpott SM: **The presence of aggressive ants is**
519 **associated with fewer insect visits to and altered microbe communities in**
520 **coffee flowers.** *Basic and Applied Ecology* 2017, **20**:62–74.
- 521 57. Tarno H, Wicaksono KP, Begliomini E: **Floral Stimulation and Behavior of**
522 **Insect Pollinators Affected by Pyraclostrobin on Arabica Coffee.** *Agrivita J*
523 *Agr Sci* 2018, **40**:161–167.
- 524 58. Albrecht M, Kleijn D, Williams NM, Tschumi M, Blaauw BR, Bommarco R,
525 Campbell AJ, Dainese M, Drummond FA, Entling MH, et al.: **The effectiveness**
526 **of flower strips and hedgerows on pest control, pollination services and**
527 **crop yield: a quantitative synthesis.** *Ecol Lett* 2020, **23**:1488–1498.
- 528 59. Schroth G, Harvey CA, Vincent G: **Complex Agroforests: Their Structure,**
529 **Diversity, and Potential Role in Landscape Conservation.** In *Agroforestry and*
530 *biodiversity conservation in tropical landscapes.* . Island Press; 2004:227–260.
- 531 60. Avelino J, Romero-Gurdián A, Cruz-Cuellar HF, Declerck FAJ: **Landscape**
532 **context and scale differentially impact coffee leaf rust, coffee berry borer,**
533 **and coffee root-knot nematodes.** *Ecological Applications* 2012, **22**:584–596.
- 534 61. Lima M de FC de, Veiga AF de SL: **Ocorrência de inimigos naturais de**
535 **Dione juno juno (CR.), Agraulis vanillae maculosa S. e Eueides isabella**
536 **dianasa (Hüb.)(Lepidoptera: Nymphalidae) em Pernambuco.** *An Soc Entomol*
537 *Bras* 1995, **24**:631–634.

- 538 62. Ribeiro BM, dos Santos ER, Trentin LB, da Silva LA, de Melo FL, Kitajima EW,
539 Ardisson-Araújo DMP: **A Nymphalid-Infecting Group I Alphabaculovirus**
540 **Isolated from the Major Passion Fruit Caterpillar Pest *Dione juno juno***
541 **(Lepidoptera: Nymphalidae). *Viruses* 2019, 11:602.**
- 542 63. Souza AR de, Giustolin TA, Querino RB, Alvarenga CD: **Natural Parasitism**
543 **of Lepidopteran Eggs by *Trichogramma* Species (Hymenoptera:**
544 ***Trichogrammatidae*) in Agricultural Crops in Minas Gerais, Brazil. *Florida***
545 ***Entomologist* 2016, 99:221–225.**
- 546 64. Wysoki M, Berg MA van den, Ish-Am G, Gazit S, Peña JE, Waite GK: **Pests**
547 **and pollinators of avocado.** In *Tropical fruit pests and pollinators: biology,*
548 *economic importance, natural enemies and control.* Edited by Peña JE, Sharp JL,
549 Wysoki M. CABI; 2002:223–293.
- 550 65. Zegada Herbas L, Lafuente Cartagena I, Naoki K, Armengot L: **Variation in**
551 **composition of cacao floral visitors (*Theobroma cacao*) among five**
552 **production systems in Sara Ana, Alto Beni, Bolivia. *Ecologia en Bolivia* 2020,**
553 **55:145–159.**
- 554 66. Kim HH, Choo HY, Kaya HK, Lee DW, Lee SM, Jeon HY: ***Steinernema***
555 ***carpocapsae* (Rhabditida: Steinernematidae) as a biological control agent**
556 **against the fungus Gnat *Bradysia agrestis* (Diptera: Sciaridae) in**
557 **propagation houses. *Biocontrol Science and Technology* 2004, 14:171–183.**
- 558 67. González-Chaves A, Jaffé R, Metzger JP, de M. P. Kleinert A: **Forest**
559 **proximity rather than local forest cover affects bee diversity and coffee**
560 **pollination services. *Landscape Ecol* 2020, 35:1841–1855.**
- 561 68. Alves D de A, Lima JE, Soares MA, Rabelo JM, Da Cruz M do CM: **Ataque de**
562 ***Trigona spinipes* (Fabr.) (Hymenoptera: Apidae: Meliponinae) em pitaia**
563 ***Hylocereus undatus* (Haw.) e *Hylocereus polyrhizus* (Weber) (Cactaceae)**
564 **em Couto de Magalhães de Minas, Minas Gerais, Brasil. *EntomoBrasilis* 2018,**
565 **11:223–225.**
- 566 69. Morris JR, Vandermeer J, Perfecto I: **A Keystone Ant Species Provides**
567 **Robust Biological Control of the Coffee Berry Borer Under Varying Pest**
568 **Densities. *PLoS ONE* 2015, 10:e0142850.**
- 569 70. Rader R, Bartomeus I, Garibaldi LA, Garratt MPD, Howlett BG, Winfree R,
570 Cunningham SA, Mayfield MM, Arthur AD, Andersson GKS, et al.: **Non-bee**

- 571 **insects are important contributors to global crop pollination.** *Proc Natl Acad*
572 *Sci USA* 2016, **113**:146–151.
- 573 71. Young AM: **The Ineffectiveness of the Stingless Bee *Trigona jaty***
574 **(Hymenoptera: Apidae: Meliponinae) as a Pollinator of Cocoa (*Theobroma***
575 ***cacao* L.).** *The Journal of Applied Ecology* 1981, **18**:149.
- 576 72. Stork NE: **How Many Species of Insects and Other Terrestrial Arthropods**
577 **Are There on Earth?** *Annu Rev Entomol* 2018, **63**:31–45.
- 578 73. <http://www.fao.org/faostat/en/#data/QCL/visualize>. 2019.
- 579 74. Web of Knowledge: *Online bibliographical database from the Institute for*
580 *Scientific Information*. 2021.
- 581 75. Spafford H, Ching A, Manley M, Hardin C, Bittenbender H: **Management of**
582 **Chinese Rose Beetle (*Adoretus sinicus*) Adults Feeding on Cacao**
583 **(*Theobroma cacao*) Using Insecticides.** *Insects* 2016, **7**:28.
- 584 76. Sarfo JE, Campbell CAM, Hall DR: **Design and placement of synthetic sex**
585 **pheromone traps for cacao mirids in Ghana.** *Int J Trop Insect Sci* 2018,
586 **38**:122–131.
- 587 77. Sarfo JE, Campbell CAM, Hall DR: **Effect of lure age and blend on sex**
588 **pheromone trap catches of the mirid *Sahlbergella singularis* on cacao in**
589 **Ghana.** *Crop Protection* 2020, **138**:105344.
- 590 78. Vanhove W, Vanhoudt N, Bhanu KRM, Abubeker S, Feng Y, Yu M, Van
591 Damme P, Zhang A: **Geometric isomers of sex pheromone components do**
592 **not affect attractancy of *Conopomorpha cramerella* in cocoa plantations.** *J*
593 *Appl Entomol* 2015, **139**:660–668.
- 594 79. Sarfo JE, Campbell CAM, Hall DR: **Optimal pheromone trap density for**
595 **mass trapping cacao mirids.** *Entomol Exp Appl* 2018, **166**:565–573.
- 596 80. Awudzi GK, Cudjoe AR, Hadley P, Hatcher PE, Daymond AJ: **Optimizing**
597 **mirid control on cocoa farms through complementary monitoring systems.**
598 *J Appl Entomol* 2017, **141**:247–255.
- 599 81. Mahot HC, Mahob JR, Hall DR, Arnold SEJ, Fotso AK, Membang G, Ewane N,
600 Kemga A, Fiaboe KKM, Bilong CFB, et al.: **Visual cues from different trap**
601 **colours affect catches of *Sahlbergella singularis* (Hemiptera: Miridae) in sex**
602 **pheromone traps in Cameroon cocoa plantations.** *Crop Protection* 2020,
603 **127**:104959.

- 604 82. Bisseleua DHB, Begoude D, Tonnang H, Vidal S: **Ant-mediated ecosystem**
605 **services and disservices on marketable yield in cocoa agroforestry**
606 **systems**. *Agriculture, Ecosystems & Environment* 2017, **247**:409–417.
- 607 83. Forbes SJ, Northfield TD: **Oecophylla smaragdina ants provide pest**
608 **control in Australian cacao**. *Biotropica* 2017, **49**:328–336.
- 609 84. Rizali A, Tscharrntke T, Buchori D, Clough Y: **Separating effects of species**
610 **identity and species richness on predation, pathogen dissemination and**
611 **resistance to invasive species in tropical ant communities: Separating**
612 **effects of identity and richness of ants**. *Agr Forest Entomol* 2018, **20**:122–130.
- 613 85. Saleh A, Armaniar, Ahmad AH: **Strategies for Controlling Cocoa Pod**
614 **Borer, Conopomorpha cramerella Snellen, on Cocoa Farmers in Langkat**
615 **District, North Sumatra, Indonesia**. In *Proceedings of the International*
616 *Conference and the 10th Congress of the Entomological Society of Indonesia*
617 *(ICCESI 2019)*. . Atlantis Press; 2020.
- 618 86. Bagny Beilhe L, Piou C, Tadu Z, Babin R: **Identifying Ant-Mirid Spatial**
619 **Interactions to Improve Biological Control in Cacao-Based Agroforestry**
620 **System**. *Environmental Entomology* 2018, **47**:551–558.
- 621 87. Mboussi SB, Ambang Z, Kakam S, Bagny Beilhe L: **Control of cocoa mirids**
622 **using aqueous extracts of Thevetia peruviana and Azadirachta indica**.
623 *Cogent Food & Agriculture* 2018, **4**:1430470.
- 624 88. Cassano CR, Silva RM, Mariano-Neto E, Schroth G, Faria D: **Bat and bird**
625 **exclusion but not shade cover influence arthropod abundance and cocoa**
626 **leaf consumption in agroforestry landscape in northeast Brazil**. *Agriculture,*
627 *Ecosystems & Environment* 2016, **232**:247–253.
- 628 89. Armengot L, Ferrari L, Milz J, Velásquez F, Hohmann P, Schneider M: **Cacao**
629 **agroforestry systems do not increase pest and disease incidence compared**
630 **with monocultures under good cultural management practices**. *Crop*
631 *Protection* 2020, **130**:105047.
- 632 90. Felicitas AC, Hervé BDB, Ekesi S, Akutse KS, Djuideu CTCL, Meupia MJ,
633 Babalola OO: **Consequences of shade management on the taxonomic**
634 **patterns and functional diversity of termites (Blattodea: Termitidae) in**
635 **cocoa agroforestry systems**. *Ecol Evol* 2018, **8**:11582–11595.

- 636 91. Riedel J, Kägi N, Armengot L, Schneider M: **Effects of rehabilitation pruning**
637 **and agroforestry on cacao tree development and yield in an older full-sun**
638 **plantation.** *Ex Agric* 2019, **55**:849–865.
- 639 92. Mulia S, McMahon PJ, Purwantara A, Bin Purung H, Djufry F, Lambert S,
640 Keane PJ, Guest DI: **Effect of organic and inorganic amendments on**
641 **productivity of cocoa on a marginal soil in Sulawesi, Indonesia.** *Ex Agric*
642 2019, **55**:1–20.
- 643 93. Muslimin, Wijayanti W, Anshary A, Basri Z, Cruz AF, Suwastika IN, Shiina T:
644 **Sulawesi cacao (*Theobroma cacao*, L.) performances under two different**
645 **agricultural system in east coast of Central Sulawesi.** *IOP Conf Ser: Earth*
646 *Environ Sci* 2018, **144**:012066.
- 647 94. Syarief M, Mudjiono G, Abadi AL, Himawan T: **Arthropods Diversity and**
648 **Population Dynamic of *Helopeltis antonii* Sign. (Hemiptera: Miridae) on**
649 **Various Cocoa Agroecosystems Management.** *Agrivita J Agr Sci* 2018,
650 **40**:350–359.
- 651 95. Salazar-Díaz R, Torres-Coto V: **Study of the dynamics of cocoa**
652 **(*Theobroma cacao*) pollinators in three production systems.** *TM* 2017,
653 **30**:90–100.
- 654 96*. Djuideu CTL, Bisseleua HDB, Kekeunou S, Ambele FC: **Rehabilitation**
655 **practices in cocoa agroforestry systems mitigate outbreaks of termites and**
656 **support cocoa tree development and yield.** *Agriculture, Ecosystems &*
657 *Environment* 2021, **311**:107324.
- 658 This study investigated how rehabilitation pruning of cocoa trees influences termite
659 outbreaks and yield in three cocoa agroforestry systems (rustic, intermediate and full
660 sun systems). They monitored cocoa and non-cocoa trees at different development
661 stages after rehabilitation pruning and assessed termite diversity and damage and
662 found that farm rehabilitation practices are associated with high beta diversity of
663 shade trees. Farm rehabilitation practices also increased biodiversity of beneficial
664 termite species which in turn improved cocoa tree vigor and yield in cocoa
665 agroforestry systems.
- 666 97. Djuideu TCL, Bisseleua DHB, Kekeunou S, Meupia MJ, Difouo FG, Ambele
667 **CF: Plant community composition and functional characteristics define**
668 **invasion and infestation of termites in cocoa agroforestry systems.**
669 *Agroforest Syst* 2020, **94**:185–201.

- 670 98. Saripah B, Nor Aizam A, Ummu Syaira Ain R: **Temporal geospatial**
671 **assessment of cocoa pollinator**. *Serranga* 2019, **24**:159–172.
- 672 99. Saripah B, Alias A: **The Use of Transferable Breeding Container for**
673 **Population Enrichment of Pollinator, *Forcipomyia* spp. in the Cocoa**
674 **Ecosystem**. *JOAAT* 2018, **5**:245–251.
- 675 100. Góngora CE, Laiton-J. LA, Gil ZN, Benavides P: **Evaluación de *Beauveria***
676 ***bassiana* para el control de *Monalonion velezangeli* (Hemiptera: Miridae) en**
677 **el cultivo del café**. *Rev Colomb Entomol* 2020, **46**:e7685.
- 678 101. De la Mora A, García-Ballinas JA, Philpott SM: **Local, landscape, and**
679 **diversity drivers of predation services provided by ants in a coffee**
680 **landscape in Chiapas, Mexico**. *Agriculture, Ecosystems & Environment* 2015,
681 **201**:83–91.
- 682 102. Gitirana Neto J, Cunha JPARD: **Spray deposition and chemical control of**
683 **the coffee leaf-miner with different spray nozzles and auxiliary boom**. *Eng*
684 *Agríc* 2016, **36**:656–663.
- 685 103. Egonyu JP, Kucel P, Kagezi G, Kovach J, Rwomushana I, Erbaugh M,
686 Wekono R, Salifu D, Kyamanywa S: ***Coffea arabica* Variety KP423 May be**
687 **Resistant to the Cerambycid Coffee Stemborer *Monochamus leuconotus*,**
688 **but Common Stem Treatments Seem Ineffective Against the Pest**. *African*
689 *Entomology* 2015, **23**:68–74.
- 690 104. Vega FE, Simpkins A, Miranda J, Harnly JM, Infante F, Castillo A, Wakarchuk
691 D, Cossé A: **A Potential Repellent Against the Coffee Berry Borer**
692 **(Coleoptera: Curculionidae: Scolytinae)**. *Journal of Insect Science* 2017, **17**.
- 693 105*. Machado ACP, Barônio GJ, Oliveira FF, Garcia CT, Rech AR: **Does a coffee**
694 **plantation host potential pollinators when it is not flowering? Bee**
695 **distribution in an agricultural landscape with high biological diversity in the**
696 **Brazilian *Campo Rupestre***. *J Sci Food Agric* 2021, **101**:2345–2354.

697 This work evaluated the distribution of bees in an agricultural landscape of coffee
698 plantation before and after coffee flowering. They collected bee species in dry and
699 rainy seasons in Minas Gerais Brazil. Few bees were recurrent in both seasons and
700 vegetation types (coffee and non-coffee). There was no temporal difference in
701 richness or abundance; however, both varied in relation to the vegetation type and
702 were higher in the coffee-native transition area. Diverging from richness or

703 abundance, the composition of the communities differed regarding season and
704 vegetation types.

- 705 106. Oliva M, Rubio K, Rivasplata E, Leiva S: **Integrated management of the**
706 **coffee berry borer: A comparison of cultural, biological, and ethological**
707 **control**. *Entomological Research* 2020, **50**:539–544.
- 708 107. Rostaman, Prakoso B: **Response of Coffee Berry Borer (*Hypothenemus***
709 ***hampaei*) to Alcohol-Based Attractants on Coffee Crops in Banjarnegara,**
710 **Indonesia**. In *Proceedings of the International Conference and the 10th*
711 *Congress of the Entomological Society of Indonesia (ICCESI 2019)*. . Atlantis
712 Press; 2020.
- 713 108. Egonyu JP, Torto B: **Responses of the ambrosia beetle *Xylosandrus***
714 ***compactus* (Coleoptera: Curculionidea: Scolytinae) to volatile constituents**
715 **of its symbiotic fungus *Fusarium solani* (Hypocreales: Nectriaceae).**
716 *Arthropod-Plant Interactions* 2018, **12**:9–20.
- 717 109. Vargas RI, Souder SK, Rendon P, Mackey B: **Suppression of Mediterranean**
718 **Fruit Fly (Diptera: Tephritidae) With Trimedlure and Biolure Dispensers in**
719 ***Coffea arabica* (Gentianales: Rubiaceae) in Hawaii**. *Journal of Economic*
720 *Entomology* 2018, **111**:293–297.
- 721 110. Leiva-Espinoza S, Oliva-Cruz M, Rubio-Rojas K, Maicelo-Quintana J, Milla-
722 Pino M: **Use of color traps and alcoholic attractants for the capture of the**
723 **coffee berry borer (*Hypothenemus hampei*) in highly infested coffee**
724 **plantations**. *Rev Colomb Entomol* 2020, **45**:e8537.
- 725 111. Perfecto I, Hajian-Forooshani Z, White A, Vandermeer J: **Ecological**
726 **complexity and contingency: Ants and lizards affect biological control of**
727 **the coffee leaf miner in Puerto Rico**. *Agriculture, Ecosystems & Environment*
728 2021, **305**:107104.
- 729 112. Rivera-Salinas IS, Hajian-Forooshani Z, Jiménez-Soto E, Cruz-Rodríguez JA,
730 Philpott SM: **High intermediary mutualist density provides consistent**
731 **biological control in a tripartite mutualism**. *Biological Control* 2018, **118**:26–
732 31.
- 733 113. Ogogol R, Egonyu JP, Bwogi G, Kyamanywa S, Erbaugh M: **Interaction of**
734 **the predatory ant *Pheidole megacephala* (Hymenoptera: Formicidae) with**
735 **the polyphagous pest *Xylosandrus compactus* (Coleoptera: Curculionidea).**
736 *Biological Control* 2017, **104**:66–70.

- 737 114. Rodrigues-Silva N, de Oliveira Campos S, de Sá Farias E, de Souza TC,
738 Martins JC, Picanço MC: **Relative importance of natural enemies and abiotic**
739 **factors as sources of regulation of mealybugs (Hemiptera:**
740 **Pseudococcidae) in Brazilian coffee plantations: Identification of potential**
741 **biological control agent of the mealybug.** *Ann Appl Biol* 2017, **171**:303–315.
- 742 115. Cancino J, Ruiz L, López E, Aguilar E, Galvez C, Montoya P, Liedo P:
743 **Suppression of *Ceratitis capitata* (Wied.) (Diptera: Tephritidae) populations**
744 **in coffee in the Mexico–Guatemala border region through the augmentative**
745 **releases of *Diachasmimorpha longicaudata* (Ashmead) (Hymenoptera:**
746 **Braconidae).** *Biocontrol Science and Technology* 2019, **29**:822–826.
- 747 116. Bizumungu G, Majer JD: **The Distribution of Ants in a Rwandan Coffee**
748 **Plantation and Their Potential to Control Pests.** *African Entomology* 2019,
749 **27**:159–166.
- 750 117. Bizumungu G, Majer JD, Fay HAC: **The potential protective value of ants in**
751 **Northern Australian coffee plantations.** *International Journal of Pest*
752 *Management* 2020, **66**:227–238.
- 753 118. Greco EB, Wright MG, Burgueño J, Jaronski ST: **Efficacy of *Beauveria***
754 ***bassiana* applications on coffee berry borer across an elevation gradient in**
755 **Hawaii.** *Biocontrol Science and Technology* 2018, **28**:995–1013.
- 756 119. Hollingsworth RG, Aristizábal LF, Shriner S, Mascarín GM, Moral R de A,
757 Arthurs SP: **Incorporating *Beauveria bassiana* Into an Integrated Pest**
758 **Management Plan for Coffee Berry Borer in Hawaii.** *Front Sustain Food Syst*
759 2020, **4**.
- 760 120. Bayman P, Mariño YA, García-Rodríguez NM, Oduardo-Sierra OF, Rehner
761 SA: **Local isolates of *Beauveria bassiana* for control of the coffee berry**
762 **borer *Hypothenemus hampei* in Puerto Rico: Virulence, efficacy and**
763 **persistence.** *Biological Control* 2021, **155**:104533.
- 764 121. Toledo J, Flores S, Campos S, Villaseñor A, Enkerlin W, Liedo P, Valle Á,
765 Montoya P: **Pathogenicity of three formulations of *Beauveria bassiana* and**
766 **efficacy of autoinoculation devices and sterile fruit fly males for**
767 **dissemination of conidia for the control of *Ceratitis capitata*.** *Entomol Exp*
768 *Appl* 2017, **164**:340–349.
- 769 122. Góngora CE, Tapias J, Jaramillo J, Medina R, Gonzalez S, Casanova H, Ortiz
770 A, Benavides P: **Evaluation of Terpene-Volatile Compounds Repellent to the**

- 771 **Coffee Berry Borer, *Hypothenemus hampei* (Ferrari) (Coleoptera:**
772 **Curculionidae).** *J Chem Ecol* 2020, **46**:881–890.
- 773 123. Zayas DV, Piñeda RR, Gómez MC, Ortiz IBM, Chaviano CD: **Efecto de**
774 **Heterorhabditis bacteriophora sobre la broca del café en la zona del**
775 **Algarrobo, Trinidad, Cuba.** *Centro Agriola* 2015, **43**:15–20.
- 776 124. Vega VJ, Mariño YA, Deynes D, Greco EB, Bright DE, Bayman P: **A Beetle in**
777 **a Haystack: Are There Alternate Hosts of the Coffee Berry Borer**
778 **(*Hypothenemus hampei*) in Puerto Rico?** *Agronomy* 2020, **10**:228.
- 779 125*. Jezeer RE, Santos MJ, Verweij PA, Boot RGA, Clough Y: **Benefits for**
780 **multiple ecosystem services in Peruvian coffee agroforestry systems**
781 **without reducing yield.** *Ecosystem Services* 2019, **40**:101033.
- 782 This study analyzes the effects of input use and shade management in coffee-based
783 agroforestry systems on coffee production and the ecosystem services of butterfly
784 pollination and soil fertility. No evidence of trade-offs or synergies between yield and
785 butterfly species richness and storage of carbon was observed. The only observed
786 effect was an improvement in carbon storage in the high-shade agroforestry systems.
- 787 126. Jonsson M, Raphael IA, Ekbohm B, Kyamanywa S, Karungi J: **Contrasting**
788 **effects of shade level and altitude on two important coffee pests.** *J Pest Sci*
789 2015, **88**:281–287.
- 790 127. Hoshino AT, Bortolotto OC, Hata FT, Ventura MU, Menezes Júnior A de O:
791 **Effect of pigeon pea intercropping or shading with leucaena plants on the**
792 **occurrence of the coffee leaf miner and on its predation by wasps in**
793 **organic coffee plantings.** *Cienc Rural* 2018, **48**:e20160863.
- 794 128. Cerda R, Allinne C, Gary C, Tixier P, Harvey CA, Krolczyk L, Mathiot C,
795 Clément E, Aubertot J-N, Avelino J: **Effects of shade, altitude and**
796 **management on multiple ecosystem services in coffee agroecosystems.**
797 *European Journal of Agronomy* 2017, **82**:308–319.
- 798 129. Fisher K, Gonthier DJ, Ennis KK, Perfecto I: **Floral resource availability from**
799 **groundcover promotes bee abundance in coffee agroecosystems.** *Ecol Appl*
800 2017, **27**:1815–1826.
- 801 130. Caudill SA, Brokaw JN, Doublet D, Rice RA: **Forest and trees: Shade**
802 **management, forest proximity and pollinator communities in southern**
803 **Costa Rica coffee agriculture.** *Renew Agric Food Syst* 2017, **32**:417–427.

- 804 131. Thapa S, Lantinga EA: **Growing Coffee in the Shade: A Strategy to**
805 **Minimize the Prevalence of Coffee White Stem Borer, *Xylocoreus***
806 ***quadripes*. *Southwestern Entomologist* 2017, **42**:357–362.**
- 807 132. Bukomeko H, Jassogne L, Kagezi GH, Mukasa D, Vaast P: **Influence of**
808 **shaded systems on *Xylosandrus compactus* infestation in Robusta coffee**
809 **along a rainfall gradient in Uganda. *Agr Forest Entomol* 2018, **20**:327–333.**
- 810 133. Vaidya C, Cruz M, Kuesel R, Gonthier DJ, Iverson A, Ennis KK, Perfecto I:
811 **Local and Landscape Constraints on Coffee Leafhopper (Hemiptera:**
812 **Cicadellidae) Diversity. *Journal of Insect Science* 2017, **17**.**
- 813 134. Liebig T, Babin R, Ribeyre F, Läderach P, van Asten P, Poehling H-M,
814 Jassogne L, Cilas C, Avelino J: **Local and regional drivers of the African**
815 **coffee white stem borer (*Monochamus leuconotus*) in Uganda: Local and**
816 **regional drivers of *M. leuconotus*. *Agr Forest Entomol* 2018, **20**:514–522.**
- 817 135. Prado SG, Collazo JA, Irwin RE: **Resurgence of specialized shade coffee**
818 **cultivation: Effects on pollination services and quality of coffee production.**
819 ***Agriculture, Ecosystems & Environment* 2018, **265**:567–575.**
- 820 136. Ennis KK, Philpott SM: **Seasonal and microhabitat differences alter ant**
821 **predation of a globally disruptive coffee pest. *Agriculture, Ecosystems &***
822 ***Environment* 2019, **284**:106597.**
- 823 137. Nesper M, Kueffer C, Krishnan S, Kushalappa CG, Ghazoul J: **Shade tree**
824 **diversity enhances coffee production and quality in agroforestry systems in**
825 **the Western Ghats. *Agriculture, Ecosystems & Environment* 2017, **247**:172–181.**
- 826 138. Mariño YA, Pérez M-E, Gallardo F, Trifilio M, Cruz M, Bayman P: **Sun vs.**
827 **shade affects infestation, total population and sex ratio of the coffee berry**
828 **borer (*Hypothenemus hampei*) in Puerto Rico. *Agriculture, Ecosystems &***
829 ***Environment* 2016, **222**:258–266.**
- 830 139. Liebig T, Jassogne L, Rahn E, Läderach P, Poehling H-M, Kucel P, Van Asten
831 P, Avelino J: **Towards a Collaborative Research: A Case Study on Linking**
832 **Science to Farmers' Perceptions and Knowledge on Arabica Coffee Pests**
833 **and Diseases and Its Management. *PLoS ONE* 2016, **11**:e0159392.**
- 834 140. Bravo-Monroy L, Tzanopoulos J, Potts SG: **Ecological and social drivers of**
835 **coffee pollination in Santander, Colombia. *Agriculture, Ecosystems &***
836 ***Environment* 2015, **211**:145–154.**

- 837 141. Márquez R, Julca O: **Organically and conventionally grown coffee in three**
838 **altitudinal floors in La Convencion Province, Cusco, Peru.** *Chilean Journal of*
839 *Agricultural and Animal Sciences* 2015, **31**:188–196.
- 840 142. Johnson MA, Manoukis NC: **Abundance of Coffee berry borer in feral,**
841 **abandoned and managed coffee on Hawaii Island.** *J Appl Entomol* 2020,
842 **144**:920–928.
- 843 143. Allinne C, Savary S, Avelino J: **Delicate balance between pest and disease**
844 **injuries, yield performance, and other ecosystem services in the complex**
845 **coffee-based systems of Costa Rica.** *Agriculture, Ecosystems & Environment*
846 2016, **222**:1–12.
- 847 144. Geeraert L, Aerts R, Jordaens K, Dox I, Wellens S, Couri M, Berecha G,
848 Honnay O: **Intensification of Ethiopian coffee agroforestry drives**
849 **impoverishment of the Arabica coffee flower visiting bee and fly**
850 **communities.** *Agroforest Syst* 2019, **93**:1729–1739.
- 851 145. Constantino LM, Gil ZN, Montoya EC, Benavides P: **Coffee Berry Borer**
852 **(*Hypothenemus hampei*) Emergence from Ground Fruits Across Varying**
853 **Altitudes and Climate Cycles, and the Effect on Coffee Tree Infestation.**
854 *Neotrop Entomol* 2021, **50**:374–387.
- 855 146. Rodríguez-Benito Y, González-Vega ME, Vázquez-López E, García-Gutiérrez
856 C, Magallanes-Tapia MA, Duarte BND, Chávez ES, Pérez-Álvarez S:
857 **Recolección de Frutos Después de la Cosecha para Reducir la Infestación**
858 **de la Broca del Café *Hypothenemus hampei* (Ferrari) en Santiago de Cuba.**
859 *Southwestern Entomologist* 2018, **43**:447–456.
- 860 147. Johnson MA, Fortna S, Manoukis NC: **Evaluation of Exclusion Netting for**
861 **Coffee Berry Borer (*Hypothenemus Hampei*) Management.** *Insects* 2020,
862 **11**:364.
- 863 148. Acevedo FE, Jiménez M, Pimentel JP, Benavides P: **Spatial Distribution of**
864 **Mealybugs (Hemiptera: Coccoomorpha: Coccoidea) in the Root System of**
865 **Pruned and Non-pruned *Coffea arabica* Trees.** *Journal of Economic*
866 *Entomology* 2019, **113**:172–184.
- 867 149. De Leijster V, Santos MJ, Wassen MW, Camargo García JC, Llorca
868 Fernandez I, Verkuil L, Scheper A, Steenhuis M, Verweij PA: **Ecosystem**
869 **services trajectories in coffee agroforestry in Colombia over 40 years.**
870 *Ecosystem Services* 2021, **48**:101246.

- 871 150. Castro AM, Tapias J, Ortiz A, Benavides P, Góngora CE: **Identification of**
872 **attractant and repellent plants to coffee berry borer, *Hypothenemus***
873 ***hampei*. *Entomol Exp Appl* 2017, **164**:120–130.**
- 874 151. Martínez-Salinas A, DeClerck F, Vierling K, Vierling L, Legal L, Vílchez-
875 Mendoza S, Avelino J: **Bird functional diversity supports pest control**
876 **services in a Costa Rican coffee farm. *Agriculture, Ecosystems & Environment***
877 **2016, **235**:277–288.**
- 878 152. Librán-Embú F, De Coster G, Metzger JP: **Effects of bird and bat exclusion**
879 **on coffee pest control at multiple spatial scales. *Landscape Ecol* 2017,**
880 ****32**:1907–1920.**
- 881 153. Armas-Quiñonez G, Ayala-Barajas R, Avendaño-Mendoza C, Lindig-Cisneros
882 R, del-Val E: **Bee diversity in secondary forests and coffee plantations in a**
883 **transition between foothills and highlands in the Guatemalan Pacific Coast.**
884 ***PeerJ* 2020, **8**:e9257.**
- 885 154. Geeraert L, Aerts R, Berecha G, Daba G, De Fruyt N, D'hollander J, Helsen K,
886 Stynen H, Honnay O: **Effects of landscape composition on bee communities**
887 **and coffee pollination in *Coffea arabica* production forests in southwestern**
888 **Ethiopia. *Agriculture, Ecosystems & Environment* 2020, **288**:106706.**
- 889 155. Berecha G, Aerts R, Muys B, Honnay O: **Fragmentation and Management of**
890 **Ethiopian Moist Evergreen Forest Drive Compositional Shifts of Insect**
891 **Communities Visiting Wild Arabica Coffee Flowers. *Environmental***
892 ***Management* 2015, **55**:373–382.**
- 893 156. Saturni FT, Jaffé R, Metzger JP: **Landscape structure influences bee**
894 **community and coffee pollination at different spatial scales. *Agriculture,***
895 ***Ecosystems & Environment* 2016, **235**:1–12.**

896
897
898

899 **Box 1. Cocoa and coffee tropical crops**

900 The genus *Coffea* (Rubiaceae) originates from tropical and subtropical Africa. Coffee
901 is a crop of global importance cultivated in approximately 80 tropical countries with
902 Brazil being the greatest producer and exporter [73]. The two main varieties cultivat-
903 ed in the tropics have different pollination requirements since in *Coffea arabica* (self-
904 compatible), animal pollination represents only 10 to 40% of total pollination while
905 animal pollination represents more than 40% in *Coffea canephora* (self-incompatible)
906 (Figure S2 in [3]). Interestingly, the absence of pollinators could decrease coffee
907 yields from 20 to 30% [20] and pest attacks harm up to 26 to 38% of productions [28].
908 The species *Theobroma cacao* is the main fruit cultivated from the genus *Theobroma*
909 (Malvaceae). Native to the rainforest regions of the northern part of South America, it
910 is currently cultivated mainly in Africa, and the Ivory Coast is the leading producer
911 country providing more than a third of global production [73]. Most of the cocoa culti-
912 vated varieties are self-incompatible and the sticky pollen is not dispersed by the
913 wind. Cocoa pollination is therefore highly dependent on animal-mediated pollination
914 [3]. We performed a systematic review on the management practices of pest control
915 and pollination services in cocoa and coffee crops, from 2015 (i.e. the establishment
916 of the IPPM concept, [23]) to May 2021 and based on two search string equations in
917 Web of Science (using the "topics" field search, [74]). Each of the equations identified
918 the scientific articles dealing with one of the two services on both crops: (*Equation 1*)
919 pest* and (damage* or control* or monitor* or manag* or suppress* or regulat*) and
920 (cocoa or cacao or coffee or coffea); (*Equation 2*) pollin* and (manag* or enhance* or
921 benefi* or success or increase* or promot* or monitor* or facilita* or support* or im-
922 prov* or favo*) and (cocoa or cacao or coffee or coffea). Out of 1224 reviewed stud-
923 ies, only 32 studies on cocoa [14,21**,37,39,48*,53,65,75–99] and 70 studies on cof-
924 fee [13,15–17,20,29*,44,45*,51*,56,57,67,69,100–156] were considered by reading
925 the abstracts and full texts and selecting that (i) address coffee and cocoa in field
926 experiments, (ii) include arthropods and (iii) did not consider the effect of a practice
927 on the ecosystem service (e.g. pest or pollinator biology, climate effect, monitoring
928 and detection). The combination of the article, the ecosystem service studied, the
929 management practice and the response metric (ex: the abundance, the diversity, the
930 richness or the infestation) was named an 'experimental unit' and used to perform an
931 analysis of the management strategies conducted on cocoa and coffee crops.

932

933 **Figure legends**

934

935 **Figure 1.** Integrated Pest and Pollinator Management (IPPM) application in coffee
936 and cocoa illustrated with a systematic review of the management strategies for
937 pollination and pest control services from 102 scientific articles. The pyramid and the
938 four categories of management from human input-based strategies (in gray) to
939 biodiversity-based strategies (in dark green) are inspired from Lundin et al. 2021
940 [19**]. The category "sampling, monitoring and threshold" was not considered in the
941 pyramid because these practices aim to monitor the level of arthropods communities
942 without trying to regulate or promote them. Icons: www.flaticon.com and
943 www.vecteezy.com

944


(A) COCOA



POLLINATION



PEST CONTROL



(B) COFFEE



POLLINATION



PEST CONTROL




Table 1: Effects of IPPM management strategies and practices on pest control and pollination services in cocoa and coffee. Studies are synthesized from January 2015 to May 2021 through a systematic review, as explained in **Box 1**. The effects presented are the result obtained on the expected ecosystem service, i.e. a practice that increases the abundance, richness and visitation rate of pollinators, or decreases their mortality, is considered to have a positive effect on the pollination service. For the pest control service, a practice that reduces the abundance or diversity of pests, or decreases their observed damage level is assumed to have a positive effect. Among the selected studies, we chose to consider only the ones that provided a conclusion on the overall effect of the practice. So, we did not include studies that aimed to compare the modalities of implementation, such as color traps or biopesticides' date of application on practice effectiveness, and studies that compared agroecosystems with many differences in term of management. N = total number of action effects; n = action effect not considered for the synthesis because they were comparisons of action modalities (ex: trap's color, biopesticide's dose) instead of showing a negative, positive or null effect. Icons: www.flaticon.com and www.vecteezy.com

Crop	Management strategy	Practice	Pollination				Pest control				References
			Negative	Null	Positive	N(n)	Negative	Null	Positive	N(n)	
Cocoa	Abiotic inputs	Pesticides	0	0	0	0	0	1	1	2(0)	[75]
		Traps	0	0	0	0	0	3	2	10(5)	[14,76–81]
	Biotic inputs	Biocontrol agent	0	0	0	0	3	4	5	12(0)	[37,82–86]
		Biopesticides	0	0	0	0	0	0	1	2(1)	[75,87]
	Crop field management	Agroforestry	1	3	1	6(1)	1	4	4	9(0)	[21**,37,39,53,65,88–91]
		Fertilization	0	0	0	0	0	2	2	4(0)	[89,92,93]
		Intensification	0	0	0	1(1)	0	0	0	2(2)	[94,95]
		Mechanical management	0	0	0	0	0	0	2	2(0)	[91,96*]
		Plot floral abundance	0	1	1	2(0)	0	0	0	0	[21**]
		Nesting habitat diversity	0	2	2	7(3)	2	1	1	5(1)	[21**,48*,75,83,96*–99]
Landscape management	Biodiversity conservation	0	0	0	0	2	2	2	6(0)	[37,88]	
	Semi-natural habitats	0	1	1	2(0)	0	0	0	0	[21**]	
Coffee	Abiotic inputs	Pesticides	0	1	1	2(0)	0	3	3	6(0)	[13,57,100–103]
		Traps	0	0	0	2(2)	0	1	0	6(5)	[104–110]
	Biotic inputs	Biocontrol agent	3	0	0	3(0)	3	11	12	26(0)	[15–17,56,69,101,111–117]
		Biopesticides	0	0	0	0	0	1	7	10(2)	[100,106,118–123]

Crop field management	Agroforestry	7	8	3	18(0)	6	15	8	30(1)	[16,29*,44,45*,101,124–140]
	Fertilization	2	2	2	6(0)	0	2	1	3(0)	[45*,136,140,141]
	Intensification	3	3	1	7(0)	1	3	2	7(0)	[20,45*,125*,128,142–144]
	Mechanical management	0	0	0	0	2	3	5	10(0)	[13,103,106,145–148]
	Plot floral abundance	2	3	8	13(0)	0	0	0	0	[20,129,130,140]
	Nesting habitat diversity	1	3	1	5(0)	2	4	4	10(0)	[29*,101,130,136,137,149,150]
Landscape management	Biodiversity conservation	0	0	0	0	1	2	3	6(0)	[29*,151,152]
	Semi-natural habitats	4	19	13	38(2)	4	5	7	16(0)	[15,17,20,29*,44,51*,67,125*,130,133,140,152–156]

Pollination



Pest control



IPPM

Application to tropical crops?



Cocoa



Coffee