



HAL
open science

Estimation of multivariate generalized gamma convolutions, application to dependence structure modelling in insurance.

Véronique Maume-Deschamps

► **To cite this version:**

Véronique Maume-Deschamps. Estimation of multivariate generalized gamma convolutions, application to dependence structure modelling in insurance.. Conference on Industrial and Applied Mathematics - ECMI 2023, Jun 2023, Wrocław, Poland. hal-04294063

HAL Id: hal-04294063

<https://hal.science/hal-04294063>

Submitted on 19 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Estimation of multivariate generalized gamma convolutions, application to dependence structure modelling in insurance.

Véronique Maume-Deschamps, joint work with Oskar Laverny,
Esterina Masiello, Didier Rullière.
Partially funded by SCOR

ECMI 2023



Plan

- 1 Introduction / motivation
- 2 Multivariate generalized Gamma convolutions
- 3 Simulations

Plan

- 1 Introduction / motivation
- 2 Multivariate generalized Gamma convolutions
- 3 Simulations

Introduction

A (re-)insurer observes many (positive) random variables that represent losses from several portfolios of risks.

A major topic is the [evaluation and modeling of the dependency](#) between those risks. Indeed, the properties of this dependency has a huge impact on the behavior of the total loss, and therefore on the capital management of the company.

Risks can be heavy-tailed, skewed, and are usually dependant with each other, including in the extremes.

The infinite divisibility may be a wanted property.


Why infinite divisibility ?

Definition (ID r.v. X)

- 1 X is n -divisible if and only if there exists X_1, \dots, X_n i.i.d. such that $X = X_1 + \dots + X_n$
- 2 X is ID iff this can be done for all n .
- 3 X_i is called a *piece* of X .

- Dependencies are largely unobservables in reinsurance;
- Experts are providing information as divisibility parameters: "20% of this is driven by the same risk factors as 10% of that.." *this means comonotonic pieces accross marginals.*

Company's Solvency



SC (Solvency Capital): regulatory capital calculated using a risk measure that the re-insurer must have to absorb potential losses. For example, in the Solvency II directive of the European Union, it is the $VaR(99.5\%)$ of the P&L distribution.

Risk factors modelisation

In the **internal modeling** point of view, the distributions of several losses X_1, \dots, X_d are supposed to be known, but the dependence structure between them must still be evaluated and taken into account to assess the variability and behavior of the total loss:

$$\sum_{i=1}^d X_i.$$

It might be achieved by a **risk factors decomposition**: $X_i = \sum_{j=1}^k X_{i,j}$,

where $X_{i,j}, X_{\ell,j}$ are either independent or comonotonic.

Divisibility allows to interpret the $X_{i,j}$ as pieces of X_i .

⇒ multivariate generalized Gamma convolutions give an answer to that question.

Plan


- 1 Introduction / motivation
- 2 Multivariate generalized Gamma convolutions
- 3 Simulations

Recall Gamma distributions

Well known class of infinite divisible distributions.

$X \sim \mathcal{G}(k, \theta)$ if its density function is:

$$\frac{x^{k-1} e^{-x/\theta}}{\Gamma(k) \theta^k}.$$



Multivariate Gamma convolutions classes $\mathcal{G}_{d,n}$ and \mathcal{G}_d

$\mathbf{x}, \mathbf{t} \in \mathbb{R}^d$, **Recall:** Cumulant generating function:

$$K(\mathbf{t}) = \ln \left(\mathbb{E} \left(e^{\langle \mathbf{t}, \mathbf{X} \rangle} \right) \right).$$

Multivariate Thorin Classes ¹

$$\mathbf{X} \sim \mathcal{G}_{d,n}(\alpha, \mathbf{s}) \Leftrightarrow K(\mathbf{t}) = - \sum_{i=1}^n \alpha_i \ln (1 - \langle \mathbf{s}_i, \mathbf{t} \rangle)$$

$$\mathbf{X} \sim \mathcal{G}_d(\nu) \Leftrightarrow K(\mathbf{t}) = - \int \ln (1 - \langle \mathbf{s}, \mathbf{t} \rangle) \nu(d\mathbf{s}).$$

Example

All gammas, log-Normals, Paretos, α -stables, and sums and/or products of these are in \mathcal{G}_1 .

No estimation procedures were available before our work.

¹ Lennart Bondesson (Nov. 2009). en. In: *Journal of Statistical Planning and Inference*

Motivation: interpretability of $\mathcal{G}_{d,n}$ models

$\mathbf{X} \in \mathcal{G}_{d,n}$ has an additive risk-factor structure: there exists gamma random variables G_1, \dots, G_n such that:

$$\begin{pmatrix} X_1 \\ \dots \\ X_d \end{pmatrix} = \begin{pmatrix} s_{1,1} & \dots & \dots & s_{1,n} \\ \dots & \dots & \dots & \dots \\ s_{d,1} & \dots & \dots & s_{d,n} \end{pmatrix} \cdot \begin{pmatrix} G_1 \\ \dots \\ G_n \end{pmatrix}$$

- 1 For all $i \in 1, \dots, n$, $G_i \sim \mathcal{G}_{1,1}(\alpha_i, 1)$
- 2 G_1, \dots, G_n are independent.

Goal: Estimate ν from observations of the random vector \mathbf{X} .

Pb: It is a deconvolution problem, which is numerically hard (can also be interpreted as a solution-less moment problem for the Thorin measure, see MFK²).

² Justin Miles, Edward Furman, and Alexey Kuznetsov (2019). In: *Variance*

The orthonormal Laguerre basis of $L_2(\mathbb{R}_+^d)$

Consider the following **orthonormal basis of $L_2(\mathbb{R}_+^d)$** called **Laguerre basis**³: let $\mathbf{p} \in \mathbb{N}^d$,

$$\phi_{\mathbf{p}}(\mathbf{x}) = \prod_{i=1}^d \phi_{p_i}(x_i), \text{ with } \phi_p(x) = \sqrt{2} \sum_{k=0}^p \binom{p}{k} \frac{(-2x)^k}{k!} e^{-x}.$$

Any square integrable distribution density f can be expanded as :

$$f(\mathbf{x}) = \sum_{\mathbf{p} \in \mathbb{N}^d} a_{\mathbf{p}} \phi_{\mathbf{p}}(\mathbf{x}) \text{ where } a_{\mathbf{p}} = \int \phi_{\mathbf{p}}(\mathbf{x}) f(\mathbf{x}) \partial \mathbf{x}$$

³ Florian Dussap (2021). In: *Journal of Statistical Planning and Inference*

Estimation

Consider a random sample $\mathbf{X}_1, \dots, \mathbf{X}_N$ of $\mathbf{X} \in \mathcal{G}_d(\alpha, \mathbf{s})$. Let

$$\widehat{\mathbf{a}}_{\mathbf{p}} = \frac{1}{N} \sum_{i=1}^N \phi_{\mathbf{p}}(\mathbf{X}_i).$$

Consider the **integrated square error loss**:

$$L(\alpha, \mathbf{s}) = \sum_{\mathbf{k} \leq \mathbf{m}} (\widehat{a}_{\mathbf{k}} - a_{\mathbf{k}}(\alpha, \mathbf{s}))^2 \text{ and}$$

$$(\widehat{\alpha}, \widehat{\mathbf{s}}) = \operatorname{argmin}_{(\alpha, \mathbf{s})} L(\alpha, \mathbf{s}).$$

Remark that the theoretical Laguerre coefficients may be computed using the formula:

$$a_{\mathbf{k}} = \langle \phi_{\mathbf{k}}, f \rangle = \mathbb{E}(\phi_{\mathbf{k}}(\mathbf{X})) = \sqrt{2}^d \sum_{\mathbf{k} \leq \mathbf{p}} \binom{\mathbf{p}}{\mathbf{k}} \frac{(-2)^{|\mathbf{k}|}}{\mathbf{k}!} \mathbb{E}\left(\mathbf{X}^{\mathbf{k}} e^{-|\mathbf{X}|}\right).$$

Consistency

We have the following consistency result ⁴ under the technical *well-behaved* assumption (which holds for univariate Gamma convolutions, independent multivariate Gamma convolution and invertible linear transformations of them e.g.).

Theorem

If \mathbf{X} is drawn from an ϵ -well-behaved density $f_{\alpha, \mathbf{s}} \in \mathcal{G}_d(\alpha, \mathbf{s})$ then,

$$\|f_{(\widehat{\alpha}, \widehat{\mathbf{s}})} - f_{\alpha, \mathbf{s}}\|_2^2 \xrightarrow[N \rightarrow \infty, \mathbf{m} \rightarrow \infty]{a.s.} 0.$$

⁴ Oskar Laverny et al. (Jan. 2021). In: *Electronic Journal of Statistics*

Dimensionality issues

The loss $L(\alpha, \mathbf{s})$ is too costly to work with when d gets large. Indeed, the number of coefficients to compute is given by

$$D(m, d) = \sum_{i=1}^m \binom{i+d-1}{d-1}.$$

which is exponentially increasing in d and therefore unusable in high dimension (ok until $d = 4$ or 5).

Improvement of the method by Oskar Laverny⁵ by using an approximation of an integrated loss function and random projections (tested for $d = 1000$).

⁵ Oskar Laverny (2022). In: *arXiv preprint arXiv:2203.13741*

Plan

- 1 Introduction / motivation
- 2 Multivariate generalized Gamma convolutions
- 3 Simulations**

Multivariate log-Normal




Figure: MLN case: $n = 10$

Multivariate log-Normal




Figure: MLN case: $n = 20$

Tail dependency: A Clayton copula




Figure: Clayton case: $n = 20$

Conclusions

- The univariate Thorin class is wide: LN, Pareto, α -stable, (some) Weibull, ...
- The Multivariate analogue provides an asymmetrical dependence structure which include tail dependency, and can take a lot of different shapes.
- Deconvolution is a hard inverse problem, and estimation of these distributions is complicated.
- The final additive risk-factor model gives easy interpretation of parameters and easy aggregation schemes. Should be shortly **used by SCOR in some parts of their internal model.**
- Improvement to go to high dimension in Oskar' work ⁶ and **the open-source Julia package ⁷ provides the implementation.**

⁶ Oskar Laverny (2022). In: *arXiv preprint arXiv:2203.13741*

⁷ Oskar Laverny (Mar. 2021). Version v0.1

References I



Bondesson, Lennart (Nov. 2009). "On Univariate and Bivariate Generalized Gamma Convolutions". en. In: *Journal of Statistical Planning and Inference* 139.11, pp. 3759–3765.



Dussap, Florian (2021). "Anisotropic multivariate deconvolution using projection on the Laguerre basis". In: *Journal of Statistical Planning and Inference* 215, pp. 23–46.



Laverny, Oskar (Mar. 2021). *ThorinDistributions.jl*. Version v0.1.



— (2022). "Estimation of high dimensional Gamma convolutions through random projections". In: *arXiv preprint arXiv:2203.13741*.



Laverny, Oskar et al. (Jan. 2021). "Estimation of Multivariate Generalized Gamma Convolutions through Laguerre Expansions.". In: *Electronic Journal of Statistics* 15.2.



Miles, Justin, Edward Furman, and Alexey Kuznetsov (2019). "Risk Aggregation: A General Approach via the Class of Generalized Gamma Convolutions". In: *Variance*.

Thank you

Thanks for your attention.