



**HAL**  
open science

# Variability and accuracy of multiple saliva pepsin measurements in laryngopharyngeal reflux patients

Jêrome René Lechien, François Bobin

## ► To cite this version:

Jêrome René Lechien, François Bobin. Variability and accuracy of multiple saliva pepsin measurements in laryngopharyngeal reflux patients. *Journal of Otolaryngology - Head and Neck Surgery*, 2023, 52 (1), 10.1186/s40463-023-00670-5 . hal-04294692

**HAL Id: hal-04294692**

**<https://hal.science/hal-04294692v1>**

Submitted on 23 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

ORIGINAL RESEARCH ARTICLE

Open Access



# Variability and accuracy of multiple saliva pepsin measurements in laryngopharyngeal reflux patients

Jerome R. Lechien<sup>1,2,3,4,5,6\*</sup> and Francois Bobin<sup>5,6</sup>

## Abstract

**Objective** To study the variability and diagnostic value of multiple salivary pepsin measurements in the detection of laryngopharyngeal reflux (LPR).

**Methods** Patients with LPR symptoms were consecutively recruited from December 2019 to Augustus 2022. Twenty-one asymptomatic individuals completed the study. The diagnosis was confirmed with hypopharyngeal–esophageal impedance-pH monitoring (HEMII-pH). Patients collected three saliva samples during the 24-h testing period. Symptoms and findings were studied with reflux symptom score-12 and reflux sign assessment. Sensitivity, specificity, positive (PPV) and negative (NPV) predictive values of pepsin measurements were calculated considering morning, post-lunch and post-dinner samples. The consistency and relationship between HEMII-pH, pepsin measurements, and clinical features were investigated.

**Results** Morning, post-lunch and post-dinner saliva pepsin concentrations were measured in 42 patients. Pepsin measurements were 64.9%, 59.5%, and 59.0% sensitive for morning, post-lunch and post-dinner collections at cut-off  $\geq 16$  ng/mL. Considering the highest concentration of the three pepsin saliva collections, the accuracy, sensitivity, specificity and PPV were 70.5%, 73.0%; 66.7% and 78.9%, respectively. Morning pepsin measurements reported higher consistency, sensitivity, and specificity than post-dinner and post-lunch pepsin measurements.

**Conclusion** The collection of several saliva pepsin samples improves the detection rate of LPR. In case of high clinical LPR suspicion and negative pepsin test, a HEMII-pH study could provide further diagnostic information.

**Keywords** Laryngopharyngeal, Reflux, Voice, pH monitoring, Laryngeal, Otolaryngology, Saliva, Pepsin

\*Correspondence:

Jerome R. Lechien

[Jerome.Lechien@umons.ac.be](mailto:Jerome.Lechien@umons.ac.be)

<sup>1</sup> Division of Laryngology and Broncho-Esophagology, Department of Otolaryngology-Head Neck Surgery, EpiCURA Hospital, Baudour, Belgium

<sup>2</sup> Department of Human Anatomy and Experimental Oncology, Faculty of Medicine, UMONS Research Institute for Health Sciences and Technology, University of Mons (UMons), Avenue du Champ de Mars, 6, 7000 Mons, Belgium

<sup>3</sup> Department of Otorhinolaryngology and Head and Neck Surgery, Foch Hospital, School of Medicine, UFR Simone Veil, Université Versailles Saint-Quentin-en-Yvelines (Paris Saclay University), Paris, France

<sup>4</sup> Department of Otorhinolaryngology and Head and Neck Surgery, CHU de Bruxelles, CHU Saint-Pierre, School of Medicine, Université Libre de Bruxelles, Brussels, Belgium

<sup>5</sup> Polyclinique Elsan de Poitiers, Poitiers, France

<sup>6</sup> Research Committee of Young Otolaryngologists of the International Federation of Oto-Rhinolaryngological Societies (YO-IFOS), Paris, France



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Introduction

Laryngopharyngeal reflux (LPR) is an inflammatory condition of the upper aerodigestive tract tissues related to direct and indirect effect of gastroduodenal content reflux, which induces morphological changes in the upper aerodigestive tract [1]. The symptoms and findings are related to the deposit of pepsin in laryngopharyngeal mucosa and the development of inflammatory reaction and mucosa injuries [2]. Currently, the LPR diagnosis may be confirmed with the 24-h hypopharyngeal–esophageal multichannel intraluminal impedance-pH monitoring (HEMII-pH), which detects pharyngeal reflux events [3]. HEMII-pH is considered as the gold standard for the LPR diagnosis, but this tool remains costly and inconvenient. HEMII-pH is uncommonly used by otolaryngologists [4, 5]. Over the past 2 decades, saliva pepsin measurement was proposed as a cost-effective and minimal invasive diagnostic tool for LPR [6]. However, studies reported inconsistencies regarding the cutoff, the collection time and the appropriate number of saliva samples for the diagnostic. These inconsistencies may be due to the variability of the saliva pepsin concentration over time [6–10].

The aim of this study was to investigate the variability of pepsin saliva concentration throughout the day and its diagnostic value for detecting laryngopharyngeal reflux.

## Material and methods

### Subjects and setting

Patients with suspected LPR regarding laryngopharyngeal symptoms and findings were consecutively recruited from two European hospitals (Saint-Pierre University Hospital of Brussels (Belgium) and Elsan Polyclinic of Poitiers (France)). The LPR diagnosis was confirmed with 24-h HEMII-pH. Gastrointestinal (GI) endoscopy was proposed to elderly individuals and those with gastroesophageal reflux disease (GERD) complaints. The exclusion criteria included the following: active smoker, alcoholic (>3 alcohol glasses daily), history of upper respiratory tract infection within the last month, neurological or psychiatric illness, head and neck malignancy, head and neck radiotherapy, active seasonal allergies, intake of inhaled corticosteroids, or asthma.

The local ethics committee approved the study protocol (CHU Saint-Pierre, n°BE076201837630). Patients consented to participate.

### Hypopharyngeal–esophageal multichannel intraluminal impedance-pH monitoring

The catheter placement and tracing analyses were described in a previous publication (Versaflex Z<sup>®</sup>, Digi-trapper pH-Z testing System, Medtronic, Europe) [3]. In sum, the catheter was placed in the morning fasting (8:00

AM) and removed the next day in the morning. The catheter was composed of 8 impedance segments and 2 pH electrodes. The six esophageal impedance segments were placed along the esophagus zones (Z1 to Z6) at 19, 17, 11, 9, 7 and 5 cm above the lower esophageal sphincter (LES). The pharyngeal impedance segments were placed 1 and 2 cm above the upper esophageal sphincter (UES) in the hypopharynx. The pH electrodes were placed 2 cm above LES and 2 cm below UES, respectively. According to recent systematic review on HEMII-pH features in healthy individuals [11], the LPR diagnosis criteria was based on the occurrence of >1 acid ( $\text{pH} \leq 4.0$ ) or non-acid ( $\text{pH} > 4.0$ ) hypopharyngeal reflux events (off proton pump inhibitors).

### Saliva pepsin measurements

The pepsin concentration was measured in the saliva samples with Peptest<sup>®</sup> device (RD Biomed Ltd., Hull, United Kingdom). Patients collected 1 to 5 mL saliva samples in the morning (fasting, after waking) and 2 h after the lunch and the dinner. The saliva was collected during the 24-h HEMII-pH period. The saliva was collected into a 30-mL universal sample collection tube containing a pre-established concentration of citric acid. Pepsin is active at acidic pH (around 2.0 to 4.0). The citric acid in the collection tube helps to maintain this low pH level acting as a buffer. It prevents the pH of the sample of increasing, which would deactivate the pepsin and compromise the accuracy of the test results.

The pepsin sample collections were stored in the refrigerator for a period of up to one week, which was found to have no significant impact on the pepsin measurement [12]. A trained lab technician analyzed the samples according to a standardized procedure [12]. The result of the pepsin test was validated when a blue line appeared under the letter C (control) of the device 15 min after applying the sample. The apparition of a blue line under the letter T (test) meant that the pepsin test was positive. The Cube Reader<sup>®</sup> was used to precisely measure the level of pepsin, the concentration ranging from 1 to 500 ng/mL.

### Clinical outcomes and control group

Demographic outcomes (e.g. age, gender, body mass index) were collected from the patient's medical record. Reflux Symptom Score-12 (RSS-12) [13] was used to rate the severity and the frequency of symptoms. A RSS-12 > 11 was suggestive of LPR. Findings were evaluated with Reflux Sign Assessment (RSA) [14], which rates oral, pharyngeal and laryngeal signs associated with LPR. An RSA > 14 may be suggestive of LPR.

RSS-12 and RSA were used to recruit 21 asymptomatic individuals who reported RSS-12 < 11 and RSA < 14.

Exclusion criteria of asymptomatic individuals were similar to those of LPR patients. They similarly collected saliva samples to measure pepsin.

**Statistical methods**

Statistical Package for the Social Sciences for Windows (SPSS version 27.0; IBM Corp, Armonk, NY, USA) was used for the statistical analyses.

A power analysis was performed, in which the ideal sample size for our study was calculated focusing on the diagnostic accuracy of pepsin tests in the previous studies. Precisely, the anticipated SE of the tests was set at 85.0, indicating an expectation of high true positive rate, while the anticipated SP was set at 40.0, acknowledging a relatively high rate of false positives. We assumed an imbalance in the distribution of healthy to diseased individuals, with a ratio (R) set to 1/5, reflecting the prevalence of LPR in the population ( $\cong 10\%$ ). The statistical power, a measure of the study’s ability to detect a true effect, was set at a standard value of 0.80. The significance level, a threshold for determining statistical significance, was set at 0.05. Based on these assumptions, a simplified R function was used to calculate the required sample size.

The sensitivity, specificity, positive (PPV) and negative predictive value (NPV) of the pepsin measurement were evaluated considering several thresholds ( $\geq 16, \geq 36, \geq 45, \geq 75$  and  $\geq 100$  ng/mL). Associations between morning, post-lunch, post-dinner pepsin saliva measurements and clinical findings was investigated through multivariate analysis. The association was defined as low ( $r_s < 0.30$ ), moderate ( $r_s = 0.30-0.60$ ) or strong ( $r_s > 0.60$ ), respectively. The consistency between pepsin measurements, HEMII-pH and clinical findings was assessed with kappa-Cohen analysis.

**Results**

**Setting**

Forty-two patients and 21 asymptomatic individuals collected saliva samples. The mean age of patients was  $48.0 \pm 19.6$  years. There were 21 females (50%). Twenty-eight patients underwent GI endoscopy, which was unremarkable in 13 (46.0%) patients (Table 1). The LPR diagnosis was positive at the HEMII-pH in 39 patients (92.9%), which was not significantly consistent with the pepsin measurements (Table 2). The mean RSS-12 was  $71.1 \pm 41.1$ . The mean RSA was  $22.9 \pm 10.9$ . The combination of  $RSS-12 > 11$  with  $RSA > 14$  was significantly consistent with the LPR diagnostic at the HEMII-pH (Table 2).

**Accuracy of pepsin test**

The detection of LPR at the pepsin measurements regarding several cutoffs was reported in Table 3. The

**Table 1** Characteristics of patients

Characteristics	
Age (mean, SD)	50.0 ± 16.6
BMI (mean, SD)	26.2 ± 5.5
Male (N, %)	21 (50)
Female (N, %)	21 (50)
<i>Gastrointestinal endoscopy (N = 28)</i>	
Normal	13 (46)
Esophagitis	3 (11)
Hiatal hernia	8 (29)
LES insufficiency	9 (32)
Gastritis	7 (25)
<i>HEMII-pH (mean, SD)</i>	
Pharyngeal acid events	17.0 ± 17.7
Pharyngeal nonacid events	16.8 ± 22.3
Pharyngeal events (total number)	33.6 ± 27.8
<i>Saliva pepsin measurements (mean, SD)</i>	
Morning pepsin test	85.5 ± 96.5
Post-lunch pepsin test	120.2 ± 132.2
Post-dinner pepsin test	104.7 ± 118.6
Mean concentration of pepsin tests	99.5 ± 86.4
Highest concentration of pepsin test	170.0 ± 136.4

BMI, body mass index; HEMII-pH, hypopharyngeal–esophageal multichannel intraluminal impedance-pH monitoring; LES, lower esophageal sphincter; N, number; SD, standard deviation

**Table 2** Consistency findings

Outcomes	HEMII-pH	
	Kappa	p value
Morning pepsin test	0.036	NS
Post-lunch pepsin test	0.024	NS
Post-dinner pepsin test	0.022	NS
Highest pepsin test	0.095	NS
RSS-12 > 11	0.050	NS
RSS-12 > 11 and RSA > 14	0.638	0.005

HEMII-pH, hypopharyngeal–esophageal multichannel intraluminal impedance-pH monitoring; NS, non-significant; RSA, reflux sign assessment; RSS-12, reflux symptom score-12

sensitivity, specificity, PPV and NPV of pepsin measurements was calculated considering a single, two or three measurements (Table 4). The selection of the highest pepsin saliva concentration of the 3 pepsin measurements was associated with the highest accuracy for detecting LPR (70.5%; cutoff  $\geq 16$  ng/mL). Considering the association of two pepsin measurements, the combination of morning and post-lunch pepsin measurements was 85.7% accurate (Table 4).

**Table 3** Accuracy of saliva pepsin test according to thresholds

Pepsin tests	Rate
Morning	
≥ 16 ng/mL	65.6
≥ 36 ng/mL	64.5
≥ 45 ng/mL	64.4
≥ 75 ng/mL	62.7
≥ 100 ng/mL	62.7
Post-lunch	
≥ 16 ng/mL	62.3
≥ 36 ng/mL	65.6
≥ 45 ng/mL	63.9
≥ 75 ng/mL	60.7
≥ 100 ng/mL	60.7
Post-dinner	
≥ 16 ng/mL	61.9
≥ 36 ng/mL	63.5
≥ 45 ng/mL	61.9
≥ 75 ng/mL	61.9
≥ 100 ng/mL	61.9

The best accuracy value was found for fasting saliva pepsin test with a cutoff of ≥ 16 ng/mL

**Table 4** Accuracy of saliva pepsin test association

Highest sample concentration	
cutoff ≥ 16 ng/mL	Rate
Morning + post-lunch peptest	65.6
Morning + post-dinner peptest	65.6
Post-lunch + post-dinner	62.3
Morning + post-lunch + post-dinner	70.5

The association of the three saliva pepsin tests (morning, post-lunch and post-dinner) reported the best accuracy rate

HEMII-pH, hypopharyngeal-esophageal multichannel intraluminal impedance-pH monitoring

**Sensitivity, specificity and predictive values of pepsin test**

The sensitivity, specificity, PPV and NPV of pepsin measurements at cutoffs ≥ 16, ≥ 36, ≥ 45, ≥ 75 and ≥ 100 ng/

mL were reported in Table 5. Morning saliva pepsin measurement was 64.9% sensitive and 66.7% specific at cutoffs ≥ 16 ng/mL. At cutoffs ≥ 36 ng/mL, the morning pepsin test was 54.1% sensitive and 80.0% specific. Morning pepsin measurement reported overall higher sensitivity, specificity, PPV and NPV than post-lunch and post-dinner measurements (Table 5). The sensitivity, specificity, PPV and NPV of the ‘highest pepsin measurement’ were 73.0% (95%IC: 69.7, 76.3), 66.7% (95%IC: 66.3, 67.1), 78.9% (95%IC: 78.1, 79.6), and 64.0% (95%IC: 61.4, 66.6) at cutoff ≥ 16 ng/mL, respectively.

**Multivariate analyses**

There was a strong association between the number of pharyngeal reflux events and the RSA ( $r_s=0.634$ ;  $p=0.006$ ). The level of morning saliva pepsin was moderately associated with the level of post-dinner pepsin ( $r_s=0.429$ ;  $p=0.007$ ) and the RSA ( $r_s=0.578$ ;  $p=0.019$ ). The post-lunch pepsin level was moderately correlated with the post-dinner pepsin level ( $r_s=0.369$ ;  $p=0.019$ ). There was no significant association between the morning and post-dinner pepsin saliva concentrations.

**Discussion**

The pepsin saliva test was developed to detect laryngopharyngeal reflux disease without the need for HEMII-pH [6]. To date, studies reported controversial results about the most appropriate time of saliva collection, and the related accuracy, sensitivity, specificity, and predictive values of pepsin saliva measurements. The accuracy and predictive values of the pepsin test were investigated in few studies, which reported controversial results (Table 6) [2, 6, 7, 9, 10, 14–18]. Overall, SE and SP ranged from 29.4 to 100% according to studies, where authors collected saliva sample in the morning, post-meals or after symptoms. Pepsin test appears to be sensitive but not specific. However, most authors included only LPR patients and the lack of control groups may significantly influence the assessment of predictive values and accuracy of pepsin test.

**Table 5** Characteristics of patients according to the reflux profiles

	Morning pepsin test				Post-lunch pepsin test				Post-dinner pepsin test			
	SE	SP	PPV	NPV	SE	SP	PPV	NPV	SE	SP	PPV	NPV
≥ 16 ng/mL	64.9	66.7	75.0	55.2	59.5	66.7	73.3	51.6	59.0	66.7	74.2	50.0
≥ 36 ng/mL	54.1	80.0	80.0	54.0	56.8	79.2	80.8	54.3	53.9	79.2	80.0	51.4
≥ 45 ng/mL	51.4	83.3	81.8	54.1	54.1	79.2	80.0	52.8	51.3	79.2	80.0	50.0
≥ 75 ng/mL	48.6	83.3	81.0	52.6	45.9	83.3	80.9	50.0	48.7	83.3	82.6	50.0
≥ 100 ng/mL	45.7	87.5	84.2	52.5	45.9	83.3	80.9	50.0	48.7	83.3	82.6	50.0

SE, sensitivity; SP, specificity; PPV, positive predictive value; NPV, negative predictive value

**Table 6** Literature studies

Authors	Ref	Years	LPR/CT	LPR diagnosis	Sample time	Pepsin analysis	Thresholds	SE	SP	PPV	NPV	Accuracy
Wang China	10	2022	112/26	HEMI-pH ≥ 1 HRE	From morning 07:00:00 to 6:00 PM	Rapid pepsin lateral flow Highest saliva concentration PepTest®	> 45 ng/mL Fasting test > 45 ng/mL Highest test ≥ 16 ng/mL	38.4 86.6 48.0	84.6 80.8 27.0	91.5 95.1 63.0	24.2 58.3 40.0	55.7 73.9 48.0
Zelenik Czech	9	2021	45/1	HEMI-pH > 1 HRE	Morning	PepTest®	≥ 16 ng/mL	76.9	25.0	87.0	14.3	87.0
Zhang Australia	7	2020	26/4	HEMI-pH ≥ 2 HRE ≥ 6 PRE	Morning 1-h post-lunch 1-h post-dinner When symptoms	PepTest®	≥ 75 ng/mL	57.7	75.0	93.8	21.4	N.P
Klimara U.S.A	2	2019	19/7	HEMI-pH > 1 HRE > 40 PRE	Morning 1-h post-lunch 1-h post-dinner 1-h post-breakfast 3 samples	ELISA Western blot Highest pepsin concentration PepTest®	> 1 ng/mL	29.4	50.0	62.5	20.0	42.0
Weitzendorfer Austria	15	2019	41/29	Restech Ryan score (> 9.4)	3 samples	Highest of 3 samples PepTest®	> 16 ng/mL > 50 ng/mL > 100 ng/mL > 150 ng/mL > 216 ng/mL	85.4 78.1 68.3 53.7 41.5	27.6 41.4 58.6 69.0 86.2	62.5 65.3 70.0 71.0 81.0	57.1 57.1 56.7 51.3 51.0	85.4
Hayat UK	6	2015	111/100	MII-pH EAT > 4.2%	Morning 1-h post-lunch 1-h post-dinner	PepTest®	> 16 ng/mL > 50 ng/mL > 100 ng/mL > 150 ng/mL > 210 ng/mL	77.6 67.2 51.7 41.4 44.2	63.2 76.3 74.5 90.8 96.3	58.4 67.2 54.5 75.0 95.7	80.4 76.8 72.3 69.9 36.5	N.P
Ocak Turkey	16	2015	18/2	Dual-probe pH testing EAT > 4.2%	N.P	PepTest®	≥ 16 ng/mL	33.0	100	100	14.2	N.P
Saritas U.S.A	17	2012	22/25	Wireless pH testing EAT > 4.2%	N.P	PepTest®	≥ 50 ng/mL	50.0	92.0	85.0	68.0	N.P
Potluri U.S.A	18	2003	3/13	Dual-probe pH testing ≥ 1 proximal acid esophageal event	When symptoms	Pepsin assay	> 1 ng/mL	100	92.3	N.P	N.P	N.P

EAT, esophageal acid exposure time; (HEMI)-pH, hypopharyngeal-esophageal multichannel-intraluminal impedance-pH monitoring; HRE, hypopharyngeal reflux event(s); LPR/CT, laryngopharyngeal reflux event(s); NP, not provided; NPV, negative predictive value; PPV, positive predictive value; PRE, proximal esophageal reflux event(s); ref., reference; SE, sensitivity; SP, specificity



Our results suggested a variability of the pepsin saliva concentration throughout the day, and the lack of significant consistency with the HEMII-pH results. The morning saliva pepsin measurement appeared to be associated with the highest sensitivity and accuracy, when compared to other measurements. In 2016, Na et al. observed that the average pepsin level upon waking was higher than that measured at any other time [19]. Wang et al. corroborated these findings in a recent study where the morning saliva pepsin measurement was associated with the highest LPR detection rate [10]. The importance of the morning saliva collection was however not supported by Weitzendorfer et al. who observed higher saliva pepsin concentrations after the dinner and the lunch compared to waking concentrations [15]. In other studies, authors reported a variability between morning, post-lunch and post-dinner pepsin saliva concentrations [6, 7] without determining the most adequate time of saliva collection.

The problem of the variability of saliva pepsin concentration and the related discrepancies across studies in accuracy, sensitivity and predictive values may be tackled by the collection of multiple saliva samples. Indeed, as suggested in the present study, recent studies supported that sensitivity, specificity and predictive values may be raised when considering the highest pepsin measurement of 2 or 3 saliva sample collections within the testing day [2, 10, 15]. Considering the highest saliva pepsin measurement, sensitivity, specificity, and PPV found in the present study corroborated those of the literature (Table 6) [2, 10, 15]. Precisely, Wang et al. [10] reported that 55.7% of the true positive cases were missed by considering a single pepsin test. Similarly, Hayat et al. and Zhang et al. supported that the accuracy, sensitivity, specificity, and predictive values of pepsin saliva measurements were improved when considering the highest pepsin saliva concentration of three or four measurements, respectively [6, 7].

To date, the variability of pepsin saliva concentration throughout the day is not fully understood. Several factors may influence the gastric pepsin secretions, the esophageal motility, the relaxation of sphincters, and the related pepsin saliva concentration. First, it has been suggested that the foods and beverages consumed during the testing day may influence the pepsin saliva concentration [16, 17]. On the one hand, foods and beverages may influence the gastric secretion of pepsin, and, therefore, the pepsin concentration into the stomach content that may refluxate into the upper aerodigestive tract tissues [18, 19]. On the other hand, acid, spicy, low-protein, and high-fat foods may increase the number of transient relaxations of esophageal sphincters, leading to an increased number of pharyngeal reflux events that contain pepsin [16, 17]. Regarding the influence of diet, the

differences across studies from different world regions should be interpreted according to the diet habits of populations.

Both esophageal sphincter tonicity and motility are known to be influenced by the autonomic nerve function [20, 21]. The activation of sympathetic nervous system may impair the esophageal antireflux barriers (sphincter tonicity and esophageal motility), leading to pharyngeal reflux events. In that way, patients with stress, anxiety or depressive findings at the time of the diagnosis/testing should have higher number of pharyngeal reflux events and, theoretically, higher pepsin saliva concentration compared to patients without autonomic nerve dysfunction [20, 21]. In addition to these factors, it is important to keep in mind that the saliva pepsin measurements highlight the extracellular pepsin concentrations, while recent studies suggested a potential internalization of pepsin into the Golgi apparatus of pharyngeal cells [22], which makes undetectable a part of refluxate pepsin.

To the best of our knowledge, the present report is the first study investigating accuracy, sensitivity, specificity and predictive values of pepsin saliva measurements according to the time of saliva collection. The primary limitation of the present study was the homogeneity of the study population, which mainly included patients with a positive diagnostic at the HEMII-pH and only 21 asymptomatic individuals. The lack of healthy individuals benefiting from HEMII-pH may be considered as a limitation but HEMII-pH is costly and inconvenience for asymptomatic patients. Future studies are needed to better understand the low SE and SP of pepsin test, and to investigate the presence of other gastroduodenal enzymes in the saliva of patients. Indeed, the presence of other enzymes, such as bile salts, should explain the mucosa injuries and related symptoms and findings without detected pepsin.

## Conclusion

The collection of several saliva pepsin samples improves the detection rate of LPR. The consideration of the highest concentration of multiple saliva pepsin collections was associated with the highest detection rate, and sensitivity. In case of high clinical LPR suspicion and negative pepsin test, a HEMII-pH study could provide further diagnostic information. Future studies are needed to confirm the most adequate number and time of saliva sample collection for the pepsin measurement.

## Abbreviations

GI	Gastrointestinal
HEMII-pH	Hypopharyngeal-esophageal multichannel intraluminal impedance-pH monitoring
LPR	Laryngopharyngeal reflux
N/PPV	Negative/positive predictive value

RSA Reflux sign assessment  
RSS-12 Reflux symptom score-12

### Acknowledgements

B. Johnson, native US speaker for the proofread.

### Author contributions

JRL: design, acquisition of data, data analysis & interpretation, drafting, final approval, and accountability for the work; final approval of the version to be published; agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. FB: design, acquisition of data, data analysis & interpretation, drafting, final approval, and accountability for the work; final approval of the version to be published; agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

### Funding

Vesale Grant (No. 2019) & IRIS-Recherche Grant (Foundation Roi Baudouin). The study is funded by the Groupement de Coopération Sanitaire ELSAN, 58 bis rue de la Boétie, 75008 Paris, France.

### Availability of data and materials

Data are available on request.

### Declarations

#### Ethics approval and consent to participate

The local ethics committee approved the study protocol (CHU Saint-Pierre, n°BE076201837630). Patients consented to participate.

#### Competing interests

Authors have no conflict of interest.

Received: 29 January 2023 Accepted: 27 September 2023  
Published online: 04 October 2023

### References

1. Lechien JR, Akst LM, Hamdan AL, et al. Evaluation and management of laryngopharyngeal reflux disease: state of the art review. *Otolaryngol Head Neck Surg.* 2019;160(5):762–82. <https://doi.org/10.1177/0194599819827488>.
2. Klimara MJ, Johnston N, Samuels TL, Visotcky AM, Poetker DM, Loehrl TA, Blumin JH, Bock JM. Correlation of salivary and nasal lavage pepsin with MII-pH testing. *Laryngoscope.* 2020;130(4):961–6. <https://doi.org/10.1002/lary.28182>.
3. Lechien JR, Bobin F, Dapri G, Eisendrath P, Salem C, Mouawad F, Horoi M, Thill MP, Dequanter D, Rodriguez A, Muls V, Saussez S. Hypopharyngeal–esophageal impedance-pH monitoring profiles of laryngopharyngeal reflux patients. *Laryngoscope.* 2021;131(2):268–76. <https://doi.org/10.1002/lary.28736>.
4. Hirano I. Review article: modern technology in the diagnosis of gastro-oesophageal reflux disease—Bilitec, intraluminal impedance and Bravo capsule pH monitoring. *Aliment Pharmacol Ther.* 2006;23(Suppl 1):12–24. <https://doi.org/10.1111/j.1365-2036.2006.02800.x>.
5. Lechien JR, Allen J, Mouawad F, et al. Do laryngologists and general otolaryngologists manage laryngopharyngeal reflux differently? *Laryngoscope.* 2020. <https://doi.org/10.1002/lary.28484>.
6. Hayat JO, Gabieta-Somnez S, Yazaki E, Kang JY, Woodcock A, Dettmar P, Mabary J, Knowles CH, Sifrim D. Pepsin in saliva for the diagnosis of gastro-oesophageal reflux disease. *Gut.* 2015;64(3):373–80. <https://doi.org/10.1136/gutjnl-2014-307049>.
7. Zhang M, Chia C, Stanley C, Phyland DJ, Paddle PM. Diagnostic utility of salivary pepsin as compared with 24-hour dual pH/impedance probe in laryngopharyngeal reflux. *Otolaryngol Head Neck Surg.* 2021;164(2):375–80. <https://doi.org/10.1177/0194599820951183>.
8. Calvo-Henríquez C, Ruano-Ravina A, Vaamonde P, Martínez-Capoccioni G, Martín-Martín C. Is pepsin a reliable marker of laryngopharyngeal reflux? A systematic review. *Otolaryngol Head Neck Surg.* 2017;157(3):385–91. <https://doi.org/10.1177/0194599817709430>.
9. Zeleník K, Hránková V, Vrtková A, Staníková L, Komínek P, Formánek M. Diagnostic value of the Peptest™ in detecting laryngopharyngeal reflux. *J Clin Med.* 2021;10(13):2996. <https://doi.org/10.3390/jcm10132996>.
10. Wang J, Li J, Nie Q, Zhang R. Are multiple tests necessary for salivary pepsin detection in the diagnosis of laryngopharyngeal reflux? *Otolaryngol Head Neck Surg.* 2022;166(3):477–81. <https://doi.org/10.1177/01945998211026837>.
11. Lechien JR, Chan WW, Akst LM, Hoppo T, Jobe BA, Chiesa-Estomba CM, Muls V, Bobin F, Saussez S, Carroll TL, Vaezi MF, Bock JM. Normative ambulatory reflux monitoring metrics for laryngopharyngeal reflux: a systematic review of 720 healthy individuals. *Otolaryngol Head Neck Surg.* 2022;166(5):802–19. <https://doi.org/10.1177/01945998211029831>.
12. Bobin F, Journe F, Lechien JR. Saliva pepsin level of laryngopharyngeal reflux patients is not correlated with reflux episodes. *Laryngoscope.* 2019. <https://doi.org/10.1002/lary.28260>.
13. Lechien JR, Bobin F, Rodriguez A, Dequanter D, Muls V, Huet K, Harmegnies B, Crevier-Buchman L, Hans S, Saussez S, Carroll TL. Development and validation of the short version of the reflux symptom score: reflux symptom score-12. *Otolaryngol Head Neck Surg.* 2021;164(1):166–74. <https://doi.org/10.1177/0194599820941003>.
14. Lechien JR, Bobin F, Muls V, et al. Validity and reliability of the reflux sign assessment (RSA). *Ann Otol Rhinol Laryngol.* 2019. <https://doi.org/10.1177/0003489419888947>.
15. Na SY, Kwon OE, Lee YC, Eun YG. Optimal timing of saliva collection to detect pepsin in patients with laryngopharyngeal reflux. *Laryngoscope.* 2016;126(12):2770–3. <https://doi.org/10.1002/lary.26018>.
16. Lechien JR, Bobin F, Muls V, Horoi M, Thill MP, Dequanter D, Finck C, Rodriguez A, Saussez S. Saliva pepsin concentration of laryngopharyngeal reflux patients is influenced by meals consumed before the samples. *Laryngoscope.* 2021;131(2):350–9. <https://doi.org/10.1002/lary.28756>.
17. Lechien JR, Bobin F, Muls V, Horoi M, Thill MP, Dequanter D, Rodriguez A, Saussez S. Patients with acid, high-fat and low-protein diet have higher laryngopharyngeal reflux episodes at the impedance-pH monitoring. *Eur Arch Otorhinolaryngol.* 2020;277(2):511–20. <https://doi.org/10.1007/s00405-019-05711-2>.
18. Luo Q, Zhan W, Boom RM, Janssen AEM. Interactions between acid and proteins under in vitro gastric condition—a theoretical and experimental quantification. *Food Funct.* 2018;9(10):5283–9. <https://doi.org/10.1039/c8fo01033a>.
19. Guldvog I, Linnestad P, Schrupf E, Berstad A. Dopaminergic and adrenergic influence on gastric acid and pepsin secretion stimulated by food. The role of vagal innervation. *Scand J Gastroenterol.* 1984;Suppl 89:113–6.
20. Wang AM, Wang G, Huang N, Zheng YY, Yang F, Qiu X, Chen XM. Association between laryngopharyngeal reflux disease and autonomic nerve dysfunction. *Eur Arch Otorhinolaryngol.* 2019;276(8):2283–7. <https://doi.org/10.1007/s00405-019-05482-w>.
21. Huang WJ, Shu CH, Chou KT, Wang YF, Hsu YB, Ho CY, Lan MY. Evaluating the autonomic nervous system in patients with laryngopharyngeal reflux. *Otolaryngol Head Neck Surg.* 2013;148(6):997–1002. <https://doi.org/10.1177/0194599813482103>.
22. Johnston N, Wells CW, Samuels TL, Blumin JH. Pepsin in nonacidic refluxate can damage hypopharyngeal epithelial cells. *Ann Otol Rhinol Laryngol.* 2009;118(9):677–85. <https://doi.org/10.1177/00034894091800913>.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.