



# **Optogenetic Globus Pallidus Stimulation Improves Motor Deficits in 6-Hydroxydopamine-Lesioned Mouse Model of Parkinson's Disease**

Sonia Di Bisceglie Caballero, Aurelia Ces, Martine Liberge, Frederic Ambroggi, Marianne Amalric, Abdel- Mouttalib Ouagazzal

## **► To cite this version:**

Sonia Di Bisceglie Caballero, Aurelia Ces, Martine Liberge, Frederic Ambroggi, Marianne Amalric, et al.. Optogenetic Globus Pallidus Stimulation Improves Motor Deficits in 6-Hydroxydopamine-Lesioned Mouse Model of Parkinson's Disease. International Journal of Molecular Sciences, 2023, 24 (9), pp.7935. <10.3390/ijms24097935>. <hal-04296470>

**HAL Id: hal-04296470**

**<https://hal.science/hal-04296470v1>**

Submitted on 20 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



Type of the Paper: Article

# Contrasting behavioral effects of optogenetic globus pallidus modulations in normal and hemi-parkinsonian mice

Sonia Di Bisceglie Caballero<sup>1</sup>, Aurelia Ces<sup>1</sup>, Martine Liberge<sup>1</sup>, Frederic Ambroggi<sup>1</sup>, Marianne Amalric<sup>1</sup>, Abdel-Mouttalib Ouagazzal<sup>1</sup>, \*

<sup>1</sup>Aix Marseille Université, CNRS, LNC (UMR 729), Marseille, France

\* Correspondence: [abdel-mouttalib.ouagazzal@univ-amu.fr](mailto:abdel-mouttalib.ouagazzal@univ-amu.fr)

**Abstract:** Excessive inhibition of the external globus pallidus (GPe) by striatal GABAergic neurons is considered as a central mechanism contributing to motor symptoms of Parkinson's disease (PD). While electrophysiological findings support this view, behavioral studies assessing beneficial effects of global GPe activations are scarce and the reported results are controversial. We used an optogenetic approach and the standard unilateral 6-hydroxydopamine nigrostriatal dopamine (DA) lesion model of PD to explore the effects of GPe photostimulation on motor deficits in mice. Global optogenetic GPe inhibition was used in normal mice to verify whether it reproduced the typical motor impairment induced by DA lesion. GPe activation improved ipsilateral circling, contralateral forelimb akinesia, locomotor hypoactivity and bradykinesia in hemi-parkinsonian mice at ineffective photostimulation parameters in normal mice. GPe photoinhibition had no effect on locomotor activity and forelimb use in normal mice but reduced directed exploration and improved working memory performances indicating that recruitment of GPe in physiological conditions may depend on the behavioral task involved. Collectively, these findings suggest that GPe plays more important role in PD as consequence of the complex functional compensatory remodeling affecting BG circuits upon DA depletion.

**Keywords:** external globus pallidus, motor behavior, mice, optogenetic, Parkinson's disease.

## 1. Introduction

Basal ganglia (BG) are a highly organized network of subcortical nuclei that controls various aspect of voluntary motor movements and were implicated in many neurodegenerative disorders, including Parkinson's disease (PD)<sup>1,2</sup>. The external globus pallidus (GPe) was traditionally considered as a simple relay nucleus within the network, connecting the striatum and subthalamic nucleus (STN) in the indirect pathway<sup>2,3</sup>. However, recent anatomical and electrophysiological studies uncovered a diversity of GABAergic neurons and anatomical connectivity suggesting that the GPe may serve a much more complex function than just a relay station<sup>4–6</sup>. GABAergic neurons of GPe comprise distinct neuronal subpopulations that can be identified based on their electrophysiological properties, molecular features, and projection targets. The two major classes are prototypic neurons that send widespread projections within and outside the BG and arky pallidal neurons that project back to the striatum<sup>4–8</sup>. Despite surge of interest over the past years, our understanding of the functional importance of GPe within the BG circuits remains incomplete. Early behavioral studies showed that pharmacological manipulations with GABAergic and glutamatergic agents that enhance GPe activity promote motor behaviors<sup>9</sup>. Unilateral GPe activations induce contralateral circling in rats and dyskinesia in monkeys while bilateral stimulations produce locomotor hyperactivity in rats<sup>10–16</sup>. Consistent with pharmacological studies, unilateral optogenetic GPe activation was also reported to induce contralateral circling and hyperkinesia in mice<sup>17</sup>. While these findings consistently show that the net effect of global GPe activation is a facilitation of motor behaviors, studies assessing the impact of cell-type specific optogenetic stimulations revealed a dissociable contribution of GABAergic neuron subtypes to motor behavior<sup>18–21</sup>.

**Citation:** To be added by editorial staff during production.

Academic Editor: Firstname  
Lastname

Received: date

Accepted: date

Published: date

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors.

Submitted for possible open access publication under the terms and conditions of the Creative Commons

Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

In PD condition, the degeneration of nigrostriatal dopaminergic neurons triggers a range of functional compensatory changes in BG circuits that lead to development of motor symptoms<sup>22</sup>. Excessive inhibition of GPe neurons by striatal GABAergic neurons of the indirect pathway is considered a critical mechanism underlying the pathological overactivity of the subthalamic nucleus and BG output structures and thus the expression of the motor symptoms<sup>1,22,623</sup>. While electrophysiological and neurochemical studies in patients and animal models of PD provide support for this view<sup>24–28</sup>, few studies have explored behavioral effects of global GPe stimulations in the context of dopamine (DA) depletion and the reported results are controversial. Pharmacological and global chemogenetic GPe activations ameliorate motor deficits in reserpine-treated rat and in mice with bilateral medial forebrain bundle (MFB) lesion, respectively<sup>29–31</sup>. By contrast, no effect was found with either global optogenetic GPe activation or inhibition in MFB-lesioned mice<sup>32</sup>. Whether the discrepancy is due to the functional heterogeneity of GABAergic projection neurons and/or the difference in neuromodulation approaches used is still an open question. To date, it remains also unclear whether GPe inhibition in physiological conditions produces motor deficits like dopaminergic lesion. While pharmacological inhibitions of GPe produce motor impairments in rats<sup>10,11,33,34</sup>, recent optogenetic studies targeting striatal indirect pathway and GPe found no effects in mice<sup>35,36</sup>, thus raising a question about GPe participation to locomotor behavior in physiological conditions.

In the present study we used an optogenetic approach and the standard unilateral 6-hydroxydopamine (6-OHDA) nigrostriatal dopamine (DA) lesion model in mice to reevaluate the effects of global GPe stimulation on motor deficits. The behavioral effects of optogenetic inhibition of GPe were also studied in normal mice under the same testing conditions to examine whether reduction of GPe activity reproduces the typical motor deficits of DA lesion. The findings show that global GPe photostimulation improves a range of motor deficits in hemi-parkinsonian mice. In normal mice, GPe photoinhibition had a little effect on motor behavior but promoted cognitive functions. These findings shed a new light into GPe function and show that this BG nucleus differentially contribute to modulation of motor functions in normal and PD conditions.


## 2. Results

### 2.1. GPe photostimulation improves ipsilateral circling behavior and contralateral forelimb akinesia in 6-OHDA-lesioned mice

To examine whether GPe activation improves motor impairments induced by unilateral DA lesion, the excitatory channelrhodopsin 2 (ChR2) variant, ChR2(H134R), which has a larger photocurrent than wildtype ChR2<sup>37</sup>, was expressed in GPe neurons under the control of the human synapsin-1 (hSyn1) promoter using an adeno-associated virus (AAV) vector (**Figures 1A**). Only mice with successful optic fiber implantation (**Figure 1B**) and opsin expression in the GPe (**Figure 1C**) were included in the study. Immunohistochemical staining of tyrosine hydroxylase (TH) confirmed the complete loss of dopaminergic fibers in the striatum of lesioned mice (**Figure 1D**), and quantitative analysis of TH immunofluorescence in the substantia nigra pars compacta (SNc) revealed over 98% loss of TH positive cells in 6-OHDA/TAG and 6-OHDA/ChR2 mice compared to Sham/controls ( $p \leq 0.005$  for both groups, Dunn's post-hoc test following a significant Kruskal-Wallis test, **Figures 1E** and **F**).

To reveal potential alterations of GPe activity, we selected an optical stimulation (532 nm, 5 Hz, 5 ms, 3 mW, 2 min application, **Figure 1G**) that was behaviorally ineffective in normal mice behavior based on pilot studies (data not shown). The illumination parameters triggered no contralateral circling (null scores) and produced no change in

spontaneous locomotor activity and forelimb use in non-lesioned control mice expressing the opsin (Sham/ChR2) or the fluorescent protein alone (Sham/TAG From **Supplementary Figure 1**). The two control groups were therefore pooled together in this study (Sham/controls, **Figure 1**). During the pre-stimulation period (first OFF, **Figure 1H**), 6-OHDA/TAG and 6-OHDA/ChR2 mice had a comparable ipsilateral circling scores that were significantly higher than those of Sham/controls ( $p \leq 0.02$ , non-parametric Dunn's post-hoc test). The photostimulation had no effect in 6-OHDA/TAG mice ( $p < 0.0005$  for all ON periods vs Sham/controls, Dunn's post-hoc test), but it reduced ipsilateral circling in 6-OHDA/ChR2 mice ( $p > 0.05$  vs Sham/controls, Dunn's post-hoc test, **Figure 1H**). The improvement immediately vanished when the optical stimulation was turned OFF. Analysis of cumulative circling scores over the OFF and ON periods revealed a significant main effect of lesion ( $H(3) \leq 16.58$  for both conditions,  $p < 0.0005$ , Kruskal-Wallis test; **Figure 1I**) and post-hoc test confirmed a beneficial effect of the photostimulation only in 6-OHDA/ChR2 mice ( $p > 0.05$  vs Sham/controls, Dunn's post-hoc test).




**Figure 1: GPe photostimulation ameliorates circling behavior and forelimb akinesia in hemiparkinsonian mice.** (A) Schematic illustration of the experimental procedure for opsin expression in GPe and nigrostriatal DA lesion. (B) Placement of individual optical fiber across the antero-posterior axis of the GPe. (C) Immunofluorescence image illustrating unilateral eYFP expression in GPe neurons. (D) DAB immunostaining for TH-positive fibers in the striatum of a lesioned mouse. (E) Immunostaining for TH+ cells in SNc of a lesioned mouse. (F) Percentage of TH+ cells loss in SNc in lesioned mice relative to Sham mice. (G) Photostimulation protocol used for the open field test. Testing was carried over 8 min with repeated sequences of light pulses stimulation (532 nm, 5 Hz, 5 ms, 3 mW) turned OFF and ON for 2 min. (H-I) Time course and total number of spontaneous ipsilateral rotations during OFF and ON periods, respectively. (J) Photostimulation protocol used for the cylinder test. Testing was carried over 6 min with repeated sequences of light pulses stimulation (532 nm, 5 Hz, 5 ms, 3 mW) turned ON and OFF for 1 min. (K) Percentage of contralateral

teral forelimb use during OFF and ON periods relative to the total number of forepaw contacts. (L) Total number of rears (forepaw contacts) during OFF and ON periods. Results are presented as mean  $\pm$  SEM. \* $p < 0.05$  and \*\* $p < 0.01$  significantly different from Sham mice. # $p < 0.05$  significantly different from 6-OHDA/TAG mice, Dunn's test following a significant non-parametric Kruskal-Wallis test for both circling behavior and forelimb use; Tukey's test following a significant two-way repeated measures ANOVA for rears.

We next verified whether GPe activation could restore contralateral forelimb akinesia in the cylinder test. Forelimb use in the cylinder test is associated with postural support when the mouse is up on its hind limbs and is a reliable index of akinesia induced by dopaminergic depletion. **Figure 1K** illustrates the global score of contralateral forelimb use in OFF and ON periods. A significant impairment was detected in all lesioned groups during the OFF periods ( $p \leq 0.01$  vs Sham/controls, Dunn's post-hoc test). Application of the optical stimulation had no effect in 6-OHDA/TAG mice ( $p < 0.001$ , Dunn's post-hoc test), but improved contralateral forelimb use deficit in 6-OHDA/ChR2 mice ( $p > 0.05$ , Dunn's post-hoc test). Total number of rearing was significantly reduced in both lesioned groups (main effect of 6-OHDA lesion,  $F_{(2, 18)} = 11.18$ ,  $p < 0.001$ , **Figure 1L**) and GPe photostimulation was ineffective ( $p < 0.05$  for both lesioned groups vs Sham/controls, Tukey's post-hoc test, **Figure 1L**).

## 2.2. GPe photostimulation improves locomotor hypoactivity and bradykinesia in 6-OHDA-lesioned mice

To further characterize the beneficial effects of GPe activation in lesioned mice we analyzed various locomotor activity parameters. **Figure 2B** illustrates the time-course of the distance traveled in the open field arena upon repeated application of the photostimulation (532 nm, 5 Hz, 5 ms, 3 mW, **Figure 2A**). In the pre-stimulation period, 6-OHDA/TAG and 6-OHDA/ChR2 mice had a significantly lower baseline locomotor activity levels than Sham/controls ( $p < 0.01$  for both groups, Tukey's post-hoc test). GPe photostimulation increased locomotor activity in 6-OHDA/ChR2 mice ( $p > 0.05$  vs Sham/controls, Tukey's post-hoc test) but not in 6-OHDA/TAG mice ( $p < 0.05$  vs Sham/controls, Tukey's post-hoc test). Repeated measures ANOVA performed on global scores (**Figure 2C**) yielded a significant main effect of 6-OHDA lesion ( $F_{(2, 19)} = 6.12$ ,  $p < 0.05$ ) and optical stimulation ( $F_{(1, 19)} = 7.16$ ,  $p < 0.01$ ). Post-hoc analysis confirmed that GPe photostimulation had a beneficial effect only in 6-OHDA/ChR2 mice ( $p > 0.05$  vs Sham/controls, Tukey's post-hoc test).



**Figure 2:** GPe photostimulation ameliorates locomotor activity measures in hemiparkinsonian mice. (A) Photostimulation protocol used for the open field test. Testing was carried over 8 min with repeated sequences of light pulses stimulation (532 nm, 5 Hz, 5 ms, 3 mW) turned OFF and ON for 2 min. (B–C) Time course and total distance traveled (cm) during OFF and ON periods, respectively. (D–E) Time course and total time spent in immobility (%) during OFF and ON periods, respectively. (F–G) Time course and global locomotor velocity during OFF and ON periods, respectively. Results are presented as mean  $\pm$  SEM. \* $p < 0.05$  and \*\* $p < 0.01$  significantly different from Sham mice or within group, Tukey's test following a significant two-way repeated measures ANOVA.

Hypokinesia assessed by the percentage of time spent in immobility (Figure 2D) was overall higher in lesioned groups compared to Sham/controls ( $F_{(2, 19)} = 2.97$ ,  $p < 0.05$ , 6-OHDA lesion effect). Though, a significant difference between 6-OHDA/TAG mice and Sham controls was only detected at the 2<sup>nd</sup> OFF period ( $p < 0.05$ , Tukey's post-hoc test). The photostimulation reduced % of the time spent in immobility in 6-OHDA/ChR2 in a reversible manner ( $F_{(6, 57)} = 2.44$ ,  $p < 0.05$ , 6-OHDA lesion  $\times$  photostimulation interaction, Figure 2D). Analysis of the global scores (Figure 2E) yielded a significant main effect of optical stimulation ( $F_{(1, 19)} = 5.06$ ,  $p < 0.05$ ) and a significant 6-OHDA lesion  $\times$  optical stimulation interaction ( $F_{(2, 19)} = 4.54$ ,  $p < 0.05$ , Figure 2E), but post-hoc test failed to reveal a significant difference between groups ( $p > 0.05$ , Tukey's test). Further analysis performed




within groups confirmed that only 6-OHDA/ChR2 mice had a significantly lower scores in the ON than the OFF period ( $p < 0.05$ , Student's *t* test, **Figure 2E**).

Lesioned mice also displayed a marked bradykinesia illustrated by a slower locomotor compared to Sham controls ( $F_{(2, 19)} = 6.12$ ,  $p < 0.01$ , 6-OHDA lesion effect, **Figure 2F**). The average velocity in the OFF periods was around 8 cm/s for Sham/controls and below 5 cm/s for lesioned mice ( $p \leq 0.04$ , Tukey's post-hoc test). Application of the photostimulation enhanced locomotor velocity in 6-OHDA/ChR2 ( $F_{(3, 57)} = 2.26$ ,  $p < 0.05$ ) at all ON periods ( $p > 0.05$  vs Sham/controls, Tukey's post-hoc test, **Figure 2F**). Analysis of the global scores (**Figure 2G**) revealed a significant main effect of 6-OHDA lesion ( $F_{(2, 19)} = 5.27$ ,  $p < 0.05$ ) and optical stimulation ( $F_{(1, 19)} = 8.83$ ,  $p < 0.01$ ) and post-hoc multiple comparisons confirmed an improvement of bradykinesia only in 6-OHDA/ChR2 ( $p > 0.05$  vs Sham/controls, Tukey's post-hoc test, **Figure 2G**).

### 2.3. GPe photoinhibition in normal mice does not mimic motor deficits of DA depletion

We next examined whether optogenetic inhibition of GPe in normal mice could reproduce the typical motor deficits of unilateral 6-OHDA lesion. To do so, we used the engineered blue-sensitive chloride-conducting channelrhodopsin, iC++, which allows more efficient optogenetic control than the hyperpolarization induced by light-activated chloride pumps<sup>38</sup>. iC++ was expressed in GPe under the control of the hSyn1 promoter (**Figure 3A**). Only mice with successful opsin expression (**Figure 3B**) and optic fiber implantation within the GPe (**Figures 3C and D**) were included in the study. We first examined whether unilateral GPe photoinhibition induces ipsilateral circling. Various photoinhibition protocols were used with pulsed and continuous illuminations (**Figure 3E**) at different frequencies (20–60 Hz) and intensities (5–12 mW) based on published studies<sup>38,39</sup>. Only a tendency to circle toward the illumination side (few incomplete ipsilateral rotations) was detected in iC++ mice. From **Figures 3F and G**, it can be seen that continuous photoinhibition with the highest illumination intensity (12 mW) produces no notable circling behavior ( $p > 0.05$ , Mann-Whitney test). Analysis of locomotor activity measures revealed no changes in distance traveled ( $p > 0.05$ , repeated measures ANOVA, **Supplementary Figures 2B and C**), % of the time spent in immobility ( $p > 0.05$ , repeated measures ANOVA, **Supplementary Figures 2D and E**) and velocity ( $p > 0.05$ , repeated measures ANOVA, **Supplementary Figures 2F and G**). In the cylinder test, the unilateral photoinhibition (12 mW, **Figure 3H**) produced no impairment in contralateral forelimb use ( $p > 0.05$ , Student's *t*-test, **Figures 3I**) and rearing behavior ( $p > 0.05$  vs TAG mice, Student's *t*-test, **Figures 3J**).




**Figure 3: Unilateral GPe photoinhibition does not reproduce motor deficits.** (A) Schematic illustration of the experimental procedure for opsin expression in GPe. (B) Immunofluorescence image illustrating bilateral viral expression in GPe neurons. (C) Cresyl violet staining showing bilateral fiber optic placement in the GPe. (D) Placement of individual optical fiber across antero-posterior GPe axis. (E) Photoinhibition protocol used in the open field test. Testing was carried over 10 min with continuous light illumination (450 nm, 12 mW). (F) Time course and total number of ipsilateral rotations over 10 min of testing, respectively. (G) Photoinhibition protocol used in the cylinder test. Testing was carried over 6 min with continuous light illumination (450 nm, 12 mW). (I) Percentage of contralateral forelimb use relative to the total number of forepaw contacts. (J) Total number of rears (forepaw contacts). Results are presented as mean  $\pm$  SEM.  $p > 0.05$ , Mann-Whitney test or unpaired t-test following a significant non-parametric Kruskal-Wallis test for circling behavior and forelimb akinesia, respectively.

#### 2.4. GPe photoinhibition improves cognitive functions in normal mice

The above findings indicate that GPe may not contribute to modulation of spontaneous locomotor behavior in physiological conditions. GPe establishes direct anatomical connections with the cortex, thalamus, and amygdala<sup>19,40–43</sup> suggesting that it may participate to modulation of motivational and cognitive aspects of motor behavior. To explore this possibility, we examined whether bilateral GPe photoinhibition (6 mW/side) could impact short-term spatial working memory performance assessed by spontaneous alternations in the Y-maze task (**Figures 4A and B**). The latency to exit the starting arm and the number of arm entries were used as an index of neophobia and general locomotor activity, respectively. TAG and iC++ mice, displayed good spontaneous alternation per-



formances that were above chance level (calculated at 22.2%,  $p < 0.05$ , One-group test **Figure 4B**). Interestingly, iC++ mice had a better score than TAG mice ( $p < 0.05$ , Student's t-test) indicating that the photoinhibition promotes short-term working memory. The photoinhibition did not change the latency to exit the starting arm ( $p < 0.05$  vs TAG mice, Student's t-test, **Figure 4C**) or the number of arm entries ( $p < 0.05$  vs TAG mice, Student's t-test, **Figures 4D and E**).



**Figure 4: Bilateral GPe photoinhibition impacts working memory and directed exploration.** (A) Photoinhibition protocol used in the Y-maze test. Testing was carried over 6 min with continuous light stimulation (450 nm, 6 mW/side). (B) Percentage of spontaneous alternations. The dashed line shows a chance level of 22.2%. (C) Latency to exit the starting arm (sec). (D-E) Time course and total number of arm entries in the Y-maze over 6 min of testing, respectively. (F) Photoinhibition protocol used in the nose-poke habituation test. Two 15-min of testing were carried one week apart, session 1 with continuous light illumination (450 nm, 6 mW/side) and session 2 with light OFF. (G-H) Time course and total number of nose-pokes per session. Results are presented as mean  $\pm$  SEM. \* $p < 0.05$ , unpaired t-test for Y-maze test and unpaired t-test following a significant repeated measures ANOVA for nose-poke exploration.

The effect of the photoinhibition was also studied on the long-term habituation to spatial cues, a simple form of non-associative learning, using one week of retention delay. Testing was conducted over 15 min in operant chambers equipped with illuminated nose-poke ports placed on the front wall. We previously demonstrated that nose-poke ports can be employed as spatial cues for evaluating directed exploration and learning performances in mice<sup>44–46</sup>. Mice received bilateral optical inhibition (6 mW/side) solely on the first session (**Figure 4F**). TAG mice displayed the expected exploration profile (**Fig-**

ure 4G). On session 1, they initially displayed a high number of nose-poking that declined over the course of the session and then remained at the same low level on the second testing session, reflecting respectively short-term (within-session) and long-term (between-session) habituation phenomenon ( $F_{(9, 81)} = 5.062$ ,  $p < 0.0001$ , main effect across days). iC++ mice had a remarkably low exploration level on session 1 compared to TAG mice ( $F_{(9, 81)} = 2.574$ ,  $p < 0.05$ , sessions  $\times$  optical inhibition interaction) and a significant difference between groups was detected in the first block ( $p < 0.05$ , Student's t-test). On session 2, they displayed a normal habituation performance illustrated by a low exploration level comparable to TAG mice ( $p > 0.05$ , Student's t-test for all blocks). Analysis of the global scores confirmed that iC++ mice had a significantly lower number of nose-poking than TAG mice only in session 1 ( $p < 0.05$ , Student's t-test, Figure 4H) indicating that they were able to efficiently process and retain novel spatial information despite their low exploration level.

### 3. Discussion

The present study shows that global GPe optogenetic activation is effective at reducing motor impairments in hemi-parkinsonian mice. As expected, unilateral SNc lesion produced the typical motor deficits in mice manifested by ipsilateral circling, locomotor hypoactivity, bradykinesia, and forelimb akinesia. Global GPe optogenetic activation improved the various motor abnormalities. The beneficial effects were reversible and only seen in 6-OHDA-lesioned mice expressing the excitatory opsin, confirming the selectivity of the optogenetic neuromodulation. The improvement of locomotor hypoactivity may be due to restoration of ipsilateral circling, and thus the balance in the motor circuit, which is essential to walk in a straight line and to travel over long distances in the arena. It may also be attributed to restoration of the exploration drive as dopaminergic lesions are well known to reduce novelty-induced exploration. The amelioration of hypokinesia and bradykinesia produced by GPe photostimulation is consistent with this conclusion. The photostimulation was also efficient at improving akinesia assessed by contralateral forelimb use in the cylinder test. Overall, our findings extend those reported in the literature using pharmacological and chemogenetic approaches<sup>29–31</sup>. However, they contrast with those of Mastro et al.<sup>32</sup> showing that cell-specific rather than global GPe optogenetic manipulations are effective in restoring motor deficits in MFB-lesioned mice, a PD model of severe DA deficiency. The discrepancy between the two studies may be explained by the differences in photostimulation protocols and PD mouse models used. The unilateral intranigral lesion model we used produces a less severe DA depletion and motor impairments than the bilateral MFB lesion that targets all midbrain DA neurons. It is therefore possible that global GPe photostimulation may more readily alleviate motor impairments induced by unilateral SNc than bilateral MFB lesion. It should be stressed that in Mastro et al. study the behavioral effects of GPe photostimulation were assessed 3–5 days post-lesion while in our study they were evaluated at a much later time point (5–6 weeks post-lesion). Such a difference in post-lesion testing time may be another possible source of the discrepant results considering that BG circuitry undergoes major functional changes several weeks after DA depletion<sup>3</sup>. An interesting study will be to assess the beneficial effects of global GPe photostimulations in hemi-parkinsonian mice at different time-points post-lesion.

Numerous electrophysiological studies demonstrated a reduction of GPe neuronal firings in animal models and PD patients<sup>3,24,26,28</sup>. GPe hypoactivity has been attributed to an abnormal elevation of extracellular GABA concentrations caused by the overactivity of the striato-pallidal pathway<sup>23,25,47</sup> and to local changes at the level of pallidal GABAergic synapses<sup>48,49</sup>. Our study provides new behavioral evidence that GPe activity is altered upon DA depletion. We show that an ineffective optical stimulation in normal

mice is sufficient for improving many aspect of motor impairments caused by DA lesion. Here, we selected low optical stimulation parameters to reveal possible functional alterations of the GPe, and we do expect that a full restoration of the motor deficits can be achieved using higher illumination frequencies and/or intensities. The greater sensitivity of hemi-parkinsonian mice to behavioral effects of the photostimulation may also reflect a functional hypersensitivity of GPe GABAergic neurons that develops following DA lesion. For instance, the number of GABAergic synapses and the expression of GABA<sub>A</sub> receptors are increased in the STN upon DA depletion leading to a strengthening of the functional connectivity between GPe and STN<sup>50–52</sup>. Hence, the motor improvement seen in lesioned mice may be attributed to restoration of the inhibitory drive of GPe prototypic neurons expressing parvalbumine (PV-neurons), which are the main source of GPe inputs to STN and BG output structures (substantia nigra pars reticulata and entopeduncular nucleus homologous to primates globus pallidus interna)<sup>4</sup>. Selective optogenetic stimulations of PV-neurons was consistently shown to alleviate motor deficits in mouse models of PD<sup>32,53,54</sup>. Interestingly, we found that a behaviorally ineffective optical stimulation in normal mice restores to normal level the motor deficits in lesioned mice<sup>54</sup>. It is worth noting that not all GPe neurons promote motor behaviors. Optogenetic silencing of prototypic neurons expressing Lim homeobox 6 (Lhx6-neurons) improves motor deficits in MFB-lesioned mice indicating that this neuronal subpopulation plays a deleterious role in the context of DA depletion<sup>32</sup>. Optical stimulations of arky pallidal neurons, namely FoxP2-expressing neurons that exclusively innervate the striatum, were also found to suppress locomotion in normal mice<sup>21,55</sup>. The fact that global GPe photostimulation produces a beneficial effect in hemi-parkinsonian mice implies that the recruitment of PV-neurons overrides deleterious influences of other GPe neuronal populations that suppress motor behavior, an action that may be exerted through a local collateral inhibition and suppression of downstream BG nuclei as recently demonstrated<sup>21</sup>.

The series of experiments conducted in normal mice show that mere reduction of GPe activity did not produce an impairment of spontaneous locomotor behavior and forelimb use, like DA lesion did. It is unlikely that the illumination parameters used are insufficient for silencing GPe neurons. We tested a range of photoinhibition protocols that were previously demonstrated to be effective for silencing neurons and triggering behavioral responses in normal mice<sup>39,56</sup>. Our results are in a good agreement with a recent study of Isett et al.<sup>36</sup> demonstrating a little effect of optogenetic inhibitions of GPe and striatal indirect pathway on locomotor activity in mice. They also extend those reported in rats and monkeys showing that neurotoxic lesions of GPe do not reproduce changes in BG discharge patterns and motor dysfunction observed in DA-depleted states<sup>28,57</sup>. Collectively, these findings suggest that GPe contribution to gross motor function may not be critical in physiological conditions. Though, in a parkinsonian patient, a lesion confined to the GPe exacerbates akinesia<sup>58</sup>. GPe ablation was also reported to worsen motor deficits in parkinsonian monkeys and to abolish the improvement induced by the dopaminergic receptor agonist, apomorphine<sup>59</sup>. It thus appears that in DA-depleted state, GPe may play a more important role in the development of motor deficits. In PD, the degeneration of DA neurons also leads to a hypoactivity of the direct striatal pathway that promotes motor behaviors. It is therefore possible that the deleterious impact of reduced GPe activity on motor function may be exacerbated when direct striatal pathway transmission is concomitantly reduced. Another possibility could be that the inhibitory drive of some GPe GABAergic neurons that suppress motor behavior is enhanced upon DA depletion. As mentioned above, optical inhibition of GPe Lhx6-neurons improves motor deficits in MFB-lesioned mice<sup>32</sup>. On the other hand, striatal inputs of Npas1<sup>+</sup>-expressing neurons, which suppress locomotor behavior, have been shown to be strengthened upon DA lesion<sup>55</sup>. Finally, it should be noted that motor symptoms of PD were linked not only to changes in local firing rates but also to an ab-

normal rhythmic activity in BG and cortex manifested by an enhanced oscillations in the beta frequency range (13–30 Hz)<sup>60,61</sup>. Pathological beta-oscillations are thought to compromise information flow and processing in the cortico-basal ganglia (CBC) network, thereby leading to the impairment of motor functions. Recent preclinical studies showed that GPe plays a central role in this context by promoting the expression and propagation of abnormal beta-oscillations throughout the CBG network<sup>62–66</sup>. A such capability of orchestrating changes in the pattern of activity across the CBG network may thus emerge as a consequence of the compensatory alterations of GPe inhibitory drive onto BG nuclei and thalamus following DA neuron loss<sup>19,50,63</sup>. The contrasting effects of optogenetic GPe modulations on motor functions in normal and hemi-parkinsonian mice is in line with this conclusion.

An important finding was the improvement of cognitive performances produced by GPe inactivation in normal mice. In the Y-maze task, the photoinhibition induced a specific improvement of short-term working memory that was manifested by an increase of spontaneous alternations without any changes in locomotor exploration and emotionality. In the nose-poke habituation task, the promnesic effect was a less clear-cut because the photoinhibition reduced nose-poking behavior. Despite the exploration deficit, iC++ mice displayed a normal long-term habituation performance like control mice, which also points to a cognitive facilitation. The reduction of nose-poking behavior induced by the photoinhibition contrasts with the null effects found in the open field and the Y-maze tests. In both cases, higher illumination intensities (8–12 vs 6 mW) produced no locomotor dysfunction in mice, supporting the view that directed exploration and locomotor activity are subserved by distinct neural substrates<sup>44–46</sup>. The low level of nose-poke exploration displayed by iC++ mice on session 1 may also be indicative of a promnesic action of the photoinhibition. An enhanced ability to process and retain spatial information should favor a rapid familiarization of mice with surrounding salient spatial cues and thus lead to a reduced investigatory activity. A such cognitive facilitation can explain the good working memory performance of iC++ mice revealed in the Y-maze task, which also requires spatial processing. The precise mechanisms underlying the inhibitory influence of GPe on cognitive processing remains to be delineated. GPe contains a diversity of GABAergic neurons that sends widespread projections within and outside BG and could each influence specific aspects of motor and cognitive behaviors. In support of this idea, it was recently shown that PV-neurons projecting to the SNr promote motor behavior while those innervating the parafascicular thalamus negatively modulate behavioral flexibility in mice<sup>19</sup>. It is therefore possible that the promnesic effects produced by the photoinhibition may in part be attributed to an inhibition of GPe inputs to the parafascicular thalamus. It may also be linked to a suppression of the GPe inhibitory drive onto the frontal cortex<sup>40</sup> that subserves executive functions. Future studies are necessary to elucidate the specific contribution of the GPe pathways to the different facets of cognitive behaviors.

In conclusion, our findings provide a new insight into the functional role of GPe in health and disease state. They suggest that GPe modulation of gross motor function may not be critical in physiological conditions as previously thought. This BG nucleus may rather participate to specific aspects of motor and cognitive behaviors. In PD condition, the GPe may however play a central role in the expression of a range motor signs of PD because of the complex functional compensatory remodeling affecting BG circuits upon DA depletion.

## 4. Materials and Methods

### 4.1. Subjects

Eight weeks old C57BL/6J (BL6) male mice were purchased from Charles River Laboratories (Saint Germain Nuelles, France). Mice were housed in groups of 3-4 in individually ventilated cages (Techniplast, Grostenquin, France) and kept in 12 h light/dark cycle (light on at 07:00am, off at 07:00pm) with water and food ad lib. Behavioral testing was performed during the light cycle between 07:00am and 07:00pm. All experimental procedures were approved by the French national ethics committee (CE071) and conducted in accordance with EEC (2010/63/UE) guidelines for care and use of laboratory animals.

#### 4.2. Viral constructions

For global optical stimulation of GPe neurons the excitatory channelrhodopsin 2 variant (ChR2(H134)) fused to a yellow fluorescent protein (eYFP) was expressed under the hSYN1 promoter using an adeno-associated virus serotype 2 vector (AAV2-hSYN1-ChR2(H134R)-eYFP,  $5.6 \times 10^{12}$  particles/ml). For optical inhibition we used the AAV2 vector encoding the engineered chloride channel, IC++, fused to eYFP under the hSYN1 promoter (AAV2-hSYN1-IC++-eYFP,  $4.1 \times 10^{12}$  particles/ml). AAV2 encoding the eYFP reporter protein only (AAV2-hSyn1-eYFP,  $2 \times 10^{12}$  particles/ml) was used as a control. All viral vectors were obtained from Vector Core (University of North Carolina, Chapel Hill, USA and MTA from Karl Deisseroth at Stanford University).

#### 4.3. Stereotaxic surgery

##### 4.3.1. Unilateral 6-OHDA lesion

Mice were pre-treated subcutaneously with buprenorphine (0.05mg/kg) for analgesia, then deeply anesthetized with isoflurane (4% induction then 1-2%) and placed in a stereotaxic frame (David Kopf instruments) on a heated pad and OcryGel was used to protect from dry eyes. Lidocaine (6mg/kg) was injected subcutaneously before incising the scalp. Mice received 1.5  $\mu$ l infusion of 6-OHDA hydrobromide (2.7  $\mu$ g/ $\mu$ l, free base, Tocris) either into the left or right SNc over 5 min at the following coordinates: AP: 3 mm, ML  $\pm$  1.3mm and DV -4.7 mm, according to the atlas of Paxinos and Watson, (2001). The injection side was randomized between experimental groups. The microinjector was slowly removed 5 min after the end of the injection to minimize backflow. Control mice (Sham) received an injection of the corresponding volume of vehicle (0.02% ascorbic acid in NaCl) in the same conditions.

##### 4.3.2. Viral infusion and fiber optic implantation

This procedure was performed either immediately after 6-OHDA lesion for studying the effects of GPe optostimulation in PD mice or in separate surgery session in normal mice for assessing behavioral effects of optogenetic manipulations of GPe activity in a baseline condition. For GPe photostimulation in PD mice or in non-lesioned control mice (Sham), 300 nl of AAV2-hSyn1-ChR2(H134R)-eYFP or AAV2-hSyn1-eYFP was unilaterally infused into the GPe over 6 min (AP: -0.3 mm, ML:  $\pm$  2.0 mm and DV: -3.8 mm from the skull surface, according to the atlas of Paxinos and Watson, 2001). The microinjector was slowly removed 6 min after the end of the injection. Following viral injection, an optical fiber-ferrule (optical fiber 200  $\mu$ m diameter, 0.22 N.A., Doric lenses, Quebec, Canada) was unilaterally implanted 0.2 mm above the injection site and cemented to the skull with Super-Bond C&B. For GPe photoinhibition in a normal condition, normal mice received bilateral infusion of 300 nl of AAV2-hSyn1-IC++-eYFP or AAV2-hSyn1-eYFP in GPe over 6 min at the same stereotaxic coordinates as above. Optical fiber-ferrules were bilaterally implanted 0.2 mm above the injection site and cemented to the skull with Super-Bond C&B.

Animals received a subcutaneous injection of carprofen (3mg/kg, diluted in NaCl 0.9% and glucose 5%) immediately after surgery and then for 2 consecutive post-operative days. They were kept for 2-3 weeks in the post-operative room and their weight monitored daily. Lesioned mice received supplemental food (diet gel, Nutella, chocolate puffed cereals) and additional NaCl 0.9% injections if needed until complete recovery.

#### 4.4. Optogenetic manipulations

Mice were connected through an optical patch cable (Doric lenses, Quebec, Canada) coupled either to a green DPSS laser (532 nm, LaserGlow Technologies, Toronto, Canada) or a blue LD Fiber Light Source (450 nm, LDFS, Doric Lenses, Quebec, Canada) via a rotary joint (Doric Lenses, Quebec, Canada). The light pulses were triggered by a programmable TTL Pulse Train Generators (custom made). The optical light power was calibrated before behavioral testing by measuring the output power at the tip of an optical fiber-ferrule using a light power meter (PM100D, Thorlabs, France).

#### 4.5. Behavioral Procedures

Behavioral testing was carried 5-6 weeks after the surgery. All experiments were performed on 20-22 weeks old mice. Mice were handled and habituated to the optical connection few days before behavioral testing. Pilot tests were carried in a viewing jar (20 cm diameter and 30 cm height Plexiglas cylinder) to establish the adequate optical stimulation parameters (light pulse frequency, duration, and intensity) before behavioral testing. For optogenetic studies in 6-OHDA-lesioned mice, a green wavelength light (DPSS laser) was used for photostimulation because the delivery of the blue LDFS was delayed due to the Covid19 pandemic, and pilot tests carried in the viewing jar showed that it is effective in normal mice (Figure S3 for behavioral comparison conducted afterwards with green DPSS laser and blue LDFS, see also<sup>56</sup>). For GPe photostimulation study, mice were first submitted to the open field test for assessing circling behavior and changes in spontaneous locomotor activity then the cylinder test for assessing forelimb use impairment and postural asymmetry. For GPe photoinhibition study, mice were tested in the following order: open field test, then the nose-poke habituation test session 1, Y-maze test, nose-poke habituation test session 2 and the cylinder test.

##### 4.5.1. Cylinder test

Mice were placed in a small Plexiglas cylinder (10 cm diameter, 20 cm height) and forepaw contacts performed against the wall during vertical exploration (rearing) were manually scored using an ethological keyboard (LabWatcher, ViewPoint, France). For studying the effects of GPe optostimulation in PD mice testing was conducted as follow: after 30 sec of familiarization, mice received light pulses for 1 min (ON period) followed by no stimulation for 1 min (OFF period), repeated three times. For GPe photoinhibition in normal mice, the illumination was applied during the whole testing session (6 min). Mice failing to reach 3 contacts during the 3 min OFF or ON period were excluded from the analyses. One mouse from the 6-OHDA/ChR2 group and one from the iC++ group were excluded. Data are expressed as a percentage of contralateral forepaw contacts relative to the total number of contacts.

##### 4.5.2. Open field test



Testing was carried in a white PVC arena (50 x 50 x 45 cm) and spontaneous locomotor activity was recorded by a video-tracking system (ViewPoint, France). Each mouse was individually placed facing a corner of the arena and allowed to freely explore the apparatus for 8 min. For GPe photostimulation in PD mice, light pulses were delivered as follows: 2-min OFF periods (baseline locomotor activity level) and 2-min ON periods, repeated twice to assess whether the behavioral effect of the photostimulation was reversible. For GPe photoinhibition in normal mice, a repeated 2-min and continuous illumination protocols were used. In each period, distance traveled, velocity, % of time spent in immobility (movement threshold < 1cm/s) were analyzed. Circling behaviors was hand-scored in parallel.

#### 4.5.3. Y-maze test

Mice were placed in a Y-maze made of grey Plexiglas with 3 identical arms (40 x 9 x 16 cm) placed at 120° from each other. Each arm had walls with specific motifs allowing to distinguish it from the others. Mice freely explored the maze for 6 min with light ON. Spontaneous alternation performance was assessed by visually recording for each mouse the pattern of entrance into each arm in the maze. Alternations were operationally defined as successive entries into each of the three arms as on overlapping triplet sets. Percent spontaneous alternation performance was defined as the ratio of actual (total alternations) to possible alternations (total arm entries - 2) x 100. Total entries were scored as an index of exploratory activity in the maze. Latency to exit the starting arm was used to assess neophobia.

#### 4.5.4. Nose-poke habituation test

Mice were tested in five-choice operant chambers (Coulbourn Instruments, Allentown, United State) dimly lit with a permanent house-light. The front panel was curved and composed of five bays filled with metal wall panels interchangeable by nose-poke modules (Model H21-10M). Each nose-poke hole was equipped with a controlled yellow LED cue light at the end and infrared photobeam across the opening that detects the nose-pokes. Two adjacent nose-poke units (spaced 4 cm apart) were presented in the right corner and two others in the left corner of the front wall (no nose-poke unit in central bay). The back wall was composed of a single bay fitted with metal panels and the plexiglass side walls were completely covered by cardboard with distinguishable geometrical motifs. The metal stainless-steel rod floor (the grid shock floor provided by the manufacturer) was covered by a grey vinyl-coated paper that was used as the standard flooring throughout the study. Testing consisted of 15 min long sessions with nose-poke holes illuminated. The bilateral photoinhibition (continuous illumination) was applied only the first day. One week retention delay was used to assess the impact of GPe inhibition on long-term habituation performances of mice. The number of nose poking was monitored and used as an index of directed exploration and habituation capacity to salient spatial cues.

### 4.6. Immunofluorescence and histological analysis

#### 4.6.1. Analysis of 6-OHDA-induced dopamine cell loss

Mice were perfused transcardially with 4% paraformaldehyde and the brains were removed, postfixed, and cryoprotected. Free floating coronal sections (40 µm thick) containing the SNc were incubated with rabbit anti-TH antibody (1/1000, Synaptic Systems, 213 102) overnight at 4°C followed by a 2h incubation with the Alexa Fluor 594 goat anti-rabbit antibody (1/400, Jackson Laboratory, 111-585-003) at room temperature (RT). Sections were mounted onto SuperFrost Plus glass slides (VWR) and coverslipped with Ro-

ti®-Mount FluorCare mounting medium (Carl Roth). Immunofluorescence was analyzed by a laser confocal microscopy (Zeiss LM710, Germany) at a 20 X magnification. TH positive cells were counted through the SNc region based on the stereotaxic mouse atlas (Paxinos & Franklin, 2001) delineating the SNc from the ventral tegmental area by the medial optic tract. Counting was done manually in five regions of interest (ROIs) per hemisphere, covering the whole medio-lateral extension of the structure (-2.70 to -3.80 mm relative to bregma, 3 sections per mouse), using cell counter plugin of FIJI software (ImageJ, National Institutes of Health).

DA depletion in the striatum was also examined. Coronal sections (40µm) were incubated overnight at 4 °C with a mouse anti-TH antibody (1/1000, Millipore, MAB318). Thereafter, they were incubated with a biotinylated secondary antibody (goat anti-mouse, 1/200; Jackson ImmunoResearch; 115-065-003) for 1 hour and then in a solution containing 0.01% DAB (3,3-diaminobenzidine) and 3% H<sub>2</sub>O<sub>2</sub> diluted in PBS 1X for 3 min. Sections were finally mounted on slides as described before. Images were acquired using a bright field microscope (Nikon Leica DMLB) at 10X magnification.

#### 4.6.2. Analysis of opsins expression

Free floating coronal sections (40 µm thick) containing the GPe were incubated a 48h with a rabbit anti-GFP antibody (1/500, Invitrogen, A-11122) at 4°C then 1h30 with Alexa Fluor 488 donkey anti-rabbit antibody (1/500, Invitrogen, R37118) at RT. Sections were mounted onto SuperFrost Plus glass slides (VWR) and coverslipped with Roti®-Mount FluorCare mounting medium (Carl Roth). Immunofluorescence was analyzed by a laser confocal microscopy (Zeiss LM710, Germany) at 10 X magnification.

#### 4.6.3. Verification of optic fiber-ferrule placement:

For confirming the placement of the optical fiber in the GPe, coronal sections of the GPe were stained with cresyl violet. Images were acquired using Leica DM LB Microscope at 2.5 X magnification equipped with Nikon digital camera DXM1200. Mice with the optical fiber-ferrule placements outside of the GPe were excluded from analysis.

#### 4.7. Statistical analysis

All data are expressed as mean group value ± standard error of the mean (s.e.m.). Analysis of forelimb akinesia and rears data in the cylinder test were carried out using a non-parametric Kruskal-Wallis test followed by Dunn's post-hoc test and a two-way repeated measures ANOVA followed by unpaired t-test, respectively. Analysis of circling behavior was performed using Kruskal-Wallis test followed by Dunn's post-hoc test or using an Mann-Whitney test. Locomotor activity data in the open field test were analyzed using two-way ANOVA with repeated measures (OFF and ON periods or time). When relevant, post-hoc comparisons were carried using Tukey's multiple comparisons tests or t-tests, respectively. The accepted level of significance was  $p < 0.05$ .

**Supplementary Materials:** The following supporting information can be downloaded at: [www.mdpi.com/xxx/s1](http://www.mdpi.com/xxx/s1),

Figure S1: Behavioral effect of GPe photostimulation in Sham controls groups

Figure S2: Unilateral GPe photoinhibition has no impact on locomotor activity measures in normal mice;

Figure S3: Similar behavioral effects of GPe photostimulation with green or blue light in normal mice ;

**Author Contributions:** Conceptualization, AM. O., M. A. and F. A.; methodology, AM. O., M. L., A. C. and S. DBC.; formal analysis, AM. O., M. A., M. L., A. C. and S. DBC.; investigation, AM. O., M. L., A. C. and S. DBC.; resources, AM. O., M. A. and F. A.; writing—review and editing, AM. O., M. A. and F. A., M. L. and S. DBC.; supervision, AM. O., M. A. and M. L.; project administration, AM. O. and M. A.; funding acquisition, AM. O., M. A. and F. A. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Centre National de la Recherche Scientifique (CNRS), Aix-Marseille University (AMU), European Union’s Horizon 2020 Research and Innovation Program (FETOPEN, Grant # 767092) and the Association France Parkinson (AFP).

**Institutional Review Board Statement:** The animal study protocol was approved by the French national ethics committee in animal experimentation (CEEA071) and the Ministère de l’Enseignement Supérieur, de la Recherche et de l’Innovation (authorization no 24894) and conducted in accordance with EEC (2010/63/UE) guidelines for care and use of laboratory animals.

**Data Availability Statement:** The data reported in this paper will be shared by the lead contact upon reasonable request.

**Acknowledgments:** Sonia Di Bisceglie Caballero was supported by French Ministry of Higher Education and Research and Aix-Marseille University Neuroschool. We would like to thank, E. Mansour, JL Fina and J. Espejo for animal care, D. Louber for technical assistance, the histology platform headed by N. Lorenzo and A. Tonetto for assistance in confocal microscopy.

**Conflicts of Interest:** All authors declare no biomedical financial interest or potential conflicts of interest.

## References

1. Lanciego, J.L., Luquin, N., and Obeso, J.A. (2012). Functional neuroanatomy of the basal ganglia. *Cold Spring Harb. Perspect. Med.* *2*, 743–746. 10.1101/cshperspect.a009621.
2. Alexander, G.E., DeLong, M.R., and Strick, P.L. (1986). Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu. Rev. Neurosci.* *VOL. 9*, 357–381. 10.1146/annurev.ne.09.030186.002041.
3. Albin, R.L., Young, A.B., and Penney, J.B. (1989). The functional anatomy of basal ganglia disorders. *Trends Neurosci.* *12*, 366–375. 10.1016/0166-2236(89)90074-X.
4. Hegeman, D.J., Hong, E.S., Hernández, V.M., and Chan, C.S. (2016). The external globus pallidus: Progress and perspectives. *Eur. J. Neurosci.* *43*, 1239–1265. 10.1111/ejn.13196.
5. Gittis, A.H., Berke, J.D., Bevan, M.D., Chan, C.S., Mallet, N., Morrow, M.M., and Schmidt, R. (2014). New roles for the external globus pallidus in basal ganglia circuits and behavior. *J. Neurosci.* *34*, 15178–15183. 10.1523/JNEUROSCI.3252-14.2014.
6. Dong, J., Hawes, S., Wu, J., Le, W., and Cai, H. (2021). Connectivity and Functionality of the Globus Pallidus Externa Under Normal Conditions and Parkinson’s Disease. *Front. Neural Circuits* *15*, 1–19. 10.3389/fncir.2021.645287.
7. Jaeger, D., and Kita, H. (2011). Functional connectivity and integrative properties of globus pallidus neurons. *Neuroscience* *198*, 44–53. 10.1016/j.neuroscience.2011.07.050.
8. Mallet, N., Micklem, B.R., Henny, P., Brown, M.T., Williams, C., Bolam, J.P., Nakamura, K.C., and Magill, P.J. (2012). Dichotomous Organization of the External Globus Pallidus. *Neuron* *74*, 1075–1086. 10.1016/j.neuron.2012.04.027.
9. Amalric, M., and Koob, G.F. (1989). Dorsal pallidum as a functional motor output of the corpus striatum. *Brain Res.* *483*, 389–394. 10.1016/0006-8993(89)90186-8.
10. Ossowska, K., Wędzony, K., and Wolfarth, S. (1984). The role of the GABA mechanisms of the globus pallidus in mediating catalepsy, stereotypy and locomotor activity. *Pharmacol. Biochem. Behav.* *21*, 825–831. 10.1016/S0091-3057(84)80060-X.
11. Chen, L., Chan, S.C.Y., and Yung, W.H. (2002). Rotational behavior and electrophysiological effects induced by GABAB

- receptor activation in rat globus pallidus. *Neuroscience* *114*, 417–425. 10.1016/S0306-4522(02)00299-3.
12. Wisniecki, A., Correa, M., Arizzi, M.N., Ishiwari, K., and Salamone, J.D. (2003). Motor effects of GABAA antagonism in globus pallidus: Studies of locomotion and tremulous jaw movements in rats. *Psychopharmacology (Berl.)* *170*, 140–149. 10.1007/s00213-003-1521-z.
  13. Grabli, D., McCairn, K., Hirsch, E.C., Agid, Y., Féger, J., François, C., and Tremblay, L. (2004). Behavioural disorders induced by external globus pallidus dysfunction in primates: I. Behavioural study. *Brain* *127*, 2039–2054. 10.1093/brain/awh220.
  14. Xue, Y., Han, X.H., and Chen, L. (2010). Effects of pharmacological block of GABAA receptors on pallidal neurons in normal and parkinsonian state. *Front. Cell. Neurosci.* *4*, 1–10. 10.3389/neuro.03.002.2010.
  15. Ikeda, H., Kotani, A., Koshikawa, N., and Cools, A.R. (2010). Differential role of GABAA and GABAB receptors in two distinct output stations of the rat striatum: Studies on the substantia nigra pars reticulata and the globus pallidus. *Neuroscience* *167*, 31–39. 10.1016/j.neuroscience.2010.01.054.
  16. Crossman, A.R., Mitchell, I.J., Sambrook, M.A., and Jackson, A. (1988). Chorea and myoclonus in the monkey induced by gamma-aminobutyric acid antagonism in the lentiform complex. The site of drug action and a hypothesis for the neural mechanisms of chorea. *Brain* *111* ( Pt 5), 1211–1233. 10.1093/brain/111.5.1211.
  17. Tian, J., Yan, Y., Xi, W., Zhou, R., Lou, H., Duan, S., Chen, J.F., and Zhang, B. (2018). Optogenetic stimulation of GABAergic neurons in the globus pallidus produces hyperkinesia. *Front. Behav. Neurosci.* *12*, 1–11. 10.3389/fnbeh.2018.00185.
  18. Cui, Q., Pamukcu, A., Cherian, S., Chang, I.Y.M., Berceau, B.L., Xenias, H.S., Higgs, M.H., Rajamanickam, S., Chen, Y., Du, X., et al. (2021). Dissociable roles of pallidal neuron subtypes in regulating motor patterns. *J. Neurosci.* *41*, 4036–4059. 10.1523/JNEUROSCI.2210-20.2021.
  19. Lilascharoen, V., Wang, E.H.J., Do, N., Pate, S.C., Tran, A.N., Yoon, C.D., Choi, J.H., Wang, X.Y., Pribiag, H., Park, Y.G., et al. (2021). Divergent pallidal pathways underlying distinct Parkinsonian behavioral deficits. *Nat. Neurosci.* *24*, 504–515. 10.1038/s41593-021-00810-y.
  20. Hernández, V.M., Hegeman, D.J., Cui, Q., Kever, D.A., Fiske, M.P., Glajch, K.E., Pitt, J.E., Huang, T.Y., Justice, N.J., and Savio Chan, C. (2015). Parvalbumin + neurons and Npas1 + neurons are distinct neuron classes in the mouse external globus pallidus. *J. Neurosci.* *35*, 11830–11847. 10.1523/JNEUROSCI.4672-14.2015.
  21. Aristieta, A., Barresi, M., Azizpour Lindi, S., Barrière, G., Courtand, G., de la Crompe, B., Guilhemsang, L., Gauthier, S., Fioramonti, S., Baufreton, J., et al. (2021). A Disynaptic Circuit in the Globus Pallidus Controls Locomotion Inhibition. *Curr. Biol.* *31*, 707–721.e7. 10.1016/j.cub.2020.11.019.
  22. McGregor, M.M., and Nelson, A.B. (2019). Circuit Mechanisms of Parkinson's Disease. *Neuron* *101*, 1042–1056. 10.1016/j.neuron.2019.03.004.
  23. O'Connor, W.T. (1998). Functional neuroanatomy of the basal ganglia as studied by dual-probe microdialysis. *Nucl. Med. Biol.* *25*, 743–746. 10.1016/S0969-8051(98)00066-3.
  24. Fillion, M., and Tremblay, L. (1991). Abnormal spontaneous activity of globus pallidus neurons in monkeys with MPTP-induced parkinsonism. *Brain Res.* *547*, 140–144. 10.1016/0006-8993(91)90585-J.
  25. Bianchi, L., Galeffi, F., Bolam, J.P., and Della Corte, L. (2003). The effect of 6-hydroxydopamine lesions on the release of amino acids in the direct and indirect pathways of the basal ganglia: A dual microdialysis probe analysis. *Eur. J. Neurosci.* *18*, 856–868. 10.1046/j.1460-9568.2003.02795.x.
  26. Starr, P.A., Rau, G.M., Davis, V., Marks, W.J., Ostrem, J.L., Simmons, D., Lindsey, N., and Turner, R.S. (2005). Spontaneous pallidal neuronal activity in human dystonia: Comparison with Parkinson's disease and normal macaque. *J. Neurophysiol.* *93*, 3165–3176. 10.1152/jn.00971.2004.
  27. Chazalon, M., Paredes-Rodriguez, E., Morin, S., Martinez, A., Cristóvão-Ferreira, S., Vaz, S., Sebastiao, A., Panatier, A., Boué-Grabot, E., Miguelez, C., et al. (2018). GAT-3 Dysfunction Generates Tonic Inhibition in External Globus Pallidus

Neurons in Parkinsonian Rodents. *Cell Rep.* **23**, 1678–1690. 10.1016/j.celrep.2018.04.014.

28. Soares, J., Kliem, M.A., Betarbet, R., Greenamyre, J.T., Yamamoto, B., and Wichmann, T. (2004). Role of external pallidal segment in primate parkinsonism: Comparison of the effects of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced parkinsonism and lesions of the external pallidal segment. *J. Neurosci.* **24**, 6417–6426. 10.1523/JNEUROSCI.0836-04.2004.

29. Maneuf, Y.P., Mitchell, I.J., Crossman, A.R., and Brotchie, J.M. (1994). On the Role of Enkephalin Cotransmission in the GABAergic Striatal Efferents to the Globus Pallidus. *Exp. Neurol.* **125**, 65–71. 10.1006/exnr.1994.1007.

30. Liu, J., Shelkar, G.P., Sarode, L.P., Gawande, D.Y., Zhao, F., Clausen, R.P., Ugale, R.R., and Dravid, S.M. (2021). Facilitation of GluN2C-containing NMDA receptors in the external globus pallidus increases firing of fast spiking neurons and improves motor function in a hemiparkinsonian mouse model. *Neurobiol. Dis.* **150**, 105254. 10.1016/j.nbd.2021.105254.

31. Assaf, F., and Schiller, Y. (2019). A chemogenetic approach for treating experimental Parkinson's disease. *Mov. Disord.* **34**, 469–479. 10.1002/mds.27554.

32. Mastro, K.J., Zitelli, K.T., Willard, A.M., Leblanc, K.H., Kravitz, A. V., and Gittis, A.H. (2017). Cell-specific pallidal intervention induces long-lasting motor recovery in dopamine-depleted mice. *Nat. Neurosci.* **20**, 815–823. 10.1038/nn.4559.

33. Chen, L., Chan, C.S., and Yung, W.H. (2004). Electrophysiological and behavioral effects of zolpidem in rat globus pallidus. *Exp. Neurol.* **186**, 212–220. 10.1016/j.expneurol.2003.11.003.

34. Turski, L., Klockgether, T., Turski, W.A., Schwarz, M., and Sontag, K.H. (1990). Blockade of excitatory neurotransmission in the globus pallidus induces rigidity and akinesia in the rat: implications for excitatory neurotransmission in pathogenesis of Parkinson's diseases. *Brain Res.* **512**, 125–131. 10.1016/0006-8993(90)91180-O.

35. Bolkan, S.S., Stone, I.R., Pinto, L., Ashwood, Z.C., Iravedra Garcia, J.M., Herman, A.L., Singh, P., Bandi, A., Cox, J., Zimmerman, C.A., et al. (2022). Opponent control of behavior by dorsomedial striatal pathways depends on task demands and internal state (Springer US) 10.1038/s41593-022-01021-9.

36. Isett, B.R., Nguyen, K.P., Schwenk, J.C., Snyder, C.N., Adegbesan, K.A., Ziausyte, U., and Gittis, A.H. (2022). The Indirect Pathway of the Basal Ganglia Promotes Negative Reinforcement, But Not Motor Suppression. *bioRxiv*, 2022.05.18.492478.

37. Lerner, T.N., Ye, L., and Deisseroth, K. (2016). Communication in Neural Circuits: Tools, Opportunities, and Challenges. *Cell* **164**, 1136–1150. 10.1016/j.cell.2016.02.027.

38. Berndt, A., Lee, S.Y., Wietek, J., Ramakrishnan, C., Steinberg, E.E., Rashid, A.J., Kim, H., Park, S., Santoro, A., Frankland, P.W., et al. (2016). Structural foundations of optogenetics: Determinants of channelrhodopsin ion selectivity. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 822–829. 10.1073/pnas.1523341113.

39. Park, S., Kramer, E.E., Mercaldo, V., Rashid, A.J., Insel, N., Frankland, P.W., and Josselyn, S.A. (2016). Neuronal Allocation to a Hippocampal Engram. *Neuropsychopharmacology* **41**, 2987–2993. 10.1038/npp.2016.73.

40. Saunders, A., Oldenburg, I.A., Berezovskii, V.K., Johnson, C.A., Kingery, N.D., Elliott, H.L., Xie, T., Gerfen, C.R., and Sabatini, B.L. (2015). A direct GABAergic output from the basal ganglia to frontal cortex. *Nature* **521**, 85–89. 10.1038/nature14179.

41. Karube, F., Takahashi, S., Kobayashi, K., and Fujiyama, F. (2019). Motor cortex can directly drive the globus pallidus neurons in a projection neuron type dependent manner in rat. *bioRxiv*, 1–25. 10.1101/677377.

42. Giovanniello, J., Yu, K., Furlan, A., Nachtrab, G.T., Sharma, R., Chen, X., and Li, B. (2020). A central amygdala-globus pallidus circuit conveys unconditioned stimulus-related information and controls fear learning. *J. Neurosci.* **40**, 9043–9054. 10.1523/JNEUROSCI.2090-20.2020.

43. Hunt, A.J., Dasgupta, R., Rajamanickam, S., Jiang, Z., Beierlein, M., Chan, C.S., and Justice, N.J. (2018). Paraventricular hypothalamic and amygdalar CRF neurons synapse in the external globus pallidus. *Brain Struct. Funct.* **223**, 2685–2698. 10.1007/s00429-018-1652-y.

44. Reiss, D., Walter, O., Bourgoin, L., Kieffer, B.L., and Ouagazzal, A.-M. (2014). New automated procedure to assess context

- recognition memory in mice. *Psychopharmacology (Berl)*. 231, 4337–4347. 10.1007/s00213-014-3577-3.
45. Boccia, M.M., and Baratti, C.M. (1999). Effects of oxytocin and an oxytocin receptor antagonist on retention of a nose-poke habituation response in mice. *Acta Physiol. Pharmacol. Ther. Latinoam*. 49, 155–160.
46. Brodtkin, J. (1999). Assessing memory in mice using habituation of nose-poke responding. *Behav. Pharmacol.* 10, 445–451. 10.1097/00008877-199909000-00002.
47. Ochi, M., Koga, K., Kurokawa, M., Kase, H., Nakamura, J., and Kuwana, Y. (2000). Systemic administration of adenosine A(2A) receptor antagonist reverses increased GABA release in the globus pallidus of unilateral 6-hydroxydopamine-lesioned rats: A microdialysis study. *Neuroscience* 100, 53–62. 10.1016/S0306-4522(00)00250-5.
48. Chazalon, M., Paredes-Rodriguez, E., Morin, S., Martinez, A., Cristóvão-Ferreira, S., Vaz, S., Sebastiao, A., Panatier, A., Boué-Grabot, E., Miguelez, C., et al. (2018). GAT-3 Dysfunction Generates Tonic Inhibition in External Globus Pallidus Neurons in Parkinsonian Rodents. *Cell Rep.* 23, 1678–1690. 10.1016/j.celrep.2018.04.014.
49. Kimberly Johnson<sup>1, 2</sup>, Jessica Barragan<sup>1</sup>, Sarah Bashiruddin<sup>1</sup>, Cody J. Smith<sup>3</sup>, Chelsea Tyrrell<sup>4</sup>, Michael J. Parsons<sup>5</sup>, Rosemarie Doris<sup>6</sup>, Sarah Kucenas<sup>3</sup>, Gerald B. Downes<sup>7</sup>, Carla M. Velez<sup>1</sup>, Caitlin Schneider<sup>1</sup>, Catalina Sakai<sup>1</sup>, Narendra Pathak<sup>1</sup>, Katrina Anders, <sup>4</sup>, and <sup>1</sup>Department (2017). 乳鼠心肌提取 HHS Public Access. *Physiol. Behav.* 176, 139–148. 10.1038/nm.2692.HCN.
50. Fan, K.Y., Baufreton, J., Surmeier, D.J., Chan, C.S., and Bevan, M.D. (2012). Proliferation of external globus pallidus-subthalamic nucleus synapses following degeneration of midbrain dopamine neurons. *J. Neurosci.* 32, 13718–13728. 10.1523/JNEUROSCI.5750-11.2012.
51. Kovalski, R.F., Callahan, J.W., Chazalon, M., Wokosin, D.L., Baufreton, J., and Bevan, M.D. (2020). Dysregulation of external globus pallidus-subthalamic nucleus network dynamics in parkinsonian mice during cortical slow-wave activity and activation. *J. Physiol.* 598, 1897–1927. 10.1113/JP279232.
52. Shen, K.-Z., and Johnson, S.W. (2005). Dopamine depletion alters responses to glutamate and GABA in the rat subthalamic nucleus. *Neuroreport* 16, 171–174. 10.1097/00001756-200502080-00021.
53. Pamukcu, A., Cui, Q., Xenias, H.S., Berceau, B.L., Augustine, E.C., Fan, I., Chalasani, S., Hantman, A.W., Lerner, T.N., Boca, S.M., et al. (2020). Parvalbumin<sup>1</sup> and Npas<sup>11</sup> pallidal neurons have distinct circuit topology and function. *J. Neurosci.* 40, 7855–7876. 10.1523/JNEUROSCI.0361-20.2020.
54. Di Bisceglie Caballero, S., Ces, A., Ambroggi, F., Amalric, M., and Ouagazzal, A.-M. (2022). Contribution of external globus pallidus projection neurons to motor deficits of Parkinson's disease: optogenetic studies in murine model. In *Movement Disorders*, p. Abstract# 1541. 10.1002/mds.29223.
55. Glajch, K.E., Kolver, D.A., Hegeman, D.J., Cui, Q., Xenias, H.S., Augustine, E.C., Hernández, V.M., Verma, N., Huang, T.Y., Luo, M., et al. (2016). Npas<sup>1+</sup> pallidal neurons target striatal projection neurons. *J. Neurosci.* 36, 5472–5488. 10.1523/JNEUROSCI.1720-15.2016.
56. Lin, J.Y., Lin, M.Z., Steinbach, P., and Tsien, R.Y. (2009). Characterization of engineered channelrhodopsin variants with improved properties and kinetics. *Biophys. J.* 96, 1803–1814. 10.1016/j.bpj.2008.11.034.
57. Hassani, O.K., Mouroux, M., and Féger, J. (1996). Increased subthalamic neuronal activity after nigral dopaminergic lesion independent of disinhibition via the globus pallidus. *Neuroscience* 72, 105–115. 10.1016/0306-4522(95)00535-8.
58. Frcs, L.E.M., Mrcp, R.G., Frcpath, W.S., Radatz, M., Silburn, P., Scott, R., Aziz, T., Sn, F., and Stein, J.F. (1999). Clinical study Lateral pallidotomy e x a c e r b a t e s akinesia in the Parkinsonian patient. 6, 474–476.
59. Zhang, J., Russo, G.S., Mewes, K., Rye, D.B., and Vitek, J.L. (2006). Lesions in monkey globus pallidus externus exacerbate parkinsonian symptoms. *Exp. Neurol.* 199, 446–453. 10.1016/j.expneurol.2006.01.006.
60. Yu, Y., Sanabria, D.E., Wang, J., Hendrix, C.M., Zhang, J., Nebeck, S.D., Amundson, A.M., Busby, Z.B., Bauer, D.L., Johnson, M.D., et al. (2021). Parkinsonism alters beta burst dynamics across the basal ganglia-motor cortical network. *J. Neurosci.* 41,



2274–2286. 10.1523/JNEUROSCI.1591-20.2021.

61. Witcher, M., Moran, R., Tatter, S.B., and Laxton, A.W. (2014). Neuronal oscillations in Parkinson's disease. *Front. Biosci. (Landmark Ed. 19)*, 1291–1299. 10.2741/4282.

62. Mallet, N., Pogosyan, A., Márton, L.F., Bolam, J.P., Brown, P., and Magill, P.J. (2008). Parkinsonian beta oscillations in the external globus pallidus and their relationship with subthalamic nucleus activity. *J. Neurosci.* 28, 14245–14258. 10.1523/JNEUROSCI.4199-08.2008.

63. Crompe, B. de la, Aristieta, A., Leblois, A., Elsherbiny, S., Boraud, T., and Mallet, N.P. (2020). The globus pallidus orchestrates abnormal network dynamics in a model of Parkinsonism. *Nat. Commun.* 11, 1–14. 10.1038/s41467-020-15352-3.

64. Tachibana, Y., Iwamuro, H., Kita, H., Takada, M., and Nambu, A. (2011). Subthalamo-pallidal interactions underlying parkinsonian neuronal oscillations in the primate basal ganglia. *Eur. J. Neurosci.* 34, 1470–1484. 10.1111/j.1460-9568.2011.07865.x.

65. Chan, C.S., Surmeier, D.J., and Yung, W.H. (2006). Striatal information signaling and integration in globus pallidus: Timing matters. *NeuroSignals* 14, 281–289. 10.1159/000093043.

66. Schwab, B.C., Heida, T., Zhao, Y., Marani, E., van Gils, S.A., and van Wezel, R.J.A. (2013). Synchrony in Parkinson's disease: Importance of intrinsic properties of the external globus pallidus. *Front. Syst. Neurosci.* 7, 1–7. 10.3389/fnsys.2013.00060.