

# High-field immiscibility of electrons belonging to adjacent twinned bismuth crystals

Yuhao Ye, Akiyoshi Yamada, Yuto Kinoshita, Jinhua Wang, Pan Nie, Liangcai Xu, Huakun Zuo, Masashi Tokunaga, Neil Harrison, Ross D. Mcdonald, et al.

### ▶ To cite this version:

Yuhao Ye, Akiyoshi Yamada, Yuto Kinoshita, Jinhua Wang, Pan Nie, et al.. High-field immiscibility of electrons belonging to adjacent twinned bismuth crystals. 2023. hal-04297785

## HAL Id: hal-04297785 https://hal.science/hal-04297785

Preprint submitted on 21 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#### High-field immiscibility of electrons belonging to adjacent twinned bismuth crystals

Yuhao Ye<sup>1</sup>, Akiyoshi Yamada<sup>2,3</sup>, Yuto Kinoshita<sup>3</sup>, Jinhua Wang<sup>1</sup>, Pan Nie<sup>1</sup>, Liangcai Xu<sup>1</sup>, Huakun Zuo<sup>1</sup>, Masashi Tokunaga<sup>3</sup>, Neil Harrison<sup>4</sup>, Ross D. McDonald<sup>4</sup>, Alexey V. Suslov<sup>5</sup>, Arzhang Ardavan<sup>6</sup>, Moon-Sun

Nam<sup>6</sup>, David LeBoeuf<sup>7</sup>, Cyril Proust<sup>7</sup>, Benoît Fauqué<sup>8</sup>, Yuki Fuseya<sup>2</sup>, Zengwei Zhu<sup>1,\*</sup> and Kamran Behnia<sup>9</sup>

(1) Wuhan National High Magnetic Field Center and School of Physics,

Huazhong University of Science and Technology,

Wuhan 430074, China

(2) Department of Engineering Science,

University of Electro-Communications,

Chofu, Tokyo 182-8585, Japan

(3)Institute for Solid State Physics,

The University of Tokyo,

Kashiwa, Chiba 277-8581, Japan

(4) MS-E536, NHMFL, Los Alamos National Laboratory,

Los Alamos, New Mexico 87545, USA

(5)National High Magnetic Field Laboratory,

1800 E. Paul Dirac Drive,

Tallahassee, FL 32310, USA

(6) Department of Physics,

Clarendon Laboratory, University of Oxford,

Oxford OX1 3PU, United Kingdom

(7) Laboratoire National des Champs Magnétiques Intenses (LNCMI-EMFL),

CNRS, UGA, UPS, INSA, Grenoble/Toulouse, France

(8) JEIP, USR 3573 CNRS,

Collège de France, PSL Research University,

11, place Marcelin Berthelot,

75231 Paris Cedex 05, France

(9) Laboratoire Physique et Etude de Matériaux (CNRS-UPMC)

ESPCI Paris, PSL Research University, 75005 Paris, France

(Dated: October 11, 2023)

Bulk bismuth has a complex Landau spectrum. The small effective masses and the large gfactors are anisotropic. Moreover, at a high magnetic field, when only the lowest Landau levels remain occupied, the chemical potential does not stay constant. An added complexity arises from the existence of twin boundaries, which, by producing extra anomalies, further complexify the interpretation of the data in the extreme quantum limit. Here, we present an extensive study of low-temperature angle-dependence of magnetoresistance up to 65 T together with measurements of Nernst effect, ultrasound, and magneto-optics in bismuth. We found that all observed anomalies can be explained in a single-particle picture of a sample consisting of two twinned crystals tilted by 108°. We show that a quantitative agreement between theory and experiment can be achieved only if one assumes that the two adjacent twinned crystals keep their own chemical potentials at a high magnetic field, despite a shift between chemical potentials as large as 68 meV at 65 T. This implies the existence of an energy barrier between adjacent twinned crystals reminiscent of a Schottky barrier between a metal and a semiconductor. We argue that this barrier is built by accumulating charge carriers of opposite signs across a twin boundary.

#### I. INTRODUCTION

Bismuth occupies a distinct place in the history of solid state physics (For reviews, see [1-4]). Its tiny Fermi surface was the first to be experimentally mapped out thanks to the easily detectable quantum oscillations it generates [5–8]. Recent attention focuses on finding topologically non-trivial states both for the surfaces and the hinges of bismuth crystals [9–18]. Recent transport studies on macroscopic crystals have revealed a non-trivial variation of the magnetoresistance [19] and the Seebeck effect [20] with the amplitude and the orientation of the magnetic field. A robust boundary conductance emerges at quantizing magnetic field, which is yet to be understood [21]. At the atomic scale, a bismuth layer on a NbSe<sub>2</sub> substrate was found to be a novel platform for detecting nanometric Turing patterns [22].

The rhombohedral structure of crystalline bismuth at ambient pressure is the outcome of lowering the cubic symmetry by pulling the cube along its body diagonal axis. Since this can be achieved along each of the four body diagonals, twin formation is common in bismuth crystals [23, 24] and twin boundaries in bismuth crystals were directly observed decades ago with a scanning tunneling microscope [25, 26].

The uncontrolled presence of such twin boundaries

misled researchers seeking the signatures of collective electronic effects in bismuth. In the extreme quantum limit, when all carriers are confined to their lower Landau levels, a variety of electronic instabilities are conceivable [27, 28]. Experiments seeking such instabilities [29, 30] explored bismuth beyond its quantum limit, achievable by applying a magnetic field exceeding 9 T along the trigonal axis, and found anomalies which, at first sight, could not be explained in the single-particle picture. Taken at face value, they were attributed to instabilities caused by electronic interactions. However, these interpretations in the case of bismuth were rejected subsequently. The computed Landau spectrum [31-34] was found to be in agreement with the experimental data and the hysteresis attributed to a first-order phase transition [30] was not reproduced by torque magnetometry [35] or magnetostriction [36] measurements.

It was later found that the additional anomalies [29] were caused by the presence of a secondary twinned crystal tilted by  $108^{\circ}$  with respect to the main crystal [34]. Thus, up to 28 T, the single-particle picture proved to be sufficient to explain the experimental data [34]. The same theoretical frame was then employed to explain how the evacuation of one or two electron pockets by a 40 T magnetic field, applied along the binary or the bisectrix axes, generates a sudden drop in magnetoresistance [37] and an anomaly in magnetization [38].

The success of the single-particle picture in bismuth pushed beyond the quantum limit is to be contrasted with the case of graphite, another low-carrier semi-metal, in which the quantum limit is accessible in a threedimensional context. A succession of field-induced phase transitions have been detected in graphite [39–42] indicating the capacity of strong Coulomb interaction to generate collective states [27, 28] in this material. The two elemental semi-metals differ in several aspects. Screening is weaker in graphite because of its lower dielectric constant. Spin-orbit coupling is much stronger in bismuth. Finally, graphite is much more anisotropic than bismuth, and its electron and hole Fermi pockets' longer axis are aligned.

In the case of bismuth-antimony alloys, possible signatures of field-induced electronic instability were recently observed [43]. They are to be distinguished from highfield anomalies due to the presence of twin boundaries seen before [44].

Here, we present an extensive study of the Shubnikovde Haas effect, Nernst effect, ultrasound, and magnetooptics in bismuth at low temperatures and a magnetic field as strong as 65 T. We map the angle dependence of the Landau spectra for the whole solid angle and find that all anomalies can be explained in the single-particle picture, invoking the presence of a secondary twinned crystal. This includes the anomaly observed around 38 T when the magnetic field is applied along the trigonal axis of the main crystal [45], whose origin has not been hitherto identified. Thus, up to 65 T, there is no signature of instability caused by electron-electron interaction in





(a)

40

FIG. 1. Magnetoresistance data in the three planes of rotation: Magnetoresistance up to 65 T at 1.6 K for a magnetic field rotating in the (a) binary-trigonal, (b) bisectrixtrigonal, and (c) binary-bisectrix planes. The quantum limit of holes, when the penultimate hole Landau level is evacuated, occurs at 9 T when the magnetic field is along the trigonal axis and shifts upward as the field is tilted towards the binary or bisectrix axis.

bismuth. However, the quantitative agreement between theory and experiment is obtained thanks to a crucial assumption: that the two adjacent twinned crystals each have their own chemical potential  $\mu$ , set by the orientation of the magnetic field respective to their crystalline axes. We will show that without this assumption, there is no agreement between the computed and the measured Landau spectrum. This unavoidable assumption leads us to conclude that electrons residing in two adjacent crys-



FIG. 2. Experimental Landau spectrum for three rotating planes: (a)-(c). Nernst voltage  $(S_{xy})$  vs magnetic field measured in a magnetic field up to 12 T (in a superconducting magnet), up to 28 T in resistive water-cooled magnet (curves in blue) along with the second derivative of the magnetoresistance measured in pulsed field experiments above 28 T up to 65 T (curves in red) for the three planes of rotation: binary-trigonal, bisectrix-trigonal, and binary-bisectrix planes, respectively. The data have been normalized to their highest amplitude for each data set. The 0° and 90° represent the field along trigonal and binary/bisectrix for the (a)/(b). The 0° and 90° represent the field along binary and bisectrix for the (c). The sketches show the direction of the orientation of the magnetic field and electrical/heat current for each plane of rotation. Note that in all the experiments, the electrical or the heat current is along the direction perpendicular to the plane of rotation.

tals do not mix up, despite a 68 meV difference in their chemical potentials at 65 T. We argue that this hypothesis points to the existence of a new type of p-n junction at the twin boundary of this semi-metal.

#### II. EXPERIMENTAL

Bismuth single crystals commercially obtained used in this study had typical dimensions of  $1 \times 2 \times 0.5$ mm<sup>3</sup> and a Room-temperature-to-Residual-resistivity-Ratio (RRR) of  $\rho(300 \text{ K})/\rho(4.2 \text{ K})) \sim 30 - 100$ . Magnetoresistance (MR) measurements in pulsed magnetic field were carried out at Wuhan National High Magnetic Field Center (WHMFC) and in National High Magnetic Field Laboratory- Pulsed Field Facility (NHMFL-PFF) in Los Alamos. The samples were rotated in three distinct planes: binary-trigonal, bisectrix-trigonal and binarybisectrix. For each set of measurements, the current was applied along the axis perpendicular to the rotating plane. Our high-field MR data complemented the Nernst data obtained below 28 T and reported previously [33, 34]. Additional angle-dependent Nernst measurements up to 45 T were also carried out in the hybrid magnet of the NHMFL DC Facility in Tallahassee. Ultrasound measurements have been collected at Laboratoire National des Champs Magnétique Intense (LNCMI-Toulouse) with pulsed magnetic field up to 54 T. Longitudinal ultrasonic waves were generated using commercial LiNbO<sub>3</sub> 36° Y-cut transducers glued on a polished surface. A standard pulse-echo technique was used to determine the change in the sound velocity and attenuation. Magneto-optical measurements involving polarized light (for more details, see reference [46]) were conducted at the International MegaGauss Science Laboratory, the Institute for Solid State Physics at the University of Tokyo. The resulting image was captured using a high-speed camera.



FIG. 3. Consistency among different probes and samples: (a) Nernst voltage  $(S_{xy})$  measured in Grenoble up to 28 T (dashed lines) compared with the Nernst voltage measured up to 45 T (solid lines) for several orientations of the magnetic field. (b) Nernst voltage (blue) compared with the second derivative of the magnetoresistance (red), and the sound velocity (black) vs. magnetic field at 0° (c) Same for 15°. Peaks in  $S_{xy}$ , and  $\frac{d^2\rho}{dB^2}$  are concomitant with a minimum in  $\Delta v$ . The arrows indicate the ~40 T anomaly observed previously [45]. The measurements presented in all panels were performed in the bisectrix-trigonal plane and zero degree corresponds to field orientation along the trigonal axis.

#### III. RESULTS

Angle-dependent magnetoresistance in three planes: Fig.1 shows magnetoresistance along three planes, namely binary-trigonal (bin.-tri.), bisectrixtrigonal (bis.-tri.), and binary-bisectrix (bin.-bis.) at1.6 K. Magnetoresistance shows oscillations which correspond to evacuations of the Landau levels with magnetic field. The different orientations for these planes are indicated. When the magnetic field is along the trigonal axis, MR shows a minimum around 9 T, which corresponds to the evacuation of the penultimate Landau level of holelike carriers [47]. As the magnetic field is tilted off the trigonal axis, the cross-section of the hole Fermi pocket perpendicular to the magnetic field increases and this minimum shifts to higher fields. In the binary-bisectrix of Fig.1(c), the drops are evident around 40 T because of the emptying of Dirac pockets [37].

Fig. 2 presents the second derivative of our magnetoresistance next to the Nernst measurements  $(S_{xy})$  for three high-symmetric planes [34]. The two types of data are consistent with each other, but the sensitivity of the Nernst results is relatively higher. Each peak in the Nernst response or in the second derivative of magnetoresistance corresponds to an intersection between an electron (or a hole) Landau level and the chemical potential. Note that the data have been normalized to the largest amplitude of each data set. Fig.2 (a), (b), and (c) show the data for the magnetic field rotating in the binary-trigonal, bisectrix-trigonal, and in the binarybisectrix planes, respectively. In the corner of the panel, the schematic diagram shows the measurement configurations for MR. The quantum oscillations pattern is relatively simple and easily understood when the field is along the trigonal axis [48]. The hole Landau peaks can be easily identified as their amplitudes are the largest, except for a sudden jump around 40 T along the trigonal axis.

**Consistency of measurements:** We verified the consistency between diverse techniques when the field was rotated in the bisectrix-trigonal plane and the data obtained on different samples. Fig.3(a) shows a comparison of the Nernst data collected in Grenoble up to 28 T (dashed lines) with the data collected in Tallahassee up to 45 T (solid lines) for several orientations of the magnetic field. In various experiments, the peak positions are close to each other. Panels (b) and (c) compare three distinct sets of data obtained in different laboratories at different times and on different samples. The Nernst data,  $S_{xy}$ , the second derivative of magnetoresistance,  $\frac{d^2\rho}{dB^2}$ , and the change in the sound velocity ( $\Delta v$ ), as the ultrasound propagated along the trigonal direction, are shown for two orientations of the magnetic field, when it is oriented almost along the trigonal axis (b) and when it is  $15^{\circ}$  (c) off the trigonal axis. There is a good consistency between the three experimental probes, indicat-



FIG. 4. The experimental and theoretical Landau spectrum: (a) The illustration of the twinned structure in bismuth. The two crystals tilt from each other by  $108^{\circ}$  and share one common binary. (b) The Landau spectrum of the main and twinned samples from holes according to theory (solid lines) and experiment (symbols) for binary-trigonal, (c) bisectrix-trigonal, and (d) binary-bisectrix planes. The red and blue symbols are for the peaks from the holes of the main and twinned crystals from the Fig. 2, respectively. The red and blue lines represent the calculated Landau levels from holes of the main and twinned domains, respectively. The magenta dashed lines are for calculated  $0_{e^-}$  electron Landau level which corresponds to the large drop in magnetoresistance as the field is along trigonal. The other electron peaks below 28 T are omitted for clarity, see the ref [34].

ing similar peak positions. Specifically, there are ~40 T anomalies in  $S_{xy}$ , in  $\frac{d^2\rho}{dB^2}$ , and in  $\Delta v$ . The slight difference in positions at  $\theta \approx 15^{\circ}$  is consistent with the experimental margin in measuring the angle between the magnetic field and the sample in three different experiments. We also notice that the anomalies in sound velocity and ultrasound attenuation occur at slightly different magnetic fields[49].

**Comparison with theory:** The angle-dependent Landau spectrum can be described by a theoretical model that considers electron pockets at the L points of the Brillouin zone in an extended Dirac Hamiltonian and the hole pocket at the T point as carriers with parabolic dispersion and highly anisotropic g-factor [34, 37, 50, 51]. In order to explain additional and unexpected anomalies, one needs to assume the presence of an additional secondary twinned domain [34, 37]. In the high field limit, when only a single Landau level is occupied, the interband coupling becomes important and plays a significant role in setting the details of the spectrum [37].

The twinned sample originates from the distortion by the symmetry-lowering operation due to Peierls transition[52]. Since the Schmid factor is 0.48 in bismuth [53], applying an external load[24] and thermal cycling[54] could relatively easily induce twins compared to other materials[53]. Thus, the strain appearing during sample cleaving by a blaze along cleavage basal trigonal (0001) plane in liquid nitrogen could introduce twinned domains on the sample and these twinned domains should mainly be on the surface. This could be a reason that a smaller RRR was found in a sample with a smaller dimension[55], since the ratio of the one plane area to the volume is smaller in a smaller sample. When the sample is twinned, besides the existing principal trigonal, a quasi-trigonal axis will be present.

Fig.4 compares the theoretical prediction with experimental peaks for three high symmetrical planes. The experimental peaks depicted by open symbols are compared with the theoretical results, which are presented by lines from both the main and a secondary tilted crystal. For clarity, we have omitted the electron peaks below 28 T, which have been identified earlier[34]. Beyond 28 T, we included all peaks associated with electrons from both domains. Fig.4(a) sketches the main and twinned crystals with their Fermi surfaces. The tilt angle between the two crystals is  $108^{\circ}[34]$ . The common binary axis is also shown. All theoretical results of the hole and twinned electron Landau level  $0_{e^-}$  are included.

The theoretical model here is fundamentally the same as the one used before [33, 34, 37], except that the tensor, which is the correction to the additional g-factor in the lowest Landau level V', was adjusted a little to fit the experimental Landau spectra better. The  $V'_3$  was adjusted to -0.0025 from -0.0625 [37]. As seen in Fig.4, whereas a slight displacement occurs, which may be due to misalignment during measurement, there is a relatively good agreement between the theory and the experiment. Some more peaks may be due to the electron Landau levels, which have not been clearly identified in the figures, see Fig.4(c). We may conclude, therefore, that the Landau spectrum of bismuth corresponds to what is expected in the single-particle theory. No collective effect due to electron-electron interaction is visible up to 65 T. Because of extreme sensitivity, nullification of the Zeeman splitting and of holes along the trigonal axis, the experimentally resolved hole peaks are split due to a small residual misalignment in the case of binary-bisectrix plane (see Fig.4(d)). The twinned crystal shares one common binary axis with the original crystal, but tilts its trigonal axis by  $108^{\circ}$  off[37]. This is demonstrated by Fig.4(d) at zero degrees where two sets of hole peaks merge. The additional peaks vanish as the two crystals are in the same configuration regarding the magnetic field and possess an identical Landau spectrum.

The twinning scenario provides an explanation for the previously reported drop in the magnetoresistance for a magnetic field of 38 T along the trigonal axis of the main crystal [45]. We can identify it as the result of emptying the electron pockets of the twinned domain. As seen in Fig.4(b) and Fig.4(c), the magneta dashed line, which represents the intersection between the  $0_{e-}$  Landau level of the twinned domain and the chemical potential happens to be at ~ 40 T when the magnetic field is along the trigonal axis of the main crystal. Thus, there is no ultra-quantum feature in the main crystal beyond the quantum limit of holes, which occurs at 9 T along the trigonal axis.

Twins observed by magneto-optics: We confirmed the presence of twin boundaries in our bismuth samples with a magneto-optical probe and found that



FIG. 5. No detectable change of twins under pulsed magnetic field: (a), (b), and (c) show photos at the specific time at 0 T, 30.8 T, and 0 T after the pulse, indicated in the (d). The crystalline orientations are marked in the (a). (d). The time pattern of the magneto-optical measurements performed at 8.1 K. The magnetic field pulse for the imaging experiment lasts  $\sim 5$  ms. Images were taken every 0.2 ms as marked by red open circles.

pulsed magnetic fields have no influence on the domain population. The *in-situ* images are shown in Fig.5. The crystal was illuminated with polarized light and the image was recorded using a high-speed camera in pulsed magnetic fields up to 30.8 T oriented along the trigonal axis. The field profile is shown in Fig.5(d). Images are taken at the magnetic field at the times represented by red circles. Three typical images taken at 0 T, 30.8 T, and 0 T after the pulse are shown in Fig.5 (a), (b), and (c), respectively. The binary axis common to the two domains is labelled in Fig.5(a). The twinned domains occupy a large area close to one-third of the surface.

Independent or common chemical potentials for main and twinned crystals: Let us now turn our attention to the most striking aspect of our results. Since the charge is conserved, any change in the density of electrons is to be compensated by an identical change in the density of holes. This leads to a shift in the position of the chemical potential with increasing magnetic field [6, 33, 34, 37]. This field-induced shift strongly depends on the orientation of the applied magnetic field. The main crystal and its twinned domain are tilted by 108 degrees. Therefore, the magnitude of the field-induced



FIG. 6. Comparison between theoretical and experimental results for independent and common chemical **potential:** The Landau spectrum in the bisectrix-trigonal plane. In (a),  $\mu$  in each crystal is independent of the other. In (b), the two crystals share a common  $\mu$ . The calculated Landau spectrum and the experimental results agree in (a), but radically differ in (b). (c) The chemical potential shift for the main and twinned sample according to the theoretical results with independent  $\mu$ . The difference becomes as large as 68 meV at 65 T.

Twinned 108° tilted

B (T)

-10

-20

10 20 30 40 50 60

chemical potential shift is different for the two twinned crystals. Thus, the Landau spectrum would be very different in two imaginable cases: a) The two crystals keep their own distinct chemical potential; b) The two crystals share the same chemical potential. Fig.6 (a) and (b) compare the experimental data on twinned hole peaks and the theoretically expected Landau spectrum in the two cases. As seen in Fig.6 (b), assuming that the two twinned crystals share a common chemical potential set by the main crystal, the theoretical expectation is incompatible with the experimental results, in sharp contrast with the agreement seen between theory and experiment

60

50

40 30

20

10

L

39 prie/

28 me\

B(T)

in the former case, see Fig.6(a).

This conclusion implies a discontinuity in chemical potential and in carrier density across the twin boundary, which is reminiscent of a Schottky barrier. Remarkably, when the magnetic field is along the trigonal axis, the difference in chemical potentials increases up to 68 meV at 65 T (see Fig.6 (c)). This is remarkable given that the twin boundary width is as narrow as 1 nm [26, 56]. This raises the following question: What impedes electrons belonging to adjacent crystals from mixing up?

Twinned

Т

8



FIG. 7. Schematic of band bending in p-n junctions, Schottky junction and newly proposed twinned p-n junction in semimetal bismuth before and after joining: Band bending in (a) the p-n junctions between two semiconductors; (b) the Schottky junction between a metal and a n-type semiconductor; (c) the twin boundary in semimetal bismuth

#### IV. DISCUSSION

We now propose a possible scenario explaining this effect. It is well known that the bands are bent near the interface between electronically different solids [57]. The typical examples are shown in Fig. 7 (a) for the p-n junction and (b) for the Schottky junction. The former is a junction between semiconductors, and the latter is a junction between a metal and a semiconductor. In each case, chemical potentials are different in each side when separated (upper panels of Fig. 7). When they are joined, the carriers are forced to move to make  $\mu$ common. For example, holes move from p-type to ntype, and electrons move from n-type to p-type in the p-n junction. The n-type band is shifted upward and the p-type band downward by the potential gradient due to the carrier transfer across the interface, resulting in the band bending with a common  $\mu$  as shown in Fig. 7 (a).

A similar band bending is also expected in the present twin boundary between Bi crystals. At zero field,  $\mu$  is shared between the main and twinned crystals. Under a strong magnetic field,  $\mu$  of the main crystal shifts differently from that of the twinned one, as shown in Fig. 6 (c). This situation can be expressed by the model where two electronically different semimetals are joined at the twin boundary as depicted in Fig. 7 (c).

Suppose that  $\mu$  of the main crystal is lower than that of the twinned one when two semimetals are separated. When they are joined, the carriers are forced to move to make  $\mu$  common. As a result, both the electron and hole bands are bent, as if two types of p-n junction coexist. By this band bending, the relative levels between  $\mu$ , the conduction and the valence bands are kept in each crystal as before joining. The Landau level spectrum of this twinned p-n junction is equivalent to that of the case where the independent chemical potentials are employed. Eventually, the mystery of the apparent jump of  $\mu$  is solved by this scenario of the twinned p-n junction.

For the twinned p-n junction scenario, we assumed the electric potential difference between the main and twinned crystals. One may question whether an inhomogeneous electric field can exist inside a (semi) metal. The standard answer would be "No" because finite current will screen the potential difference. However, Landauer argued against this simplistic view [58, 59]. Even in a metal, a point-like scattering center inside a metal will create an unscreened dipolar field, i.e., the electric field is inhomogeneous near the point-like defects. In the present case, the twin boundary, a line or plane defect, will play a similar role to the point-like scattering center of Landauer's theory.

In this scenario, the Landau eigenvalues curve in the vicinity of a twin boundary in the same way as they bend near the edges of a 2D electron gas [60]. Any time of such Landau level bending results in edge currents, the continuity equation causes the edge charge and currents to remain close to the edge but distributed in real space in a 2D case[60]. In the present 3D case, the continuity equation strictly confines the currents and any associated charge variations to the vicinity of the twin boundary. The currents flow along the twin boundaries and essentially account for small differences in magnetization between twins, which should be present due to differences in Landau level filling factors related to the difference in chemical potential. The equilibrium distribution

of currents and charge ultimately depends on a balance between the Lorentz force of the current and the twin boundary potential that causes the Landau level eigenstates to curve in real space. These currents flowing along the twin boundary result from a novel form of extended bound state associated with the twin junction. This line of inquiry deserves further investigations.

In summary, we have carried out angle-dependence of magnetoresistance along three high symmetric planes, and mapped out their Landau spectra up to 65 T. All anomalies including the one observed at  $\sim 40$  T for a magnetic field oriented along the trigonal axis could be explained by assuming the presence of a second twinned crystal and the two crystals keep separate chemical potentials. We suggested an interpretation of this result by a scenario in which a p-n junction builds up at the twin boundary-areas. This implies the existence of an inhomogeneous electric field in the vicinity of twin boundaries, which can be checked by future experiments.

This work was supported by The National Key Research and Development Program of China (Grants No.2022YFA1403500 and 2016YFA0401704), the National Science Foundation of China (Grant No. 11574097 and No. 51861135104), and by directors funding grant number 20120772 at LANL. N.H. and RMcD acknowledge support from the US-DOE BES 'Science of 100T' B. F. acknowledges support from Jeunes program. Equipes de l'Institut de Physique du Collége de France (JEIP). K. B. was supported by the Agence Nationale de la Recherche (ANR-19-CE30-0014-04). Y. F. was supported by JSPS KAKENHI grants 23H04862, 23H00268. and 18KK0132. Work performed at National High Magnetic Field Laboratory was supported by National Science Foundation Cooperative Agreements DMR-0654118 and DMR-2128556 and the State of Florida. M. T. acknowledges support from the JSPS KAKENHI grant No. 23H04862.

\* zengwei.zhu@hust.edu.cn

- M. Dresselhaus, Electronic properties of the group V semimetals, J Phys Chem Solids 32, Suppl.1, 3 (1971).
- [2] V. S. Edel'man, Properties of electrons in bismuth, Soviet Physics Uspekhi 10, 819 (1977).
- [3] J. P. Issi, Low temperature transport properties of the group V semimetals, Australian Journal of Physics 32, 585 (1979).
- [4] Y. Fuseya, M. Ogata, and H. Fukuyama, Transport properties and diamagnetism of Dirac electrons in bismuth, Journal of the Physical Society of Japan 84, 012001 (2015).
- [5] D. Shoenberg and P. L. Kapitza, The magnetic properties of bismuth, III. further measurements on the de Haasvan Alphen effect, Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences 170, 341 (1939).
- [6] G. E. Smith, G. A. Baraff, and J. M. Rowell, Effective g factor of electrons and holes in bismuth, Physical Review 135, A1118 (1964).
- [7] R. N. Bhargava, de Haas-van Alphen and galvanomagnetic effect in Bi and Bi-Pb alloys, Physical Review 156, 785 (1967).
- [8] Y. Liu and R. E. Allen, Electronic structure of the semimetals Bi and Sb, Physical Review B 52, 1566 (1995).
- [9] L. Fu, C. L. Kane, and E. J. Mele, Topological insulators in three dimensions, Physical Review Letters 98, 106803 (2007).
- [10] I. K. Drozdov, A. Alexandradinata, S. Jeon, S. Nadj-Perge, H. Ji, R. J. Cava, B. Andrei Bernevig, and A. Yazdani, One-dimensional topological edge states of bismuth bilayers, Nature Physics 10, 664 (2014).
- [11] H. M. Benia, C. Straßer, K. Kern, and C. R. Ast, Surface band structure of  $\text{Bi}_{1-x}\text{Sb}_x(111)$ , Phys. Rev. B **91**, 161406 (2015).
- [12] I. Aguilera, C. Friedrich, and S. Blügel, Electronic phase transitions of bismuth under strain from relativistic self-

consistent GW calculations, Phys. Rev. B **91**, 125129 (2015).

- [13] S. Ito, B. Feng, M. Arita, A. Takayama, R.-Y. Liu, T. Someya, W.-C. Chen, T. Iimori, H. Namatame, M. Taniguchi, C.-M. Cheng, S.-J. Tang, F. Komori, K. Kobayashi, T.-C. Chiang, and I. Matsuda, Proving nontrivial topology of pure bismuth by quantum confinement, Phys. Rev. Lett. **117**, 236402 (2016).
- [14] A. Murani, A. Kasumov, S. Sengupta, Y. A. Kasumov, V. T. Volkov, I. I. Khodos, F. Brisset, R. Delagrange, A. Chepelianskii, R. Deblock, H. Bouchiat, and S. Guéron, Ballistic edge states in bismuth nanowires revealed by SQUID interferometry, Nature Communications 8, 15941 (2017).
- [15] Y. Fuseya and H. Fukuyama, Analytical solutions for the surface states of  $\text{Bi}_{1-x}\text{Sb}_x$  ( $0 \le x \le 0.1$ ), J. Phys. Soc. Jpn. 87, 044710 (2018).
- [16] F. Schindler, Z. Wang, M. G. Vergniory, A. M. Cook, A. Murani, S. Sengupta, A. Y. Kasumov, R. Deblock, S. Jeon, I. Drozdov, H. Bouchiat, S. Guéron, A. Yazdani, B. A. Bernevig, and T. Neupert, Higher-order topology in bismuth, Nature Physics 14, 918 (2018).
- [17] A. K. Nayak, J. Reiner, R. Queiroz, H. Fu, C. Shekhar, B. Yan, C. Felser, N. Avraham, and H. Beidenkopf, Resolving the topological classification of bismuth with topological defects, Science Advances 5, eaax6996 (2019).
- [18] C.-H. Hsu, X. Zhou, T.-R. Chang, Q. Ma, N. Gedik, A. Bansil, S.-Y. Xu, H. Lin, and L. Fu, Topology on a new facet of bismuth, Proceedings of the National Academy of Sciences **116**, 13255 (2019).
- [19] A. Collaudin, B. Fauqué, Y. Fuseya, W. Kang, and K. Behnia, Angle dependence of the orbital magnetoresistance in bismuth, Phys. Rev. X 5, 021022 (2015).
- [20] F. Spathelf, B. Fauqué, and K. Behnia, Magneto-seebeck effect in bismuth, Phys. Rev. B 105, 235116 (2022).
- [21] W. Kang, F. Spathelf, B. Fauqué, Y. Fuseya, and K. Behnia, Boundary conductance in macroscopic bis-

muth crystals, Nature Communications 13, 189 (2022).

- [22] Y. Fuseya, H. Katsuno, K. Behnia, and A. Kapitulnik, Nanoscale turing patterns in a bismuth monolayer, Nature Physics 17, 1031 (2021).
- [23] V. S. Savenko, V. V. Uglov, O. M. Ostrikov, and A. P. Khodoskin, Twinning of bismuth single crystals bombarded by boron ions, Technical Physics Letters 24, 287 (1998).
- [24] V. I. Bashmakov and T. S. Chikova, Evolution of twinboundary geometry in bismuth crystals with an increase of load, Crystallography Reports 47, 489 (2002).
- [25] V. S. Edelman, D. Y. Sharvin, I. N. Khlyustikov, and A. M. Troyanovskii, STM revealing of twin microlayers with quantized width on cleaved bismuth surface, Europhysics Letters (EPL) 34, 115 (1996).
- [26] V. S. Edel'man, Scanning tunnel microscopy and spectroscopy of an atomically clean bismuth surface, Phys. Usp. 48, 1057 (2005).
- [27] B. I. Halperin, Possible states for a three-dimensional electron gas in a strong magnetic field, Japanese Journal of Applied Physics 26, 1913 (1987).
- [28] A. H. MacDonald and G. W. Bryant, Strong-magneticfield states of the pure electron plasma, Phys. Rev. Lett. 58, 515 (1987).
- [29] K. Behnia, L. Balicas, and Y. Kopelevich, Signatures of electron fractionalization in ultraquantum bismuth, Science **317**, 1729 (2007).
- [30] L. Li, J. G. Checkelsky, Y. S. Hor, C. Uher, A. F. Hebard, R. J. Cava, and N. P. Ong, Phase transitions of Dirac electrons in bismuth, Science **321**, 547 (2008).
- [31] J. Alicea and L. Balents, Bismuth in strong magnetic fields: Unconventional Zeeman coupling and correlation effects, Phys. Rev. B 79, 241101 (2009).
- [32] Y. V. Sharlai and G. P. Mikitik, Origin of the peaks in the Nernst coefficient of bismuth in strong magnetic fields, Phys. Rev. B 79, 081102 (2009).
- [33] Z. Zhu, B. Fauqué, Y. Fuseya, and K. Behnia, Angleresolved landau spectrum of electrons and holes in bismuth, Physical Review B 84, 115137 (2011).
- [34] Z. Zhu, B. Fauqué, L. Malone, A. B. Antunes, Y. Fuseya, and K. Behnia, Landau spectrum and twin boundaries of bismuth in the extreme quantum limit, Proceedings of the National Academy of Sciences 109, 14813 (2012).
- [35] B. Fauqué, H. Yang, I. Sheikin, L. Balicas, J. P. Issi, and K. Behnia, Hall plateaus at magic angles in bismuth beyond the quantum limit, Physical Review B 79, (2009).
- [36] R. Kuchler, L. Steinke, R. Daou, M. Brando, K. Behnia, and F. Steglich, Thermodynamic evidence for valleydependent density of states in bulk bismuth, Nature Materials 13, 461 (2014).
- [37] Z. Zhu, J. Wang, H. Zuo, B. Fauqué, R. D. McDonald, Y. Fuseya, and K. Behnia, Emptying Dirac valleys in bismuth using high magnetic fields, Nature Communications 8, 15297 (2017).
- [38] A. Iwasa, A. Kondo, S. Kawachi, K. Akiba, Y. Nakanishi, M. Yoshizawa, M. Tokunaga, and K. Kindo, Thermodynamic evidence of magnetic-field-induced complete valley polarization in bismuth, Scientific Reports 9, 1672 (2019).
- [39] H. Yaguchi and J. Singleton, A high-magnetic-fieldinduced density-wave state in graphite, Journal of Physics: Condensed Matter 21, 344207 (2009).
- [40] B. Fauqué, D. LeBoeuf, B. Vignolle, M. Nardone, C. Proust, and K. Behnia, Two phase transitions induced

by a magnetic field in graphite, Phys. Rev. Lett. 110, 266601 (2013).

- [41] J. Wang, P. Nie, X. Li, H. Zuo, B. Fauqué, Z. Zhu, and K. Behnia, Critical point for Bose-Einstein condensation of excitons in graphite, Proceedings of the National Academy of Sciences 117, 30215 (2020).
- [42] Z. Zhu, P. Nie, B. Fauqué, B. Vignolle, C. Proust, R. D. McDonald, N. Harrison, and K. Behnia, Graphite in 90 t: Evidence for strong-coupling excitonic pairing, Phys. Rev. X 9, 011058 (2019).
- [43] Y. Kinoshita, T. Fujita, R. Kurihara, A. Miyake, Y. Izaki, Y. Fuseya, and M. Tokunaga, Field-induced reentrant insulator state of a gap-closed topological insulator  $(Bi_{1-x}Sb_x)$  in quantum-limit states, Phys. Rev. B 107, 125140 (2023).
- [44] A. Banerjee, B. Fauqué, K. Izawa, A. Miyake, I. Sheikin, J. Flouquet, B. Lenoir, and K. Behnia, Transport anomalies across the quantum limit in semimetallic Bi<sub>0.96</sub>Sb<sub>0.04</sub>, Phys. Rev. B **78**, 161103 (2008).
- [45] B. Fauqué, B. Vignolle, C. Proust, J.-P. Issi, and K. Behnia, Electronic instability in bismuth far beyond the quantum limit, New Journal of Physics 11, 113012 (2009).
- [46] I. Katakura, M. Tokunaga, A. Matsuo, K. Kawaguchi, K. Kindo, M. Hitomi, D. Akahoshi, and H. Kuwahara, Development of high-speed polarizing imaging system for operation in high pulsed magnetic field, Review of Scientific Instruments 81, 043701 (2010).
- [47] S. G. Bompadre, C. Biagini, D. Maslov, and A. F. Hebard, Unambiguous determination of the g factor for holes in bismuth at high B/T, Phys. Rev. B 64, 073103 (2001).
- [48] K. Behnia, M.-A. Méasson, and Y. Kopelevich, Oscillating Nernst-Ettingshausen effect in bismuth across the quantum limit, Phys. Rev. Lett. 98, 166602 (2007).
- [49] K. Akiba, A. Miyake, H. Sakai, K. Katayama, H. Murakawa, N. Hanasaki, S. Takaoka, Y. Nakanishi, M. Yoshizawa, and M. Tokunaga, Quantitative evaluation of Dirac physics in PbTe, Phys. Rev. B 98, 115144 (2018).
- [50] Z. Zhu, H. Yang, A. Banerjee, L. Malone, B. Fauqué, and K. Behnia, Nernst quantum oscillations in bulk semi-metals, Journal of Physics: Condensed Matter 23, 094204 (2011).
- [51] Y. Fuseya, Z. Zhu, B. Fauqué, W. Kang, B. Lenoir, and K. Behnia, Origin of the large anisotropic g factor of holes in bismuth, Physical Review Letters 115, 216401 (2015).
- [52] R. Peierls, More Surprises in Theoretical Physics (Princeton Univ Press, Princeton, 1991).
- [53] O. M. Ostrikov, Effect of carbon ion bombardment and surface oxidation on the twinning rate of single-crystal bismuth, Technical Physics 44, 597 (1999).
- [54] O. M. Ostrikov, Twinning in bismuth single crystals induced by thermal cycling, Technical Physics 46, 1199 (2001).
- [55] V. Galev, V. Kozlov, N. Kolomoets, S. Y. Skipidarov, and N. Tsvetkova, Exponential temperature dependence of the coefficient of the Nernst-Ettingshausen transverse effect in bismuth, Sov. Phys.-JETP Lett. 33, 106 (1981).
- [56] V. S. Edelman, Tunneling spectra at terrace boundaries on the bismuth surface, Journal of Experimental and Theoretical Physics 107, 251 (2008).
- [57] C. Kittel, Introduction to solid state physics (John Wiley & Sons, Inc., New York, 2004).

nal of Research and Development 32, 306 (1988).

- [60] A. H. MacDonald, T. M. Rice, and W. F. Brinkman, Hall voltage and current distributions in an ideal twodimensional system, Phys. Rev. B 28, 3648 (1983).
- [58] R. Landauer, Residual resistivity dipoles, Zeitschrift f
  ür Physik B Condensed Matter 21, 247 (1975).
- [59] R. Landauer, Spatial variation of currents and fields due to localized scatterers in metallic conduction, IBM Jour-