

# FLUID SHEAR STRESS ON OSTEOCYTE UNDER ULTRASOUND STIMULATION: FINITE-ELEMENT MODEL

Cécile Baron, Elise Doveri, Philippe Lasaygues, Carine Guivier-Curien

## ► To cite this version:

Cécile Baron, Elise Doveri, Philippe Lasaygues, Carine Guivier-Curien. FLUID SHEAR STRESS ON OSTEOCYTE UNDER ULTRASOUND STIMULATION: FINITE-ELEMENT MODEL. International Symposium on Computers Methods in Biomechanics and Biomedical Engineering, May 2023, Paris, France. hal-04298154

## HAL Id: hal-04298154 https://hal.science/hal-04298154

Submitted on 21 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



3 - 5 May 2023, Paris, France



Arts et Métiers Institute of Technology Institut de Biomécanique Humaine Georges Charpak 155 boulevard de l'Hôpital 75013 Paris France

www.cmbbe-symposium.com

## FLUID SHEAR STRESS ON OSTEOCYTE UNDER ULTRASOUND STIMULATION: FINITE-ELEMENT MODEL

#### Cécile Baron (1), Elise Doveri (2), Philippe Lasaygues (2), Carine Guivier-Curien (1)

1. Aix-Marseille Univ, CNRS, Centrale Marseille, IRPHE UMR 7342, Marseille, France; 2. Aix-Marseille Univ, CNRS, Centrale Marseille, LMA UMR 7031, Marseille France.

#### 1. Introduction

Ultrasound (US) stimulation of bone remodeling remains an open question as the underlying mechanisms involved are still poorly understood. Among the possible hypotheses, the propagation of ultrasonic waves in bone tissue could generate acoustic streaming (AS) resulting in fluid shear stress exerted on the wall of osteocytes known as the mechanosensitive conductors of bone remodeling. Osteocytes are dendritic cells, ubiquituous in the bone ECM forming a complex micrometric 3D network called the lacuno-canalicular network (LCN). A first finite-element (FE) model is proposed to investigate the effect of US stimulation on a network of 3 osteocytes aligned in the thickness of an osteon, surrounded by a fluid layer and embedded in a rigid extra-cellular matrix (ECM). The influence of a wavy ECM/fluid interface [1] on the wall shear stress (WSS) level is considered.

#### 2. Materials and Methods

In order to implement relevant boundary conditions, we focus on an osteon unit delimited by the cement line (C line) as the outer boundary and the Haversian canal (H canal) as the inner one. Between these two boundaries, we consider a network of 3 osteocytes whose cell body is housed in an ellipsoidal lacuna and connected by one process inside a canaliculus full of fluid considered as salt water at 37°C. A 2D axisymmetric FE model is implemented on Comsol Multiphysics (v. 6.0) to estimate the WSS at the osteocyte surface induced by the AS in the fluid [2]. The boundary conditions are summarized in Table 1. A first order oscillating velocity  $U_1$  is imposed normal to the ECM/fluid interface to model a harmonic acoustic wave at 1MHz with an amplitude of displacement  $d_0$  equal to 0.4nm. Two geometries of canaliculus are compared:

straight and wavy.

|             | Laminar flow   | Acoustics  |
|-------------|----------------|--|
| Cement line | Wall (no slip) | Wall (no slip)   |
| H canal     | Open boundary  | Z continuity   |
| ECM/Fluid   | Wall (no slip) | $\mathbf{U}_{1}(\omega) = i\omega d_{0}\mathbf{e}_{r}$ |
| Fluid/cell  | Wall (no slip) | Z continuity   |

**Table 1**: Boundary conditions. Z: normal acoustic impedance.

#### 3. Results

The geometry of the canliculus has a strong influence on the WSS.



Figure 1: WSS along the wall of the 3 osteocytes.

#### 4. Discussion and Conclusions

This preliminary study shows that harmonic ultrasound stimulation can induce WSS level greater than 0.8 Pa, known as the physiological threshold to trigger bone remodeling [3]. The value of  $d_0$  needs to be further investigated. Moreover, the model can be improved by integrating the osteocyte elasticity and by adding processes.

#### 5. References

- 1. McNamara L et al. Anat. Rec.; 292(3):355-363 (2009).
- 2. Muller P et Bruus H, Phys. Rev E; 90(4):043016 (2014).
- 3. Weinbaum S et al., JoB; 27(3):339-360 (1994).