



HAL
open science

Macroscopic model for unsteady slip flow in porous media

Didier Lasseux, Francisco J. Valdès-Parada, Alessandro Bottaro

► **To cite this version:**

Didier Lasseux, Francisco J. Valdès-Parada, Alessandro Bottaro. Macroscopic model for unsteady slip flow in porous media. *Rarefied Gas Dynamics (RGD32)*, Jul 2022, Séoul, South Korea. hal-04299449

HAL Id: hal-04299449

<https://hal.science/hal-04299449>

Submitted on 22 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Macroscopic model for unsteady slip flow in porous media

[oral]

D. Lasseux¹, F. Valdés-Parada², A. Bottaro³

¹ I2M, UMR5295, CNRS, Univ. Bordeaux, 351, Cours de la Libération, 33405 Talence CEDEX, France

² Universidad Autónoma Metropolitana-Iztapalapa, Av. San Rafael Atlixco 186 09340 Mexico D.F., Mexico

³ Scuola Politecnica, Università di Genova, Via Montallegro 1 Genova 16145, Italy

In this work, a macroscopic model of unsteady gas flow in a homogeneous porous medium in the slip regime is derived. The starting point is the unsteady creeping, isothermal slip flow model at the microscale (the pore-scale), considering a first-order slip condition at the solid-fluid interface [1,2]. An adjoint method with a Green's function formulation [3], coupled to a simplified version of the volume averaging technique [4], is employed to obtain the macroscopic model using the Laplace transform. Under the assumption of slightly compressible flow, the macroscopic mass conservation equation keeps a form similar to its counterpart at the pore-scale. Moreover, the macroscopic momentum balance is a modified version of Darcy's law that does not correspond to a heuristic model that has been often postulated for this type of flow, as already highlighted in [5]. It involves the convolution product of the time-derivative of the apparent permeability tensor (\mathbf{K}_{st} , function of the macroscopic Knudsen number, $\xi\bar{\lambda}^*$) with the (time-dependent) macroscopic pressure gradient. In addition, the effect of the initial flow condition is accounted for by an effective memory term, α_s . Both \mathbf{K}_{st} (which symmetry and positiveness are analyzed) and α_s are determined by a single unsteady closure problem, that depends on the Knudsen number, to be solved in a representative elementary volume (periodic unit cell) of the porous structure under consideration [6].

The dependences of \mathbf{K}_{st} and α_s upon $\xi\bar{\lambda}^*$ are analyzed and the validity of the macroscopic model is illustrated through a comparison with pore-scale direct numerical simulations (DNS). This is carried out in a two-dimensional model structure made of a square pattern of parallel cylinders of uniform circular cross section, with flow orthogonal to the cylinder axes (*i.e.*, in the x -direction). An oscillatory macroscopic forcing along the x -axis is considered at $t \geq 0$, following a constant pressure gradient in the same direction for $t < 0$. As shown in Fig.1, where the evolution of the x -component of the average velocity is represented, the prediction of the macroscopic model is in excellent agreement with the pore-scale DNS. Moreover, it is further evidenced that the heuristic model is inadequate, in general.




Fig.1: x -component evolution of the average velocity within a square pattern of parallel cylinders resulting from an oscillatory pressure gradient in the x -direction, corresponding to a lattice axis orthogonal to the cylinder axes [6].

REFERENCES

- [1] D. Einzel, P. Panzer, M. Liu, *Phys. Rev. Let.*, **64**(19), 2269–2272 (1990)
- [2] D. Lasseux, F. J. Valdés-Parada, M. L. Porter, *J. Fluid Mech.*, **805**, 118-146 (2016)
- [3] A. Bottaro, *J. Fluid Mech.*, **877**, P1 (2019)
- [4] S. Whitaker, *The Method of Volume Averaging*. Springer (1999)
- [5] D. Lasseux, F. J. Valdés-Parada, *J. Fluid Mech.*, **862**, 283-311 (2019)
- [6] D. Lasseux, F.J. Valdés-Parada, A. Bottaro, *J. Fluid Mech.*, **923**, A37 (2021)