



HAL
open science

Harris Recurrent Markov Chains and Nonlinear Monotone Cointegrated Models

Patrice Bertail, Cécile Durot, Carlos Fernández

► **To cite this version:**

Patrice Bertail, Cécile Durot, Carlos Fernández. Harris Recurrent Markov Chains and Nonlinear Monotone Cointegrated Models. 2023. hal-04299698

HAL Id: hal-04299698

<https://hal.science/hal-04299698>

Preprint submitted on 22 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HARRIS RECURRENT MARKOV CHAINS AND NONLINEAR MONOTONE COINTEGRATED MODELS

BY PATRICE BERTAIL^{1,a}  CÉCILE DUROT^{1,b}  AND CARLOS FERNÁNDEZ^{1,2,c} 

¹MODAL'X, UMR CNRS 9023, Université Paris Nanterre, ^apatrice.ber tail@parisnanterre.fr; ^bcecile.durot@parisnanterre.fr

²LTCI, Telecom Paris, Institut Polytechnique de Paris, ^cfernandez@telecom-paris.fr

In this paper, we study a nonlinear cointegration-type model of the form $Z_t = f_0(X_t) + W_t$ where f_0 is a monotone function and X_t is a Harris recurrent Markov chain. We use a nonparametric Least Square Estimator to locally estimate f_0 , and under mild conditions, we show its strong consistency and obtain its rate of convergence. New results (of the Glivenko-Cantelli type) for localized null recurrent Markov chains are also proved.

1. Introduction and motivations.

1.1. *Linear and nonlinear cointegration models.* *Linear cointegration* introduced by [16] and developed by [14] and [21, 22], is a concept used in statistics and econometrics to describe a long-term relationship between two or more time series. In general, these time series are non-stationary, integrated of order 1, that is, they behave roughly as random walks. In traditional linear cointegration analysis, variables are assumed to have a linear relationship, which means their long-term equilibrium, as time grows, is characterized by a constant linear combination. This concept has since been extensively studied, particularly in the field of econometrics [14, 21, 22, 30, 31]. Notice that, when there is indeed a significant linear relationship, the link is monotone in each of the variables.

However, in some cases, the relationship between variables may exhibit nonlinear behavior, which cannot be adequately captured by linear cointegration models. The incorporation of nonlinearities allows for a more comprehensive understanding of long-term relationships between variables. [20] have developed an approach for analyzing nonlinear cointegration through threshold cointegration models. These models assume that the linear relationship between variables differs after some changepoints, leading to different long-run equilibrium states (for instance according to some latent regimes). Threshold cointegration models provide a framework for capturing nonlinearities in the data and estimating the changepoints. Refer to [32] for examples and discussions on the importance to introduce nonlinearities in cointegration applications and for further references.

Another method for analyzing nonlinear cointegration is through the use of smooth transition cointegration models introduced by [17]. These models assume a ECM (Error Correction Model) form and allow for smooth transitions between different regimes in the data. Most estimators of non-linear cointegration may be seen as Nadaraya-Watson estimators of the link function. For instance, Wang and Phillips [35–37] show that it is possible to estimate and perform asymptotic inference in specific nonparametric cointegration regression models using kernel regression techniques. Furthermore, they established that the self-normalized kernel regression estimators converge to a standard normal distribution limit, even when the explanatory variable is integrated. These findings indicate that the estimators can consistently

MSC2020 subject classifications: Primary 62G05, 62M05; secondary 62G30.

Keywords and phrases: monotone regression, isotonic regression, nonlinear cointegration, nonparametric estimation, null recurrent Markov chain.

capture the underlying relationship between variables, even in cases where the explanatory variable exhibits non-stationary behavior. The problem of estimating f_0 under the Markovian assumptions has also been tackled using local linear M-type estimators in [8, 26] using smoothing techniques.

These results have been partially extended in the framework of general β -recurrent Markov chain and not just integrated $I(1)$ time series by [7, 23]. Consider the simple framework where we observe two Markov chains, Z_t and X_t . [23] are essentially interested in the study of *nonlinear cointegrated* models such as,

$$(1) \quad Z_t = f_0(X_t) + W_t,$$

where f_0 is a nonlinear function. X_t and W_t are independent processes, and X_t is a positive or β -null recurrent Markov chain. Despite the fact that there is no stationary probability measure for X_t , they apply the Nadaraya-Watson method to estimate f_0 and established the asymptotic theory of the proposed estimator. The rate of convergence of the estimators at some point x is essentially linked to the local properties of the β -null recurrent chain X_t and typically of the order of the square root of the number of visits of the chain in a neighborhood of the point x .

1.2. *Monotone cointegration models: motivations.* Monotonicity in cointegration is a rather natural assumption in many economic applications, for instance for modeling demand as a function of income or prices (see for instance [11]) or other variables. Suppose, for example, we are interested in analyzing the long-term relationship between ice cream sales and the average monthly temperature. These two non-stationary variables may be modeled by some β -recurrent Markov chains. We hypothesize that as the average monthly temperature increases, the demand for ice cream also increases: however the rate of increase may vary according to the season. In that case, the nonlinear relationship between the two variables will be monotone. In microeconomics, the same phenomenon is expected for Engel curves, describing how real expenditure varies with household income (see [11]). Expenditures and income (or their log in most models) may be considered as non-stationary variables. However considering a linear cointegration between them may be misleading, since the relationship may change along the life cycle. By Engle's law, the relationship between the two variables should be monotone. Other types of examples of monotone non-linear cointegration phenomenon may be found in [32].

The purpose of this paper is to propose a simple estimator that is automatically monotone, does not require strong smoothness assumptions (we only require continuity of the link function), and operates under general Markovian assumptions. We establish a nonparametric estimation theory of the nonparametric least squares estimator (LSE) for the function f_0 in the model (1) under the constraints that f_0 is monotone non-increasing. Here, $\{W_t\}$ is an unobserved process such that $E(W_t|X_t) = 0$ to ensure identifiability of f_0 . Since a minimal condition for undertaking asymptotic analysis on $f_0(x_0)$ at a given point x_0 is that, as the number of observations on $\{X_t\}$ increases, there must be infinitely many observations in the neighborhood of x_0 , the process $\{X_t\}$ will be assumed to be a Harris recurrent Markov chain (cf section 2). We consider at the same time the stationary and β null recurrent non-stationary framework. To our knowledge, it is the first time that such an estimator is proposed in the literature in such a large framework.

1.3. *The estimator.* Let C be a set whose interior contains our point of interest x_0 . Having observed $(X_t, Z_t)_{t=0}^n$, we denote by $T_n(C)$ the number of times that \mathbf{X} visited C up to time n and by $\sigma_C(i)$ the time of the i -th visit. Then, we consider the nonparametric LSE defined as the minimizer of

$$(2) \quad f \mapsto \sum_{i=1}^{T_n(C)} (Z_{\sigma_C(i)} - f(X_{\sigma_C(i)}))^2$$

over the set of non-increasing functions f on \mathbb{R} . The nonparametric LSE \widehat{f}_n has a well known characterization, as follows. Let m be the number of unique values of $X_{\sigma_C(1)}, \dots, X_{\sigma_C(T_n(C))}$, and $Y_1 < \dots < Y_m$ be the corresponding order statistics. Then, $\widehat{f}_n(Y_k)$ is the left-hand slope at $\sum_{i=1}^{T_n(C)} \mathbb{I}\{X_{\sigma_C(i)} \leq Y_k\}$ of the least concave majorant of the set of points

$$(3) \quad \left\{ (0, 0), \left(\sum_{i=1}^{T_n(C)} \mathbb{I}\{X_{\sigma_C(i)} \leq Y_k\}, \sum_{i=1}^{T_n(C)} Z_{\sigma_C(i)} \mathbb{I}\{X_{\sigma_C(i)} \leq Y_k\} \right), k = 1, \dots, m \right\},$$

and it can be computed using simple algorithms as discussed in [3]. Thus, the constrained LSE is uniquely defined at the observation points, however, it is not uniquely defined between these points: any monotone interpolation of these values is a constrained LSE. As is customary, we consider in the sequel the piecewise-constant and left-continuous LSE that is constant on every interval $(Y_{k-1}, Y_k]$, $k = 2, \dots, m$ and also on $(-\infty, Y_1]$ and on $[Y_m, \infty)$.

The use of a localized estimator is due to the fact that we need to control the behavior of the chain around x_0 , and, to do this, we need to estimate the asymptotic "distribution" of \mathbf{X} in a vicinity of x_0 . For Harris recurrent Markov chains, the long-term behavior of the chain is given by its invariant measure (see Section 2). In the positive recurrent case, the invariant measure is finite and it can be estimated by simply considering the empirical cumulative distribution function of the X_t . However, in the null recurrent case, the invariant measure is only σ -finite, hence, we need to localize our analysis in a set big enough such that the chain visits it infinitely often, but small enough that the restriction of the invariant measure to it is finite. Moreover, contrary to the bandwidth in kernel type estimators, C does not depend on n , and the rate of convergence of the estimator does not depend on C .

1.4. *Outline.* Since our paper draws quite heavily on the theory of Harris recurrent Markov chains, we have added a small introduction to the subject as well as the main results that we use throughout the paper in Section 2. In Section 3, we show that under very general assumptions, our estimator \widehat{f}_n is strongly consistent, while its rate of convergence is presented in Section 4. In Section 5, we present new results concerning the localized empirical process of Harris recurrent Markov chains that have emerged during our investigation and we believe are interesting in their own right. Section 6 contains the proofs of our main results.

2. Markov chain theory and notation. In this section, we present the notation and main results related to Markov chains that are needed to present our main results. For further details, we refer the reader to the first section of the Supplementary Material [4] and the books [12, 27, 29].

Consider a time-homogeneous irreducible Markov Chain, denoted as $\mathbf{X} = X_0, X_1, X_2, \dots$, defined on a probability space $(E, \mathcal{E}, \mathbb{P})$, where \mathcal{E} is countably generated. The irreducibility measure of the chain is represented by ψ . The transition kernel of the chain is denoted as $P(x, A)$ and its initial distribution is represented by λ . If the initial measure of the chain is specified, we use \mathbb{P}_λ (and \mathbb{E}_λ) to denote the probability (and the expectation) conditioned on the law of the initial state $\mathcal{L}(X_0) = \lambda$.

For any set $C \in \mathcal{E}$, we will denote by σ_C and τ_C , respectively, the times of first visit and first return of the chain to the set C , i.e. $\tau_C = \inf \{n \geq 1 : X_n \in C\}$ and $\sigma_C = \inf \{n \geq 0 : X_n \in C\}$. The subsequent visit and return times $\sigma_C, \tau_C(k)$, $k \geq 1$ are defined inductively.

Given that our methods will only deal with the values of \mathbf{X} in a fixed set C , if A is a measurable set, we will write $\mathbb{I}_C\{X_t \in A\}$ instead of $\mathbb{I}\{X_t \in A \cap C\}$ and if $A = E$, then we will simply write $\mathbb{I}_C(X_t)$. We will use $T_n(C)$ to denote the random variable that counts the number of times the chain has visited the set C up to time n , that is $T_n(C) = \sum_{t=0}^n \mathbb{I}_C(X_t)$. Similarly, we will write $T(C)$ for the total of numbers of visits the chain \mathbf{X} to C . The set C is called *recurrent* if $\mathbb{E}_x T(C) = +\infty$ for all $x \in C$ and the chain \mathbf{X} is recurrent if every set $A \in \mathcal{E}$ such that $\psi(A) > 0$ is recurrent. A recurrent chain is called *Harris recurrent* if for all $x \in E$ and all $A \in \mathcal{E}$ with $\psi(A) > 0$ we have $\mathbb{P}(X_n \in A \text{ infinitely often} | X_0 = x) = 1$.

Denote by \mathcal{E}^+ the class of nonnegative measurable functions with positive ψ support. A function $s \in \mathcal{E}^+$ is called *small* if there exists an integer $m_0 \geq 1$ and a measure $\nu \in \mathcal{M}(\mathcal{E})_+$ such that

$$(4) \quad P^{m_0}(x, A) \geq s(x) \nu(A) \quad \forall x \in E, A \in \mathcal{E}.$$

When a chain possesses a small function s , we say that it satisfies the *minorization inequality* $M(m_0, s, \nu)$. A set $A \in \mathcal{E}$ is said to be *small* if the function \mathbb{I}_A is small. Similarly, a measure ν is *small* if there exist m_0 , and s that satisfy (4). By Theorem 2.1 in [29], every irreducible Markov chain possesses a small function and Proposition 2.6 of the same book shows that every measurable set A with $\psi(A) > 0$ contains a small set. In practice, finding such a set consists in most cases in exhibiting an accessible set, for which the probability that the chain returns to it in m steps is uniformly bounded below. Moreover, under quite wide conditions a compact set will be small, see [15].

An irreducible chain possesses an accessible atom, if there is a set $\alpha \in \mathcal{E}$ such that for all x, y in α : $P(x, \cdot) = P(y, \cdot)$ and $\psi(\alpha) > 0$. When an accessible atom exists, the *stochastic stability* properties of \mathbf{X} amount to properties concerning the speed of return time to the atom only. Moreover, it follows from the *strong Markov property* that the sample paths may be divided into independent blocks of random length corresponding to consecutive visits to α . The sequence $\{\tau_\alpha(j)\}_{j \geq 1}$ defines successive times at which the chain forgets its past, called *regeneration times*. Similarly, the sequence of i.i.d. blocks $\{\mathcal{B}_j\}_{j \geq 1}$ are named *regeneration blocks*. The random variable $T(n) = T_n(\alpha) - 1$ counts the number of i.i.d. blocks up to time n . This term is called *number of regenerations up to time n* .

If \mathbf{X} does not possess an atom but is Harris recurrent (and therefore satisfies a minorization inequality $M(m_0, s, \nu)$), a *splitting technique*, introduced in [28, 29], allows us to extend in some sense the probabilistic structure of \mathbf{X} in order to artificially construct an atomic chain (named the *split chain* and denoted by $\tilde{\mathbf{X}}$) that inherits the communication and stochastic stability properties from \mathbf{X} . One of the main results derived from this construction is the fact that every Harris recurrent Markov chain admits a unique (up to multiplicative constant) invariant measure (see Proposition 10.4.2 in [27]), that is, a measure π such that

$$\pi(B) = \int P(x, B) d\pi(x). \quad \forall B \in \mathcal{E}.$$

The invariant measure is finite if and only if $\mathbb{E}_\alpha \tau_\alpha < +\infty$, in this case we say the chain is *positive recurrent*, otherwise, we say the chain is *null recurrent*. A null recurrent chain is called β -null recurrent (c.f. Definition 3.2 in [24]) if there exists a small nonnegative function h , a probability measure λ , a constant $\beta \in (0, 1)$ and a slowly varying function L_h such that

$$\mathbb{E}_\lambda \left(\sum_{t=0}^n h(X_t) \right) \sim \frac{1}{\Gamma(1+\beta)} n^\beta L_h(n) \quad \text{as } n \rightarrow \infty.$$

As argued in [24], is not a too severe restriction to assume $m_0 = 1$. Therefore, throughout this paper we assume that \mathbf{X} satisfies the minorization inequality $M(1, s, \nu)$, i.e, there exist a measurable function s and a probability measure ν such that $0 \leq s(x) \leq 1$, $\int_E s(x) d\nu(x) > 0$ and

$$(5) \quad P(x, A) \geq s(x) \nu(A).$$

REMARK 2.1. The extensions to the case where $m_0 > 1$ of the results that will be presented in this paper can be carried out (although they involve some complicated notations/proofs) using the m -skeleton or the resolvent chains, as described in [9, 10] and Chapter 17 of [27]. However, they are not treated in this paper.

The following theorem is a compendium of the main properties of Harris's recurrent Markov chains that will be used throughout the paper. Among other things, it shows that the asymptotic behavior of $T(n)$ is similar to the function $u(n)$ defined as

$$(6) \quad u(n) = \begin{cases} n, & \text{if } \mathbf{X} \text{ is positive recurrent} \\ n^\beta L(n), & \text{if } \mathbf{X} \text{ is } \beta\text{-null recurrent} \end{cases}.$$

THEOREM 2.1. *Suppose \mathbf{X} is a Harris recurrent, irreducible Markov chain, with initial measure λ , that satisfies the minorization condition (5). Let $T(n)$ be the number of complete regenerations until time n of the split chain $\check{\mathbf{X}}$, let $C \in \mathcal{E}$ be a small set and π be an invariant measure for \mathbf{X} . Then,*

1. $0 < \pi(C) < +\infty$.
2. $\frac{T(n)}{T_n(C)}$ converges almost surely to a positive constant.
3. $\frac{T(n)}{u(n)}$ converges almost surely to a positive constant if \mathbf{X} is positive recurrent and converges in distribution to a Mittag-Leffler¹ random variable with index β if \mathbf{X} is β -null recurrent.

3. Consistency. The aim of the section is to show that for an arbitrary x_0 in the support of f_0 , the LSE $\hat{f}_n(x_0)$ is consistent. We make the following assumptions on the processes $\mathbf{X} = \{X_t\}$ and $\mathbf{W} = \{W_t\}$.

(A1) \mathbf{X} is a Harris recurrent Markov chain whose kernel $P(x, A)$ satisfies the minorization condition (5).

Let $\mathcal{F}_n = \sigma(\{X_0, \dots, X_n\})$ be sigma algebra generated by the chain \mathbf{X} up to time n .

(A2) For each n , the random variables W_1, \dots, W_n are conditionally independent given \mathcal{F}_n , $\mathbb{E}(W_t | \mathcal{F}_n) = 0$ and $\text{Var}(W_t | \mathcal{F}_n) \leq \sigma^2$ for some $\sigma > 0$.

It follows from Assumption (A1) that the Markov Chain \mathbf{X} admits a unique (up to a multiplicative constant) σ -finite invariant measure π . Let C be a set such that $0 < \pi(C) < \infty$ and $x_0 \in C$. We denote by F_n the process defined by

$$(7) \quad F_n(y) = \frac{1}{T_n(C)} \sum_{i=1}^{T_n(C)} \mathbb{I}\{X_{\sigma_C(i)} \leq y\} = \frac{1}{T_n(C)} \sum_{t=0}^n \mathbb{I}_C\{X_t \leq y\}$$

¹The Mittag-Leffler distribution with index β is a non-negative continuous distribution, whose moments are given by

$$\mathbb{E}\left(M_\beta^m(1)\right) = \frac{m!}{\Gamma(1+m\beta)} \quad m \geq 0.$$

See (3.39) in [24] for more details.

for all $y \in \mathbb{R}$, which is a localized version of the empirical distribution function of the X_t 's. It is proved in Lemma 5.1 that F_n converges almost surely to the distribution function F supported on C and defined by

$$(8) \quad F(y) = \frac{\pi(C \cap (-\infty, y])}{\pi(C)}.$$

Our next two assumptions guarantee that there is a compact C , that is a small set and contains x_0 as an interior point. Sets like this can be found under very wide conditions (cf [15]).

(A3) There is $\delta = \delta(x_0)$ such that the set $C = [x_0 - \delta, x_0 + \delta]$ is *small*.

(A4) x_0 belongs to the interior of the support of X_t .

Notice that by part 1 of Theorem 2.1, (A3) guarantees that $\pi(C)$ is finite and positive, and hence, F is properly defined.

In addition to the assumptions on the processes $\{X_t\}$ and $\{W_t\}$, we need smoothness assumptions on F and on f_0 . In particular, we will assume that F and f_0 are continuous and strictly monotone in C . This implies that f_0 and F are invertible in C , so we can find neighborhoods of $f_0(x_0)$ and $F(x_0)$ respectively, over which the inverse functions are uniquely defined. We denote by f_0^{-1} and F^{-1} respectively the inverses of f_0 and F over such a neighborhood of $f_0(x_0)$ and $F(x_0)$ respectively. The function f_0 is assumed to be monotone on its whole support.

(A5) F is locally continuous and strictly increasing in the sense that for all x' in C , for all $\varepsilon > 0$, there exists $\gamma > 0$ such that $|F^{-1}(u) - x'| > \gamma$ for all u such that $|u - F(x')| \geq \varepsilon$.

(A6) f_0 is non-increasing, and f_0 is locally strictly decreasing in the sense that for all x' in C , for all $\varepsilon > 0$, there exists $\gamma > 0$ such that $|f_0(x') - f_0(y)| > \gamma$ for all y such that $|y - x'| \geq \varepsilon$.

(A7) f_0 continuous in C .

Assumptions (A1), (A3) and (A5) ensure that X_t visits infinitely many times any small enough neighborhood of x_0 with probability 1, and guarantee that x_0 is not at the boundary of the recurrent states. Assumptions (A1) and (A3) and Lemma 3.2 in [24] imply that $T_n(C) \rightarrow \infty$ almost surely.

THEOREM 3.1. *Suppose that assumptions (A1)-(A7) are satisfied. Then, as $n \rightarrow \infty$, one has*

$$(9) \quad \widehat{f}_n(x_0) = f_0(x_0) + o_P(1),$$

and

$$(10) \quad \widehat{f}_n^{-1}(f_0(x_0)) = x_0 + o_P(1).$$

4. Rates of convergence. To compute rates of convergence, we need stronger assumptions than for consistency. We replace assumption (A1) for the following stronger version

(B1) $\{X_t\}$ is a positive or β -null recurrent, aperiodic and irreducible Markov Chain whose kernel $P(x, A)$ satisfies the minorization condition (5).

We replace, (A5), (A6) and (A7), for the following slightly more restrictive assumption

(B2) The function f_0 is non-increasing, the functions f_0 and F_C are differentiable in C , and the derivatives F'_C and f'_0 are bounded, in absolute value, above and away from zero in C .

Let λ be the initial measure of \mathbf{X} . Our next hypothesis imposes some control on the behavior of the chain in the first regenerative block.

(B3) There exists a constant K and a neighborhood V of 0, such that

$$\mathbb{E}_\lambda \left(\sum_{t=0}^{\tau_\alpha} (\mathbb{I}_C \{X_t \leq x_0 + \gamma\} - \mathbb{I}_C \{X_t \leq x_0 - \gamma\}) \right) \leq K\gamma \quad \forall \gamma \in V.$$

Assumption (B3) is satisfied if we assume that the initial measure of the chain is the small measure used for the construction of the split chain (see equation 4.16c in [29]). In the positive recurrent case, taking λ equal to the unique invariant probability measure of the chain also satisfies (B3).

And finally, we need to control the number of times the chain visits C in a regeneration block.

(B4) $\ell_C(\mathcal{B}_1) = \sum_{t \in \mathcal{B}_1} \mathbb{I}_C \{X_t\}$ has finite second moment.

THEOREM 4.1. *Assume that (A2), (A3), (A4), (B1), (B2), (B3) and (B4) hold. Then, as $n \rightarrow \infty$, one has*

$$(11) \quad \hat{f}_n(x_0) = f_0(x_0) + O_P(u(n)^{-1/3}),$$

with $u(n)$ as defined in (6).

The rate $u(n)$ comes from Lemmas 5.3 and 6.7, and as it can be seen from Theorem 2.1, it is a deterministic approximation of $T(n)$. Note that in the positive recurrent case, $u(n) = n$, hence we obtain the same rate $n^{-1/3}$ as in the i.i.d. case [18, Chapter 2]. In the β -null recurrent case, however, the rate of convergence is $n^{\beta/3} L^{1/3}(n)$ which is slower than the usual rate. This is due to the null-recurrence of the chain because it takes longer for the process to return to a neighborhood of the point x_0 and it is these points in the neighborhood of x_0 which are used in nonparametric estimation.

5. Localized Markov chains. Given the localized nature of our approach, in this section, we present some results that are particularly useful in this scenario. These results are well known for positive recurrent chains but are new in the null recurrent case. The detailed proofs of these results can be found in Section 2 of the Supplementary material [4].

The first result can be viewed as an extension of the Glivenko-Cantelli theorem to the localized scenario.

LEMMA 5.1. *Assume that (A1) and (A3) hold. Then, there exists a stationary σ -finite measure π , and F defined by (8), such that,*

$$(12) \quad \sup_{y \in \mathbb{R}} |F_n(y) - F(y)| \rightarrow 0 \quad a.s.$$

as $n \rightarrow \infty$. If (A5) is also satisfied, then, for all sufficiently small $\varepsilon > 0$, as $n \rightarrow \infty$ we have

$$(13) \quad \sup_{|p - F(x_0)| \leq \varepsilon} |F_n^{-1}(p) - F^{-1}(p)| \rightarrow 0 \quad a.s.$$

Our next result (Lemma 5.2), which is an extension of Lemma 2 in [5] to the localized β -null recurrent case, deals with the properties of classes of functions defined over the regeneration blocks. Before presenting the result, we need some machinery.

Recall that $E \subseteq \mathbb{R}$ denotes the state space of X . Define $\widehat{E} = \bigcup_{k=1}^{\infty} E^k$ (i.e. the set of finite subsets of E) and let the *localized occupation measure* M_C be given by

$$M_C(B, dy) = \sum_{x \in B \cap C} \delta_x(y), \quad \text{for every } B \in \widehat{E}.$$

The function that gives the size of the localized blocks is $\ell_C : \widehat{E} \rightarrow \mathbb{N}$

$$\ell_C(B) = \int M_C(B, dy), \quad \text{for every } B \in \widehat{E}.$$

Let $\widehat{\mathcal{E}}$ denote the smallest σ -algebra formed by the elements of the σ -algebras \mathcal{E}^k , $k \geq 1$, where \mathcal{E}^k stands for the classical product σ -algebra. Let \widehat{Q} denote a probability measure on $(\widehat{E}, \widehat{\mathcal{E}})$. If $B(\omega)$ is a random variable with distribution Q' , then $M_C(B(\omega), dy)$ is a random measure, i.e., $M_C(B(\omega), dy)$ is a (counting) measure on (E, \mathcal{E}) , almost surely, and for every $A \in \mathcal{E}$, $M_C(B(\omega), A) = \int_A M_C(B(\omega), dy)$ is a measurable random variable (valued in \mathbb{N}). Henceforth $\ell(B(\omega)) \times \int f(y) M_C(B(\omega), dy)$ is a random variable and, provided that $\widehat{Q}(\ell^2) < \infty$, the map Q_C , defined by

$$(14) \quad Q_C(A) = E_{\widehat{Q}} \left(\ell_C(B) \times \int_A M_C(B, dy) \right) / E_{\widehat{Q}}(\ell_C^2), \quad \text{for every } A \in \mathcal{E},$$

is a probability measure on (E, \mathcal{E}) . The notation E_{Q_C} stands for the expectation with respect to the underlying measure Q_C . Introduce the following notations: for any function $g : E \rightarrow \mathbb{R}$, let $\widehat{g}_C : \widehat{E} \rightarrow \mathbb{R}$ be given by

$$(15) \quad \widehat{g}_C(B) = \int g(y) M_C(B, dy) = \sum_{x \in B \cap C} g(x) = \sum_{x \in B} g_C(x),$$

and for any class \mathcal{G} of real-valued functions defined on E , denote the localized version of the sums on the blocks by $\widehat{\mathcal{G}}_C = \{\widehat{g}_C : g \in \mathcal{G}\}$.

Notice that, for any function g ,

$$(16) \quad \mathbb{E}_{Q_C}(g) = \frac{\mathbb{E}_{\widehat{Q}}(\ell_C(B) \times \int g(y) M_C(B, dy))}{\mathbb{E}_{\widehat{Q}}(\ell_C^2)} = \frac{\mathbb{E}_{\widehat{Q}}(\ell_C(B) \widehat{g}_C(B))}{\mathbb{E}_{\widehat{Q}}(\ell_C^2)}.$$

LEMMA 5.2. *Let \widehat{Q} be a probability measure on $(\widehat{E}, \widehat{\mathcal{E}})$ such that $0 < \|\ell_C\|_{L^2(\widehat{Q})} < \infty$ and \mathcal{G} be a class of measurable real-valued functions defined on (E, \mathcal{E}) . Then we have, for every $0 < \varepsilon < \infty$,*

$$\mathcal{N}(\varepsilon \|\ell_C\|_{L^2(\widehat{Q})}, \widehat{\mathcal{G}}_C, L^2(\widehat{Q})) \leq \mathcal{N}(\varepsilon, \mathcal{G}, L^2(Q)),$$

where Q is given in (14). Moreover, if \mathcal{G} belongs to the Vapnik–Chervonenkis (VC) class of functions with constant envelope U and characteristic (C, v) , then $\widehat{\mathcal{G}}_C$ is VC with envelope $U\ell_C$ and characteristic (C, v) .

REMARK 5.1. For a probability measure μ , and a class of functions \mathcal{H} , the covering number $\mathcal{N}(\varepsilon, \mathcal{H}, L^r(\mu))$ is the minimum number of $L^r(\mu)$ ε -balls needed to cover \mathcal{H} . For more details about this concept and the VC class of functions, see [25].

To put into perspective Lemma 5.2, consider a class of bounded functions \mathcal{G} that is VC with finite envelope. Lemma 5.2 tells us that the class of unbounded functions $\widehat{\mathcal{G}}_C$ is also VC. If we also have that (B4) holds, then Theorem 2.5 in [25] tells us that $\widehat{\mathcal{G}}_C$ is a Donsker class. A reasoning like this is used in the proof of the following result, which is a stronger version of Lemma 5.1 under assumptions (B1) and (B2) and has some interest on its own.

LEMMA 5.3. *Assume that (B1), (B2), (A3), (A4) and (B4) hold. Then, for all sufficiently small $\varepsilon > 0$ we have,*

$$(17) \quad T_n(C) \sup_{|y-x_0| \leq \varepsilon} |F_n(y) - F(y)|^2 = O_p(1)$$

when n goes to $+\infty$. If (B2) is also satisfied, as $n \rightarrow \infty$ we have

$$(18) \quad T_n(C) \sup_{|p-F(x_0)| \leq \varepsilon} |F_n^{-1}(p) - F^{-1}(p)|^2 = O_p(1).$$

6. Proofs. In this section, we provide the proof of Theorems 3.1 and 4.1. These proofs make use of several intermediate lemmas, whose proofs can be found in Sections 7 and 8.

6.1. *Proof of Theorem 3.1.* Recall that we consider the piecewise constant and left-continuous LSE \widehat{f}_n , that is constant on every interval $(Y_{k-1}, Y_k]$, $k = 2, \dots, m$ and also on $(-\infty, Y_1]$ and on $[Y_m, \infty)$. With $\delta > 0$ fixed, we denote by $T_n(C)$ the number of times the Markov Chain \mathbf{X} visits the set $C := [x_0 - \delta, x_0 + \delta]$ until time n :

$$(19) \quad T_n(C) = \sum_{t=0}^n \mathbb{I}\{X_t \in C\}.$$

Let $l_k = \sum_{t=1}^n \mathbb{I}_C\{X_t \leq Y_k\}$ for all $k \in \{1, \dots, m\}$ and $l_0 = 0$.

Our aim is to provide a characterization of $\widehat{f}_n(x_0)$. Recall from (7) that the localized empirical distribution function F_n is defined as

$$F_n(y) = \frac{1}{T_n(C)} \sum_{i=0}^{T_n(C)} \mathbb{I}\{X_{\sigma_C(i)} \leq y\} = \frac{1}{T_n(C)} \sum_{t=0}^n \mathbb{I}_C\{X_t \leq y\}$$

for $y \in \mathbb{R}$. F_n is 0 on $(-\infty, Y_1)$, so, with an arbitrary random variable $Y_0 < Y_1$ we have $F_n(y) = F_n(Y_0) = 0$ for all $y < Y_1$. Let \mathcal{K} be the set

$$(20) \quad \mathcal{K} := \{F_n(Y_k), k = 0, \dots, m\}$$

and let Λ_n be the continuous piecewise-linear process on $[F_n(Y_0), F_n(Y_m)]$ with knots at the points in \mathcal{K} and values

$$(21) \quad \Lambda_n(F_n(Y_k)) = \frac{1}{T_n(C)} \sum_{t=0}^n Z_t \mathbb{I}_C\{X_t \leq Y_k\}$$

at the knots. The characterization of \widehat{f}_n in Lemma 6.2 involves the least concave majorant of Λ_n . Note that we use $T_n(C)$ as a normalization in the definitions of the processes F_n and Λ_n since this choice ensures that F_n and Λ_n converge to fixed functions, see Lemma 5.1.

LEMMA 6.1. *For all $y \in [F_n(Y_0), F_n(Y_m)]$,*

$$\Lambda_n(y) = L_n(y) + M_n(y),$$

where,

$$(22) \quad L_n(y) = \int_0^y f \circ F_n^{-1}(u) du,$$

and M_n is a piece-wise linear processes with knots at $F_n(Y_k)$ for $k \in \{0, \dots, m\}$ such that

$$M_n(F_n(Y_k)) = \frac{1}{T_n(C)} \sum_{t=0}^n W_t \mathbb{I}_C \{X_t \leq Y_k\}.$$

Moreover, M_n can be written as

$$(23) \quad M_n(y) = \begin{cases} 0 & , \text{ if } y = 0 \\ R_n^j(y) + M_n^j & , \text{ otherwise} \end{cases}$$

where,

$$(24) \quad M_n^j = M_n(F_n(Y_j)) = \frac{1}{T_n(C)} \sum_{t=0}^n W_t \mathbb{I}_C \{X_t \leq Y_j\},$$

$$(25) \quad R_n^j(y) = \frac{\sum_{t=0}^n W_t \mathbb{I}_C \{X_t = Y_{j+1}\}}{l_{j+1} - l_j} (y - F_n(Y_j)),$$

and j is such that $Y_{j+1} = F_n^{-1}(y)$.

In the next lemma, we give an alternative characterization of the monotone nonparametric LSE \hat{f}_n at the observation points Y_1, \dots, Y_m .

LEMMA 6.2. *Let $C = [x - \delta, x + \delta]$ for some fixed $\delta > 0$. Let $\hat{\lambda}_n$ be the left-hand slope of the least concave majorant of Λ_n . Then,*

$$(26) \quad \hat{f}_n(Y_k) = \hat{\lambda}_n \circ F_n(Y_k), \quad \forall k \in \{1, \dots, m\}.$$

with probability 1 for n big enough.

We consider below the generalized inverse function of \hat{f}_n since it has a more tractable characterization than \hat{f}_n itself. To this end, let us define precisely the generalized inverses of all processes of interest. Since $\hat{\lambda}_n$ is a non-increasing left-continuous step function on $(F_n(Y_0), F_n(Y_m)]$ that can have jumps only at the points $F_n(Y_k)$, $k \in \{1, \dots, m\}$, we define its generalized inverse $\hat{U}_n(a)$, for $a \in \mathbb{R}$, as the greatest $y \in (F_n(Y_0), F_n(Y_m)]$ that satisfies $\hat{\lambda}_n(y) \geq a$, with the convention that the supremum of an empty set is $F_n(Y_0)$. Then for every $a \in \mathbb{R}$ and $y \in (F_n(Y_0), F_n(Y_m)]$, one has

$$(27) \quad \hat{\lambda}_n(y) \geq a \text{ if and only if } \hat{U}_n(a) \geq y.$$

Likewise, since \hat{f}_n is a left-continuous non-increasing step function on \mathbb{R} that can have jumps only at the observation times $Y_1 < \dots < Y_m$, we define the generalized inverse $\hat{f}_n^{-1}(a)$, for $a \in \mathbb{R}$, as the greatest $y \in [Y_0, Y_m]$ that satisfies $\hat{f}_n(y) \geq a$, with the convention that the supremum of an empty set is Y_0 . We then have

$$(28) \quad \hat{f}_n(y) \geq a \text{ if and only if } \hat{f}_n^{-1}(a) \geq y$$

for all $a \in \mathbb{R}$ and $y \in (Y_0, Y_m]$. On the other hand, since F_n is a right-continuous non-decreasing step function on \mathbb{R} with range $[F_n(Y_0), F_n(Y_m)]$, we define the generalized inverse $F_n^{-1}(a)$, for $a \leq F_n(Y_m)$, as the smallest $y \in [Y_0, Y_m]$ which satisfies $F_n(y) \geq a$. Note that the infimum is achieved for all $a \leq F_n(Y_m)$. We then have

$$(29) \quad F_n(y) \geq a \text{ if and only if } F_n^{-1}(a) \leq y$$

for all $a \leq F_n(Y_m)$ and $y \in [Y_0, Y_m]$, and thanks to Lemma 6.2 we have

$$(30) \quad \widehat{f}_n^{-1} = F_n^{-1} \circ \widehat{U}_n$$

on \mathbb{R} . Moreover, one can check that

$$(31) \quad \widehat{U}_n(a) = \operatorname{argmax}_{p \in [F_n(Y_0), F_n(Y_m)]} \{\Lambda_n(p) - ap\}, \text{ for all } a \in \mathbb{R},$$

where argmax denotes the greatest location of maximum (which is achieved on the set \mathcal{K} in (20)). Thus, the inverse process \widehat{U}_n is a location process that is more tractable than \widehat{f}_n and $\widehat{\lambda}_n$ themselves. A key idea in the following proofs is to derive properties of \widehat{U}_n from its argmax characterization (31), then, to translate these properties to \widehat{f}_n^{-1} thanks to (30), and finally to translate them to \widehat{f}_n thanks to (28).

To go from \widehat{U}_n to \widehat{f}_n^{-1} using (30) requires to approximate F_n^{-1} by a fixed function. Hence, in the sequel, we are concerned by the convergence of the process F_n given in (7), where $\delta > 0$ is chosen sufficiently small, and by the convergence of the corresponding inverse function F_n^{-1} .

It is stated in Lemma 5.1 that under (A1) and (A3), F_n converges to a fixed distribution function F that depends on C , hence on δ . If, moreover, F is strictly increasing in C , then we can find a neighborhood of $F(x_0)$ over which the (usual) inverse function F^{-1} is uniquely defined, and F_n^{-1} converges to F^{-1} .

In the following lemma, we show that $F(x_0)$ belongs to the domain of Λ_n with probability tending to one as $n \rightarrow \infty$.

LEMMA 6.3. *Assume that (A1), (A3), (A4) and (A5) hold. Then, we can find $\varepsilon > 0$ such that the probability that $Y_1 + \varepsilon \leq x_0 \leq Y_m - \varepsilon$ tends to one as $n \rightarrow \infty$. Moreover, the probability that $F_n(Y_1) \leq F(x_0) \leq F_n(Y_m)$ tends to one as $n \rightarrow \infty$.*

We will also need to control the noise $\{W_t\}$. The following lemma shows that the noise is negligible under our assumptions.

LEMMA 6.4. *Assume that (A1) and (A2) hold. Let $\mathcal{F}_n = \sigma(\{X_1, \dots, X_n\})$. Then,*

$$\sum_{t=0}^n W_t \mathbb{I}_C \{X_t = A_n\} = o_P(T_n(C)),$$

and

$$\sup_{u > A_n} \left| \sum_{t=0}^n W_t \mathbb{I}_C \{X_t \in (A_n, u]\} \right| = o_P(T_n(C)).$$

for any sequence of random variables A_n , independent of the process $\{W_t\}$, that is adapted to the filtration $\{\mathcal{F}_n\}$.

With the above lemmas, we can prove convergence of \widehat{U}_n to $F(x_0)$ given by (31).

LEMMA 6.5. *Suppose that assumptions (A1)-(A7) are satisfied. Then, as $n \rightarrow \infty$, one has*

$$(32) \quad \widehat{U}_n(f_0(x_0)) = F(x_0) + o_P(1).$$

Now we proceed to the proof of (10). Fix $\varepsilon > 0$ arbitrarily small. It follows from (30) and (29) that

$$\begin{aligned} \mathbb{P}\left(\widehat{f}_n^{-1}(a) > x_0 + \varepsilon\right) &\leq \mathbb{P}\left(F_n^{-1} \circ \widehat{U}_n(a) > x_0 + \varepsilon\right) \\ &\leq \mathbb{P}\left(\widehat{U}_n(a) \geq F_n(x_0 + \varepsilon)\right) \\ &\leq \mathbb{P}\left(\widehat{U}_n(a) \geq F(x_0 + \varepsilon) - K_n\right), \end{aligned}$$

where

$$K_n = \sup_{|y-x_0| \leq \varepsilon} |F_n(y) - F(y)|.$$

With $\nu := F(x_0 + \varepsilon) - F(x_0)$, we obtain

$$\mathbb{P}\left(\widehat{f}_n^{-1}(a) > x_0 + \varepsilon\right) \leq \mathbb{P}\left(\widehat{U}_n(a) \geq F(x_0) + \nu - K_n\right),$$

and ν is strictly positive since F is strictly increasing in the neighborhood of x_0 . Hence, it follows from (12) that for sufficiently small $\varepsilon > 0$ one has

$$\mathbb{P}\left(\widehat{f}_n^{-1}(a) > x_0 + \varepsilon\right) \leq \mathbb{P}\left(\widehat{U}_n(a) \geq F(x_0) + \nu/2\right) + o(1),$$

so it follows from (32) that the probability that $\widehat{f}_n^{-1}(a) > x_0 + \varepsilon$ tends to zero as $n \rightarrow \infty$. Similarly, the probability that $\widehat{f}_n^{-1}(a) < x_0 - \varepsilon$ tends to zero as $n \rightarrow \infty$ so we conclude that the probability that $|\widehat{f}_n^{-1}(a) - x_0| > \varepsilon$ tends to zero as $n \rightarrow \infty$. This completes the proof of (10).

To prove (9), fix $\varepsilon > 0$ sufficiently small so that F and f_0 are continuous and strictly increasing in the neighborhood of $x' := f_0^{-1}(f_0(x_0) + \varepsilon)$. Equation (10) shows that

$$(33) \quad \widehat{f}_n^{-1}(f_0(x_0) + \varepsilon) = f_0^{-1}(f_0(x_0) + \varepsilon) + o_P(1),$$

as $n \rightarrow \infty$. Now, it follows from the switch relation (27) that

$$(34) \quad \begin{aligned} \mathbb{P}\left(\widehat{f}_n(x_0) > f_0(x_0) + \varepsilon\right) &\leq \mathbb{P}\left(\widehat{f}_n^{-1}(f_0(x_0) + \varepsilon) \geq x\right) \\ &\leq \mathbb{P}\left(\widehat{f}_n^{-1}(f_0(x_0) + \varepsilon) \geq f_0^{-1}(f_0(x_0) + \varepsilon) + \nu\right), \end{aligned}$$

where $\nu := x - f_0^{-1}(f_0(x_0) + \varepsilon) > 0$. It follows from (33) that the probability on the right-hand side tends to zero as $n \rightarrow \infty$. Hence, the probability on the left-hand side tends to zero as well as $n \rightarrow \infty$.

Similarly, the probability that $\widehat{f}_n(x_0) < f_0(x_0) - \varepsilon$ tends to zero as $n \rightarrow \infty$ so we conclude that the probability that $|\widehat{f}_n(x_0) - f_0(x_0)| > \varepsilon$ tends to zero as $n \rightarrow \infty$. This completes the proof of Theorem 3.1.

6.2. Proof of Theorem 4.1. The proof of Theorem 4.1, uses similar ideas as the ones used in the proof of Theorem 3.1 but under stronger assumptions (and therefore using stronger lemmas).

The first intermediate result is the following stronger version of Lemma 6.4.

LEMMA 6.6. *Assume that (A2), (A3), (A4), (B1), (B2) and (B3) hold. Then, there exists $K > 0$, $\gamma_0 > 0$ that do not depend on n and $N_{\gamma_0} \in \mathbb{N}$, such that for all $\gamma \in [0, \gamma_0]$ and $n \geq N_{\gamma_0}$ one has*

$$(35) \quad \mathbb{E}_\lambda \left(\sup_{|y-x_0| \leq \gamma} \left| \sum_{t=0}^n W_t (\mathbb{I}_C\{X_t \leq y\} - \mathbb{I}_C\{X_t \leq x_0\}) \right|^2 \right) \leq Ku(n)\gamma$$

$$(36) \quad \mathbb{E}_\lambda \left(\sup_{|y-x_0| \leq \gamma} \left| \sum_{t=0}^n W_t \mathbb{I}_C\{X_t = y\} \right|^2 \right) \leq Ku(n)\gamma$$

Then, we need to quantify how well we can approximate $T_n(C)$ by $u(n)$.

LEMMA 6.7. *Assume that (B1) and (A3) hold. Then we have*

a) *As $n \rightarrow \infty$ we have*

$$\frac{u(n)}{T_n(C)} = O_P(1).$$

b) *Let α and η be positive constants, then there positive exists constants N_η , \underline{c}_η and \bar{c}_η , such that*

$$\mathbb{P} \left(\left(\frac{T_n(C)}{a(n)} \right)^\alpha \in [\underline{c}_\eta, \bar{c}_\eta] \right) \geq 1 - \eta, \quad \forall n \geq N_\eta.$$

With the above lemmas (including Lemma 5.3 and the ones used in Section 6.1), we can obtain the rate of convergence of \widehat{U}_n given by (31).

LEMMA 6.8. *Assume that (A2), (A3), (A4), (B1), (B2), (B3) and (B4) hold. Then, as $n \rightarrow \infty$, one has*

$$(37) \quad \widehat{U}_n(f_0(x_0)) = F(x_0) + O_P \left(u(n)^{-1/3} \right),$$

and

$$(38) \quad \widehat{f}_n^{-1}(f_0(x_0)) = x + O_P \left(u(n)^{-1/3} \right).$$

Inspecting the proof of Lemma 6.8, one can see that the convergences in (37) and (38) hold in a uniform sense in the neighborhood of x_0 . More precisely, there exists $\gamma > 0$, independent on n , such that for all $\eta > 0$ we can find $K_1 > 0$ such that

$$\sup_{|a-f_0(x_0)| \leq \gamma} \mathbb{P} \left(\left| \widehat{U}_n(a) - F \circ f_0^{-1}(a) \right| > K_1 u(n)^{-1/3} \right) \leq \eta$$

and

$$\sup_{|a-f_0(x_0)| \leq \gamma} \mathbb{P} \left(\left| \widehat{f}_n^{-1}(a) - f_0^{-1}(a) \right| > K_1 u(n)^{-1/3} \right) \leq \eta.$$

Let $\varepsilon = K_1 u(n)^{-1/3}$ where $K_1 > 0$ does not depend on n , and recall (34) where $\nu = x - f_0^{-1}(f_0(x_0) + \varepsilon) > 0$. It follows from the assumption (B2) that f_0^{-1} has a derivative that is bounded in sup-norm away from zero in a neighborhood of $f_0(x_0)$. Hence, it follows from the Taylor expansion that there exists $K_2 > 0$ that depends only on f_0 such that $\nu \geq K_2 \varepsilon$,

provided that n is sufficiently large to ensure that $f_0(x_0) + \varepsilon$ belongs to this neighborhood of $f_0(x_0)$. Hence,

$$\begin{aligned} \mathbb{P}\left(\widehat{f}_n(x_0) > f_0(x_0) + \varepsilon\right) &\leq \mathbb{P}\left(\widehat{f}_n^{-1}(f_0(x_0) + \varepsilon) \geq f_0^{-1}(f_0(x_0) + \varepsilon) + K_2\varepsilon\right) \\ &\leq \sup_{|a-f_0(x_0)| \leq \gamma} \mathbb{P}\left(\left|\widehat{f}_n^{-1}(a) - f_0^{-1}(a)\right| > K_2 K_1 u(n)^{-1/3}\right), \end{aligned}$$

provided that n is sufficiently large to ensure that $f_0(x_0) + \varepsilon$ belongs to the above neighborhood of $f_0(x_0)$, and that $\gamma \geq C u(n)^{-1/3}$. For fixed $\eta > 0$ one can choose $K_2 > 0$ such that the probability on the right-hand side of the previous display is smaller than or equal to η and therefore,

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(\widehat{f}_n(x_0) > f_0(x_0) + K_2 u(n)^{-1/3}\right) \leq \eta.$$

Similarly, for all fixed $\eta > 0$, one can find K_3 that does not depend on n such that

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(\widehat{f}_n(x_0) < f_0(x_0) - K_3 u(n)^{-1/3}\right) \leq \eta.$$

Hence, for all fixed $\eta > 0$, there exists $K > 0$ that independent of n such that

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(\left|\widehat{f}_n(x_0) - f_0(x_0)\right| > K u(n)^{-1/3}\right) \leq \eta.$$

This completes the proof of Theorem 4.1.

7. Technical proofs for Section 6.1. Proof of Lemma 6.1. Combining (21) and (1) yields

$$\Lambda_n(F_n(Y_k)) = \frac{1}{T_n(C)} \sum_{t=0}^n f_0(X_t) \mathbb{I}_C\{X_t \leq Y_k\} + \frac{1}{T_n(C)} \sum_{t=0}^n W_t \mathbb{I}_C\{X_t \leq Y_k\}.$$

The first term on the right-hand side of the previous display can be rewritten as follows:

$$\begin{aligned} \frac{1}{T_n(C)} \sum_{t=0}^n f_0(X_t) \mathbb{I}_C\{X_t \leq Y_k\} &= \frac{1}{T_n(C)} \sum_{j=1}^m f_0(Y_j) (l_j - l_{j-1}) \mathbb{I}_C\{Y_j \leq Y_k\} \\ &= \sum_{j=1}^k \int_{l_{j-1}/T_n(C)}^{l_j/T_n(C)} f_0 \circ F_n^{-1}(u) du, \end{aligned}$$

using that $F_n^{-1}(u) = Y_j$ for all $u \in (l_{j-1}/T_n(C), l_j/T_n(C)]$. Hence, for all k in $\{0, \dots, m\}$

$$(39) \quad \Lambda_n(F_n(Y_k)) = \int_0^{l_k/T_n(C)} f_0 \circ F_n^{-1}(u) du + \frac{1}{T_n(C)} \sum_{t=0}^n W_t \mathbb{I}_C\{X_t \leq Y_k\}.$$

Combining (39) with the piece-wise linearity of Λ_n yields

$$\Lambda_n(F_n(Y_k)) = L_n(F_n(Y_k)) + M_n(F_n(Y_k)),$$

where L_n and M_n are piece-wise linear processes with knots at $F_n(Y_k)$ for k in $\{0, \dots, m\}$ and such that

$$L_n(F_n(Y_k)) = \int_0^{l_k/T_n(C)} f_0 \circ F_n^{-1}(u) du$$

and

$$M_n(F_n(Y_k)) = \frac{1}{T_n(C)} \sum_{t=0}^n W_t \mathbb{I}_C\{X_t \leq Y_k\}.$$

In order to ease the notation, we will write $F_n^i = F_n(Y_i)$, $L_n^i = L_n(F_n(Y_i))$ and $M_n^i = M_n(F_n(Y_i))$. Let $y \in (F_n(Y_0), F_n(Y_m)]$, take j such that $Y_{j+1} = F_n^{-1}(y)$, then $F_n(Y_j) < y \leq F_n(Y_{j+1})$. With this notation,

$$L_n(y) = \frac{L_n^{j+1} - L_n^j}{F_n^{j+1} - F_n^j} (y - F_n^j) + L_n^j,$$

$$M_n(y) = \frac{M_n^{j+1} - M_n^j}{F_n^{j+1} - F_n^j} (y - F_n^j) + M_n^j.$$

Notice that

$$L_n^{j+1} - L_n^j = \int_{\frac{l_j}{T_n(C)}}^{\frac{l_{j+1}}{T_n(C)}} f_0 \circ F_n^{-1}(u) du = \frac{l_{j+1} - l_j}{T_n(C)} f(Y_{j+1}),$$

$$F_n^{j+1} - F_n^j = \frac{l_{j+1} - l_j}{T_n(C)},$$

therefore,

$$L_n(y) = f_0(Y_{j+1})(y - F_n^j) + L_n^j = \int_{\frac{l_j}{T_n(C)}}^y f_0 \circ F_n^{-1}(u) du + L_n^j = \int_0^y f_0 \circ F_n^{-1}(u) du,$$

which proves (22).

For M_n we have,

$$M_n^{j+1} - M_n^j = \frac{1}{T_n(C)} \sum_{t=0}^n W_t \mathbb{I}_C\{X_t = Y_{j+1}\},$$

then,

$$M_n(y) = \frac{\sum_{t=1}^n W_t \mathbb{I}_C\{X_t = Y_{j+1}\}}{l_{j+1} - l_j} (y - F_n^j) + M_n^j = R_n^j(y) + M_n^j.$$

and this completes the proof.

Proof of Lemma 6.2. By definition, with $l_0 = 0$, and $l_k = \sum_{t=0}^n \mathbb{I}_C\{X_t \leq Y_k\}$ for all $k \in \{1, \dots, m\}$, we have $F_n(Y_k) = al_k$ for all $k \in \{0, \dots, m\}$, where $a = 1/T_n(C)$ and does not depend on k . Moreover,

$$\Lambda_n(F_n(Y_k)) = a \sum_{t=0}^n Z_t \mathbb{I}_C\{X_t \leq Y_k\}$$

Since $\widehat{f}_n(Y_k)$ is the left-hand slope at l_k of the least concave majorant of the set of points in (3), the equality in (26) follows from Lemma 2.1 in [13].

Proof of Lemma 6.3. The first assertion follows from Assumption (A4) and the second immediately follows from the first one by (12) combined with the strict monotonicity of F in C .

Proof of Lemma 6.4. Let $\mathcal{F}_n = \sigma(\{X_0, \dots, X_n\})$ be sigma algebra generated by the chain $\{X_t\}$ up to time n . Denote by $\mathbb{P}_{\mathcal{F}_n}$ the probability conditioned to \mathcal{F}_n . Take $\varepsilon > 0$.

By Chebyshev's inequality, we have

$$\mathbb{P}_{\mathcal{F}_n} \left(\left| \frac{\sum_{t=0}^n W_t \mathbb{I}_C \{X_t = A_n\}}{T_n(C)} \right| > \varepsilon \right) \leq \frac{\sigma^2 \sum_{t=0}^n \mathbb{I}_C \{X_t = A_n\}}{\varepsilon^2 T_n^2(C)} \leq \frac{\sigma^2}{\varepsilon^2 T_n(C)},$$

which implies the first part of the Lemma because $T_n(C) \rightarrow \infty$ with probability 1.

For the second part, let $\gamma_n(u)$ be the number of times the chain visits $(A_n, u] \cap C$ up to time n and $A_n(u) = \{t \leq n : X_t \in (A_n, u] \cap C\} = \{a_1, \dots, a_{\gamma_n(u)}\}$ the times of those visits. Using that $\gamma_n = \sup_{u > A_n} \gamma_n(u) \leq T_n(C)$ and Kolmogorov's inequality (Th 3.1.6, pp 122 in [19]) we obtain,

$$\begin{aligned} \mathbb{P}_{\mathcal{F}_n} \left(\sup_{u > A_n} \left| \frac{\sum_{t=0}^n W_t \mathbb{I}_C \{X_t \in (A_n, u]\}}{T_n(C)} \right| > \varepsilon \right) &= \mathbb{P}_{\mathcal{F}_n} \left(\sup_{u > A_n} \left| \sum_{i=1}^{\gamma_n(u)} \frac{W_{t_{a_i}}}{T_n(C)} \right| > \varepsilon \right) \\ &\leq \mathbb{P}_{\mathcal{F}_n} \left(\sup_{1 \leq k \leq \gamma_n} \left| \sum_{i=1}^k \frac{W_{t_{a_i}}}{T_n(C)} \right| > \varepsilon \right) \\ &\leq \frac{\sigma^2}{\varepsilon^2 T_n(C)}. \end{aligned}$$

which by the same argument as before, implies the second part of the Lemma.

Proof of Lemma 6.5. In the sequel, we set $a = f_0(x_0)$. We begin with the proof of (32).

Fix $\varepsilon > 0$ arbitrarily, and let $\nu > 0$ and $\gamma > 0$ be such that $|F^{-1}(u) - x_0| > \nu$ for all u such that $|u - F(x_0)| \geq \varepsilon/2$, and $|f_0(x_0) - f_0(y)| > \gamma$ for all y such that $|y - x_0| \geq \nu/2$. Note that existence of ν and γ is ensured by assumptions (A5) and (A6).

By Lemma 6.3, we can assume without loss of generality that $F(x_0)$ belongs to the domain $[F_n(Y_1), F_n(Y_m)]$ of Λ_n , since this occurs with probability tending to one. Therefore, we can find $j(x_0)$ such that $Y_{j(x_0)} = F_n^{-1}(F(x_0))$. It follows from the characterization in (31) that the event $E_n^1 := \{\widehat{U}_n(a) > F(x_0) + \varepsilon\}$ is contained in the event that there exists $p \in \mathcal{K}$ such that $p > F(x_0) + \varepsilon$ and

$$\Lambda_n(p) - ap \geq \Lambda_n(F(x_0)) - aF(x_0),$$

where we recall that $a = f_0(x_0)$.

By Lemma 6.1, E_n^1 is contained in the event that there exists $p \in \mathcal{K}$ such that $p > F(x_0) + \varepsilon$ and

$$(40) \quad L_n(p) + M_n(p) - ap \geq L_n(F(x_0)) + M_n(F(x_0)) - aF(x_0)$$

Using (22) in (40) we obtain that E_n^1 is contained in the event that there exists $p \in \mathcal{K}$ such that $p > F(x_0) + \varepsilon$ and

$$\int_{t_0/T_n(C)}^p f_0 \circ F_n^{-1}(u) du + S_n - ap \geq \int_{t_0/T_n(C)}^{F(x_0)} f_0 \circ F_n^{-1}(u) du - aF(x_0),$$

where

$$S_n = \sup_{p > F(x_0) + \varepsilon, p \in \mathcal{K}} \{M_n(p) - M_n(F(x_0))\}.$$

Let j and k such that $Y_{j+1} = F_n^{-1}(F(x_0))$ and $p = F_n(Y_k)$. By equation (23) we have $M_n(p) - M_n(F(x_0)) = M_n^k - M_n^j - R_n^j(F(x_0))$, therefore,

$$\begin{aligned} S_n &= \sup_{\substack{p > F(x_0) + \varepsilon \\ p \in \mathcal{K}}} \left\{ M_n^k - M_n^j \right\} - R_n^j(F(x_0)) \\ &\leq \sup_{\substack{p > F(x_0) + \varepsilon \\ p \in \mathcal{K}}} \left| \frac{1}{T_n(C)} \sum_{t=0}^n W_t (\mathbb{I}_C\{X_t \leq F_n^{-1}(p)\} - \mathbb{I}_C\{X_t \leq F_n^{-1}(F(x_0))\}) \right| \\ &\quad + |R_n^j(F_n(Y_{j+1}))| \\ &\leq \sup_{\substack{p > F(x_0) + \varepsilon \\ p \in \mathcal{K}}} \left| \frac{1}{T_n(C)} \sum_{t=0}^n W_t \mathbb{I}_C\{X_t \in (F_n^{-1}(F(x_0)); F_n^{-1}(p))\} \right| \\ &\quad + \left| \frac{\sum_{t=0}^n W_t \mathbb{I}_C\{X_t = F_n^{-1}(F(x_0))\}}{T_n(C)} \right|. \end{aligned}$$

Hence,

$$\begin{aligned} T_n(C)S_n &\leq \sup_{\substack{p > F(x_0) + \varepsilon \\ p \in \mathcal{K}}} \left| \sum_{t=0}^n W_t \mathbb{I}_C\{X_t \in (F_n^{-1}(F(x_0)); F_n^{-1}(p))\} \right| \\ &\quad + \left| \sum_{t=0}^n W_t \mathbb{I}_C\{X_t = F_n^{-1}(F(x_0))\} \right|. \end{aligned}$$

Therefore, the event E_n^1 is contained in the event that there exists $p > F(x_0) + \varepsilon$ such that

$$\int_{F(x_0)}^p f_0 \circ F_n^{-1}(u) du + S_n \geq a(p - F(x_0)).$$

Now, let E_n^2 be the event that

$$\sup_{|u - F(x_0)| \leq \varepsilon} |F_n^{-1}(u) - F^{-1}(u)| \leq \eta$$

where $\eta \in (0, \nu/4)$ is such that $|f_0(y) - f_0(x_0)| \leq \gamma/2$ for all y such that $|x_0 - y| \leq \eta$. Note that the existence of η is ensured by assumption (A7). Then, it follows from the monotonicity of f_0 and F_n that on E_n^2 ,

$$\int_{F(x_0)}^p f_0 \circ F_n^{-1}(u) du \leq \int_{F(x_0)}^{F(x_0) + \varepsilon/2} f_0(F^{-1}(u) - \eta) du + \int_{F(x_0) + \varepsilon/2}^p f_0(F_n^{-1}(F(x_0) + \varepsilon/2)) du.$$

Hence, it follows from the definitions of η , ν and γ that on E_n^2 ,

$$\begin{aligned} \int_{F(x_0)}^p f_0 \circ F_n^{-1}(u) du &\leq \frac{\varepsilon}{2} f_0(x_0) + \frac{\gamma\varepsilon}{4} + (p - F(x_0) - \varepsilon/2) f_0(F^{-1}(F(x_0) + \varepsilon/2) - \eta) \\ &\leq \frac{\varepsilon}{2} f_0(x_0) + \frac{\gamma\varepsilon}{4} + (p - F(x_0) - \varepsilon/2) f_0(x_0 + \nu/2) \\ &\leq \frac{\varepsilon}{2} f_0(x_0) + \frac{\gamma\varepsilon}{4} + (p - F(x_0) - \varepsilon/2) (f_0(x_0) - \gamma). \end{aligned}$$

This implies that on E_n^2 ,

$$\begin{aligned} \int_{F(x_0)}^p f_0 \circ F_n^{-1}(u) du &\leq a(p - F(x_0)) - (p - F(x_0) - 3\varepsilon/4)\gamma \\ &\leq a(p - F(x_0)) - \varepsilon\gamma/4 \end{aligned}$$

for all $p > F(x_0) + \varepsilon$. Hence, the event $E_n^1 \cap E_n^2$ is contained in the event $\{S_n \geq \varepsilon\gamma/4\}$. Now, on E_n^2 , for all $p > F(x_0) + \varepsilon$ we have

$$\begin{aligned} F_n^{-1}(p) &\geq F_n^{-1}(F(x_0) + \varepsilon) \\ &\geq F^{-1}(F(x_0) + \varepsilon) - \eta \\ &\geq x + \nu - \eta \\ &\geq F_n^{-1}(F(x_0)) + \nu - 2\eta \\ &\geq F_n^{-1}(F(x_0)) + \nu/2, \end{aligned}$$

since $\nu > 4\eta$. Therefore,

$$\begin{aligned} T_n(C)S_n &\leq \sup_{u > F_n^{-1}(F(x_0)) + \nu/2} \left| \sum_{t=0}^n W_t \mathbb{I}_C\{X_t \in (F_n^{-1}(F(x_0)), u]\} \right| + \\ &\quad + \left| \sum_{t=0}^n W_t \mathbb{I}_C\{X_t = F_n^{-1}(F(x_0))\} \right|. \end{aligned}$$

Hence, it follows from Lemma 6.4 that S_n converges in probability to zero as $n \rightarrow \infty$, so that the probability of the event $\{S_n \geq \varepsilon\gamma/4\}$ tends to zero as $n \rightarrow \infty$. It follows from Lemma 5.1 that for ε sufficiently small, the probability of the event E_n^2 tends to one as $n \rightarrow \infty$, so we conclude that the probability of E_n^1 tends to zero as $n \rightarrow \infty$. Similarly, the probability of the event $\{\widehat{U}_n(a) < F(x_0) - \varepsilon\}$ tends to zero as $n \rightarrow \infty$, so that

$$\lim_{n \rightarrow \infty} \mathbb{P}(|\widehat{U}_n(a) - F(x_0)| > \varepsilon) = 0$$

for all $\varepsilon > 0$. This completes the proof of (32).

8. Technical proofs for Section 6.2. Proof of Lemma 6.6. Let $\mathcal{F}_n = \sigma(\{X_0, \dots, X_n\})$ be sigma algebra generated by the chain \mathbf{X} up to time n . Denote by $\mathbb{E}_{\mathcal{F}_n}$ the expected value conditioned to \mathcal{F}_n . Take $0 < \gamma \leq \delta$ and define $I_0 = [x_0 - \gamma, x_0]$, $I_1 = [x_0, x_0 + \gamma]$ and

$$\begin{aligned} S_0(\gamma) &= \sup_{y \in I_0} \left| \sum_{t=0}^n W_t (\mathbb{I}\{X_t \leq y\} - \mathbb{I}\{X_t \leq x_0\}) \right|^2 \\ S_1(\gamma) &= \sup_{y \in I_1} \left| \sum_{t=0}^n W_t (\mathbb{I}\{X_t \leq y\} - \mathbb{I}\{X_t \leq x_0\}) \right|^2 \end{aligned}$$

then,

$$\begin{aligned} S(\gamma) &= \sup_{|y-x_0| \leq \gamma} \left| \sum_{t=0}^n W_t (\mathbb{I}\{X_t \leq y\} - \mathbb{I}\{X_t \leq x_0\}) \right|^2 = \max(S_0(\gamma), S_1(\gamma)), \\ (41) \quad &\leq S_0(\gamma) + S_1(\gamma) \end{aligned}$$

Following the notation of section 2, let

$$\alpha_n^{(0)}(\gamma) = \sup_{y \in I_0} T_n([y, x_0]) \quad , \quad \alpha_n^{(1)}(\gamma) = \sup_{y \in I_1} T_n([x_0, y]) ,$$

with this notation, $S_0 = \sup_{y \in I_0} \left| \sum_{i=1}^{T_n([y, x_0])} W_{\sigma_{[y, x_0]}(i)} \right|^2$ and $S_1 = \sup_{y \in I_1} \left| \sum_{i=1}^{T_n([x_0, y])} W_{\sigma_{[x_0, y]}(i)} \right|^2$.

By Doob's maximal inequality (Th 10.9.4 in [19]), we have, for $j = 0, 1$,

$$\begin{aligned} \mathbb{E}_{\mathcal{F}_n} S_j(\gamma) &\leq 4 \mathbb{E}_{\mathcal{F}_n} \left(\sum_{i=1}^{\alpha_n^{(j)}} W_{t_i} \right)^2 \\ &\leq 4 \sigma^2 T_n([x_0 - \gamma, x_0 + \gamma]) \\ (42) \quad &\leq 4 \sigma^2 \sum_{t=0}^n (\mathbb{I}\{X_t \leq x_0 + \gamma\} - \mathbb{I}\{X_t < x_0 - \gamma\}). \end{aligned}$$

Therefore, by (41) and (42)

$$(43) \quad \mathbb{E}_{\mathcal{F}_n} S(\gamma) \leq 8 \sigma^2 \sum_{t=0}^n (\mathbb{I}\{X_t \leq x_0 + \gamma\} - \mathbb{I}\{X_t < x_0 - \gamma\}).$$

Define,

- $h(y, \gamma) = \mathbb{I}\{y \in [x_0 - \gamma, x_0 + \gamma]\}$,
- $h(\mathcal{B}_j, \gamma) = \begin{cases} \sum_{t=0}^{\tau_\alpha} h(X_t, \gamma) & , \quad j = 0 \\ \sum_{t=\tau_A(j)+1}^{\tau_A(j+1)} h(X_t, \gamma) & , \quad j \geq 1 \end{cases}$
- $Z_n(\gamma) = \sum_{t=0}^n h(X_t, \gamma)$
- $\ell(\mathcal{B}_j) = \begin{cases} \tau_\alpha & , \quad j = 0 \\ \tau_\alpha(j+1) - \tau_\alpha(j) & , \quad j \geq 1 \end{cases}$
- $\tilde{T}(n) = \min \left\{ k : \sum_{i=0}^k \ell(\mathcal{B}_i) \geq n \right\}$.
- $\mathcal{G}_k = \sigma \left(\{(h(\mathcal{B}_j, \gamma), \ell(\mathcal{B}_j))\}_{j=0}^k \right)$ for $k \geq 0$.

By the Strong Markov property, $\{(h(\mathcal{B}_j, \gamma), \ell(\mathcal{B}_j))\}_{j=1}^{+\infty}$ is an i.i.d. sequence which is independent of $(h(\mathcal{B}_0, \gamma), \ell(\mathcal{B}_0))$ (and, therefore, of the initial measure λ). For n fixed, the random variable $\tilde{T}(n)$ is a stopping time for the sequence $\{(h(\mathcal{B}_j, \gamma), \ell(\mathcal{B}_j))\}_{j=0}^{+\infty}$, in effect

$$\begin{aligned} \{\tilde{T}(n) = 0\} &= \{\ell(\mathcal{B}_0) \geq n\} \in \mathcal{G}_0, \\ \{\tilde{T}(n) = k\} &= \bigcap_{j=0}^{k-1} \left\{ \sum_{i=0}^j \ell(\mathcal{B}_i) < n \right\} \cap \left\{ \sum_{i=0}^k \ell(\mathcal{B}_i) \geq n \right\} \in \mathcal{G}_k \quad \forall k \geq 1. \end{aligned}$$

For each n and γ we have that

$$Z_n(\gamma) = \sum_{t=0}^{\tau_\alpha} h(X_t, \gamma) + \sum_{j=1}^{T(n)} h(\mathcal{B}_j, \gamma) + \sum_{t=t_\alpha(T(n))+1}^n h(X_t, \gamma)$$

$$(44) \quad \leq h(\mathcal{B}_0, \gamma) + \sum_{j=1}^{\tilde{T}(n)} h(\mathcal{B}_j, \gamma).$$

where the last inequality is justified by the fact that, $T(n) \leq \tilde{T}(n)$ and $h(y, \gamma)$ is a nonnegative function. Because $\ell(\mathcal{B}_j) \geq 1$ for all j , we have that,

$$\sum_{j=1}^{\tilde{T}(n)} h(\mathcal{B}_j, \gamma) = \sum_{j=1}^n h(\mathcal{B}_j, \gamma) \mathbb{I}\{\tilde{T}(n) \geq j\},$$

then,

$$(45) \quad \mathbb{E} \left(\sum_{j=1}^{\tilde{T}(n)} h(\mathcal{B}_j, \gamma) \right) = \sum_{j=1}^n \mathbb{E} \left(h(\mathcal{B}_j, \gamma) \mathbb{I}\{\tilde{T}(n) \geq j\} \right).$$

For each j we have,

$$\mathbb{E}_\lambda \left(h(\mathcal{B}_j, \gamma) \mathbb{I}\{\tilde{T}(n) \geq j\} \right) = \mathbb{E}_\lambda \left(\mathbb{E} \left(h(\mathcal{B}_j, \gamma) \mathbb{I}\{\tilde{T}(n) \geq j\} \mid \mathcal{G}_{j-1} \right) \right)$$

Notice that $\mathbb{I}\{\tilde{T}(n) \geq j\} = 1 - \mathbb{I}\{\tilde{T}(n) \leq j-1\} \in \mathcal{G}_{j-1}$ and $h(\mathcal{B}_j, \gamma)$ is independent of \mathcal{G}_{j-1} , therefore,

$$\mathbb{E}_\lambda \left(h(\mathcal{B}_j, \gamma) \mathbb{I}\{\tilde{T}(n) \geq j\} \right) = \mathbb{E}_\lambda \left(\mathbb{I}\{\tilde{T}(n) \geq j\} \right) \mathbb{E} \left(h(\mathcal{B}_j, \gamma) \right).$$

Plugging this into equation (45) we get,

$$\mathbb{E}_\lambda \left(\sum_{j=1}^{\tilde{T}(n)} h(\mathcal{B}_j, \gamma) \right) = \sum_{j=1}^n \mathbb{E} \left(h(\mathcal{B}_j, \gamma) \right) \mathbb{P}_\lambda \left(\tilde{T}(n) \geq j \right) \leq \mathbb{E} \left(h(\mathcal{B}_1, \gamma) \right) \mathbb{E}_\lambda \tilde{T}(n).$$

Then, by taking expectation in (44) we obtain

$$(46) \quad \begin{aligned} \mathbb{E}_\lambda Z_n(\gamma) &\leq \mathbb{E}_\lambda h(\mathcal{B}_0, \gamma) + \mathbb{E} \left(h(\mathcal{B}_1, \gamma) \right) \mathbb{E}_\lambda \tilde{T}(n) \\ &\leq \mathbb{E}_\lambda h(\mathcal{B}_0, \gamma) + \mathbb{E} \left(h(\mathcal{B}_1, \gamma) \right) \mathbb{E}_\lambda (T(n) + 1). \end{aligned}$$

By Theorem 1.1 and the fact that F is Lipschitz we can find K_1 independent of γ such that,

$$(47) \quad \begin{aligned} \mathbb{E} \left(h(\mathcal{B}_1, \gamma) \right) &= \int h(t, \gamma) d\pi(t) = K_\pi \pi(C) (F(x_0 + \gamma) - F(x_0 - \gamma)) \\ &\leq K_1 \gamma. \end{aligned}$$

If \mathbf{X} is positive recurrent, by Theorem 1.1, $\frac{T(n)}{u(n)}$ converges almost surely to a positive constant

$K_2 > 0$. Moreover, $\frac{T(n)}{u(n)} \leq 1$ therefore, by the Dominated Convergence Theorem we obtain that $\mathbb{E}_\lambda T(n) \sim \frac{u(n)}{K_2}$. If \mathbf{X} is β -null recurrent, by Lemma 3.3 in [24], $\mathbb{E}_\lambda T(n) \sim \frac{u(n)}{\Gamma(1+\beta)}$, hence, for both positive and β -null recurrent chains, we can find K_2 and N , both independent of γ , such that $\mathbb{E}_\lambda T(n) \leq K_2 u(n)$ for all $n \geq N$. Using this with (46) and (47) we get,

$$(48) \quad \frac{\mathbb{E}_\lambda Z_n(\gamma)}{u(n)\gamma} \leq \frac{\mathbb{E}_\lambda h(\mathcal{B}_0, \gamma)}{u(n)\gamma} + K_1 K_2 \quad \forall n \geq N, \forall \gamma \in (0, \delta].$$

Combining (48) with assumption (B3) and the fact that $Z_n(0) \equiv 0$ we obtain that there exist positive constants K_3 and γ_0 such that

$$\mathbb{E}_\lambda Z_n(\gamma) \leq u(n)\gamma \quad \forall n \geq N, \forall \gamma \in (0, \gamma_0].$$

Equation (35) now follows after taking expectation in (43). The proof of (36) follows the same reasoning, but using

$$S_j(\gamma) = \sup_{y \in I_j} \left| \sum_{t=0}^n W_t (\mathbb{I}_C \{X_t = y\}) \right|^2.$$

Proof of Lemma 6.7. a) If \mathbf{X} is positive recurrent, Theorem 1.1 implies that there exists a positive constant K such that $\frac{T_n(C)}{u(n)}$ converges almost surely to $K\pi(C)$, which is not zero by (A3).

On the other hand, if \mathbf{X} is β -null recurrent, Theorem 1.1 and Slutsky's Theorem implies that there exists a constant $K > 0$ such that $\frac{T_n(C)}{u(n)}$ converges in distribution to $K M_\beta(1)$ where $M_\beta(1)$ denotes a Mittag-Leffler distribution with parameter β . This distribution is continuous and strictly positive with probability 1, then, by the Continuous Mapping Theorem, $\frac{u(n)}{T_n(C)}$ converges in distribution to a multiple of $\frac{1}{M_\beta}$, therefore, $\frac{u(n)}{T_n(C)}$ is bounded in probability by Theorem 2.4 in [33].

b) Let \mathbf{X} be positive recurrent, then, we can find N_η such that

$$\mathbb{P} \left(\left| \left(\frac{T_n(C)}{u(n)} \right)^\alpha - K^\alpha \pi(C)^\alpha \right| \leq \left(\frac{K\pi(C)}{2} \right)^\alpha \right) \geq 1 - \eta, \quad \forall n \geq N_\eta.$$

hence,

$$\mathbb{P} \left(\left(\frac{T_n(C)}{u(n)} \right)^\alpha \in \left[\frac{K^\alpha \pi(C)^\alpha}{2}, \frac{3K^\alpha \pi(C)^\alpha}{2} \right] \right) \geq 1 - \eta, \quad \forall n \geq N_\eta.$$

Now let \mathbf{X} be β -null recurrent. Let $Z = (K M_\beta(1))^\alpha$, This random variable is continuous and positive, therefore, we can find positive constants \underline{c}_η and \bar{c}_η such that

$$(49) \quad \mathbb{P}(Z \in [\underline{c}_\eta, \bar{c}_\eta]) \geq 1 - \frac{\eta}{2}.$$

By the Continuous Mapping Theorem, $\left(\frac{T_n(C)}{u(n)} \right)^\alpha$ converges in distribution to Z , therefore, we can find $N_\eta \in \mathbb{N}$ such that

$$(50) \quad \left| \mathbb{P} \left(\left(\frac{T_n(C)}{u(n)} \right)^\alpha \in [\underline{c}_\eta, \bar{c}_\eta] \right) - \mathbb{P}(Z \in [\underline{c}_\eta, \bar{c}_\eta]) \right| \leq \frac{\eta}{2}, \quad \forall n \geq N_\eta,$$

Combining (49) and (50) we obtain that

$$(51) \quad \mathbb{P} \left(\left(\frac{T_n(C)}{u(n)} \right)^\alpha \in [\underline{c}_\eta, \bar{c}_\eta] \right) \geq 1 - \eta, \quad \forall n \geq N_\eta.$$

Proof of Lemma 6.8. Fix $\varepsilon \in (0, 1)$ small enough so that F' and $|f'_0|$ are bounded from above and away from zero on $[F^{-1}(F(x_0) - 2\varepsilon), F^{-1}(F(x_0) + 2\varepsilon)]$, see the assumption (B2). Then, the proper inverse functions of F and f_0 are well defined on $[F(x_0) - 2\varepsilon, F(x_0) + 2\varepsilon]$ and

$$[f_0 \circ F^{-1}(F(x_0) - 2\varepsilon), f_0 \circ F^{-1}(F(x_0) + 2\varepsilon)]$$

respectively. We denote the inverses on that intervals by F^{-1} and f_0^{-1} respectively. Let

$$(52) \quad U_n(a) = \operatorname{argmax}_{|p - F(x_0)| \leq \varepsilon} \{\Lambda_n(p) - ap\}$$

where $a = f_0(x_0)$ and where the supremum is restricted to $p \in [F_n(Y_0), F_n(Y_m)]$. We will show below that

$$(53) \quad U_n(a) = F(x_0) + O_P(u(n)^{-1/3}),$$

as $n \rightarrow \infty$. Combining (31) to Lemma 6.5 ensures that $\widehat{U}_n(a)$ coincides with $U_n(a)$ with a probability that tends to one as $n \rightarrow \infty$, so (37) follows from (53).

We turn to the proof of (53). Fix $\eta > 0$ arbitrarily and let

$$(54) \quad \gamma_n = K_0 u(n)^{-1/3}$$

for some $K_0 \geq 1$ sufficiently large so that

$$(55) \quad \gamma_n \geq \frac{1}{\sqrt{u(n)}}.$$

Then, by part ii) of Lemma 6.7, we can find positive constants \underline{c}_η , \bar{c}_η and N_η such that

$$(56) \quad \mathbb{P}\left(T_n(C)^{2/3} \gamma_n u(n)^{-1/3} \in [K_0 \underline{c}_\eta, K_0 \bar{c}_\eta]\right) \geq 1 - \eta/2 \quad \forall n \geq N_\eta,$$

Let $\underline{c} = K_0 \underline{c}_\eta$ and $\bar{c} = K_0 \bar{c}_\eta$. It follows from (18) that for sufficiently small $\varepsilon > 0$, we can find $K_1 > 0$ such that

$$\mathbb{P}\left(T_n(C) \sup_{|p-F(x_0)| \leq 2\varepsilon} |F_n^{-1}(p) - F^{-1}(p)|^2 \leq K_1\right) \geq 1 - \eta/2$$

for all n . Hence for $n \geq N_\eta$,

$$\mathbb{P}(\mathcal{E}_n) \geq 1 - \eta,$$

where \mathcal{E}_n denotes the intersection of the events

$$(57) \quad T_n(C)^{2/3} \gamma_n u(n)^{-1/3} \in [\underline{c}, \bar{c}]$$

and

$$(58) \quad T_n(C) \sup_{|p-F(x_0)| \leq 2\varepsilon} |F_n^{-1}(p) - F^{-1}(p)|^2 \leq K_1.$$

Combining equations (57) and (58), we obtain that, in \mathcal{E}_n ,

$$(59) \quad \sup_{|p-F(x_0)| \leq 2\varepsilon} |F_n^{-1}(p) - F^{-1}(p)|^2 \leq K_2 a(n)^{-1}$$

where $K_2 = K_1 \left(\frac{K_0}{\underline{c}}\right)^{3/2}$ is independent of n and K_0 .

By Lemma 6.3, we can assume without loss of generality that $F(x_0)$ belongs to $[F_n(Y_0), F_n(Y_m)]$, since this occurs with probability that tends to one. Hence, by (52), the event $\{|U_n(a) - F(x_0)| \geq \gamma_n\}$ is contained in the event that there exists $p \in [F_n(Y_0), F_n(Y_m)]$ with $|p - F(x_0)| \leq \varepsilon$, $|p - F(x_0)| \geq \gamma_n$ and

$$(60) \quad \Lambda_n(p) - ap \geq \Lambda_n(F(x_0)) - aF(x_0).$$

Obviously, the probability is equal to zero if $\gamma_n > \varepsilon$ so we assume in the sequel that $\gamma_n \leq \varepsilon$. For all $p \in [F(x_0) - \varepsilon, F(x_0) + \varepsilon]$ define

$$\Lambda(p) = \int_{F(x_0)}^p f_0 \circ F^{-1}(u) du.$$

Let $c > 0$ such that $|f'_0|/F' > 2c$ on the interval $[F^{-1}(F(x_0) - 2\varepsilon), F^{-1}(F(x_0) + 2\varepsilon)]$. Since $\Lambda'(F(x_0)) = a$ and $\Lambda'' = f'_0 \circ F^{-1}/F' \circ F^{-1}$, it then follows from Taylor's expansion that

$$\Lambda(p) - \Lambda(F(x_0)) \leq (p - F(x_0))a - c(p - F(x_0))^2$$

for all $p \in [F(x_0) - \varepsilon, F(x_0) + \varepsilon]$ and therefore, (60) implies that

$$\Delta_n(p) - \Delta_n(F(x_0)) - c(p - F(x_0))^2 \geq 0$$

for all such p 's, where we set $\Delta_n := \Lambda_n - \Lambda$. Hence, for all $n \geq N_\eta$,

$$\begin{aligned} & \mathbb{P}(|U_n(a) - F(x_0)| \geq \gamma_n) \\ & \leq \eta + \mathbb{P}\left(\sup_{|p-F(x_0)| \in [\gamma_n, \varepsilon]} \{\Delta_n(p) - \Delta_n(F(x_0)) - c(p - F(x_0))^2\} \geq 0 \text{ and } \mathcal{E}_n\right) \\ & \leq \eta + \sum_j \mathbb{P}\left(\sup_{|u| \in [\gamma_n 2^j, \gamma_n 2^{j+1}]} \{\Delta_n(F(x_0) + u) - \Delta_n(F(x_0))\} \geq c(\gamma_n 2^j)^2 \text{ and } \mathcal{E}_n\right) \\ (61) \quad & \leq \eta + \sum_j \mathbb{P}\left(\sup_{|u| \leq \gamma_n 2^{j+1}} |\Delta_n(F(x_0) + u) - \Delta_n(F(x_0))| \geq c(\gamma_n 2^j)^2 \text{ and } \mathcal{E}_n\right) \end{aligned}$$

where the sums are taken over all integers $j \geq 0$ such that $\gamma_n 2^j \leq \varepsilon$. Recall that we have (39) for all $k \in \{0, \dots, m\}$. Since Λ_n is piecewise-linear with knots at $F_n(Y_0), \dots, F_n(Y_m)$, by (22) and (23) we get that for every j in the above sum,

$$\begin{aligned} & \sup_{|u| \leq \gamma_n 2^{j+1}} |\Delta_n(F(x_0) + u) - \Delta_n(F(x_0))| \\ & \leq \sup_{|u| \leq \gamma_n 2^{j+1}} \left| \int_{F(x_0)}^{F(x_0)+u} (f_0 \circ F_n^{-1}(y) - f_0 \circ F^{-1}(y)) dy \right| \\ (62) \quad & + \sup_{|u| \leq \gamma_n 2^{j+1}} |M_n(F(x_0) + u) - M_n(F(x_0))|. \end{aligned}$$

Moreover, $|f'_0|$ is bounded above on $[F^{-1}(F(x_0) - 2\varepsilon), F^{-1}(F(x_0) + 2\varepsilon)]$, so we obtain that for every j with $\gamma_n 2^j \leq \varepsilon$, the first term on the right-hand side of (62) satisfies

$$\begin{aligned} & \sup_{|u| \leq \gamma_n 2^{j+1}} \left| \int_{F(x_0)}^{F(x_0)+u} (f_0 \circ F_n^{-1}(p) - f_0 \circ F^{-1}(p)) dp \right| \\ & \leq \int_{F(x_0) - \gamma_n 2^{j+1}}^{F(x_0) + \gamma_n 2^{j+1}} |f_0 \circ F_n^{-1}(p) - f_0 \circ F^{-1}(p)| dp \\ & \leq K_3 \gamma_n 2^j \sup_{|p-F(x_0)| \leq 2\varepsilon} |F_n^{-1}(p) - F^{-1}(p)|, \end{aligned}$$

for some $K_3 > 0$ that does not depend on n . Hence, it follows from the previous display and (59) that

$$\begin{aligned} & \mathbb{E} \left(\sup_{|u| \leq \gamma_n 2^{j+1}} \left| \int_{F(x_0)}^{F(x_0)+u} (f_0 \circ F_n^{-1}(p) - f_0 \circ F^{-1}(p)) dp \right|^2 \mathbb{I}(\mathcal{E}_n) \right) \\ & \leq K_3^2 \gamma_n^2 2^{2j} \mathbb{E} \left(\sup_{|p-F(x_0)| \leq 2\varepsilon} |F_n^{-1}(p) - F^{-1}(p)|^2 \mathbb{I}(\mathcal{E}_n) \right) \\ & \leq K_3^2 \gamma_n^2 2^{2j} K_2 u(n)^{-1}. \end{aligned}$$

Hence, taking $K_4 = K_3^2 K_2$ we get that for all j with $\gamma_n 2^j \leq \varepsilon \leq 1$.

$$(63) \quad \mathbb{E} \left(\sup_{|u| \leq \gamma_n 2^{j+1}} \left| \int_{F(x_0)}^{F(x_0)+u} (f_0 \circ F_n^{-1}(p) - f_0 \circ F^{-1}(p)) dp \right|^2 \mathbb{I}(\mathcal{E}_n) \right) \leq K_4 \gamma_n 2^j u(n)^{-1}.$$

By equations (23) and (24) in Lemma 6.1, the second term on the right-hand side of (62) satisfies,

$$(64) \quad \sup_{|u| \leq \gamma_n 2^{j+1}} |M_n(F(x_0) + u) - M_n(F(x_0))| \leq I_1^{n,j} + I_2^{n,j},$$

where $I_1^{n,j}$ and $I_2^{n,j}$ are given by

$$I_1^{n,j} = \frac{1}{T_n(C)} \sup_{|u| \leq \gamma_n 2^{j+1}} \left| \sum_{t=0}^n W_t \left(\mathbb{I}_C\{X_t \leq F_n^{-1}(F(x_0) + u)\} - \mathbb{I}_C\{X_t \leq F_n^{-1}(F(x_0))\} \right) \right|,$$

$$I_2^{n,j} = \frac{2}{T_n(C)} \sup_{|u| \leq \gamma_n 2^{j+1}} \left| \sum_{t=0}^n W_t \left(\mathbb{I}_C\{X_t = F_n^{-1}(F(x_0) + u)\} \right) \right|.$$

For $I_1^{n,j}$, it follows from the triangle inequality that

$$I_1^{n,j} \leq \frac{2}{T_n(C)} \sup_{|u| \leq \gamma_n 2^{j+1}} \left| \sum_{t=0}^n W_t \left(\mathbb{I}_C\{X_t \leq F_n^{-1}(F(x_0) + u)\} - \mathbb{I}_C\{X_t \leq x_0\} \right) \right|.$$

Combining (59) and the fact that F^{-1} is Lipschitz in $[F(x_0) - 2\varepsilon, F(x_0) + 2\varepsilon]$ we can find $K_5 = \max(\sqrt{K_2}, \sup(F^{-1}))$ independent of n such that, on \mathcal{E}_n ,

$$\sup_{|p-F(x_0)| \leq 2\varepsilon} |F_n^{-1}(p) - F^{-1}(p)| \leq \frac{K_5}{\sqrt{u(n)}}$$

and $|F^{-1}(F(x_0) + u) - x| = |F^{-1}(F(x_0) + u) - F^{-1}(F(x_0))| \leq K_5|u|/2$ for all u with $|u| \leq 2\varepsilon$. Hence, on \mathcal{E}_n

$$I_1^{n,j} \leq \frac{2}{T_n(C)} \sup_{|y-x_0| \leq K_5 \gamma_n 2^j + K_5/\sqrt{u(n)}} \left| \sum_{t=0}^n W_t \left(\mathbb{I}_C\{X_t \leq y\} - \mathbb{I}_C\{X_t \leq x_0\} \right) \right|,$$

$$I_2^{n,j} \leq \frac{2}{T_n(C)} \sup_{|y-x_0| \leq K_5 \gamma_n 2^j + K_5/\sqrt{u(n)}} \left| \sum_{t=0}^n W_t \left(\mathbb{I}_C\{X_t = y\} \right) \right|.$$

It follows from (55) that $\gamma_n 2^j \geq \gamma_n \geq 1/\sqrt{u(n)}$ for all $j \geq 0$, then, on \mathcal{E}_n

$$I_1^{n,j} \leq \frac{2}{T_n(C)} \sup_{|y-x_0| \leq 2K_5 \gamma_n 2^j} \left| \sum_{t=0}^n W_t \left(\mathbb{I}_C\{X_t \leq y\} - \mathbb{I}_C\{X_t \leq x_0\} \right) \right|,$$

$$I_2^{n,j} \leq \frac{2}{T_n(C)} \sup_{|y-x_0| \leq 2K_5 \gamma_n 2^j} \left| \sum_{t=0}^n W_t \left(\mathbb{I}_C\{X_t = y\} \right) \right|.$$

By Lemma 6.6, we conclude that there exists $K_6 > 0$ and N'_η such that, for $n \geq N'_\eta$

$$(65) \quad \mathbb{E} \left(\left(I_1^{n,j} + I_2^{n,j} \right)^2 \mathbb{I}(\mathcal{E}_n) \right) \leq K_6 \gamma_n 2^j u(n)^{-1}$$

Combining (62), (63), (64) and (65), we conclude that there exists $K_7 > 0$, independent of n and K_0 , such that for all $n \geq N'_\eta$ and $j \geq 0$ where $\gamma_n 2^j \leq \varepsilon$, one has

$$\mathbb{E} \left(\sup_{|u| \leq \gamma_n 2^{j+1}} |\Delta_n(F(x_0) + u) - \Delta_n(F(x_0))|^2 \mathbb{I}(\mathcal{E}_n) \right) \leq K_7 \gamma_n 2^j u(n)^{-1}.$$

Combining this with (61) and the Markov inequality, we conclude that there exist $K_8 > 0$ and N''_η , that do not depend on n nor K_0 , such that, for all $n \geq N''_\eta$,

$$\begin{aligned} \mathbb{P}(|U_n(a) - F(x_0)| \geq \gamma_n) &\leq \eta + K_8 \sum_{k \geq 0} \frac{\gamma_n 2^k u(n)^{-1}}{(\gamma_n 2^k)^4} \\ &\leq \eta + K_8 \gamma_n^{-3} u(n)^{-1} \sum_{j \geq 0} 2^{-3j}. \end{aligned}$$

The sum on the last line is finite, so there exists $K > 0$, independent of n and K_0 , such that for n bigger than N''_η

$$(66) \quad \mathbb{P}(|U_n(a) - F(x_0)| \geq \gamma_n) \leq \eta + K \gamma_n^{-3} u(n)^{-1} = \eta + \frac{K}{K_0^3}.$$

The above probability can be made smaller than 2η by setting (54) for some sufficiently large K_0 independent of n . This proves (53) and completes the proof of (37).

Now, we turn to the proof of (38). It follows from (30) combined to (32) and Lemma 5.3 that

$$\widehat{f}_n^{-1}(f_0(x_0)) = F^{-1} \circ \widehat{U}_n(f_0(x_0)) + T_n(C)^{-1/2} O_P(1).$$

Hence, by Lemma 6.7 we have

$$\widehat{f}_n^{-1}(f_0(x_0)) = F^{-1} \circ \widehat{U}_n(f_0(x_0)) + O_P(u(n)^{-1/2}).$$

Now, it follows from the assumption (B2) that F^{-1} has a bounded derivative in the neighborhood of $F(x_0)$, to which $\widehat{U}_n(f_0(x_0))$ belongs with probability that tends to one. Hence, it follows from Taylor's expansion that

$$\begin{aligned} \widehat{f}_n^{-1}(f_0(x_0)) &= F^{-1} \circ F(x_0) + O(|\widehat{U}_n(f_0(x_0)) - F(x_0)|) + O_P(u(n)^{-1/2}) \\ &= x + O_P(u(n)^{-1/3}) + O_P(u(n)^{-1/2}), \end{aligned}$$

where we used (37) for the last equality. This proves (38) and completes the proof of Lemma 6.8.

SUPPLEMENTARY MATERIAL

1. Markov chain theory and notation. This section extends Section 2 of the main paper, presenting a more detailed exposition of the Markov chain theory required in the proofs. For further details, we refer the reader to [12, 27, 29].

Let $\mathbf{X} = X_0, X_1, X_2, \dots$ be a time-homogeneous Markov Chain defined on a probability space $(E, \mathcal{E}, \mathbb{P})$ where \mathcal{E} is countably generated. Let $P(x, A)$ denote its transition kernel, i.e. for $x \in E$, $A \in \mathcal{E}$ we have

$$P(x, A) = \mathbb{P}(X_{i+1} \in A | X_i = x), \quad i = 0, 1, \dots$$

Let $P^n(x, A)$ denote the n -step transition probability, i.e.

$$P^n(x, A) = \mathbb{P}(X_{i+n} \in A | X_i = x) \quad \forall i.$$

If λ is a probability measure in (E, \mathcal{E}) such that $\mathcal{L}(X_0) = \lambda$, then λ is called the *initial measure* of the chain \mathbf{X} . A homogeneous Markov chain is uniquely identified by its kernel and initial measure.

When the initial measure of the chain is given, we will write \mathbb{P}_λ (and \mathbb{E}_λ) for the probability (and the expectation) conditioned on $\mathcal{L}(X_0) = \lambda$. When $\lambda = \delta_x$ for some $x \in E$ we will simply write \mathbb{P}_x and \mathbb{E}_x .

An homogeneous Markov chain is *irreducible* if there exists a σ -finite measure ϕ on (E, \mathcal{E}) such that for all $x \in E$ and all $A \in \mathcal{E}$ with $\phi(A) > 0$ we have $P^n(x, A) > 0$ for some $n \geq 1$. In this case, there exists a maximal irreducibility measure ψ (all other irreducibility measures are absolutely continuous with respect to ψ). In the following, all Markov chains are supposed to be irreducible with maximal irreducibility measure ψ .

For any set $C \in \mathcal{E}$, we will denote by σ_C and τ_C , respectively, the times of first visit and first return of the chain to the set C , i.e. $\tau_C = \inf \{n \geq 1 : X_n \in C\}$ and $\sigma_C = \inf \{n \geq 0 : X_n \in C\}$. The subsequent visit and return times $\sigma_C, \tau_C(k)$, $k \geq 1$ are defined inductively as follows:

$$(67) \quad \tau_C(1) = \tau_C \quad , \quad \tau_C(k) = \min \{n > \tau_C(k-1) : X_n \in C\},$$

$$(68) \quad \sigma_C(1) = \sigma_C \quad , \quad \sigma_C(k) = \min \{n > \sigma_C(k-1) : X_n \in C\}.$$

Given that our methods will only deal with the values of \mathbf{X} in a fixed set C , if A is a measurable set, we will write $\mathbb{I}_C\{X_t \in A\}$ instead of $\mathbb{I}\{X_t \in A \cap C\}$ and if $A = E$, then we will simply write $\mathbb{I}_C(X_t)$.

We will use $T_n(C)$ to denote the random variable that counts the number of times the chain has visited the set C up to time n , that is $T_n(C) = \sum_{t=0}^n \mathbb{I}_C(X_t)$. Similarly, we will write $T(C)$ for the total of numbers of visits the chain \mathbf{X} to C . The set C is called *recurrent* if $\mathbb{E}_x T(C) = +\infty$ for all $x \in C$. The chain \mathbf{X} is considered recurrent if every set $A \in \mathcal{E}$, such that $\psi(A) > 0$, is recurrent.

Although recurrent chains possess many interesting properties, a stronger type of recurrence is required in our analysis. An irreducible Markov chain is *Harris recurrent* if for all $x \in E$ and all $A \in \mathcal{E}$ with $\psi(A) > 0$ we have

$$\mathbb{P}(X_n \in A \text{ infinitely often} | X_0 = x) = 1.$$

An irreducible chain possesses an accessible atom, if there is a set $\alpha \in \mathcal{E}$ such that for all x, y in α : $P(x, \cdot) = P(y, \cdot)$ and $\psi(\alpha) > 0$. Denote by \mathbb{P}_α and $\mathbb{E}_\alpha(\cdot)$ the probability and the expectation conditionally to $X_0 \in \alpha$. If \mathbf{X} possesses an accessible atom and is Harris recurrent, the probability of returning infinitely often to the atom α is equal to one, no matter the starting point, i.e. $\forall x \in E, \mathbb{P}_x(\tau_\alpha < \infty) = 1$. Moreover, it follows from the *strong Markov property* that the sample paths may be divided into independent blocks of random length corresponding to consecutive visits to α :

$$\begin{aligned} \mathcal{B}_0 &= (X_0, X_1, \dots, X_{\tau_\alpha(1)}) \\ \mathcal{B}_1 &= (X_{\tau_\alpha(1)+1}, \dots, X_{\tau_\alpha(2)}) \\ &\dots \\ \mathcal{B}_n &= (X_{\tau_\alpha(n)+1}, \dots, X_{\tau_\alpha(n+1)}) \\ &\dots \end{aligned}$$

taking their values in the torus $\mathbb{T} = \cup_{n=1}^{\infty} E^n$. Notice that the distribution of \mathcal{B}_0 depends on the initial measure, therefore it does not have the same distribution as \mathcal{B}_j for $j \geq 1$. The sequence $\{\tau_\alpha(j)\}_{j \geq 1}$ defines successive times at which the chain forgets its past, called *regeneration*

times. Similarly, the sequence of i.i.d. blocks $\{\mathcal{B}_j\}_{j \geq 1}$ are named *regeneration blocks*. The random variable $T(n) = T_n(\alpha) - 1$ counts the number of i.i.d. blocks up to time n . This term is called *number of regenerations up to time n* .

Notice that for any function defined on E , we can write $\sum_{t=0}^n f(X_t)$ as a sum of independent random variables as follows:

$$(69) \quad \sum_{t=0}^n f(X_t) = f(\mathcal{B}_0) + \sum_{j=1}^{T(n)} f(\mathcal{B}_j) + f(\mathcal{B}_{(n)}),$$

where, $f(\mathcal{B}_0) = \sum_{t=0}^{\tau_\alpha} f(X_t)$, $f(\mathcal{B}_j) = \sum_{t=\tau_\alpha(j)+1}^{\tau_\alpha(j+1)} f(X_t)$ for $j = 1, \dots, T(n)$ and $f(\mathcal{B}_{(n)})$ denotes the last incomplete block, i.e. $f(\mathcal{B}_{(n)}) = \sum_{t=\tau_\alpha(T(n)+1)+1}^n f(X_t)$.

When an accessible atom exists, the *stochastic stability* properties of \mathbf{X} amount to properties concerning the speed of return time to the atom only. For instance, the measure π_α given by:

$$(70) \quad \pi_\alpha(B) = \mathbb{E}_\alpha \left(\sum_{n=1}^{\tau_\alpha} \mathbb{I}\{X_n \in B\} \right), \quad \forall B \in \mathcal{E}$$

is invariant, i.e.

$$\pi_\alpha(B) = \int P(x, B) d\pi_\alpha(x).$$

Denote by \mathcal{E}^+ the class of nonnegative measurable functions with positive ψ support. A function $s \in \mathcal{E}^+$ is called *small* if there exists an integer $m_0 \geq 1$ and a measure $\nu \in \mathcal{M}(\mathcal{E})_+$ such that

$$(71) \quad P^{m_0}(x, A) \geq s(x) \nu(A) \quad \forall x \in E, A \in \mathcal{E}.$$

When a chain possesses a small function s , we say that it satisfies the *minorization inequality* $M(m_0, s, \nu)$. As pointed out in [29], there is no loss of generality in assuming that $0 \leq s(x) \leq 1$ and $\int_E s(x) d\nu(x) > 0$.

A set $A \in \mathcal{E}$ is said to be *small* if the function \mathbb{I}_A is small. Similarly, a measure ν is *small* if there exist m_0 , and s that satisfy (71). By Theorem 2.1 in [29], every irreducible Markov chain possesses a small function and Proposition 2.6 of the same book shows that every measurable set A with $\psi(A) > 0$ contains a small set. In practice, finding such a set consists in most cases in exhibiting an accessible set, for which the probability that the chain returns to it in m steps is uniformly bounded below. Moreover, under quite wide conditions a compact set will be small, see [15].

If \mathbf{X} does not possess an atom but is Harris recurrent (and therefore satisfies a minorization inequality $M(m_0, s, \nu)$), a *splitting technique*, introduced in [28, 29], allows us to extend in some sense the probabilistic structure of \mathbf{X} in order to artificially construct an atom. The general idea behind this construction is to expand the sample space so as to define a sequence $(Y_n)_{n \in \mathbb{N}}$ of Bernoulli r.v.'s and a bivariate chain $\check{\mathbf{X}} = \{(X_n, Y_n)\}_{n=0}^{+\infty}$, named *split chain*, such that the set $\check{\alpha} = (E, 1)$ is an atom of this chain. A detailed description of this construction can be found in [29].

The whole point of this construction consists in the fact that $\check{\mathbf{X}}$ inherits all the communication and stochastic stability properties from \mathbf{X} (irreducibility, Harris recurrence,...). In particular, the marginal distribution of the first coordinate process of $\check{\mathbf{X}}$ and the distribution of the original \mathbf{X} are identical. Hence, the splitting method enables us to establish all the results known for atomic chains to general Harris chains, for example, the existence of an invariant measure which is unique up to multiplicative constant (see Proposition 10.4.2 in [27]).

The invariant measure is finite if and only if $\mathbb{E}_{\tilde{\alpha}}\tau_{\tilde{\alpha}} < +\infty$, in this case we say the chain is *positive recurrent*, otherwise, we say the chain is *null recurrent*. A null recurrent chain is called β -null recurrent (c.f. Definition 3.2 in [24]) if there exists a small nonnegative function h , a probability measure λ , a constant $\beta \in (0, 1)$ and a slowly varying function L_h such that

$$\mathbb{E}_{\lambda} \left(\sum_{t=0}^n h(X_t) \right) \sim \frac{1}{\Gamma(1+\beta)} n^{\beta} L_h(n) \quad \text{as } n \rightarrow \infty.$$

As argued in [24], is not a too severe restriction to assume $m_0 = 1$. Therefore, throughout this paper we assume that \mathbf{X} satisfies the minorization inequality $M(1, s, \nu)$, i.e, there exist a measurable function s and a probability measure ν such that $0 \leq s(x) \leq 1$, $\int_E s(x) d\nu(x) > 0$ and

$$(72) \quad P(x, A) \geq s(x) \nu(A).$$

A measurable and positive function L , defined in $[a, +\infty)$ for some $a \geq 0$, is called *slowly varying at $+\infty$* if it satisfies $\lim_{x \rightarrow +\infty} \frac{L(xt)}{L(x)} = 1$ for all $t \geq a$. See [6] for a detailed compendium of these types of functions.

It was shown in Theorem 3.1 of [24] that if the chain satisfies the minorization condition (5), then it is β -null recurrent if and only if

$$(73) \quad \mathbb{P}(\tau_{\tilde{\alpha}} > n) \sim \frac{1}{\Gamma(1-\beta) n^{\beta} L(n)},$$

where L is a slowly varying function.

The following theorem is a compendium of the main properties of Harris's recurrent Markov chains that will be used throughout the proofs. Among other things, it shows that the asymptotic behavior of $T(n)$ is similar to the function $u(n)$ defined as

$$(74) \quad u(n) = \begin{cases} n, & \text{if } \mathbf{X} \text{ is positive recurrent} \\ n^{\beta} L(n), & \text{if } \mathbf{X} \text{ is } \beta\text{-null recurrent} \end{cases}.$$

The Mittag-Leffler distribution with index β is a non-negative continuous distribution, whose moments are given by

$$\mathbb{E}(M_{\beta}^m(1)) = \frac{m!}{\Gamma(1+m\beta)} \quad m \geq 0.$$

See (3.39) in [24] for more details.

THEOREM 1.1. *Suppose \mathbf{X} is a Harris recurrent, irreducible Markov chain, with initial measure λ , that satisfies the minorization condition (72). Let $T(n)$ be the number of complete regenerations until time n of the split chain $\tilde{\mathbf{X}}$, let $C \in \mathcal{E}$ be a small set and π be an invariant measure for \mathbf{X} . Then,*

1. $0 < \pi(C) < +\infty$.
2. For any function f , defined on E , the decomposition (69) holds. Moreover, there is a constant K_{π} , that only depends on π , such that if $f \in L^1(E, \pi)$, then $\mathbb{E}_{\lambda} f(\mathcal{B}_1) = K_{\pi} \int_E f d\pi$.
3. $\frac{T(n)}{T_n(C)}$ converges almost surely to a positive constant.
4. $\frac{T(n)}{u(n)}$ converges almost surely to a positive constant if \mathbf{X} is positive recurrent and converges in distribution to a Mittag-Leffler random variable with index β if \mathbf{X} is β -null recurrent.

REMARK 1.1. The Mittag-Leffler distribution with index β is a non-negative continuous distribution, whose moments are given by

$$\mathbb{E}(M_\beta^m(1)) = \frac{m!}{\Gamma(1+m\beta)} \quad m \geq 0.$$

See (3.39) in [24] for more details.

REMARK 1.2. Part 1 of Theorem 1.1 is Proposition 5.6.ii of [29], part 2 is equation (3.23) of [24] and part 3 is an application of the Ratio Limit Theorem (Theorem 17.2.1 of [27]). For the positive recurrent case, part 4 also follows by the aforementioned Ratio Limit Theorem while the claim for the null recurrent case appears as Theorem 3.2 in [24].

2. Technical proofs for Section 5. Proof of Lemma 5.1. Equation (12) follows from Corollary 2 in [2] and part 2 of Theorem 1.1.

Now, we turn to the proof of (13). To do this, we adapt some of the ideas presented in the proof of Lemma 21.2 in [33].

Let V be a normal random variable independent of the X_i 's, and Φ its distribution function. It follows from (12) that conditionally on the X_i 's, $F_n(V)$ converges almost surely to $F(V)$. Thus, denoting by \mathbb{P}_X the conditional probability given the X_i 's, it follows from (29) that $\Phi(F_n^{-1}(u)) = \mathbb{P}_X(F_n(V) < u)$ converges almost surely to $\mathbb{P}_X(F(V) < u) = \Phi(F^{-1}(u))$ at every u at which the limit function is continuous. Since F is strictly increasing in C , one can find $\varepsilon > 0$ such that F^{-1} is continuous on $[F(x_0) - \varepsilon, F(x_0) + \varepsilon]$, so the above limit function is continuous at every $u \in [F(x_0) - \varepsilon, F(x_0) + \varepsilon]$. By continuity of Φ^{-1} on $(0, 1)$, $F_n^{-1}(u)$ converges almost surely to $F^{-1}(u)$ for every such u . By monotonicity, the convergence is uniform, hence

$$\sup_{|p-F(x_0)| \leq \varepsilon} |F_n^{-1}(p) - F^{-1}(p)| = o(1) \quad a.s.$$

as $n \rightarrow \infty$.

Proof of Lemma 5.2. This proof is an adaptation to the localized case of the proof of Lemma 2 in [5]. Let $f'_C \in \mathcal{F}'_C$, i.e., there exists $f \in \mathcal{F}$ such that $f'_C(B) = \int f(y) M_C(B, dy)$. By Cauchy–Schwarz inequality,

$$\left(\int f(y) M_C(B, dy) \right)^2 \leq \ell_C(B) \left(\int f^2 M_C(B, dy) \right),$$

then

$$\mathbb{E}_{Q'}(f'^2_C) \leq \mathbb{E}_{Q'} \left(\ell_C(B) \left(\int f(y)^2 M_C(B, dy) \right) \right) = \mathbb{E}_{Q_C}(f^2) \mathbb{E}_{Q'}(\ell^2_C),$$

where the last equality follows from (16). Applying this to the function

$$f'_C(B) - f'_k(B) = \int (f(y) - f_k(y)) M_C(B, dy),$$

when each f_k is the center of an ε -cover of the space \mathcal{F} and $\|f - f_k\|_{L_2(Q_C)} \leq \varepsilon$ gives the first assertion of the lemma. To obtain the second assertion, note that $U'_C = U \ell_C$ is an envelope for \mathcal{F}'_C . In addition, we have that

$$\|U'_C\|_{L_2(Q')} = U \|\ell_C\|_{L_2(Q')}.$$

From this, we derive that, for every $0 < \varepsilon < 1$,

$$\mathcal{N}(\varepsilon \|U'_C\|_{L_2(Q')}, \mathcal{U}'_C, L_2(Q')) = \mathcal{N}(\varepsilon U \|\ell_C\|_{L_2(Q')}, \mathcal{U}', L_2(Q')).$$

Then using the first assertion of the lemma, we obtain for every $0 < \varepsilon < 1$,

$$\mathcal{N}(\varepsilon \|U'_C\|_{L_2(Q')}, \mathcal{F}'_C, L_2(Q')) \leq \mathcal{N}(\varepsilon U, \mathcal{F}, L_2(Q_C)),$$

which implies the second assertion of the Lemmaz whenever the class \mathcal{F} is VC with envelope U .

Proof of Lemma 5.3. Let $B \in \widehat{E}$ and $g : E \times \mathbb{R} \rightarrow \mathbb{R}_+$. For each $y \in \mathbb{R}$ we define $g_y(x) = g(x, y)$, then, using the notation of section 6.2 we will have $\widehat{g}_y(B) = \sum_{x \in B \cap C} g(x, y)$. Finally, for any function $h : \mathbb{R} \rightarrow \mathbb{R}$, we define

$$\widetilde{g}_y^h(B) = (g_y - \widehat{h}(y))(B) = \sum_{x \in B \cap C} (g(x, y) - h(y)) = \widehat{g}_y(B) - \ell_C(B) h(y).$$

Let $g(x, y) = \mathbb{I}\{x \leq y\}$, and $h = F$ as defined in (8). Then, $\widehat{g}_y(B) = \sum_{x \in B} \mathbb{I}_C\{x \leq y\}$ and

$$\widetilde{g}_y^F(B) = \sum_{x \in B \cap C} (\mathbb{I}\{x \leq y\} - F(y)) = \widehat{g}_y(B) - \ell_C(B) F(y).$$

From now on, we'll remove the superindex from \widetilde{g}_y^F to ease the notation. By the definition of F_n and F ((7) and (8)), we have that

$$\begin{aligned} F_n(y) - F(y) &= \frac{1}{T_n(C)} \sum_{i=1}^{T_n(C)} (\mathbb{I}\{X_{\sigma_C(i)} \leq y\} - F(y)) \\ &= \frac{1}{T_n(C)} \sum_{i=0}^n (\mathbb{I}_C\{X_t \leq y\} - \mathbb{I}_C\{X_i\} F(y)) \\ &= \frac{1}{T_n(C)} \left(\widetilde{g}_y(\mathcal{B}_0) + \sum_{i=1}^{T(n)} \widetilde{g}_y(\mathcal{B}_i) + \widetilde{g}_y(\mathcal{B}_{(n)}) \right), \end{aligned}$$

therefore,

$$\sqrt{T_n(C)} (F_n(y) - F(y)) = \frac{\widetilde{g}_y(\mathcal{B}_0)}{\sqrt{T_n(C)}} + \frac{\sum_{i=1}^{T(n)} \widetilde{g}_y(\mathcal{B}_i)}{\sqrt{T_n(C)}} + \frac{\widetilde{g}_y(\mathcal{B}_{(n)})}{\sqrt{T_n(C)}}.$$

Notice that $|\widetilde{g}_y(\mathcal{B}_0)| \leq 2\ell_C(\mathcal{B}_0) < +\infty$ and $T_n(C) \rightarrow +\infty$ almost surely, therefore, the first term in the last equation converges almost surely to 0 uniformly in y . For the last term, we have that

$$\frac{|\widetilde{g}_y(\mathcal{B}_{(n)})|}{\sqrt{T_n(C)}} \leq \frac{2\ell_C(\mathcal{B}_{T(n)})}{\sqrt{T_n(C)}} = 2\sqrt{\frac{T(n)}{T_n(C)} \frac{\ell_C(\mathcal{B}_{T(n)})}{\sqrt{T(n)}}},$$

by (B4), the expectation of $\ell_C^2(\mathcal{B}_1)$ is finite, then, Lemma 1 in [2] shows that $\frac{\ell_C^2(\mathcal{B}_n)}{n} \rightarrow 0$ a.s. which implies that $\frac{\ell_C(\mathcal{B}_n)}{\sqrt{n}}$ also converges to 0 a.s. Since $T(n) \rightarrow +\infty$ a.s., by Theorem 6.8.1 in [19] we have $\frac{\ell_C(\mathcal{B}_{T(n)})}{\sqrt{T(n)}} \rightarrow 0$ almost surely. Joining this with the almost sure convergence of $\frac{T(n)}{T_n(C)}$ to a positive constant (see Theorem 1.1) we obtain that $\frac{|\widetilde{g}_y(\mathcal{B}_{(n)})|}{\sqrt{T_n(C)}}$ converges almost surely to 0 uniformly in y . Therefore,

$$(75) \quad \sqrt{T_n(C)} (F_n(y) - F(y)) = \frac{\sum_{i=1}^{T(n)} \widetilde{g}_y(\mathcal{B}_i)}{\sqrt{T(n)}} + o_P(1).$$

where we have used that $\frac{T_n(C)}{T(n)}$ converges almost surely to a positive constant to use $T(n)$ instead of $T_n(C)$.

Then, (17) will be proved if we show that, for ε small enough

$$(76) \quad \sup_{|y-x_0| \leq \varepsilon} \frac{\left| \sum_{i=1}^{T(n)} \tilde{g}_y(\mathcal{B}_i) \right|}{\sqrt{T(n)}} = O_p(1).$$

Fix $\eta > 0$ arbitrarily. By Lemma 6.7 and Slutsky's theorem, we can find positive numbers $\underline{a}_\eta, \bar{a}_\eta$ and an integer N_η such that $P(\mathcal{E}_n) \geq 1 - \frac{\eta}{2}$ for all $n \geq N_\eta$, where

$$(77) \quad \mathcal{E}_n = \{ \underline{a}_\eta u(n) \leq T(n) \leq \bar{a}_\eta u(n) \}.$$

Define $W_n(\varepsilon) = \sup_{|y-x_0| \leq \varepsilon} \left| \sum_{i=1}^n \tilde{g}_y(\mathcal{B}_i) \right|$ and let M_η be a fixed positive number. Then, for all $n \geq N_\eta$

$$(78) \quad \begin{aligned} \mathbb{P} \left(\frac{1}{\sqrt{T(n)}} W_{T(n)} > M_\eta \right) &< \mathbb{P} \left(\left\{ \frac{1}{\sqrt{T(n)}} W_{T(n)} > M_\eta \right\} \cap \mathcal{E}_n \right) + 1 - \mathbb{P}(\mathcal{E}_n) \\ &< \mathbb{P} \left(\left\{ \frac{1}{\sqrt{T(n)}} W_{T(n)} > M_\eta \right\} \cap \mathcal{E}_n \right) + \frac{\eta}{2}. \end{aligned}$$

On \mathcal{E}_n , $\underline{a}_\eta u(n) \leq T(n) \leq \bar{a}_\eta u(n)$, therefore for all $n \geq N_\eta$

$$(79) \quad \begin{aligned} \mathbb{P} \left(\left\{ \frac{1}{\sqrt{T(n)}} W_{T(n)} > M_\eta \right\} \cap \mathcal{E}_n \right) &< \mathbb{P} \left(\left\{ \frac{1}{\sqrt{\underline{a}_\eta u(n)}} \max_{1 \leq k \leq \bar{a}_\eta u(n)} W_k > M_\eta \right\} \cap \mathcal{E}_n \right), \\ &< \mathbb{P} \left(\frac{1}{\sqrt{\underline{a}_\eta u(n)}} \max_{1 \leq k \leq \bar{a}_\eta u(n)} W_k > M_\eta \right). \end{aligned}$$

The random variables $\{\tilde{g}_{(\cdot)}(\mathcal{B}_k)\}_{k=1}^{\bar{a}_\eta u(n)}$ are i.i.d., therefore, by Montgomery-Smith's inequality (Lemma 4 in [1]), there exists a universal constant K such that for all $n \geq N_\eta$,

$$(80) \quad \begin{aligned} \mathbb{P} \left(\frac{1}{\sqrt{\underline{a}_\eta u(n)}} \max_{1 \leq k \leq \bar{a}_\eta u(n)} W_k > M_\eta \right) &< K \mathbb{P} \left(\frac{1}{\sqrt{\underline{a}_\eta u(n)}} W_{\bar{a}_\eta u(n)} > \frac{M_\eta}{K} \right), \\ &< K \mathbb{P} \left(\frac{1}{\sqrt{\underline{a}_\eta u(n)}} \sup_{|y-x_0| \leq \varepsilon} \left| \sum_{i=1}^{\bar{a}_\eta u(n)} \tilde{g}_y(\mathcal{B}_i) \right| > \frac{M_\eta}{K} \right). \end{aligned}$$

For an arbitrary set T , let $\ell^{+\infty}(T)$ be the space of all uniformly bounded, real functions on T , equipped with the uniform norm. Weak convergence to a tight process in this space is characterized by asymptotic tightness plus convergence of marginals (see Chapter 1.5 in [34]).

The class of functions $\mathcal{G} - F = \{g_y(\cdot) - F(y)\}_{y \in \mathbb{R}}$ is VC with constant envelope 2, hence, by Lemma 5.2, the class of functions $\widehat{\mathcal{G} - F}$ is also VC and has $2\ell_C$ as envelope. $\mathbb{E}\ell_C^2(\mathcal{B}_1)$ is finite (by (B4)), therefore, by Theorem 2.5 in [25], $\widehat{\mathcal{G} - F}$ is Donsker. Then, the process $\frac{1}{\sqrt{\underline{a}_\eta u(n)}} \left| \sum_{i=1}^{\bar{a}_\eta u(n)} \tilde{g}_y(\mathcal{B}_i) \right|$ converges weakly in $\ell^\infty[\widehat{\mathcal{G} - F}]$ to a tight process Z . The map

$y \mapsto \|y\|_\infty$ from $\ell^\infty \left[\widehat{\mathcal{G}} - F \right]$ to \mathbb{R} is continuous with respect to the supremum norm (cf. pp 278 of [33]), therefore, $\frac{1}{\sqrt{\underline{a}_\eta u(n)}} \sup_{|y-x_0| \leq \varepsilon} \left| \sum_{i=1}^{\bar{a}_\eta u(n)} \tilde{g}_y(\mathcal{B}_i) \right|$ converges in distribution to $\sup_{|y-x_0| \leq \varepsilon} Z(y)$, hence, we can find V_η and N'_η such that

$$(81) \quad \mathbb{P} \left(\frac{1}{\sqrt{\underline{a}_\eta u(n)}} \sup_{|y-x_0| \leq \varepsilon} \left| \sum_{i=1}^{\bar{a}_\eta u(n)} \tilde{g}_y(\mathcal{B}_i) \right| > V_\eta \right) < \frac{\eta}{2K}, \quad \forall n > N'_\eta.$$

Choosing $M_\eta = KV_\eta$ in 81 and joining (80), (79) and (78), completes the proof of (17).

Now we proceed to prove (18). Let η be fixed, by (17) and Lemma 6.7, we can find ε' , M'_η and N'_η such that

$$(82) \quad \mathbb{P} \left(\sqrt{T_n(C)} \sup_{|y-x_0| \leq \varepsilon'} |F_n(y) - F(y)| > M'_\eta \right) < \frac{\eta}{4} \quad \forall n \geq N'_\eta$$

$$(83) \quad \mathbb{P}(\mathcal{D}_n) \geq 1 - \frac{\eta}{2} \quad \forall n \geq N'_\eta$$

where $\mathcal{D}_n = \{\underline{a}_\eta u(n) \leq T_n(C) \leq \bar{a}_\eta u(n)\}$. Define the sets

$$U_n = \left\{ \sqrt{T_n(C)} \sup_{|p-F(x_0)| \leq \varepsilon} |F_n^{-1}(p) - F^{-1}(p)| > M_\eta \right\},$$

$$U_n^1 = \left\{ \exists p \in [F(x_0) - \varepsilon, F(x_0) + \varepsilon] : F_n^{-1}(p) - F^{-1}(p) > \frac{M_\eta}{\sqrt{T_n(C)}} \right\},$$

$$U_n^2 = \left\{ \exists p \in [F(x_0) - \varepsilon, F(x_0) + \varepsilon] : F^{-1}(p) - F_n^{-1}(p) > \frac{M_\eta}{\sqrt{T_n(C)}} \right\}.$$

where ε and M_η are constants that will be specified later.

On $U_n^1 \cap \mathcal{D}_n$, $F_n^{-1}(p) > \frac{M_\eta}{\sqrt{T_n(C)}} + F^{-1}(p) > \frac{M_\eta}{\sqrt{\bar{a}_\eta u(n)}} + F^{-1}(p)$, hence,

$$(84) \quad F_n \left(\frac{M_\eta}{\sqrt{\bar{a}_\eta u(n)}} + F^{-1}(p) \right) \leq F_n(F_n^{-1}(p)) \leq p + \frac{1}{T_n(C)} \leq p + \frac{1}{\underline{a}_\eta u(n)}.$$

Assumption (B2) indicates that F has bounded derivative in C , take K_1 as the maximum value of this derivative in C , then, the Mean Value Theorem implies that

$$p = F(F^{-1}(p)) = F \left(\frac{M_\eta}{\sqrt{\bar{a}_\eta u(n)}} + F^{-1}(p) \right) - \frac{F'(\theta_p) M_\eta}{\sqrt{\bar{a}_\eta u(n)}} \leq F \left(\frac{M_\eta}{\sqrt{\bar{a}_\eta u(n)}} + F^{-1}(p) \right) - \frac{K_1 M_\eta}{\sqrt{\bar{a}_\eta u(n)}}.$$

After plugging this into (84) we get

$$F \left(\frac{M_\eta}{\sqrt{\bar{a}_\eta u(n)}} + F^{-1}(p) \right) - F_n \left(\frac{M_\eta}{\sqrt{\bar{a}_\eta u(n)}} + F^{-1}(p) \right) \geq \frac{K_1 M_\eta}{\sqrt{\bar{a}_\eta u(n)}} - \frac{1}{\underline{a}_\eta u(n)}.$$

Because $u(n) \rightarrow +\infty$, we can find N_1 such that $\sqrt{\frac{\bar{a}_n}{u(n)}} \frac{1}{\underline{a}_n K_1} < 1$ for all $n \geq N_1$, taking M_η bigger than $\frac{M'_\eta}{K_1} \sqrt{\frac{\bar{a}_n}{\underline{a}_n}} + 1$ and using that $T_n(C) \leq \underline{a}_\eta u(n)$ on \mathcal{D}_n , we obtain, for all $n \geq N_1$

$$(85) \quad F\left(\frac{M_\eta}{\sqrt{\bar{a}_\eta u(n)}} + F^{-1}(p)\right) - F_n\left(\frac{M_\eta}{\sqrt{\bar{a}_\eta u(n)}} + F^{-1}(p)\right) > \frac{M'_\eta}{\sqrt{T_n(C)}}.$$

Let $N_{2,\eta}$ be such that $\frac{M_\eta}{\sqrt{\bar{a}_\eta u(n)}} < \frac{\varepsilon'}{2}$ for $n \geq N_{2,\eta}$. By the continuity of F^{-1} in $F(x_0)$ there exists $\varepsilon > 0$ such that $|F^{-1}(p) - x_0| \leq \frac{\varepsilon'}{2}$ for all p in $[F(x_0) - \varepsilon, F(x_0) + \varepsilon]$, therefore, the triangular inequality implies that $\frac{M_\eta}{\sqrt{\bar{a}_\eta u(n)}} + F^{-1}(p)$ lies in the interval $[x_0 - \varepsilon', x_0 + \varepsilon']$ for all $n \geq N_\eta = \max(N_1, N_{2,\eta})$. This, alongside (85), shows that for all $n \geq N_\eta$

$$\begin{aligned} U_n^1 \cap \mathcal{D}_n &\subseteq \left\{ \exists y \in [x_0 - \varepsilon', x_0 + \varepsilon'] : F(y) - F_n(y) > \frac{M'_\eta}{\sqrt{T_n(C)}} \right\} \\ &\subseteq \left\{ \sqrt{T_n(C)} \sup_{|y-x_0| \leq \varepsilon'} |F_n(y) - F(y)| > M'_\eta \right\}. \end{aligned}$$

By a similar argument, it can be shown that

$$U_n^2 \cap \mathcal{D}_n \subseteq \left\{ \exists y \in [x_0 - \varepsilon', x_0 + \varepsilon'] : F_n(y) - F(y) > \frac{M'_\eta}{\sqrt{T_n(C)}} \right\} \quad \forall n \geq N_\eta.$$

Using (82) and $U_n = U_n^1 \cup U_n^2$ we obtain that $\mathbb{P}(U_n \cap \mathcal{D}_n) \leq \frac{\eta}{2}$ for all $n \geq N_\eta$. Equation (18) now follows by (83).

Supplement of “Harris recurrent Markov chains and nonlinear monotone cointegrated models”

This is the supplementary material associated with the present article.

REFERENCES

[1] ADAMCZAK, R. (2008). A tail inequality for suprema of unbounded empirical processes with applications to Markov chains. *Electronic Journal of Probability* **13** 1000 – 1034.

[2] ATHREYA, K. B. and ROY, V. (2016). General Glivenko–Cantelli theorems. *Stat* **5** 306–311.

[3] BARLOW, R. E., BARTHOLOMEW, D. J., BREMNER, J. and BRUNK, H. D. (1972). *Statistical inference under order restrictions: The theory and application of isotonic regression*. Wiley New York.

[4] BERTAIL, P., DUROT, C. and FERNÁNDEZ, C. (2023). Supplement to “Harris recurrent Markov chains and nonlinear monotone cointegrated models.”. [https://doi.org/10.1214/\[providedbytypesetter\]](https://doi.org/10.1214/[providedbytypesetter])

[5] BERTAIL, P. and PORTIER, F. (2019). Rademacher complexity for Markov chains: Applications to kernel smoothing and Metropolis–Hastings. *Bernoulli* **25** 3912–3938.

[6] BINGHAM, N. H., GOLDIE, C. M. and TEUGELS, J. L. (1987). *Regular variation. Encyclopedia of mathematics and its applications* **27**. Cambridge University Press.

[7] CAI, B. and TJØSTHEIM, D. (2015). Nonparametric Regression Estimation for Multivariate Null Recurrent Processes. *Econometrics* **3** 265–288.

[8] CAI, Z. and OULD-SAIID, E. (2003). Local M-estimator for nonparametric time series. *Statistics and Probability Letters* **65** 433–449.

[9] CHEN, X. (1999). How Often Does a Harris Recurrent Markov Chain Recur? *The Annals of Probability* **27**.

[10] CHEN, X. (2000). On the limit laws of the second order for additive functionals of Harris recurrent Markov chains. *Probability Theory and Related Fields* **116**.

[11] DEATON, A. and MUELLBAUER, J. (1980). *Economics and Consumer Behavior*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511805653>

[12] DOUC, R., MOULINES, E., PRIOURET, P. and SOULIER, P. (2018). *Markov chains. Springer Series in Operations Research and Financial Engineering*. Springer.

- [13] DUROT, C. and TOCQUET, A.-S. (2003). On the distance between the empirical process and its concave majorant in a monotone regression framework. In *Annales de l'Institut Henri Poincaré (B) Probability and Statistics* **39** 217–240. Elsevier.
- [14] ENGLE, R. F. and GRANGER, C. W. J. (1987). Co-Integration and Error Correction: Representation, Estimation, and Testing. *Econometrica* **55** 251–276.
- [15] FEIGIN, P. D. and TWEEDIE, R. L. (1985). Random Coefficient Autoregressive Processes: a Markov Chain Analysis of Stationarity and Finiteness of Moments. *Journal of Time Series Analysis* **6**.
- [16] GRANGER, C. W. J. (1981). Some properties of time series data and their use in econometric model specification. *Journal of Econometrics* **16** 121–130.
- [17] GRANGER, C. and TERÄSVIRTA, T. (1993). *Modelling Non-Linear Economic Relationships*. Oxford University Press.
- [18] GROENEBOOM, P. and JONGBLOED, G. (2014). *Nonparametric Estimation under Shape Constraints: Estimators, Algorithms and Asymptotics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.
- [19] GUT, A. (2013). *Probability : a graduate course*, 2nd ed ed. Springer texts in statistics. Springer.
- [20] HANSEN, B. E. and SEO, B. (2002). Testing for two-regime threshold cointegration in vector error-correction models. *Journal of Econometrics* **110** 293–318. Long memory and nonlinear time series. [https://doi.org/10.1016/S0304-4076\(02\)00097-0](https://doi.org/10.1016/S0304-4076(02)00097-0)
- [21] JOHANSEN, S. (1988). Statistical analysis of cointegrating vectors. *Journal of Economic Dynamics and Control* **12** 231.
- [22] JOHANSEN, S. (1991). Estimation and hypothesis testing of cointegration vectors in Gaussian vector autoregressive models. *Econometrica* **59** 1551.
- [23] KARLSEN, H. A., MYKLEBUST, T. and TJØSTHEIM, D. (2007). Nonparametric estimation in a nonlinear cointegration type model. *The Annals of Statistics* **35** 252–299.
- [24] KARLSEN, H. A. and TJØSTHEIM, D. (2001). Nonparametric estimation in null recurrent time series. *The Annals of Statistics* **29**.
- [25] KOSOROK, M. R. (2008). *Introduction to empirical processes and semiparametric inference*, 1 ed. Springer Series in Statistics. Springer.
- [26] LIN, Z., LI, D. and CHEN, J. (2009). Local linear M-estimators in null recurrent times series. *Statistica Sinica* **19** 1683–1703.
- [27] MEYN, S., TWEEDIE, R. and GLYNN, P. (2009). *Markov chains and stochastic stability*, 2 ed. Cambridge Mathematical Library. Cambridge University Press.
- [28] NUMMELIN, E. (1978). A splitting technique for Harris recurrent Markov chains. *Probability Theory and Related Fields* **43**.
- [29] NUMMELIN, E. (1984). *General Irreducible Markov Chains and Non-Negative Operators*. Cambridge Tracts in Mathematics 83. Cambridge University Press.
- [30] PHILLIPS, P. C. B. (1991). Optimal inference in cointegrated systems. *Econometrica* **59** 283.
- [31] PHILLIPS, P. C. B. and SOLO, V. (1992). Asymptotics for linear processes. *The Annals of Statistics* **20** 971.
- [32] STIGLER, M. (2020). Threshold cointegration: overview and implementation in R. *Handbook of Statistics*, **42** 229–264.
- [33] VAN DER VAART, A. W. (2000). *Asymptotic statistics*. Cambridge university press.
- [34] VAN DER VAART, A. W. and WELLNER, J. A. (1996). *Weak Convergence and Empirical Processes*. Springer.
- [35] WANG, Q. (2015). *Limit Theorems for Nonlinear Cointegrating Regression*. World Scientific.
- [36] WANG, Q. and PHILLIPS, P. C. B. (2009). Asymptotic Theory for Local Time Density Estimation and Nonparametric Cointegrating Regression. *Econometric Theory* **25** 710–738.
- [37] WANG, Q. and PHILLIPS, P. C. B. (2009). Structural Nonparametric Cointegrating Regression. *Econometrica* **77** 1901–1948.