

A one-domain approach for flow near porous media boundaries

Didier Lasseux, Francisco J. Valdès-Parada

▶ To cite this version:

Didier Lasseux, Francisco J. Valdès-Parada. A one-domain approach for flow near porous media boundaries. InterPore 2022, InterPore, May 2022, Abu Dhabi, United Arab Emirates. pp.273-274. hal-04300288

HAL Id: hal-04300288

https://hal.science/hal-04300288

Submitted on 22 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A One-domain approach for flow near porous media boundaries

Authors: Francisco J. Valdés-Parada¹; Didier Lasseux²

Corresponding Authors: didier.lasseux@u-bordeaux.fr, iqfv@xanum.uam.mx

Momentum transport near porous media boundaries has been the subject of intense work for more than half a century since the pioneering work of Beavers and Joseph in 1967 [1,2]. Currently, there are two modeling strategies to study this subject: 1) A one-domain approach (ODA), where the spatial variations of average properties are accounted for in the transition zone and 2) a two-domain approach (TDA), where these variations are collapsed in a boundary condition [3]. The TDA has received much more attention than the ODA due to the practical use of jump conditions, whereas a closed-form of the ODA has been typically made obscure and dependent of pore-scale solutions in the entire system. Nevertheless, the derivation of a jump condition for the TDA requires that the ODA be developed first. In this work, a practical formulation of the ODA is presented, that is valid for a porous medium sharing boundaries with a free fluid [4], another porous medium or a solid material. It is has the simple structure of a Darcy-like model involving a position-dependent permeability tensor that is predicted from the solution of an ancillary closure problem. The performance of the model is exemplified in the vicinity of a fluid channel, near the boundary of two porous media (either in direct contact or separated by a fracture) and for flow between a porous medium and an impervious wall [5]. In all cases, the model predictions are validated with pore-scale simulations showing an excellent agreement. Furthermore, in the case of a fluid-porous medium boundary, the model is also validated with experimental data [4]. The simplicity and versatility of the new ODA model provided here make it an interesting alternative to existing approaches in the literature.

References

- G. Beavers and D. Joseph, Boundary conditions at a naturally permeable wall J. Fluid Mech. 30, 197 (1967).
- [2] D. A. Nield, The Beavers–Joseph boundary condition and related matters: A historical and critical note, Transp. Porous Media 78, 537–540 (2009).
- [3] M. Chandesris and D. Jamet, Boundary conditions at a planar fluid–porous interface for a Poiseuille flow, Int. J. Heat Mass Transfer 49, 2137–2150 (2006).

Page 273

InterPore2022 / Book of Abstracts

[4] F. J. Valdés-Parada and D. Lasseux, A novel one-domain approach for modeling flow in a fluid-porous system including inertia and slip effects, Phys. Fluids 33, 022106 (2021).
[5] F. J. Valdés-Parada and D. Lasseux, Flow near porous media boundaries including inertia and slip: A one-domain approach, Phys. Fluids 33, 073612 (2021).

Acceptance of the Terms & Conditions:

Click here to agree

MDPI Energies Student Poster Award:

No, do not submit my presenation for the student posters award.

Country:

France

References:

Time Block Preference:

Time Block A (09:00-12:00 CET)

Participation:

Online

¹ Universidad Autónoma Metropolitana-Iztapalapa

² CNRS - Univ. Bordeaux