

Measurements of hydroperoxy and total organic peroxy radicals in an Indiana forest using laser induced fluorescence

C.M.F. Rosales, Ahmad Lahib, Brandon Bottorff, Emily K. Reidy, V. Kumar, Sébastien Dusanter, Alexandre Tomas, Philip Stevens


▶ To cite this version:

C.M.F. Rosales, Ahmad Lahib, Brandon Bottorff, Emily K. Reidy, V. Kumar, et al.. Measurements of hydroperoxy and total organic peroxy radicals in an Indiana forest using laser induced fluorescence. AGU Fall Meeting, Dec 2019, San francisco, USA, United States. hal-04300879


HAL Id: hal-04300879 https://hal.science/hal-04300879v1

Submitted on 24 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



MEASUREMENTS OF HYDROPEROXY & TOTAL ORGANIC PEROXY RADICALS IN AN INDIANA FOREST USING LASER INDUCED-FLUORESCENCE



Colleen Marciel F. Rosales^{1*}, Ahmad Lahib^{1,2}, Brandon P. Bottorff³, Emily K. Reidy³, Vinay Kumar¹, Sebastien Dusanter², Alexandre Tomas², and Philip Stevens^{1,3}

¹O'Neill School of Public and Environmental Affairs, and ³Department of Chemistry, Indiana University

²Atmospheric Sciences and Environmental Engineering, IMT Lille Douai, University of Lille, France

*cfrosale@iu.edu

Poster no. A11K-2702

AMBIENT CHEMISTRY OF HYDROPEROXY (HO₂) AND PEROXY (RO₂) RADICALS

The OH radical initiates the oxidation of volatile organic compounds (VOCs), leading to HO2 and RO2 radicals whose chemistry, in the presence of nitrogen oxides (NOx), can lead to the formation of ozone and secondary organic aerosols in the environment. Previous measurements of these radicals in forest environments characterized by high mixing ratios of NO_x have shown serious discrepancies with modeled predictions, bringing into question our understanding of isoprene oxidation in the atmosphere.

LASER-INDUCED FLUORESCENCE-FLUORESCENCE ASSAY BY GAS EXPANSION (LIF-FAGE)

ROxLIF configuration

• LIF-FAGE inlet was modified by attaching a 45-cm reactor flow tube (Fig.3, right)

to convert RO_x and HO_x species to HO_2 , which is detected in the detection cell

• RO_x and HO_x modes are achieved by addition of CO and NO (Fig. 2), under low

• Detection limits for HO_2 and RO_2 were $(2-4)\times10^7$ cm⁻³, and $(1.5-3)\times10^7$ cm⁻³, re-

pressure conditions (~30 Torr) before reaching the detection cell.

spectively, for S/N = 1 and 5-minute measurements.

- Measures OH & HO₂ by excitation and fluorescence of OH at 308 nm. Fluorescence is detected by a microchannel plate PMT (Fig. 1).
- Air is sampled via an inlet and is expanded under vacuum (~3.5 to 4 Torr).
- HO₂ in the detection cell is measured by addition of NO to produce OH within the cell: $HO_2 + NO \rightarrow OH + NO_2$
- A low NO concentration (\sim 3.6 to 7 x 10¹¹ cm⁻³) is used in the detection cell for HO₂ measurements to minimize interference from RO₂ radicals (Lew, 2018). HO₂ to OH conversion efficiency in the detection cell is ~14%.

(Fuchs, Holland et al. 2008).




Figure 2. Chemistry in the reactor flow tube. Propagation reactions are shown in red, while termination reactions are shown in gray. Loss reactions are mostly wall losses or recombination with

 OH is converted to HO₂ by the addition of CO: $OH + CO \rightarrow H + CO_2$ $H + O_2 \rightarrow HO_2$

 RO₂ is measured by adding NO into the reactor: $RO_2 + NO \rightarrow RO + NO_2$

 $RO + O_2 \rightarrow R'O + HO_2$

Figure 4. (a, top) Calibration source used to

right) Calculated RO₂ and HO₂ sensitivity as func-

generate HO₂ and RO₂ radicals (Lew, 2018) (b,

tion of water %.

UDT 555-UV Quartz window 7////// 185 nm filter $a_{2} = -1.3 \times 10^{-7} \times + 5.7 \times 10^{-9}$

 $|C_{RO_2} = -1.5 \times 10^{-7} \times + 7.2 \times 10^{-5}|$ Water, %

Figure 1. Diagram of the base LIF-FAGE detection cell. When the reactor is attached, the external C_3F_6 ring is removed (Dusanter,

RO_x mode: NO + CO $10^{\hat{}}_{x}$ mode: CO + N

Figure 3. (Left) ROxLIF deployed to the IURTP site. (Right) Diagram of the ROxLIF attachment. The top inlet is a

Conversion Efficiency and Sensitivity of ROxLIF pinhole while the bottom inlet is conical. From here, the sam • Known concentrations of OH and HO₂ were generated using the calibration source ple is directed to the detection cell shown in Fig.1. shown in Fig 4a. To generate RO₂, ethane was added to the calibration source and re-

act with OH and introduced into the reactor. The RO₂ concentration was assumed to be equal to the initial OH concentration. • Sensitivity to RO₂, C_{RO2}, was calculated from ethane-RO₂ measurements (Fig 4b).


• Chemistry within the reactor have been modeled using Master Chemical Mechanism v3.3.1, wall loss rate constants from experiments and literature (Fuchs, 2008; Mihele, 1999; Sander, 2003). Residence times of RO2 species generated from methane, toluene and isoprene have been found to be within 10% of the residence time for ethane-RO2. Thus, it is reasonable to assume a generic C_{RO2} for these RO₂ species.

◆ Experimental ••• 0.4 s -2.0 0.0 0.2 0.4 0.6 0.8 CO concentration, % NO concentration, ppm

Figure 5. Measured (points) and modeled (dotted lines) instrument response to RO_x as function of (a) NO and (b) CO concentration added to reactor. Models were done for different residence times in the reactor. For the measured RO₂, ethane was used as source of RO₂.

PREVIOUS VS. CURRENT MEASUREMENTS AT THE IURTP SITE

- The Indiana University Research and Teaching Preserve (IURTP) is a mixed deciduous forest consisting of sugar maple, sycamore, tulip polar, ash and hickory trees (Lew et al., 2019).
- It is located ~2.5 km NE of the center of the Indiana University campus, and 1 km from the IN 45/46 bypass at the northern perimeter.










Figure 7. Total HO₂+RO₂ measurements at the IURTP site in 2017 and 2019 compared with HO_2 * measurements by the LIF-FAGE in 2015. HO_2 * is HO_2 + αRO_2 , where α is a fraction of alkene and aromatic hydrocarbons.

- Isoprene concentrations in 2019 were greater than that measured in 2017, but similar to that measured in 2015 (Fig. 8c). NO₂ concentrations in 2019 were lower than measurements in both 2015 and 2017 (Fig. 8b) suggesting that NO concentrations were also lower compared to 2015 and 2017.
- The higher HO₂ and RO₂ concentrations observed in 2019 may be consistent with lower NO concentrations at this site compared with 2017 and 2015 (Figs. 6 and 7).

Supporting Measurements: Isoprene, Ozone and NO_x








Figure 8. Concentrations of (a) NO, (b) NO₂, (c) Isoprene and (d) ozone at the IURTP site in 2015, 2017 and 2019. NO concentrations in 2019 are below LOD and are not shown in (a). Shaded areas, when visible, stand for standard error.

REFERENCES

Dusanter, S., D. Vimal, P. S. Stevens, R. Volkamer and L. T. Molina (2009). Atmos. Chem. Phys. 9: 1665-1685. Lew, M. L., P. S. Rickly, B. P. Bottorff, S. Sklaveniti, T. Leonardis, N. Locoge, S. Dusanter, S. Kundu, E. Wood and P. S. Stevens (2019). Atmos. Chem. Phys. Discuss. 2019: 1-35.

rkholder, J.B., Sander, S.P., Abbatt, J., Barker, J.R., Huie, R.E., Kolb C.E., Kurylo, M.J., Orkin, V.L. Wilmouth, D.M. and Wine, P.H., (2015) JPL Publication 15-10, Jet Propulsion Laboratory, Pasadena, http://jpldataeval.jpl.nasa.gov. Wolfe, G. M., M. R. Marvin, S. J. Roberts, K. R. Travis and J. Liao (2016). Geosci. Model Dev. 9(9): 3309-3319.

DAILY MEASUREMENTS & MODEL COMPARISONS





Figure 9. Time series of HO_2 and RO_2 measurements during the campaign.



the supporting measurements used as model constraints.

Modeling (FOAM) (Wolfe, 2016). The model underestimates HO₂ concentrations on the 23rd, but seem to reproduce them well on

HO₂ and RO₂ show a strong diur-

nal trend that is characteristic of

=RO₂ except for July 24th (Fig 9.),

To constrain a zero-dimensional

box model based on the Master

v.3.3.1, supporting measurements

of photolysis rates, VOCs, O₃ and

Calculations were done using the

Framework for O-D Atmospheric

Chemical Mechanism (MCM)

NO₂ (Fig. 10a-e) were used.

This day, isoprene was lower than

photolytically-driven reactions.

For most days, measured HO₂

the previous days (Fig 10a).

 On the other hand, RO₂ is underestimated on the 23rd but overestimated on the 24th especially dur-

Figure 10. Measured and modeled HO₂ and RO₂ concentrations are shown on (f), while the top panels (a-e) show

ing the night.

CONCLUSIONS

This work presents the first successful ambient measurements of HO2 and RO2 in a forested environment by the newly constructed IU ROxLIF-FAGE system. The conversion efficiency for different RO2 species within the flow tube have been estimated to be within ~10% of each other and have been detected using the ROxLIF method. Ambient concentrations of HO₂ and RO₂ were found to be around $4x10^8$ to $2x10^9$ molecules cm⁻³, similar to previous measurements at this site. Modeled HO₂ concentrations using MCM v.3.3.1 agree or underestimate the measurements on two days, while modeled RO2 concentrations either under- or overestimate the measurements by a factor of the 24th.