

Associations Among Menstrual Cycle Length, Coronavirus Disease 2019 (COVID-19), and Vaccination

Alexandra Alvergne, PhD, Emily Boniface, MPH, Blair Darney, PhD, MPH, Amanda Shea, PhD, Kirsten Weber, PhD, Cécile Ventola, PhD, Virginia J. Vitzthum, PhD, and Alison Edelman, MD, MPH

OBJECTIVE: To assess whether coronavirus disease 2019 (COVID-19) is associated with menstrual cycle length changes and, if so, how that compares with those undergoing vaccination or no event (control).

METHODS: We conducted a retrospective cohort analysis in which we analyzed prospectively tracked cycle-

length data from users of a period tracker application who also responded to a survey regarding COVID-19 symptoms and vaccination. We restricted our sample to users aged 16–45 years, with normal cycle lengths (24–38 days) and regular tracking behavior during the five cycles around COVID-19 symptoms or vaccination or a similar time period for those experiencing no event (control group). We calculated the within-user change in cycle length (days) from the three consecutive cycles preevent average (either vaccination, disease, or neither; cycles 1–3) to the event (cycle 4) and postevent (cycle 5) cycles. We used mixed-effects models to estimate the age- and country-adjusted difference in change in cycle length across the groups.

RESULTS: We included 6,514 users from 110 countries representing 32,570 cycles (COVID-19 symptoms: 1,450; COVID-19 vaccination: 4,643; control: 421). The COVID-19 cohort experienced a 1.45-day adjusted increase in cycle length during cycle 4 (COVID-19) compared with their three preevent cycles (95% CI 0.86–2.04). The vaccinated group experienced a 1.14-day adjusted increase in cycle length during cycle 4 (COVID-19 vaccine) compared with their preevent average (95% CI 0.60–1.69). The control group (neither vaccine nor disease) experienced a 0.68-day decrease (95% CI –1.18 to –0.19) in a similar time period. Post hoc tests showed no significant differences in the magnitude of changes between the COVID-19 and vaccination cohorts. In both cohorts, cycle length changes disappeared in the postevent cycle.

CONCLUSION: Experiencing COVID-19 is associated with a small change in cycle length similar to COVID-19 vaccination. These changes resolve quickly within the next cycle.

(*Obstet Gynecol* 2023;00:1–9)

DOI: 10.1097/AOG.0000000000005343

The coronavirus disease 2019 (COVID-19) pandemic has revealed many significant knowledge gaps, notably a lack of information regarding the

From the Institute for Evolutionary Sciences, Montpellier University, Montpellier, France; the School of Anthropology and Museum Ethnography, University of Oxford, Oxford, United Kingdom; the Department of Obstetrics and Gynecology, Oregon Health & Science University, and the OHSU-PSU School of Public Health, Portland, Oregon; the National Institute of Public Health (INSP), Center for Population Health (CISP), Cuernavaca, Morelos, Mexico; and Clue by BioWink GmbH, Berlin, Germany.

Research reported in this publication was funded by the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) and the National Institutes of Health Office of Research on Women's Health - NIH NICHD089957 Supplement.

The authors thank all the participants for their time in answering the surveys.

Each author has confirmed compliance with the journal's requirements for authorship.

Corresponding author: Alexandra Alvergne, PhD, Institute for Evolutionary Sciences, Montpellier University, Montpellier, France; alexandra.alvergne@umontpellier.fr.

Financial Disclosure

Kirsten Weber works for Clue by BioWink, and they received funding from Alison Edelman's grant (NIH NICHD089957 Supplement) for project support. Virginia J. Vitzthum is a paid consultant for Clue by BioWink, GmbH, Berlin, Germany, during the previous 36 months (and since 2017). Blair Darney's institution receives research support from Merck/Organon. Blair Darney reports honoraria and travel funding from SFP and ACOG. Alison Edelman reports receiving honoraria and travel reimbursement from ACOG, WHO, and the CDC for guideline development group activities. Alison Edelman also receives royalties from Up to Date, Inc. Oregon Health & Science University (OHSU) receives research funding from OHSU Foundation, Merck, HRA Pharma, the Bill & Melinda Gates Foundation, and NIH for which Alison Edelman is the principal investigator. The other authors did not report any potential conflicts of interest.

Copyright © 2023 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

ISSN: 0029-7844/23

potential effects of vaccines and infection on the menstrual cycle.^{1–3} Menstruation is increasingly being recognized as a critically important patient-reported outcome,^{4–6} in part due to the initial scrutiny placed on the association of COVID-19 vaccines and menstrual cycle disturbances.^{7–15} Yet, as the prevalence of COVID-19 has increased, growing reports of menstrual disturbances after severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection^{16,17} and with long-COVID¹⁸ are occurring.

The immune and reproductive systems are known to interact with each other,^{19,20} and temporary disruption of the menstrual cycle is seen with acute infection and febrile episodes. However, previous literature on the association between COVID-19 and menstrual cycle changes is scarce and inconsistent, limited by small samples, recall bias, lack of comparison (unexposed) groups, or small subgroups of participants with COVID-19.^{21–23}

Menstrual cycles have their own normal inherent variability,²⁴ which makes it particularly challenging to determine whether an exposure causes a change without access to prospectively collected population-level data before and after the exposure. Similar to prior work,^{8–10} we present an analysis of prospectively collected menstrual cycle tracking data to assess whether COVID-19 is associated with changes in cycle length. We compared within-individual changes in cycle length among groups: 1) a COVID-19 group, 2) a COVID-19–vaccinated group, and 3) a control group of unvaccinated participants reporting no history of COVID-19.

METHODS

We conducted a retrospective cohort analysis of menstrual cycle data collected prospectively by users of the period tracker application (app) Clue, linked to survey data on COVID-19 vaccination and disease status. The study received clearance from the French Research Institute for Development Ethic Board (IRD CCERP), Montpellier University Ethic Board, and the Oregon Health & Science University IRBs.

Only users aged 16–58 years with a registered period tracker app account and who gave consent for their pseudonymized data to be used for research purposes were sent an in-app message to take part in the survey, “Period and the Pandemic.” The survey distributed to period tracker app users in the United States, the United Kingdom, Canada, and Australia between November 29, 2021, and February 8, 2022, and then to all period tracker app users with their app set to English-language between May 19, 2022, and August 10, 2022. We stopped data collection when no

new entries were recorded. The link to the survey was seen by 3,310,221 users, and 443,134 clicked on the link, leading to a clickthrough rate of 13.4%, consistent with typical response rates for in-app surveys.

After giving consent to link their survey and their prospectively collected menstrual cycle data (starting in 2019), users completed the survey questions regarding their COVID-19 vaccination status and dates of vaccination and COVID-19 history (month and year of onset of first symptoms or date of a positive test result), as well as age, body mass index (BMI, calculated as weight in kilograms divided by height in meters squared), and country of location. To evaluate the independent associations of disease and vaccination with cycle length, we created three groups: 1) a control group, including participants with no history of either COVID-19 vaccination or having COVID-19; 2) a vaccinated group, including participants with a history of COVID-19 vaccination but no history of having COVID-19 and for whom the first vaccine occurred in the first 38 days of the cycle and the second vaccine occurred at least 38 days after the first vaccine (to avoid the potential effects of a second vaccine dose); and 3) a COVID-19 group, including participants with a history of having COVID-19 and reporting being either unvaccinated or vaccinated at least 76 days (ie, two cycles of 38 days) after the onset of COVID-19 symptoms.

We excluded individuals who reported hormonal contraceptive use at any time between 2019 and the time of the survey. Data on contraception were taken either from the survey when available or from data tracked by users within the app during the study period. We also excluded users older than age 45 years, those for whom no cycle data were available, those who did not report their COVID-19 and vaccination status, and those who gave inconsistent or no dates for COVID-19 symptoms or vaccination. We removed all cycles flagged by users as abnormal ($n=10,788$).

We included users with at least five consecutive cycles. For the control group, we selected the last three cycles of 2020 (cycles 1–3) as artificial preevent cycles and two post-first artificial event cycles, inclusive of the event cycle (cycle 4). The artificial event was the first cycle of 2021 that contained January 31, 2021, to reduce bias from asymptomatic cases and align with the timing of most vaccinations. We used alternative dates (October 31, 2019, January 31, 2020, or October 31, 2020) in sensitivity analyses. For the vaccinated group, we included three prevaccine cycles (cycles 1–3) and two post-first vaccine cycles, inclusive of the vaccination cycle (cycle 4). For the

COVID-19 group, we included three pre-COVID cycles (cycles 1–3) and two post-first COVID cycles, inclusive of the COVID cycle (cycle 4). When the month but not the day of COVID-19 symptom onset was recorded, we took the cycle that overlapped the most with the month during which COVID-19 symptoms occurred. For all three groups, we excluded all participants with average preevent cycles outside the 24–38 day range for cycle length.²⁵ To reduce the possibility that missing data increased the length of the event cycle, we excluded all participants who did not track at least one symptom of any type every 38 days in the 90 days after the start of cycle 4, so that an absence of bleeding in this period cannot be attributed to nonadherence with tracking (Fig. 1).

Our primary exposure was self-reported vaccination and disease status (ie, group). Our primary outcome was the within-user change in cycle length (days) from the three-cycle preevent average (cycle 1–3) to the event cycle (cycle 4). Our secondary outcome was the within-user change in cycle length (days) from the three-cycle preevent average (cycle 1–3) to the first postevent (cycle 5). Our third outcome was the proportion of users who experienced a clinically significant change in cycle length (more than 8 days) in cycles 4 and 5.²⁵

Assuming an SD of 4 days and a minimum sample size of 421 per group, we had more than 90% power to detect an unadjusted 1-day within-individual difference in cycle length given three groups, lowering the significance level from 0.05 to 0.0125 to adjust for multiple outcomes (Appendix 1, available online at <http://links.lww.com/AOG/D312>). All analyses were conducted using R 4.2.1.²⁶ We compared within-individual changes in cycle length between the three-cycle preevent average (cycles 1–3) and cycles 4 and 5 using a two-sided *t* test. We determined outliers using the Cook's distance method and excluded them for the analysis (Appendix 2, available online at <http://links.lww.com/AOG/D312>). We created histograms to compare the distributions of changes in cycle length and compared the proportion of individuals who experienced a clinically significant change in cycle length (8 days or more)²⁵ using Pearson's χ^2 tests. We used longitudinal multivariable mixed-effects models to determine the adjusted difference in the change in cycle length among groups and plotted predicted values. Models contained random intercepts and slopes at the individual level and interaction term between time (preevent cycle average, event cycle, and postevent cycle) and group (control, COVID-19, COVID-19 vaccination). All estimates were adjusted for age and country, and

P values were adjusted for multiple comparisons using a Bonferroni–Holm correction to control the family-wise error rate. A post hoc Tukey honestly significant difference (HSD) test was used to assess the significance of differences between pairs of group means.

Age and country data were available for the full selected sample, but information on BMI (categorized as underweight, normal weight, overweight, or obesity) was missing for most participants (94.6%; Appendix 3, available online at <http://links.lww.com/AOG/D312>). To avoid reducing statistical power, our complete cases analysis did not include BMI as a covariate. We then imputed data on BMI using a multiple imputation approach,²⁷ combining random forest imputation with predictive mean matching (Appendix 3, <http://links.lww.com/AOG/D312>), and conducted a sensitivity analysis.

We also explored whether vaccination before the onset of COVID-19 symptoms mediated any association between COVID-19 and cycle length. We created an additional group including participants reporting vaccination at least 3 months before the onset of COVID-19 symptoms and reran all steps of the analysis, including removing outliers and adjusting for age and country (Appendices 4–6, available online at <http://links.lww.com/AOG/D312>).

We conducted three sensitivity analyses to confirm the robustness of our findings: 1) we excluded individuals who reported COVID-19 symptoms after November 2021 to rule out any effect of the Omicron wave (Appendix 7, available online at <http://links.lww.com/AOG/D312>), 2) we performed a sensitivity analysis including imputed data on BMI (Appendix 8, available online at <http://links.lww.com/AOG/D312>), and 3) we changed the timing of the artificial COVID cycle in the control group (Appendix 9, available online at <http://links.lww.com/AOG/D312>).

RESULTS

Of 39,884 eligible individuals, 6,514 met inclusion criteria (Fig. 1). The final study sample included 6,514 participants representing 32,570 cycles (five cycles per individual), including 421 individuals in the control group, 1,450 in the COVID-19 group, and 4,643 in the COVID-19 vaccinated group (Table 1). Initial COVID-19 symptoms occurred between January 1, 2020, and June 28, 2022, and initial COVID-19 vaccinations occurring between December 11, 2020, and July 26, 2022 (Appendix 10, available online at <http://links.lww.com/AOG/D312>). Included vaccine types were Pfizer-BioNTech, Oxford-AstraZeneca, CoronaVac/Sinovac, Covishield, Johnson & Johnson/




Fig. 1. STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) flow diagram. COVID-19, coronavirus disease 2019.

Alvergne. Menstrual Cycle Length and COVID-19. Obstet Gynecol 2023.

Janssen, Moderna, Sinopharm, and Sputnik V. Participants were from 110 countries (Appendix 11, available online at <http://links.lww.com/AOG/D312>), with most participants coming from the United States (3,237, 49.7%), the United Kingdom (922, 14.1%), Germany (324, 5.0%), Canada (235, 3.6%), France (206, 3.2%), and Australia (206, 3.2%)

Individuals in the control group experienced a 0.68-day decrease in cycle length between cycles 1–3 and cycle 4 (95% CI -1.18 to -0.19 , $P=.007$). Indi-

viduals in the vaccinated group experienced a 1.14-day unadjusted increase in cycle length during the first vaccination cycle compared with the prevaccination average (Table 2, 95% CI 0.63–1.66, $P<.001$). Individuals in the COVID-19 cohort experienced a 1.45-day unadjusted increase in cycle length during the first COVID cycle compared with the average of their three pre-COVID cycles (Table 2, 95% CI 0.89–2.02, $P<.001$). A Tukey HSD test revealed no significant differences between the vaccinated and COVID-

Table 1. Sample Characteristics

Variable	Group				P*
	Overall (N=6,514)	COVID-19 (n=1,450)	Control (n=421)	Vaccinated (n=4,643)	
Age (y)					<.001
17–20	1,420 (22)	374 (26)	121 (29)	925 (20)	
21–25	1,252 (19)	289 (20)	83 (20)	880 (19)	
26–30	1,214 (19)	266 (18)	76 (18)	872 (19)	
31–35	1,146 (18)	225 (16)	61 (14)	860 (19)	
36–40	879 (13)	175 (12)	50 (12)	654 (14)	
41–45	603 (9.3)	121 (8.3)	30 (7.1)	452 (9.7)	
Country					<.001
Other	2,355 (36)	561 (39)	94 (22)	1,700 (37)	
United Kingdom	922 (14)	242 (17)	53 (13)	627 (14)	
United States	3,237 (50)	647 (45)	274 (65)	2,316 (50)	
Cycle length preevent average	28.7 (27.0–31.3)	28.7 (27.0–31.3)	29.0 (27.0–31.3)	28.7 (27.0–31.0)	.53

Data are n (%) or median (interquartile range) unless otherwise specified.

* Kruskal-Wallis rank sum test or Pearson’s χ^2 test.

19 groups (diff=0.31, 95% CI –0.05 to 0.67, $P=.11$), although they were both different from the control group. The overlaid histogram shows a cycle length change distribution in individuals in the COVID-19 and vaccinated groups that is roughly equivalent to that in the control group, although slightly right-skewed relative to the control distribution (Fig. 2A). The proportion of individuals who experienced a clinically significant change in cycle length of more than 8 days between the event cycle and the preevent average was 6.9% in the control group, 9.7% in the COVID-19 group, and 6.3% in the vaccinated group. After applying a Bonferroni correction, this proportion remained higher for the COVID-19 group ($P<.001$; Appendix 12, available online at <http://links.lww.com/AOG/D312>). As compared with the control group, after adjusting for age and country, the change in cycle length remains at 1.45 days for the COVID-19 cohort (95% CI 0.86–2.04, $P<.001$) and 1.14 days for the vaccinated cohort (95% CI 0.60–1.69, $P<.001$, Table 2, Fig. 3A).

Individuals in the COVID-19 cohort did not experience an increase in unadjusted cycle length cycle during the first post-COVID cycle compared with the three pre-COVID cycles (Table 3, 95% CI –0.12 to 0.96, $P=.3$), nor did individuals in the control group (95% CI –0.83 to 0.12, $P=.3$) or the vaccination group (95% CI –0.08 to 0.92, $P=.3$). The overlaid histogram shows a cycle length change distribution in individuals in the COVID-19 and vaccinated groups that is roughly equivalent to that in the control group (Fig. 2B). The proportion of individuals who experienced a clinically significant change in cycle length of more than 8 days was 4.7% in the

control group, 8.1% in the COVID-19 group, and 6.9% in the vaccinated group. After applying a Bonferroni correction, these proportions were not statistically different across groups ($P=.33$; Appendix 13, available online at <http://links.lww.com/AOG/D312>). Adjusted changes postcycle were not different from unadjusted changes (Table 3, Fig. 3B).

We conducted a subanalysis in a cohort of individuals vaccinated at least 3 months before the onset of COVID-19 ($n=2,335$). Individuals in the cohort experienced a 1.02-day unadjusted increase in cycle length from the three prevaccination cycles compared with the first post-COVID cycle (95% CI 0.50–1.54, $P<.001$) (Appendix 5, <http://links.lww.com/AOG/D312>), which is lower than the change experienced by those in the unvaccinated cohort who had COVID-19 (1.44, 95% CI 0.90–1.99, $P<.001$) (Appendix 5, <http://links.lww.com/AOG/D312>).

A Tukey HSD test revealed a significant difference between the group of individuals who had COVID-19 after vaccination and the COVID-only group (diff=–0.42, 95% CI –0.85 to 0.0031, adjusted $P=.05$) but no differences between the vaccinated-only group and the COVID-19-after-vaccination group (diff=–0.17, 95% CI –0.15 to 0.49, adjusted $P=.52$). The overlaid histogram shows a cycle length change distribution in individuals in the COVID-19 and vaccinated groups that is roughly equivalent to that in the control group (Appendix 4, <http://links.lww.com/AOG/D312>). As compared with the control group, after adjusting for age and country, the change in cycle length is 1.02 days (95% CI 0.47–1.57, $P<.001$) (Appendix 5, <http://links.lww.com/AOG/D312>) for the cohort that was vaccinated before experiencing

Table 2. Unadjusted and Adjusted Estimates for Changes in Cycle Length Between the Three Preevent Cycle Average (Cycles 1–3) and the Event Cycle (Cycle 4) (n=6,396)*

Group	Within-Individual Unadjusted Change in Cycle Length From 3 Preevent Cycle Average to Event Cycle			Adjusted Difference in Change (Event Cycle) vs Unvaccinated Individuals		
	Estimate	95% CI	P [†]	Estimate	95% CI	P [†]
Control (n=401)	-0.68	-1.18 to -0.19	.007	—	—	—
Vaccinated (n=4,597)	1.14	0.63–1.66	<.001	1.14	0.60–1.69	<.001
COVID-19 (n=1,398)	1.45	0.89–2.02	<.001	1.45	0.86–2.04	<.001

COVID-19, coronavirus disease 2019.

* The event cycle is the first vaccination cycle for the vaccinated cohort, the cycle during which COVID-19 symptoms first appeared for the COVID-19 cohort, or the cycle including January 31, 2021, for the control cohort. All cycles are consecutive.

† Adjusted P-values based on the Holm-Bonferroni method.

COVID-19. There is no significant increase in cycle length between the average of the three cycles before vaccination and the first cycle post-COVID (95% CI -0.52 to 0.50, $P=.98$) (Appendix 6, <http://links.lww.com/AOG/D312>).

After removing cases that occurred during the Omicron variant dominance (n=293), the estimates remained the same (Appendix 7, <http://links.lww.com/AOG/D312>). When imputed data on BMI were included in the adjusted model, estimates remained unchanged (Appendix 8, <http://links.lww.com/AOG/D312>). Using alternative timings for the arti-

cial COVID cycle in the control group lead to significant and similar (0.82, 1.08 and 1.09-day increase) changes for the COVID-19 group but no changes for the control group and similar changes for the vaccinated group in some but not all cases (Appendix 9, <http://links.lww.com/AOG/D312>).

DISCUSSION

We evaluated 32,570 cycles from 6,514 individuals to evaluate whether COVID-19 is associated with changes in menstrual cycle length and how any such changes might compare with COVID-19 vaccination




Fig. 2. Distribution of changes in cycle length between the three preevent cycles average and the event cycle (A) and the postevent cycle (B). This includes data from people who tracked any symptom at least every 38 days for 90 days from the start of the event cycle and excludes outliers (n=6,396). The proportion of individuals with a cycle change more than 15 days is 1.6% in the event cycle (A) and 2.1% in the postevent cycle (B). COVID-19, coronavirus disease 2019.

Alvergne. Menstrual Cycle Length and COVID-19. *Obstet Gynecol* 2023.




Fig. 3. Adjusted within-individual change in cycle length from the average length of three cycles preevent and the event cycle (A) or the postevent cycle (B) across groups and age groups. *Alvergne. Menstrual Cycle Length and COVID-19. Obstet Gynecol 2023.*

or a control group. Among unvaccinated participants reporting having COVID-19, we found a 1.45-day increase in menstrual cycle length as compared with their three preevent cycle-length average. This change quickly resolved in the post-COVID cycle. We found a similar small increase in menstrual cycle length for the COVID-19–vaccinated cohort as compared with their three preevent cycle-length average, which also quickly resolved in the postvaccination cycle. Although the differences for the COVID-19 and vaccinated groups were each statistically different from the control group, they were not significantly different from one another. Changes in cycle length associated with COVID-19 and vaccination during the event cycle were small in magnitude and not clinically significant at the population level, but the proportion of individuals who experienced a clinically significant change of more than 8 days was higher for individuals with COVID-19 than for those in the vaccination or control groups. We also found that COVID-19 vaccination at least 3 months before the onset of COVID-19 symptoms was protective for COVID-19–associated changes in cycle length.

Existing literature on the effects of COVID-19 on the menstrual cycle is scant and somewhat variable, but the overall signal appears to be consistently small in magnitude. The Nurses’ Health Study 3 showed no associations between COVID-19 and cycle length changes in a prospective study of 3,858 premeno-

pausal U.S. health care professionals.²³ This study’s use of self-reported data collected 5–10 years apart¹ likely hinders the resolution needed to detect small changes. Previous studies among individuals who tested positive for SARS-CoV-2 infection found larger effects, with 15–25% of individuals reporting changes to their menstrual cycles after SARS-CoV-2 infection.^{16,28} These studies likely overestimated the effects due to a lack of control group and a higher proportion of severely ill patients in the study populations.¹ Yet, a biological effect is suggested by a small study of 73 mostly Black and Hispanic women showing a significant association between immunoglobulin G antibodies and the perception of menstrual irregularities.²² Our results are strengthened by the use of prospectively tracked menstrual cycles and a control group, the exclusion of contraceptive users, and the power to detect a 1-day difference in cycle length.

Several limitations exist in our study. First, we relied on app users’ self-report of COVID-19 symptoms or a positive test result, as well as dates of infection or illness or vaccination. However, an individual’s self-report of COVID-19 symptoms earlier in the pandemic, when we performed data collection, was shown to be highly correlated with COVID-19, and most individuals have easy access to dates of their COVID-19 vaccinations.^{29,30} Our control cohort likely included asymptomatic individuals with

Table 3. Unadjusted and Adjusted Estimates for Changes in Cycle Length Between the Three Preevent (COVID-19 or Vaccination) Cycle Average (Cycles 1–3) and the Postevent Cycle (Cycle 5) (n=6,396)*

Group	Within-Individual Unadjusted Change in Cycle Length From Three Preevent Cycle Average to Postevent Cycle			Adjusted Difference in Change (Postevent Cycle) vs Unvaccinated Individuals		
	Estimate	95% CI	P [†]	Estimate	95% CI	P [†]
Control (n=401)	−0.36	−0.83 to 0.12	.305	—	—	—
Vaccinated (n=4,597)	0.42	−0.08 to 0.92	.305	0.42	−0.15 to 0.99	1
COVID-19 (n=1,398)	0.42	−0.12 to 0.96	.305	0.42	−0.20 to 1.04	1

COVID-19, coronavirus disease 2019.

* The postevent cycle is the cycle after the first vaccination cycle for the vaccinated cohort, after the cycle during which COVID-19 symptoms first appeared for the COVID-19 cohort, or after the cycle including January 31, 2021, for the control cohort. All cycles are consecutive.

† Adjusted P-values based on the Holm-Bonferroni method.

COVID-19, who may be similar to controls in that they likely do not experience menstrual cycle disturbances. A recent study supports this assumption, showing a positive association between the number of COVID-19 symptoms and menstrual cycle changes.²⁸ Second, SARS-CoV-2 variants may act differently on menstrual cycles, and we have no biological data to assess this possibility, although removing the more virulent Omicron wave from our data did not change the results. Third, our menstrual data rely on tracking behavior, and any heterogeneity in tracking behavior makes it difficult to distinguish between a real biological effect and a change in tracking behavior due to being ill. We addressed this issue by including only *regular trackers* (ie, participants who tracked any symptom in the app in the 90 days after the start of the cycle during which symptoms or the vaccination occurred). Although this prevents us from including individuals who had simply forgotten to track their cycles, it excludes people who stopped tracking altogether because of being too symptomatic from COVID-19.

We also do not have additional information about this cohort's education, income, or ethnicity, but prior research using a cohort of period tracker app users from the United States shows that the demographics for respondents to surveys are reflective of the broader U.S. population.³¹ We cannot account for pandemic stress, but adjusting for pandemic stress has previously been found not to alter associations between cycle length and COVID-19 vaccination.²³ We are unable to comment on how COVID-19 or vaccination affects hormonal contraceptive users or individuals with gynecologic diseases. Previous studies on vaccination and menstrual cycle length have shown that associations may be small to no increase in individuals with gynecologic disorders.^{7–9,15}

In summary, although the proportion of individuals who experienced a clinically significant change in cycle length of more than 8 days was higher for those with COVID-19, at the population level, experiencing COVID-19 was associated with a small and temporary change in cycle length similar to that with COVID-19 vaccination.

REFERENCES

1. Male V. COVID-19 vaccination and menstruation. *Science* 2022;378:704–6. doi: 10.1126/science.ade1051
2. Male V. Menstrual changes after covid-19 vaccination. *BMJ* 2021;374:n2211. doi: 10.1136/bmj.n2211
3. Sharp GC, Fraser A, Sawyer G, Kountourides G, Easey KE, Ford G, et al. The COVID-19 pandemic and the menstrual cycle: research gaps and opportunities. *Int J Epidemiol* 2022; 51:691–700. doi: 10.1093/ije/dyab239
4. Critchley HOD, Maybin JA, Armstrong GM, Williams ARW. Physiology of the endometrium and regulation of menstruation. *Physiol Rev* 2020;100:1149–79. doi: 10.1152/physrev.00031.2019
5. Critchley HOD, Babayev E, Bulun SE, Clark S, Garcia-Grau I, Gregersen PK, et al. Menstruation: science and society. *Am J Obstet Gynecol* 2020;223:624–64. doi: 10.1016/j.ajog.2020.06.004
6. Hillard PJA. Menstruation in adolescents: what do we know? And what do we do with the information? *J Pediatr Adolesc Gynecol* 2014;27:309–19. doi: 10.1016/j.jpag.2013.12.001
7. Alvergne A, Woon EV, Male V. Effect of COVID-19 vaccination on the timing and flow of menstrual periods in two cohorts. *Front Reprod Health* 2022;4:952976. doi: 10.3389/frph.2022.952976
8. Edelman A, Boniface ER, Male V, Cameron ST, Benhar E, Han L, et al. Association between menstrual cycle length and covid-19 vaccination: global, retrospective cohort study of prospectively collected data. *BMJ Med* 2022;1:e000297. doi: 10.1136/bmjmed-2022-000297
9. Edelman A, Boniface ER, Benhar E, Han L, Matteson KA, Favaro C, et al. Association between menstrual cycle length and coronavirus disease 2019 (COVID-19) vaccination. *Obstet Gynecol* 2022;139:481–9. doi: 10.1097/AOG.0000000000004695

10. Darney BG, Boniface ER, Van Lamsweerde A, Han L, Matteson KA, Cameron S, et al. Impact of coronavirus disease 2019 (COVID-19) vaccination on menstrual bleeding quantity: an observational cohort study. *BJOG: Int J Obstet Gynaecol* 2023;130:803–12. doi: 10.1111/1471-0528.17471
11. Gibson EA, Li H, Fruh V, Gabra M, Asokan G, Jukic AMZ, et al. Covid-19 vaccination and menstrual cycle length in the Apple Women's Health Study. *NPJ Digit Med* 2022;5:165. doi: 10.1038/s41746-022-00711-9
12. Costeira R, Lee KA, Murray B, Christiansen C, Castillo-Fernandez J, Ni Lochlainn M, et al. Estrogen and COVID-19 symptoms: associations in women from the COVID symptom study. *PLoS One* 2021;16:e0257051. doi: 10.1371/journal.pone.0257051
13. Lee KMN, Junkins EJ, Luo C, Fatima UA, Cox ML, Clancy KBH. Investigating trends in those who experience menstrual bleeding changes after SARS-CoV-2 vaccination. *Sci Adv* 2022;8:eabm7201. doi: 10.1126/sciadv.abm7201
14. Muhaidat N, Alshrouf MA, Azzam MI, Karam AM, Al-Nazer M, Al-Ani A. Menstrual symptoms after COVID-19 vaccine: a cross-sectional investigation in the MENA region. *Int J Women's Health* 2022;14:395–404. doi: 10.2147/IJWH.S352167
15. Alvergne A, Kountourides G, Argentieri MA, Agyen L, Rogers N, Knight D, et al. A retrospective case-control study on menstrual cycle changes following COVID-19 vaccination and disease. *iScience* 2023;26:106401. doi: 10.1016/j.isci.2023.106401
16. Al-Najjar MAA, Al-Alwany RR, Al-Rshoud FM, Abu-Farha RK, Zawiah M. Menstrual changes following COVID-19 infection: a cross-sectional study from Jordan and Iraq. *PLoS One* 2022;17:e0270537. doi: 10.1371/journal.pone.0270537
17. Carp-Veliscu A, Mehedintu C, Frincu F, Bratila E, Rasu S, Iordache I, et al. The effects of SARS-CoV-2 infection on female fertility: a review of the literature. *Int J Environ Res Public Health* 2022;19:984. doi: 10.3390/ijerph19020984
18. Lebar V, Laganà AS, Chiantera V, Kunič T, Lukanović D. The effect of COVID-19 on the menstrual cycle: a systematic review. *J Clin Med* 2022;11:3800. doi: 10.3390/jcm11133800
19. Klein SL, Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol* 2016;16:626–38. doi: 10.1038/nri.2016.90
20. Bhatia A, Sekhon HK, Kaur G. Sex hormones and immune dimorphism. *Scientific World J* 2014;2014:1–8. doi: 10.1155/2014/159150
21. Li K, Chen G, Hou H, Liao Q, Chen J, Bai H, et al. Analysis of sex hormones and menstruation in COVID-19 women of child-bearing age. *Reprod BioMedicine Online* 2021;42:260–7. doi: 10.1016/j.rbmo.2020.09.020
22. Cherenack EM, Salazar AS, Nogueira NF, Raccamarich P, Rodriguez VJ, Mantero AM, et al. Infection with SARS-CoV-2 is associated with menstrual irregularities among women of reproductive age. *PLoS One* 2022;17:e0276131. doi: 10.1371/journal.pone.0276131
23. Wang S, Mortazavi J, Hart JE, Hankins JA, Katuska LM, Farland LV, et al. A prospective study of the association between SARS-CoV-2 infection and COVID-19 vaccination with changes in usual menstrual cycle characteristics. *Am J Obstet Gynecol* 2022;227:739.e11–739.e11. doi: 10.1016/j.ajog.2022.07.003
24. Li K, Urteaga I, Wiggins CH, Druet A, Shea A, Vitzthum VJ, et al. Characterizing physiological and symptomatic variation in menstrual cycles using self-tracked mobile-health data. *NPJ Digit Med* 2020;3:79. doi: 10.1038/s41746-020-0269-8
25. Munro MG, Critchley HOD, Fraser IS. The two FIGO systems for normal and abnormal uterine bleeding symptoms and classification of causes of abnormal uterine bleeding in the reproductive years: 2018 revisions. *Int J Gynecol Obstet* 2018;143:393–408. doi: 10.1002/ijgo.12666
26. R Core Team: A language and environment for statistical computing. 2022. <https://www.r-project.org/>
27. Mayer M. MissRanger: fast imputation of missing value. <https://cran.r-project.org/package=missRanger>
28. Khan SM, Shilen A, Heslin KM, Ishimwe P, Allen AM, Jacobs ET, et al. SARS-CoV-2 infection and subsequent changes in the menstrual cycle among participants in the Arizona CoVHORT study. *Am J Obstet Gynecol* 2022;226:270–3. doi: 10.1016/j.ajog.2021.09.016
29. Adorni F, Prinelli F, Bianchi F, Giacomelli A, Pagani G, Bernacchia D, et al. Self-reported symptoms of SARS-CoV-2 infection in a nonhospitalized population in Italy: cross-sectional study of the EPICoVID19 web-based survey. *JMIR Public Health Surveill* 2020;6:e21866. doi: 10.2196/21866
30. Gibbons C, Hussain M, O'Keeffe DT, Simpkin AJ. An analysis of patient self-reported COVID-19 symptoms during the first wave of the pandemic in Ireland. *Irish J Med Sci.* 2022;191:543–6. doi: 10.1007/s11845-021-02598-z
31. Shea AA, Wever F, Ventola C, Thornburg J, Vitzthum VJ. More than blood: app-tracking reveals variability in heavy menstrual bleeding construct. *BMC Womens Health* 2023;23:170. doi: 10.1186/s12905-023-02312-4

PEER REVIEW HISTORY

Received March 31, 2023. Received in revised form June 8, 2023. Accepted June 15, 2023. Peer reviews and author correspondence are available at <http://links.lww.com/AOG/D313>.