



HAL
open science

Genetic variation for growth at one and two summers of age in the common carp (*Cyprinus carpio* L.): Heritability estimates and response to selection

Marc Vandeputte, Martin Kocour, Stéphane Mauger, Marek Rodina, Amandine Launay, David Gela, Mathilde Dupont-Nivet, Martin Hulak, Otomar Linhart

► To cite this version:

Marc Vandeputte, Martin Kocour, Stéphane Mauger, Marek Rodina, Amandine Launay, et al.. Genetic variation for growth at one and two summers of age in the common carp (*Cyprinus carpio* L.): Heritability estimates and response to selection. *Aquaculture*, 2008, 277 (1-2), pp.7-13. 10.1016/j.aquaculture.2008.02.009 . hal-04302485

HAL Id: hal-04302485

<https://hal.science/hal-04302485>

Submitted on 23 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

1 **Genetic variation for growth at one and two summers of age in**
2 **the common carp (*Cyprinus carpio* L.): Heritability estimates**
3 **and response to selection**

4
5 Marc Vandeputte^{a, b, *}, Martin Kocour^c, Stéphane Mauger^a, Marek Rodina^c, Amandine
6 Launay^a, David Gela^c, Mathilde Dupont-Nivet^a, Martin Hulak^c and Otomar Linhart^c

7
8 ^a INRA UR544 Unité de Génétique des Poissons, F-78350 Jouy-en-Josas, France

9 ^b IFREMER, chemin de Maguelone, F-34250 Palavas les Flots, France

10 ^c University of South Bohemia, Research Institute of Fish Culture and Hydrobiology, Zátiší 728/II, 389 25
11 Vodňany, Czech Republic

12

13 *: Corresponding author : Vandeputte M., email address : Marc.Vandeputte@jouy.inra.fr

14

15 **Abstract:**

16

17 We estimated the heritability of growth-related traits (weight and length at ages one summer, first spring and
18 two summers) in a synthetic mirror carp strain (HSM) in the Czech Republic. The four generation pedigree was
19 obtained from parentage assignment of three factorial mating designs with microsatellite markers, and included
20 195 fish without phenotypes (48 G0, 147 G1) and 1321 fish with phenotypes (674 G2, 647 G3). Animal model
21 heritability estimates over generations were in the range
22 0.21–0.33 for length and in the range 0.31–0.44 for weight. The genetic correlation between length and weight
23 was high (0.97). The correlations between growth measurements in the first and in the second summer of age
24 were moderate to low (0.34–0.67). Divergent selection for length at two
25 summers of age was performed on G2 fish, and response to selection was evaluated by comparing
26 the offspring of the selected sires in G3, in a communal test where genetic groups were identified by microsatellite
27 parentage assignment. The response to upwards selection was moderate and indicated a realized heritability value
28 of 0.24 to 0.34. No response to downwards selection was observed. We conclude that although selection for growth
29 seems to be feasible in common carp, it would be a long- term process before results are visible.

30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Keywords: Common carp; *Cyprinus carpio*; Heritability; Growth; Selection; Selection response; Realized heritability; Parentage assignment; Microsatellites; Performance; Quantitative genetics

1. Introduction

The common carp, *Cyprinus carpio*, is a major fish species in world aquaculture production (FAO, 2003) cultured exclusively under traditional extensive or semi-intensive pond management, where fish growth is highly dependent (50 % and more) on natural food (plankton and benthos). Hence, environmental conditions have a major impact on the performance of common carp, and much attention has been given to the optimization of pond management (see Horvath et al., 1992). Genetic improvement could be a complementary way to increase the effectiveness of carp culture.

Genetic improvement of quantitative traits in carp has proceeded through selection (Moav and Wohlfarth, 1976), genome manipulations (Cherfas et al., 1996; Kocour et al., 2003) and cross-breeding (e.g. Bakos and Gorda, 1995; Bialowas, 1991; Gross and Wohlfarth, 1994; Wohlfarth, 1993, see review by Hulata, 1995). Presently, the only widely applied method in common carp culture is intra-specific crossbreeding (Kocour et al., 2005). Concerning within-breed selection, a well documented experiment (but still controversial – see Kinghorn, 1983) found that mass selection for growth was ineffective (Moav and Wohlfarth, 1976). Estimated heritabilities for growth traits range from

0.0 to 0.5 (Vandeputte, 2003, for a review), but some of those past heritability estimates could have been biased due to the small number of breeders used, and/or the inability to separate common environment and genetic effects.

In a previous study with common carp, we used microsatellite markers for parentage assignment of 240 families produced from 10 dams and 24 sires (Vandeputte et al., 2004). Heritability estimates for weight, standard length and Fulton’s condition coefficient were found to range from 0.33 to 0.37. However, these estimates were obtained with 8-week-old juveniles. Since the results were encouraging and showed a potential for mass selection on growth in common carp, we undertook a first generation of selection to test the effectiveness of selective breeding for growth improvement. We also set up new crosses to estimate the heritability of growth traits in larger fish. As in our previous experiments, we used microsatellite parentage assignment to access family information and avoid environmental effects common to full-sib families and the divergent selected genetic groups.

61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93

2. Materials and methods

The reproduction and culture of common carp was carried out at the experimental facility of the University of South Bohemia, Research Institute of Fish Culture and Hydrobiology (Vodňany, Czech Republic). The traits of interest were recorded in generation 2 (G2) and generation 3 (G3) fish.

2.1. Production of G1

G1 offspring were produced in May 2001 using a full factorial mating between 24 G0 males and 10 G0 females of the HSM (Hungarian Synthetic Mirror) line (Vandeputte et al., 2004). Fin samples (1 cm²) were taken on all 34 G0 parents and kept in tubes filled with 98% ethanol. A first attempt to produce selected fish from the G1 was done by selecting the fish on length (up, down and control) in July 2001 and April 2002. However, floods that occurred in Vodňany in September 2002 resulted in mixing of the selected batches which were kept in different ponds. We decided to re-start the experiment from the males of this mixed population, as we could assign most of them to their G0 parents, thus allowing one more pedigreed generation compared with re-starting from the base HSM population.

2.2. Production of G2

In May 2003, 147 spermiating males were collected at random from the G1 population, and mated simultaneously with eight suitable G0 HSM females in a full factorial design to establish the G2. Fin samples were collected on all 155 parents (147 G1 males and 8 G0 females), and stored in 98 % ethanol. Only one G0 female was common to both 2001 and 2003 crosses, so the generations can be considered discrete. The details of the mating methodology are given in Vandeputte et al. (2004) and Kocour et al. (2007). The G2 population was reared as a single batch in one pond under semi-intensive pond management. In September 2003 (one summer), a random sample of 2000 yearlings of G2 population (mean weight 31g, standard deviation 10.3g) was individually PIT tagged and fin sampled. These G2 tagged fish were reared

94 in one batch in an earthen pond until September 2004 when the 1458 survivors were measured for body length
95 (from the tip of the nose to the end of the caudal peduncle) and body weight.

96
97

98 **2.3. Production of G3 by divergent selection**

99
100 Among the 1458 G2 fish, it happened that 42% had a deformed mouth, which has a negative impact on growth
101 (Kocour et al., 2006). Only the 848 fish without mouth deformity were then considered as potential candidates for
102 selection. Using September 2004 data, potential male broodfish were identified and pre-selected based on
103 their body length. Body length was chosen as it is highly genetically correlated with body weight (Vandeputte
104 et al., 2004), but easier to measure on many fish in field conditions. The proportion of fish in the Up group was
105 9.2%, 16.2% in the Control group (average length) and 12% in the Down group.

106 The numbers (and hence provisional selection intensities) per group were different, as we feared that less
107 spermiating males would be found among smaller fish in the next spring. In April 2005, 833 fish with normal
108 mouth had survived, and 99 spermiating males (30 U, 36C, 33D) were identified among
109 the three pre-selected groups. Only males were selected in order to shorten the whole experiment: in
110 the climatic conditions of the Czech Republic, males start to spermiate at the age of 2 or 3 years and they can be
111 used at that age for artificial reproduction, but females mature at the age of 3-4 years and can be used for
112 reproduction at the earliest at the age of 4 years, optimally at 5 years of age. This is why, for this experiment again,
113 we used females from the G0 generation.

114 On May 19th, 2005, the 99 spermiating males were stripped and 96 of them gave enough sperm (30U,
115 34C, 32D). They were mated in a full factorial design with 8 G0 HSM females (none of which had been used as
116 parent of the G1 and G2 fish), using our usual methodology (Kocour et al., 2007; Vandeputte et al., 2004). The
117 eggs from the three Up, Control and Down groups were mixed and reared as a
118 single batch in one Zuger jar, and then transferred to a resorption trough. At 8 days post-fertilization
119 (dpf), larvae were randomly split in two equal groups of 20,000 larvae and stocked in two ponds of
120 0.16 ha. Both ponds were harvested in September 2005, and 750 fish were randomly chosen from each pond,
121 measured, PIT-tagged and DNA-sampled. After that, the 1500 tagged fish were put in the same over-wintering
122 pond, and were then treated as a single batch until the end of the second growing season. All surviving fish
123 were individually measured in April 2006 and November 2006.

124
125

126 2.4. Parentage assignment

127

128 The following fish were genotyped:

129 the 34 G0 parents (10 females, 24 males);

130 the 147 G1 males and the 8 G0 females used as parents for the G2;

131 812 G2 fish randomly sampled in Sept. 2003;

132 the 96 G2 males and the 8 G0 females used for producing the G3;

133 797 randomly sampled G3 fish (of the total 2x750 fish randomly sampled and measured in Sept.

134 2005);

135 All these fish were genotyped for six to eleven microsatellites (Crooijmans et al., 1997): MFW7, MFW9,

136 MFW11, MFW16, MFW18 and MFW26 for all fish, MFW3, MFW12, MFW20, MFW29 and

137 MFW40 for some fish only. Parentage assignment was performed by exclusion with one or two

138 mismatches tolerated using VITASSIGN (Vandeputte et al., 2006). Only fish assigned to a single parental pair

139 were considered for the analysis.

140

141

142 2.5. Statistical analyses

143

144 The significance of fixed and random effects was evaluated using SAS-Glm in the G3 population. Heritability

145 values were computed for length and weight at each stage (1st summer, 1st spring and 2nd summer), using VCE

146 5.1.2 (Kovac and Groeneveld, 2003) with an animal model:

$$147 \underline{Y} = \underline{X}\underline{\beta} + \underline{Z}\underline{u} + \underline{e}$$

148 where \underline{Y} is the vector of observations, $\underline{\beta}$ is the vector of fixed effects, including year, pond (in G3) and mouth

149 deformity score (in G2), \underline{u} is the vector of random additive genetic effects, and \underline{e} is the vector of random residual

150 effects. \underline{X} and \underline{Z} are known incidence matrices.

151 The whole dataset (pedigree from G0 to G3, phenotypes in G2 and G3) was used to estimate heritabilities across

152 generations, while the whole pedigree but only the phenotypes in G2 or in G3 were used to estimate within-

153 generation heritabilities.

154 Animals from generations G1 and G2 whose parents could not be identified due to imperfect parentage

155 assignment were considered belonging to the base (G0) population. Univariate models were used to estimate

156 heritabilities, bivariate models were used to estimate genetic correlations between length and weight at each

157 stage, and a trivariate model was used to assess genetic correlation between weights (respectively lengths) at

158 different stages (first summer, spring and second summer). Estimated breeding values for all animals in the
159 pedigree were obtained from the solutions for the animal additive genetic values in VCE, using the whole dataset.

160 The divergence between the selected groups was tested with the following model in SAS-Glm:

$$161 \quad Y_{ij} = \mu + G_i + e_{ij}$$

162 Where Y_{ij} is the estimated breeding value of fish j from genetic group i , μ is the population mean, G_i is the fixed
163 effect of genetic group (G0, G1, G2, G3-U, G3-C, G3-D and e_{ij} is the random residual. Comparison of
164 means between genetic groups was done with the Tukey-Kramer test for multiple comparisons.

165 Realized response to upwards selection was calculated in two ways, first as twice the difference between the mean
166 breeding values of G3-U and G2, and then as twice the difference between the mean breeding values of G3-U and
167 G3-C. The differences were doubled as in the G3 groups, only the males were selected and then the observed
168 response is half the true selection response. The same type of calculation was done for downwards selection
169 response. Realized heritability was calculated as the ratio of these realized responses (standardized in phenotypic
170 standard deviation units) on the selection differentials.

171

172

173 **3. Results**

174

175

176 **3.1. Selection process**

177 The coefficients of variation of length and weight at the time of G2 selection (2nd summer) were moderately
178 high (24% for weight and 8.3% for length). Selection was performed on spermiating males in April 2005 (ca. 2
179 years of age), based on their length performance after the second summer (in September 2004). The phenotypic
180 means of the three selected groups are given in Table 1. As suspected, mouth deformities had an impact on
181 the size of the fish, and only fish with normal mouth were selected. Then, the selection differentials were also
182 calculated from fish with normal mouth only, and were equivalent to a mass selection intensity of 13% ($i=$
183 1.63) downwards and 11% ($i=1.71$) upwards.

184

185

186 **3.2. Parentage assignment**

187

188 Parentage assignment results were as follows:

189 126 of the 147 G1 males (85.7%) could be assigned to a single pair (7 of them with one or two mismatches allowed),
190 the rest being assigned to two or more pairs;
191 615 individuals in the random sample of 812 G2 fish (75.7%) could be assigned to a single parental pair (all
192 with perfect match), the rest being assigned to two or more pairs;
193 63 of the 96 G2 males used for producing the G3 offspring (65.6%) could be assigned to a single pair
194 (11 of them with one or two mismatches allowed), the rest being assigned to two or more pairs;
195 647 of the 797 G3 fish (81.2%) could be assigned to a single pair (7 of them with one or two mismatches allowed),
196 2 could not be assigned to any pair and the rest was assigned to two or more pairs.
197 These results allowed us to set up a pedigree including animals from all four generations (G0 to G3). The pedigree
198 included 195 fish without phenotypes (48 G0, 147 G1) and 1321 fish with phenotypes (674 G2, 647 G3).

199
200

201 3.3. Heritabilities and correlations

202
203 The heritabilities calculated using univariate animal models are given in Table 2. They ranged from
204 0.31 to 0.44 for body weight and 0.21 to 0.33 for body length when the whole dataset (phenotypes from G2
205 and G3) was used. At any given age, the genetic correlation between length and weight was
206 0.97 ± 0.01 . The genetic correlation between body weights measured at 1st summer and at spring was high
207 (0.96 ± 0.01), but it was much lower (0.34-0.41) between spring or 1st summer and 2nd summer (Table 3).
208 Genetic correlation between length at first summer and at spring was almost unity (0.996 ± 0.004), and the
209 genetic correlations between length at 2nd summer and at the two younger ages were higher than those seen for
210 weight (0.64 ± 0.07 for 1st summer, 0.67 ± 0.06 for spring). When we considered weight and length recorded at the
211 same age but in different years (generations) as different traits, we obtained different figures for heritability, higher
212 in G2 than in G3 (Table2).

213
214

215 3.4. Response to selection

216
217 Phenotypic means of the groups are given in Table 4, and breeding values in the different genetic groups and
218 generations are given in Table 5. There was no significant variation of the breeding values for any trait, as expected,
219 in the (unselected) G0, G1 and G2 generations. In the G3 generation a difference in breeding values could be seen
220 between the three groups ($U > C > D$, $P < 0.05$) at one summer of age. At spring and at second summer, the Control
221 and Down-selected groups were equivalent, and were outperformed by the Up-selected group ($P < 0.05$). When

222 breeding values were compared between G3 and G2, the Down and Control group had a genetic level equivalent
223 to G2, while the Up-selected group had a higher genetic level ($P < 0.05$). This was true for both length and weight,
224 and at all ages. When genetic gain was estimated as the difference between the breeding values of the Up, Down
225 and Control lines, realized heritability estimates were 0.24, for both upwards and downwards selection. When
226 upwards selection response was estimated as the difference in breeding values between the Up line and G2, the
227 corresponding realized heritability estimate was
228 0.34. When the same was done for downwards selection, realized heritability was 0.14 only.

229
230
231

232 **4. Discussion**

233
234

235 **4.1. Parentage assignment**

236

237 A total of 1451 out of the 1852 fish genotyped could be assigned to a single parent pair (78.3%). This figure is
238 relatively low compared to other parentage assignment studies in fish (usually in the 90-99% range, e.g. Fishback
239 et al., 2002; Norris and Cunningham, 2004; Vandeputte et al., 2004; Wesmajervi et al., 2006). However, we used
240 large mating schemes in G2 and G3 (147 males x 8 females and 96 males x 8 females, respectively). In such cases
241 with many potential parents, it is not unusual to have lower assignment rates (e.g. 73.5% in red sea bream with 250
242 potential parents, Perez-Enriquez et al.,
243 1999). It must be noted also that the number of non assigned fish is very low (2 fish in G3), and that the number
244 of fish for which mismatches are needed to achieve unique assignment is also low (25 fish in total), which is
245 indicative of a low genotyping error rate (Vandeputte et al., 2006). Therefore, the lack of assignment power is
246 essentially due to an insufficient resolution of the microsatellite loci set in crosses with many parents and
247 combinations, possibly combined with the fact that domesticated common carp strains commonly have a low allelic
248 richness, due to their long empirical breeding history (Kohlmann et al., 2005). Therefore, the pedigree used
249 should comprise only a limited proportion of mis-assigned fish, and therefore be suitable for quantitative genetics
250 studies. If present, a lack of accuracy in the molecular pedigree should anyway lead to an under-estimation of
251 heritabilities and larger standard errors (Milner et al., 2000). If microsatellites were to be used for family based
252 selection in common carp however, it would be necessary to improve the marker set (more loci, more variable), as

253 more than 20% unassigned fish would not be economically acceptable.

254
255

256 **4.2. Heritabilities and correlations**

257
258 The heritabilities for growth estimated in the present study with the whole dataset (0.21-0.44) are in the usual
259 range for common carp, (Nagy et al., 1980; Nenashev, 1969; Nenashev, 1966; Smisek,
260 1981; Tanck et al., 2001 – see review by Vandeputte, 2003). They are very similar to those we found for 8-weeks
261 weight in the same HSM population (0.33, Vandeputte et al., 2004), but much lower than the one we estimated for
262 weight at 3 summers of age (0.70, Kocour et al., 2007). However, it must be noted that heritability of weight
263 estimated solely in the G2 generation gives values comparable to this latter one (0.48-0.67). The fish used by
264 Kocour et al. (2007) were fish from the G2 generation, so this similarity could be expected. This global heritability
265 estimate of 0.21-0.44, confirmed over generations, places common carp in the usual range of heritability for growth
266 in commercial aquaculture species (reviewed by Gjedrem and Olesen, 2005). This value clearly leaves room for
267 selective breeding for growth in the common carp.

268 We could see that the genetic correlation between growth measurements (weight or length) in first and second
269 summer was not high (0.3-0.4 for weight, 0.6-0.7 for length). When we consider traits in G2 and in G3 as
270 different traits, the heritability estimates are never as low as in first summer in G3, where two environments (ponds)
271 were used for first summer growth. The low heritability could be explained if the offspring from the different sires
272 were ranked differently in both ponds. Unfortunately, the small family size (6.8 offspring/sire on average, or 3.4
273 in each pond) did not allow us to test for a sire*pond interaction with enough power. Nevertheless, genotype by
274 environment (GxE) interactions are well documented in common carp, at least at the population level, where it has
275 been shown that different populations may be re-ranked when tested in different environments (Gross and
276 Wohlfarth, 1994; Moav *et al.*, 1975; Wohlfarth et al., 1983). Therefore, the existence of such interactions at the
277 family level could be a possibility, although here, we did not try to generate GxE effects, as we used most of the
278 time only one pond with all genotypes mixed within.

279 Correlations between sizes at different ages were low to moderate. Then, pre-selection of young fish might not be
280 effective and might bring only moderate improvement to the weight at commercial size. A second argument for
281 this is the possibility of low survival (due to severe overwintering, predation, diseases etc.) of fish in ponds:
282 here in G3, at the end of the 2nd summer, only 479 of the 1500 fish tagged at 1st summer were still alive – only
283 32 % survival. It would seem more rational to wait for older fish to be selected when less inadvertent mortalities
284 occur. However, pre-selection of younger fish

285 might be still cost-effective considering the difficulties connected to later selection (feeding costs, higher
286 number of fish to be reared, manipulation and transport problems with bigger fish, number and size of ponds needed
287 etc.). Overall, in Central European conditions we recommend selection of the common carp at two years of age
288 (300-600 g).

289 The genetic correlation between length and weight at a given age was close to unity, as usual in fish species, so
290 selection for increased weight can be performed on length, which is much easier to measure in the field on a large
291 number of fishes. However, heritability of length (except in G3) seems to be slightly lower than heritability of
292 weight, and therefore it might be more advantageous, even if less practical, to select on weight rather than on
293 length.

294
295

296 **4.3. Response to selection**

297

298 The absolute response to selection observed was low. No significant response to downwards selection was
299 observed, when compared with the genetic level of the G2 generation. This is the exact opposite of the results
300 obtained in the first generation of the Israeli divergent selection experiment (Moav and Wohlfarth, 1976). One
301 problem about the 2nd summer G3 data is that they were obtained on a small fish sample (62 to 72 genotyped
302 fish per group), due to the mortality from 1st to 2nd summer. Therefore, the response estimates were quite
303 imprecise in the 2nd summer. The realized heritability values for upwards selection (0.24-0.34) are in the same
304 range as the animal model values (0.21-0.33 for length). Besides this, we have to keep in mind that the present
305 results for response to selection were obtained in communal rearing, with all three genetic groups (U, C, D)
306 mixed in the same pond.

307 However, it was shown that communal testing could magnify the differences between genetic groups in common
308 carp (Moav and Wohlfarth, 1974), and so the differences observed here (although moderate) could be over-
309 estimated.

310 The advantage of rearing the different groups in the same pond from the beginning and retrieving their origin with
311 microsatellites is that it allowed us to test for selection response in the first generation of selection, without common
312 environmental effects, and with some chances to see the small differences generated. If we were to test the
313 difference between genotypes in separate ponds, being able to detect differences in weight around 10% of the
314 mean may require more than 20 replicated ponds, a number that may be lowered to 4-5 replicates using an
315 internal control line (Gross and Wohlfarth,
316 1994; Vandeputte et al., 2002).

317 With a mean heritability value around 0.25, mass selection for growth in common carp should lead to a gain of
318 12% weight and 4% length by generation, when selecting the best 10% of the populations. The practical generation
319 interval of common carp in the Czech Republic is 4.5 years (4 years for males, 5
320 years for females): even if males spermiate from 2-3 years of age, they are routinely used for
321 reproduction from the age of 4 years (rarely from 3 years), and females are usually used from the age of 5 years
322 (rarely from 4 years). Then, the annual genetic gain in weight could be around 2.7-3 %. Due to the moderate
323 heritability value, family selection could also be considered, but it cannot be done easily in separate ponds due to
324 pond effects, and doing it in mixed families with genotyping would be an excessive cost. Therefore, mass selection
325 could be a reasonable option with complementary selection on quality traits from measurements on live animals,
326 such as fat-meter value to control the level of muscular fat, and selection for reduced head length to increase fillet
327 yield (Kocour et al., 2007).

328 Still, this would be a long-term project of 10-15 years with sustained investment necessary before a significant
329 improvement (30-40%) could be seen by an individual farmer, because of the large year-to-year and pond-to-pond
330 variability. Still, the global impact averages on many farms would be immediately positive. The precise impact of
331 GxE interactions on the genetic gain would also have to be more precisely evaluated, using larger samples in
332 different rearing conditions.

333 Finally, considering the possibility to genetically improve growth in common carp, the following question
334 remains: the total weight of fish harvested from a pond depends on the combination of the natural pond
335 productivity, supplemental food given and survival rate of stocked fish (e.g. Szumiec,
336 1990). Is it then sure that carps with improved growth would increase the global pond productivity under
337 semi-intensive pond management? It will be certain only if there is a correlated response in feed efficiency.
338 However, this correlation seems to be vary a lot among species: non-existent in brown trout (Sanchez et al., 2001),
339 moderate in rainbow trout (Kause et al., 2006), and higher in Atlantic salmon (Thodesen et al, 1999). So, carp with
340 improved growth will need to be accompanied with adapted management procedures (e.g. lower density and/or
341 higher supplemental food) to express their potential. Another possibility would be to improve flesh yield, which
342 seems feasible on the basis of recent results (Kocour et al., 2007). In this case, the objective would not be to improve
343 pond productivity, but to improve the quantity of edible flesh from a given pond production.

344

345

346

347 **Acknowledgements**

348

349

350 The authors wish to thanks the staff of the hatchery at USB RIFCH in the Czech Republic for active participation
351 in the production of the experimental fish. This work was supported by the project of National Agency for
352 Agricultural Research in the Czech Republic, no. QF4117, by Research plan of USB RIFCH no. MSM6007665809
353 and funds from INRA Animal Genetics Department, as well as funding from France-Czech Republic bilateral
354 programs BARRANDE n° 03218RF, 07508SA and
355 2004-044-2.

356

357

358

359

360

361 **References**

362

363

364 Bakos, J., Gorda, S., 1995. Genetic improvement of common carp strains using intraspecific hybridization.
365 Aquaculture 129, 183-186.

366 Bialowas, H., 1991. Possibilities of application of the heterosis effect in commercial production of common carp
367 (Cyprinus carpio L.). 1. Production of fingerlings. Acta Hydrobiol. 33, 319-334.

368 Cherfas, N.B., Gomelsky, B., Ben-Dom, N., Joseph, D., Cohen, S., Israel, I., Kabessa, M., Zohar, G.,

369 Peretz, Y., Mires, D., Hulata, G., 1996. Assessment of all-female common carp progenies for fish culture. Isr. J.
370 Aquacult. -Bamidgeh 48, 149-157.

371 Crooijmans, R.P.M.A., Bierbooms, V.A.F., Komen, J., Van der Poel, J.J., Groenen, M.A.M., 1997. Microsatellite
372 markers in common carp (Cyprinus carpio L.). Anim. Genet. 28, 129-134.

373 FAO, 2003. Aquaculture production 2001. FAO Yearbook of Fisheries Statistics 92/2.

374 Fishback, A.G., Danzmann, R.G., Ferguson, M.M., Gibson, J.P., 2002. Estimates of genetic parameters and
375 genotype by environment interactions for growth traits of the rainbow trout (Oncorhynchus mykiss) as inferred
376 using molecular pedigrees. Aquaculture 206, 137-150.

377 Gjedrem, T., Olesen, I., 2005. Basic statistical parameters. In: Gjedrem, T. (Ed.), Selection and

378 Breeding Programs in Aquaculture. Springer, Dordrecht, The Netherlands, pp. 45-72.

379 Gross, R., Wohlfarth, G.W., 1994. Use of genetic markers in growth testing of common carp, Cyprinus carpio L.,

380 carried out over 2 or 3 year cycles. *Aquacult. Fish. Manage.* 25, 585-599.

381 Horvath, L., Tamas, G., Seagrave, C., 1992. *Carp and pond fish culture*. Blackwell Scientific
382 Publications Ltd., UK, 158 pp.

383 Hulata, G., 1995. A review of genetic improvement of the common carp (*Cyprinus carpio* L.) and other cyprinids
384 by crossbreeding, hybridization and selection. *Aquaculture* 129, 143-155.

385 Kause, A., Tobin, D., Houlihan, D.F., Martin, S.A.M., Mantysaari, E.A., Ritola, O., Ruohonen, K., 2006. Feed
386 efficiency of rainbow trout can be improved through selection: Different genetic potential on alternative diets. *J.*
387 *Anim. Sci.* 84, 807-817.

388 Kinghorn, B.P., 1983. A review of quantitative genetics in fish breeding. *Aquaculture* 31, 283-304.

389 Kocour, M., Gela, D., Rodina, M., Linhart, O., 2005. Testing of performance in common carp *Cyprinus carpio* L.
390 under pond husbandry conditions I: top-crossing with Northern mirror carp. *Aquacult. Res.*
391 36, 1207-1215.

392 Kocour, M., Linhart, O., Gela, D., 2003. Results of comparative growing test of all-female and bisexual
393 population in two-year-old common carp (*Cyprinus carpio* L.). *Aquacult. Int.* 11, 369-378.

394 Kocour, M., Linhart, O., Vandeputte, M., 2006. Mouth and fin deformities in common carp: is there a genetic
395 basis? *Aquacult. Res.* 37, 419-422.

396 Kocour, M., Mauger, S., Rodina, M., Gela, D., Linhart, O., Vandeputte, M., 2007. Heritability estimates
397 for processing and quality traits in common carp (*Cyprinus carpio* L.) using a molecular pedigree.
398 *Aquaculture* 270, 43-50.

399 Kohlmann, K., Kersten, P., Flajshans, M., 2005. Microsatellite-based genetic variability and differentiation of
400 domesticated, wild and feral common carp (*Cyprinus carpio* L.) populations. *Aquaculture* 247, 253-266.

401 Kovac, M., Groeneveld, E., 2003. VCE5 user's guide and manual version 5.1. Department of Animal
402 Sciences, University of Ljubljana, Ljubljana, 68 pp.

403 Lynch, M., Walsh, B., 1998. *Genetics and analysis of quantitative traits*. Sinauer Associates,
404 Sunderland, MA, 980 pp.

405 Milner, J.M., Pemberton, J.M., Brotherstone, S., Albon, S.D., 2000. Estimating variance components and
406 heritabilities in the wild: a case study using the 'animal model' approach. *J. Evol. Biol.* 13, 804-
407 813.

408 Moav, R., Hulata, G., Wohlfarth, G.W., 1975. Genetic differences between the Chinese and European races of
409 the common carp. I. Analysis of genotype-environment interactions for growth rate. *Heredity*
410 34, 323-340.

411 Moav, R., Wohlfarth, G.W., 1974. Magnification through competition of genetic differences in yield

412 capacity in carp. *Heredity* 33, 181-202.

413 Moav, R., Wohlfarth, G.W., 1976. Two way selection for growth rate in the common carp (Cyprinus carpio
414 L.). *Genetics* 82, 83-101.

415 Nagy, A., Csanyi, V., Bakos, J., Horvath, L., 1980. Development of a short-term laboratory system for
416 the evaluation of carp growth in ponds. *Bamidgeh* 32, 6-15.

417 Nenashev, G.A., 1969. Heritability of some selective characters in Ropsha carp (Russian). *Izvestija*
418 Gosud. Nauchno-issled. Inst. Ozern. Recn. Rybn. Kos. (GosNIORKh) 65, 185-195.

419 Nenashev, G.A., 1966. The determination of heritability of different characters in fishes. *Genetika* 11,
420 100-108.

421 Norris, A.T., Cunningham, E.P., 2004. Estimates of phenotypic and genetic parameters for flesh colour traits in
422 farmed Atlantic salmon based on multiple trait animal model. *Livest. Prod. Sci.* 89, 209-222. Perez-Enriquez, R.,
423 Takagi, M., Taniguchi, N., 1999. Genetic variability and pedigree tracing of a
424 hatchery-reared stock of red sea bream (Pagrus major) used for stock enhancement, based on
425 microsatellite DNA markers. *Aquaculture* 173, 413-423.

426 Sanchez, M.P., Chevassus, B., Labbe, L., Quillet, E., Mambrini, M., 2001. Selection for growth of brown
427 trout (Salmo trutta) affects feed intake but not feed efficiency. *Aquat. Liv. Resour.* 14, 41-48. Smisek, J., 1981.
428 The effect of gene pool on the performance and conformation of filial generations of carp fry from line crossing.
429 *Bull. VURH Vodnany* 17, 3-11.

430 Szumiec, M.A., 1990. Stochastic model of carp fingerling growth. *Aquaculture* 91, 87-99.

431 Tanck, M.W.T., Vermeulen, H., Bovenhuis, H., Komen, J., 2001. Heredity of stress-related cortisol response
432 in androgenetic common carp (Cyprinus carpio L.). *Aquaculture* 199, 283-294.

433 Thodesen, J., Grisdale-Helland, B., Stale, J., Gjerde, B., 1999. Feed intake, growth and feed utilization
434 of offspring from wild and selected Atlantic salmon (Salmo salar). *Aquaculture* 180, 237-246. Vandeputte, M.,
435 2003. Selective breeding of quantitative traits in the common carp (Cyprinus carpio L.): a review. *Aquat. Liv.*
436 *Resour.* 16, 399-407.

437 Vandeputte, M., Kocour, M., Mauger, S., Dupont-Nivet, M., De Guerry, D., Rodina, M., Gela, D., Vallod,
438 D., Chevassus, B., Linhart, O., 2004. Heritability estimates for growth-related traits using microsatellite parentage
439 assignment in juvenile common carp (Cyprinus carpio L.). *Aquaculture* 235,
440 223-236.

441 Vandeputte, M., Mauger, S., Dupont-Nivet, M., 2006. An evaluation of allowing for mismatches as a way to
442 manage genotyping errors in parentage assignment by exclusion. *Mol. Ecol. Notes* 6, 265-267. Vandeputte, M.,
443 Peignon, E., Vallod, D., Haffray, P., Komen, J., Chevassus, B., 2002. Comparison of

- 444 growth performances of three French strains of common carp (Cyprinus carpio) using hemi-isogenic
445 scaly carp as internal control. *Aquaculture* 205, 19-36.
- 446 Wesmajervi, M.S., Westgaard, J.I., Delghandi, M., 2006. Evaluation of a novel pentaplex microsatellite marker
447 system for paternity studies in Atlantic cod (Gadus morhua L.). *Aquacult. Res.* 37, 1195-1201.
- 448 Wohlfarth, G.W., 1993. Heterosis for growth rate in common carp. *Aquaculture* 113, 31-46.
- 449 Wohlfarth, G.W., Moav, R., Hulata, G., 1983. A genotype-environment interaction for growth rate in the
450 common carp, growing in intensively manured ponds. *Aquaculture* 33, 187-195.

Tables

Table1. Numbers and size of G2 carp at two summers, before and after divergent selection. S.D.= standard deviation

	Number of fish	Length at 2 nd summer (mm (S.D.))	Selection differential (Length phenotypic S.D. units)
Fish with deformed mouth	610	234.0 (20.7)	-
Fish with normal mouth	848	249.4 (19.3)	-
Spermiating Up fish	30	282.4 (7.5)	+1.71
Spermiating Control fish	36	250.6 (5.3)	+0.06
Spermiating Down fish	33	218.1 (9.4)	-1.63

Table 2. Heritabilities (\pm S.E.) for body weight and length at different ages in common carp, using univariate animal models in VCE5, four generations of pedigree and two generations of phenotypes (G2 and G3, whole dataset) or just one generation of phenotypes (G2 only or G3 only).

Age	Whole dataset		G2 only		G3 only	
	Weight	Length	Weight	Length	Weight	Length
1 st summer	0.31 \pm 0.04	0.21 \pm 0.03	0.61 \pm 0.03	0.39 \pm 0.04	0.18 \pm 0.06	0.25 \pm 0.07
Spring	0.33 \pm 0.04	0.26 \pm 0.03	0.67 \pm 0.03	0.46 \pm 0.04	0.20 \pm 0.07	0.27 \pm 0.08
2 nd summer	0.44 \pm 0.05	0.33 \pm 0.04	0.48 \pm 0.03	0.36 \pm 0.05	0.37 \pm 0.13	0.40 \pm 0.13

Table 3. Heritabilities \pm S.E. (bold), genetic correlations \pm S.E. (italics) from trivariate animal models, and phenotypic correlations (plain) for body weight and body length after 1st summer, at spring and after second summer in common carp, using phenotypes from G2 and G3 and the pedigree from G0 to G3.

		1 st summer	spring	2 nd summer
Weight	1 st summer	0.32\pm0.03	<i>0.96\pm0.01</i>	<i>0.41\pm0.08</i>
	spring	0.95	0.29\pm0.03	<i>0.34\pm0.08</i>
	2 nd summer	0.65	0.71	0.46\pm0.04
Length	1 st summer	0.20\pm0.03	<i>1.00\pm0.00</i>	<i>0.64\pm0.07</i>
	spring	0.97	0.22\pm0.03	<i>0.67\pm0.06</i>
	2 nd summer	0.68	0.71	0.28\pm0.04

Table 4. Phenotypic data (mean values with standard deviation in brackets) of genotyped offspring from selected groups of carp in G3.

	1 st summer			Spring			2 nd summer		
	<u>n</u>	Weight (g)	Length (mm)	<u>n</u>	Weight (g)	Length (mm)	<u>n</u>	Weight (g)	Length (mm)
Up selected	202	32.1 (11.4)	93.0 (12.7)	103	32.4 (9.7)	97.0 (9.9)	66	703 (173)	306 (26)
Control	225	30.7 (10.5)	92.0 (11.2)	112	30.8 (8.4)	95.6 (9.1)	72	659 (152)	300 (22)
Down selected	220	30.4 (9.7)	91.5 (11.8)	110	31.1 (9.4)	95.9 (9.7)	62	670 (161)	299 (23)

Table 5. Differences between genetic groups for Estimated Breeding Values from the animal model with the whole dataset, estimated for body weight (BW) and body length (BL) at 1 summer, spring and 2 summers of age. Within each line, means with the same superscript are not significantly different (Tukey-Kramer test, $P > 0.05$).

Variable	Genetic group					
	G0 (n=48)	G1 (n=147)	G2 (n=670)	G3-Up (n=66-201 [*])	G3-Control (n=72-225 [*])	G3-Down (n=62-219 [*])
1 sum BL (mm)	-0.5 ^{ab}	-0.4 ^{ab}	-0.8 ^{ab}	1.9 ^c	0.1 ^b	-1.0 ^a
1 sum BW (g)	-0.8 ^{ab}	-0.9 ^{ab}	-1.3 ^{ab}	2.2 ^c	-0.3 ^b	-1.5 ^a
Spring BL (mm)	-0.5 ^a	-0.5 ^a	-0.5 ^a	2.8 ^b	0.6 ^a	-1.3 ^a
Spring BW(g)	-1.0 ^a	-1.2 ^a	-1.1 ^a	4.1 ^b	0.2 ^a	-2.1 ^a
2 sum BL (mm)	-0.4 ^a	0.1 ^a	0.0 ^a	6.9 ^b	2.3 ^a	-2.6 ^a
2 sum BW (g)	-6.4 ^a	-5.6 ^a	-5.8 ^a	53.7 ^b	12.2 ^a	-17.2 ^a

The larger n is for the 1st summer, the lower one for the 2nd summer.