



HAL
open science

A Journey into the Total Synthesis of Polycyclic Cytochalasines

Oscar Gayraud, Bastien Nay

► **To cite this version:**

Oscar Gayraud, Bastien Nay. A Journey into the Total Synthesis of Polycyclic Cytochalasines. Journée de l'Ecole Doctorale Interfaces, Ecole Polytechnique, Apr 2018, Palaiseau, France. hal-04303667

HAL Id: hal-04303667

<https://hal.science/hal-04303667>

Submitted on 23 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Journey into the Total Synthesis of Polycyclics Cytochalasines



Oscar Gayraud, Bastien Nay*


Laboratoire de Synthèse Organique (UMR 7652 CNRS), Ecole Polytechnique, 91128 Palaiseau
oscar.gayraud@polytechnique.edu

1. Introduction

Trichoderme and Trichoderone A have been isolated from the endophytic fungus *Trichoderma gamsii*.^[1, 2] Unprecedented **7/6/6/5/5** and **7/5/6/5** polycyclics cores possibly arised from a common isolated intermediate (**1**).




Panax Notoginseng





2. Retrosynthesis

Starting from key intermediates **2** and **3**, Trichoderone A and Trichoderme might be achieved through biomimetic late stage cyclization and oxidation.




Cycloheptanone will be functionalized using oxidation, coupling and rearrangement reaction. Then an Intramolecular Diels-Alder (IMDA) reaction will allow access to the **7/6/6/5** core.




3. Synthesis of 9 and 11

Inspired by Lallemand's work,^[3] diene **9** has been synthesized through a Sonogashira coupling followed by a hydroboration of the alkyne.




Lactame **11** has been access from the protected amino acid N-Boc Leucine through a sequence involving formation of a carbamate followed by a reduction then a cyclization and a protection.^[4]




4. Synthesis of the key intermediate 18

Fully functionalized 7-membered ring **5** has been synthesized using Sharpless dihydroxylation and a Suzuki coupling.




Protected key intermediates **18** has been access through an Ireland-Claisen rearrangement and an IMDA reaction enabled by the in situ generation of **17**.



5. Late Stage Oxidation

First try of the biomimetic late stage oxidation afforded the wrong regioisomer **19**. Work are in progress to access **20**.



References

- [1] Ding, G.; Wang, H.; Li, L.; Song, B.; Chen, H.; Zhang, H.; Liu, X.; Zou, Z. *J. Nat. Prod.* **2014**, *77*, 164-167.
- [2] Ding, G.; Wang, H.; Li, L.; Chen, A. J.; Chen, L.; Chen, H.; Zhang, H.; Liu, X.; Zou, Z. *Eur. J. Org. Chem.* **2012**, *13*, 2516-2519.
- [3] Ohanessian, G.; Six, Y.; Lallemand, J.-Y. *Bull. Soc. Chim. Fr.* **1996**, *133*, 1143-1148.
- [4] Smrcina, M.; Majer, P.; Majerová, E.; Guerassina, T. A.; Eissenstat, M. A. *Tetrahedron* **1997**, *53*, 12867-12874.