



HAL
open science

RivFishTIME: A global database of fish time-series to study global change ecology in riverine systems

Lise Comte, Juan Carvajal-Quintero, Pablo A Tedesco, Xingli Giam, Ulrich Brose, Tibor Erős, Ana F Filipe, Marie-josée Fortin, Katie Irving, Claire Jacquet, et al.

► **To cite this version:**

Lise Comte, Juan Carvajal-Quintero, Pablo A Tedesco, Xingli Giam, Ulrich Brose, et al.. RivFishTIME: A global database of fish time-series to study global change ecology in riverine systems. *Global Ecology and Biogeography*, 2020, 30 (1), pp.38-50. <10.1111/geb.13210>. <hal-04303700>

HAL Id: hal-04303700

<https://hal.science/hal-04303700v1>

Submitted on 28 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 **DATA PAPER**

2

3 **RivFishTIME: A global database of fish time-series to study global change**
4 **ecology in riverine systems**

5

6

7 **Running title:** Time-series database of riverine fish communities

8

9 Lise Comte^{1,*}, Juan Carvajal-Quintero^{2,3}, Pablo A. Tedesco², Xingli Giam¹, Ulrich Brose^{4,5}, Tibor
10 Erős⁶, Ana Filipa Filipe^{7,8}, Marie-Josée Fortin⁹, Katie Irving^{10,11}, Claire Jacquet^{12,13}, Stefano
11 Larsen^{14,15}, Sapna Sharma¹⁶, Albert Ruhi¹⁷, Fernando G. Becker¹⁸, Lilian Casatti¹⁹, Giuseppe
12 Castaldelli²⁰, Renato B. Dala-Corte²¹, Stephen R. Davenport²², Nathan R. Franssen²³, Emili García-
13 Berthou²⁴, Anna Gavioli²⁰, Keith B. Gido²⁵, Luz Jimenez-Segura²⁶, Rafael P. Leitão²⁷, Bill
14 McLarney²⁸, Jason Meador²⁸, Marco Milardi²⁹, David B. Moffatt³⁰, Thiago V. T. Occhi³¹, Paulo S.
15 Pompeu³², David L. Propst³³, Mark Pyron³⁴, Gilberto N. Salvador³⁵, Jerome A. Stefferud³⁶, Tapio
16 Sutela³⁷, Christopher Taylor³⁸, Akira Terui³⁹, Hirokazu Urabe⁴⁰, Teppo Vehanen³⁷, Jean R. S.
17 Vitule⁴¹, Jaqueline O. Zeni^{19,42}, Julian D. Olden⁴³

18

19 ¹Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, Knoxville,
20 TN 37996, USA

21 ²Research Unit 5174, Laboratoire Evolution et Diversité Biologique, CNRS, Institut de Recherche
22 pour le Développement, Université Paul Sabatier, F-31062 Toulouse, France

23 ³Laboratorio de Macroecología Evolutiva, Red de Biología Evolutiva, Instituto de Ecología, A.C., El
24 Haya, 91070 Xalapa, Veracruz, Mexico

25 ⁴German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig,
26 Germany

27 ⁵Institute of Biodiversity, Friedrich-Schiller-University Jena, 07743 Jena, Germany

28 ⁶Centre for Ecological Research, Balaton Limnological Institute, Klebelsberg K. u. 3. 8237 Tihany,
29 Hungary

30 ⁷CIBIO/InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos da Universidade do
31 Porto, 4485-661 Vairão, Portugal

32 ⁸Instituto Superior de Agronomia, Universidade de Lisboa, 1349-017 Lisboa, Portugal

33 ⁹Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario, M5S
34 3B2, Canada

35 ¹⁰Department of Ecosystem Research, Leibniz Institute of Freshwater Ecology and Inland Fisheries,
36 12587 Berlin, Germany

37 ¹¹Department of Biology, Chemistry and Pharmacy, Freie Universität Berlin, 14195 Berlin, Germany

38 ¹²Department of Aquatic Ecology, Swiss Federal Institute of Aquatic Science and Technology,
39 Eawag, Dübendorf, Switzerland

40 ¹³Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zürich,
41 Switzerland

42 ¹⁴Department of Civil, Environmental and Mechanical Engineering, University of Trento, Trento,
43 Italy

44 ¹⁵Computational Biology Unit, Research and Innovation Centre, Fondazione Edmund Mach, San
45 Michele all'Adige, Italy

46 ¹⁶Department of Biology, York University, 4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada

47 ¹⁷Department of Environmental Science, Policy, and Management, University of California,
48 Berkeley, Berkeley, CA 94720, USA

49 ¹⁸Department of Ecology, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

50 ¹⁹Department of Zoology and Botany, UNESP – São Paulo State University, São José do Rio Preto,
51 SP, Brazil

52 ²⁰ Department of Life Sciences and Biotechnology, University of Ferrara, via Luigi Borsari 46,
53 44121 Ferrara, Italy

54 ²¹Department of Ecology, Universidade Federal de Goiás, Goiânia, GO, Brazil

55 ²²New Mexico Fish and Wildlife Conservation Office, 3800 Commons NE, Albuquerque, NM
56 87109, USA

57 ²³U.S. Fish and Wildlife Service, 2105 Osuna Rd NE, Albuquerque, NM 87113, USA

58 ²⁴GRECO, Institute of Aquatic Ecology, University of Girona, 17003 Girona, Catalonia, Spain

59 ²⁵Division of Biology, Kansas State University, Manhattan, KS 66506, USA

60 ²⁶Instituto de Biologia, Universidad de Antioquia, Calle 67 No.53-108 Medellin, Colombia

61 ²⁷Laboratório de Ecologia de Peixes, Department of Genetics, Ecology and Evolution, Federal
62 University of Minas Gerais (UFMG), 31270-901, Belo Horizonte, MG, Brazil

63 ²⁸Mainspring Conservation Trust, Franklin, NC 28734, USA

64 ²⁹Fisheries New Zealand - Tini a Tangaroa, Ministry for Primary Industries - Manatū Ahu Matua, 34-
65 38 Bowen Street, Pipitea 6011, Wellington, New Zealand

66 ³⁰Department of Environment and Science, Ecosciences Precinct, Dutton Park QLD 4102, Australia

67 ³¹Department of Ecology and Conservation, Federal University of Paraná, Jardim das Américas,
68 81530-900 Curitiba, Brazil

69 ³²Department of Ecology and Conservation, University of Lavras (UFLA), 37200-900 Lavras, MG,
70 Brazil

71 ³³Department of Biology, Museum of Southwestern Biology, University of New Mexico,
72 Albuquerque, NM 87131, USA

73 ³⁴Department of Biology, Ball State University, Muncie, IN 47306, USA

74 ³⁵Ecology and Conservation Laboratory (LABECO), Federal University of Pará (UFPA), 66075-110,
75 Belém, PA, Brazil

76 ³⁶USDA Forest Service (Retired), Phoenix, AR 85012, USA

77 ³⁷Natural Resources Institute Finland, Latokartanonkaari 9, 00790 Helsinki, Finland

78 ³⁸Department of Biology and School of Earth, Environmental, and Marine Sciences, University of
79 Texas Rio Grande Valley, Edinburg, TX 78539, USA

80 ³⁹Department of Biology, University of North Carolina at Greensboro, Greensboro, NC 27412, USA

81 ⁴⁰Salmon and Freshwater Fisheries Research Institute, Hokkaido Research Organization, Eniwa,
82 Hokkaido, Japan

83 ⁴¹Department of Environmental Engineering, Federal University of Paraná, Curitiba, Jardim das
84 Américas, 81530-900, Brazil

85 ⁴²Department of Ecology, UNESP – São Paulo State University, Rio Claro, SP, Brazil

86 ⁴³School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98105, USA

87

88 * **Corresponding author:** Lise Comte, Department of Ecology and Evolutionary Biology, The

89 University of Tennessee, Knoxville, TN, USA

90 lgcomte@gmail.com

91

92 **Running title:** Time-series database of riverine fish communities

93

94 **ABSTRACT**

95 **Motivation:** We compiled a global database of long-term riverine fish surveys from 46 regional
96 and national monitoring programs as well as individual academic research efforts upon which
97 numerous basic and applied questions in ecology and global change research can be explored.
98 Such spatially- and temporally-extensive datasets have been lacking for freshwater systems
99 compared to terrestrial ones.

100 **Main types of variables contained:** The database includes 11,386 time-series of riverine fish
101 community catch data, including 646,270 species-specific abundance records together with
102 metadata related to geographic location and sampling methodology of each time-series.

103 **Spatial location and grain:** The database contains 11,072 unique sampling locations (stream
104 reach), spanning 19 countries, 5 biogeographic realms, and 402 hydrographic basins worldwide.

105 **Time period and grain:** The database encompasses the period 1951–2019. Each time-series is
106 composed of a minimum of two yearly surveys (mean = 8 years) and represents a minimum time
107 span of 10 years (mean = 19 years).

108 **Major taxa and level of measurement:** The database includes 944 species of ray-finned fishes
109 (Class Actinopterygii).

110 **Software format:** .csv

111 **Main conclusion:** Our collective effort provides the most comprehensive long-term community
112 database of riverine fishes to date. This unique database should interest ecologists who seek to
113 understand the impacts of human activities on riverine fish biodiversity, and model and predict

114 how fish communities will respond to future environmental change. Together, we hope it will
115 promote advances in macroecological research in the freshwater realm.

116 **KEYWORDS:** species abundance; biodiversity; conservation; freshwater streams and rivers;
117 Actinopterygii; temporal trends; worldwide

118 1| INTRODUCTION

119 Increasing awareness of the ongoing biodiversity crisis has motivated global initiatives to
120 compile large-scale datasets of population and community abundance records that have been
121 consistently sampled through recent times (Pereira & Cooper, 2006). Included among these are
122 the Global Population Dynamics Database (Inchausti & Halley, 2001), the Living Planet Index
123 database (Loh *et al.*, 2005), and more recently, the BioTIME database (Dornelas *et al.*, 2018).
124 These databases have proven extremely useful and allowed major advancements in ecological
125 research (e.g. Kendall *et al.*, 1998; Sibly *et al.*, 2005; Butchart *et al.*, 2010; Dornelas *et al.*,
126 2014); however, they remain highly biased towards terrestrial and marine assemblages (e.g. only
127 0.50% of the records concern riverine fishes in BioTIME, the most recent of these initiatives).
128 This is unfortunate as effective strategic plans for conserving water resources that support human
129 well-being and ecosystem integrity rely on access to comprehensive, pertinent, quantitative
130 information regarding the status and trends of riverine biodiversity over regional to continental
131 scales (Tickner *et al.*, 2020).

132 Long-term studies of riverine species are limited because they require highly specialized
133 and time-consuming sampling methods. Furthermore, rivers in remote areas are often difficult to
134 access (Olden *et al.*, 2010; Radinger *et al.*, 2019). Nevertheless, over the past few decades, large-
135 scale policies have been enacted in response to the rapid degradation of freshwater resources,
136 such as the Water Framework Directive in the EU (Hering *et al.*, 2004) and the Clean Water Act
137 in the USA (Paulsen *et al.*, 2008), which require countries to monitor and evaluate the biological
138 integrity of surface waters through time to adopt quality standards that restore and maintain
139 ecological integrity (Kuehne *et al.*, 2017). Beyond these official national and regional monitoring
140 programs, the temporal dynamics of riverine systems and their fish communities have also been

141 assessed through various independent, though often local in extent, academic research programs
142 (e.g. Gido, 2017; Matthews & Matthews, 2017). All of these institutional and academic
143 monitoring efforts have produced considerable freshwater fish temporal data that remain largely
144 inaccessible to the broader scientific community due to the inherent difficulty in gathering and
145 harmonizing field data from disparate institutions and sampling protocols (Buss *et al.*, 2015).

146 To fill this important gap, we here present RivFishTIME, a compiled and curated
147 database of long-term (≥ 10 years) surveys of riverine fish communities at a fine spatial (stream
148 reach) and taxonomic (species) resolution, using data mining approaches to harmonize existing
149 but currently fragmented biomonitoring data sets. Riverine fish are extremely diverse in spite of
150 the small surface they inhabit on Earth: they represent about 40% of all known fish species while
151 occupying $<1\%$ of available aquatic habitat (“the freshwater fish paradox” sensu Lévêque *et al.*,
152 2008 and Tedesco *et al.*, 2017). However, they are also among the most threatened taxonomic
153 groups on Earth because of the convergence between the high concentration of biodiversity and
154 the many pressures resulting from human uses of freshwater resources and habitat change (Reid
155 *et al.*, 2019; Tickner *et al.*, 2020). The RivFishTIME database provides a unique opportunity to
156 understand the rate, magnitude, and geography of biodiversity trends, and to identify
157 opportunities to mitigate human impacts on riverine systems (Pereira & Cooper, 2006; Anderson,
158 2018). Due to the paucity of spatially- and temporally-extensive datasets in freshwater compared
159 to terrestrial systems (Heino, 2011), RivFishTIME should also help ecologists close the gap
160 between these two systems and to address a wider range of taxa in unraveling large-scale spatio-
161 temporal biodiversity patterns.

162

163 **2| METHODS**

164 **2.1| Data acquisition**

165 We gathered time-series of fish community abundance data for riverine (lotic) ecosystems,
166 broadly defined as freshwater bodies that are continually or intermittently flowing. We tried to
167 the extent possible to exclude wetlands and brackish habitats (salinity > 0.5 ‰). Note, however,
168 that due to the complex nature of the datasets, we do not guarantee that sites are located on free-
169 flowing river segments (i.e. natural condition without impoundment, diversion, or other
170 modification of the waterway). We used the following criteria for data inclusion: (1) the location
171 of the sampling sites is known and consistent through time, (2) the sampling protocol is known
172 and consistent through time, (3) the sampling survey sought to quantify all species in the fish
173 community according to well-established protocols, (4) species-specific abundances are available
174 for each survey, (5) surveys at a given site were conducted over a period of at least 10 years, and
175 (6) at least two yearly surveys with non-null abundance are available. We considered abundance
176 measures derived from direct fish counts, catch-effort indexes such as relative abundances (%)
177 and catch per unit effort (CPUE), abundance classes, as well as statistically estimated
178 abundances (e.g. Leslie method; Ricker, 1975).

179 To identify potential datasets, we used Google Search, Google Scholar and Dataset
180 Search with different combinations of the keywords “time series”, “fish”, “abundance”,
181 “stream”, “river”, “freshwater”, “community”, “temporal”, and “monitoring” or “monitoring
182 program”. We screened the scientific as well as the grey literature to identify studies involving
183 temporal datasets of fish communities and conducted similar searches in data repositories such as
184 Dryad (<https://datadryad.org/stash>) and FigShare (<https://figshare.com/>). We also conducted
185 targeted searches for national and regional monitoring programs by adding country names to the

186 previous keywords. For the European Union, we further used the EuMon database as a reference
187 to identify fish monitoring databases (available at http://eumon.ckff.si/about_daeumon.php).

188 We contacted all the authors and monitoring program coordinators to request and obtain
189 permission to publish the data and/or ensure that the license under which the data were publicly
190 released allowed their inclusion in our global effort (e.g. Open Government License, CC0 1.0
191 Universal). We excluded the datasets for which we did not receive permission, unless the
192 reusability of data was clearly stated on the online repositories where the data were released.

193

194 2.2| Quality control

195 *Taxonomy.* We validated species scientific names using the online database Fishbase (Froese &
196 Pauly, 2019). We used the R package *rfishbase* (as of December 2019; Boettiger *et al.*, 2012)
197 and confirmed names with no match manually using the Catalog of Fishes (Fricke *et al.*, 2018).
198 We then selected only records involving ray-finned fishes (Class Actinopterygii), excluding rays
199 and lampreys, and unidentified species.

200 *Coordinates.* We harmonized the coordinate system by projecting (if necessary) the
201 coordinates of the individual datasets using the World Geodetic System (WGS84) as reference
202 geographic coordinate system. We visually inspected the spatial distribution of the sites with
203 respect to their respective country, region, or state borders as given in the original data sources
204 and discarded sites with dubious coordinates (e.g. sites located in the ocean). We also removed
205 sites whose coordinates were located outside of any hydrographic basin using the global major
206 river basin GIS layer in HydroSHEDS (Lehner *et al.*, 2008).

207

208 *Consistent sampling methods.* We excluded surveys lacking information on sampling
209 methods and selected only time-series collected using a consistent sampling protocol through
210 time. The latter evaluation was dataset-specific as dictated by the complexity of the monitoring
211 scheme and the available metadata. For instance, surveys were deemed consistent if they did not
212 experience any major deviation in sampling protocol, and disregarded minor variations (e.g.
213 number of anodes or traps, area sampled) due to survey-specific constraints (e.g. water depth,
214 habitat complexity). By contrast, several monitoring programs implemented alternate sampling
215 protocols to compare the efficiency of different gears (e.g. seining versus electrofishing) or
216 sampling methods (e.g. continuous versus point electrofishing); these time-series conducted at
217 the same sites but using different sampling protocols were kept separate in the database.

218

219 *Duplicates.* We removed duplicates within individual datasets based on the coordinates of the
220 sites, date of the survey, and species collected (e.g. due to different name attribution for the same
221 site). We also identified potential duplicates among datasets (e.g. overlap between state-level and
222 national databases) based on the coordinates of the sites rounded to three digits to account for
223 different post-processing of the individual datasets.

224

225 **2.3| Database formatting**

226 Each entry (species abundance record) was assigned a unique (1) site, (2) survey, and (3) time-
227 series identifier. The site ID corresponds to a given pair of coordinates, the survey ID to a
228 sampling campaign, and the time-series ID to a combination of site \times sampling protocol. We

229 extracted the names of the sampled water bodies (e.g. creek, stream, river) from the available
230 metadata associated with each individual dataset, that we cross-referenced against several
231 continental and national geospatial river networks in GIS (e.g. Australian Hydrological
232 Geospatial Fabric, Ordnance Survey Open Rivers). Additionally, each site ID was assigned to a
233 biogeographic realm following Olson *et al.* (2001), hydrographic basin following HydroSheds
234 (Lehner *et al.*, 2008), and administrative units (country, region and province) based on its
235 coordinates. For each sampling ID, we aggregated abundance records if they were given
236 separately for individuals, size classes or sub-species for each validated species name or if
237 different sampling passes, hauls, or sub-sampling areas were considered. We also converted
238 time-series species abundances to densities or CPUE whenever possible. The different surveys
239 were kept independent when conducted on different occasions within the same calendar year. We
240 provided the year together with the quarter of the survey (1: January-March; 2: April-June; 3:
241 July-September, 4: October-December). We also provided the associated unit (abundance class,
242 count, CPUE, individuals/100m², Leslie index, relative abundance) for each species abundance
243 record. Finally, we extracted basic information regarding the sampling protocol, including details
244 on electrofishing (backpack, shore-based or boat mounted electrofishers), netting (dip nets, gill
245 nets, beach or pelagic seines), trapping (minnow traps, fyke nets or hoop nets), and trawling
246 techniques. Many survey protocols involve a combination of sampling approaches, rendering
247 challenging the inclusion of detailed information about the sampling effort in a standardized
248 way. We therefore encourage the data user to refer to each data source for more information on
249 the sampling methods.

250 The database is organized in three tables (.csv format): the time-series table, the survey
251 table, and the information source table. The tables can be linked using the unique dataset source

252 ID and time-series ID. The time-series table contains: (1) source ID, (2) site ID, (3) time-series
253 ID, (4) sampling method, (5) latitude (WGS 84), (6) longitude (WGS 84), (7) biogeographic
254 realm, (8) hydrographic basin, (9) country (ISO code), (10) region, (11) province, and (12) water
255 body. The survey table contains: (1) time-series ID, (2) survey ID, (3) sampling year, (4)
256 sampling quarter, (5) species scientific name, (6) abundance, and (7) abundance unit. The
257 information source table contains the full citation(s), online link to the raw data when publicly
258 available, as well as the name(s) and contact of the data responsible(s) for each individual
259 dataset. Data curation was performed in the *R* (3.6.0) programming environment (R Core Team,
260 2019).

261 A list of the data sources is found in Appendix 1; for further information consult the
262 metadata. We provide a static version of the database with this article (1951-2019), but we aim to
263 continue interacting with data contributors to update and add new time-series datasets to be
264 released through the iDiv portal (<https://idata.idiv.de/idiv/Content/Databases>) and the more
265 specialized Freshwater Biodiversity Data Portal (<https://data.freshwaterbiodiversity.eu/>).

266

267 **3| RESULTS**

268 Our database includes 11,386 time-series of riverine fish compiled from 46 individual source
269 datasets, representing a total of 106,785 surveys and 646,270 individual species abundance
270 records at 11,072 unique sites. Survey-specific species richness across all time-series ranges
271 from 1 to 50 species, and covers 944 ray-finned fish species. The surveyed sites display a wide
272 distribution along longitudinal and latitudinal gradients, spanning 19 countries, 402 hydrographic
273 basins, and 5 biogeographic realms (Fig. 1a). Despite broad geographical coverage, we note a

274 clear spatial bias towards the Palearctic (European Union) and, to a lesser extent, Nearctic (North
275 America) and Australasia realms. The abundance time-series are largely represented by
276 individual counts, followed by densities (individuals/100m²) and CPUE (Fig. 1b). Abundance
277 classes, Leslie index and relative abundance represent < 1% of the time-series. Electrofishing is
278 by far the main sampling technique used to record the time-series, although variations are
279 noticeable among biogeographic realms (Fig. 1c). For instance, dipnetting sampling techniques
280 are only represented in the Neotropics, whereas gillnetting is the most common gear in the
281 Afrotropics.

282 The time-series cover a time period from 1951 to 2019 and are mainly concentrated over
283 the last two decades (average first year = 1996; Fig. 2a). Surveys have been conducted primarily
284 in the 3rd (July-September) and 4th (October-December) quarters of the year, especially in the
285 Palearctic and Nearctic realms (corresponding to periods of low flows), but all quarters are
286 represented in the different biogeographic realms (Fig. 2b). The mean time span of the time-
287 series is of 19 years and ranges from 10 to 68 years, with the longest time-series located in the
288 Palearctic (Fig. 2c). The sites were sampled from (non-necessarily consecutive) 2 to 52 years,
289 with an average number of yearly surveys of 8 years (Fig 2d). Again, the highest number of
290 yearly surveys was found in the Palearctic. The completeness of the time-series (i.e. ratio of
291 number of yearly surveys to the overall time span) ranges from 4 to 100%, with a mean value of
292 45% (Figure 2e). Importantly, the degree of completeness is largely uncorrelated to the time span
293 of the time-series ($r = 0.05$).

294

295 4| CONCLUSIONS

296 Our collective effort provides the most comprehensive long-term community database of riverine
297 fishes to date, spanning large biogeographic, climatic and hydrographic gradients. Almost all
298 biogeographic realms are represented but it is important to note that our database is not exempt
299 from spatial bias. For instance, less than 1% of the time-series belong to the Afrotropic or
300 Neotropic realms, whereas 84% belong to the Palearctic realm. These spatial gaps often present
301 in biodiversity-rich regions (tropical areas, southeast Asia) are likely to mirror the current
302 networks of freshwater monitoring programs (Buss *et al.*, 2015; Radinger *et al.*, 2019) as well as
303 biodiversity research efforts (Martin *et al.*, 2012), and hence will be prioritized in future updates
304 of RivFishTIME. We also warn data users that species abundance may not be directly
305 comparable across sites without a full understanding of the specifics of sampling approach and
306 effort, with respect to their selectivity and efficiency (Goffaux *et al.*, 2005; Portt *et al.*, 2006;
307 Oliveira *et al.*, 2014; Benejam *et al.*, 2012), and refer to the original data sources for more
308 information about the sampling protocols.

309 Despite these unavoidable limitations associated with secondary datasets collected for
310 multiple purposes, we are confident that RivFishTIME will stimulate new research in the fields
311 of global change ecology and macroecology. First and foremost, it will provide the needed
312 baseline information for conservation and restoration efforts to bend the curve of freshwater
313 biodiversity loss (Tickner *et al.*, 2020). For instance, the fish abundance time-series could be
314 used to assess population or community trends in different rivers of the world, broadening the
315 taxonomic and spatial representation of existing indicators of the status of global biodiversity
316 (e.g. Living Planet Index). Coupled with high-resolution environmental time-series, this unique
317 database could also help to decipher the underlying drivers of biodiversity changes in riverine
318 systems, including (but not limited to) habitat fragmentation and destruction, invasive species,

319 pollution, hydrologic alteration and climate change (e.g. Chen & Olden, 2020, Erős *et al.*, 2020).
320 In turn, this knowledge could be integrated into ecological models used to forecast how fish
321 communities will respond to future environmental change, paving the way to mitigate those
322 impacts. RivFishTIME could also offer new macroecological insights into the implications of
323 river network complexity on community structure and assembly processes across extensive
324 environmental gradients (e.g. community composition, population persistence, spatial synchrony
325 in community dynamics) – questions that have long fascinated ecologists but have been so far
326 primarily explored through theoretical approaches.

327

328 REFERENCES

- 329 Anderson, C.B. (2018) Biodiversity monitoring, earth observations and the ecology of scale.
330 *Ecology Letters*, **21**, 1572–1585.
- 331 Benejam, L., Alcaraz, C., Benito, J., Caiola, N., Casals, F., Maceda-Veiga, A., ... García-
332 Berthou, E. (2012). Fish catchability and comparison of four electrofishing crews in
333 Mediterranean streams. *Fisheries Research*, **123-124**: 9–15.
- 334 Boettiger, C., Lang, D.T. & Wainwright, P.C. (2012) rfishbase: exploring, manipulating and
335 visualizing FishBase data from R. *Journal of fish biology*, **81**, 2030–2039.
- 336 Buss, D.F., Carlisle, D.M., Chon, T.S., Culp, J., Harding, J.S., Keizer-Vlek, ... Hughes, R.M.
337 (2015) Stream biomonitoring using macroinvertebrates around the globe: a comparison of
338 large-scale programs. *Environmental Monitoring and Assessment*, **187**.
- 339 Butchart, S.H.M., Walpole, M., Collen, B., van Strien, A., Scharlemann, J.P.W., Almond,
340 R.E.A., Baillie, J.E.M., ... Watson, R. (2010) Global biodiversity: Indicators of recent
341 declines. *Science*, **328**, 1164–1168.
- 342 Dornelas, M., Antão, L.H., Moyes, F., Bates, A.E., Magurran, A.E., Adam, D., Akhmetzhanova,
343 A.A., ... Zettler, M.L. (2018) BioTIME: A database of biodiversity time series for the
344 Anthropocene. *Global Ecology and Biogeography*, **27**, 760–786.
- 345 Dornelas, M., Gotelli, N.J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C. & Magurran, A.E.
346 (2014) Assemblage time series reveal biodiversity change but not systematic loss. *Science*,
347 **344**, 296–299.
- 348 Chen, K., & Olden, J.D. (2020). Threshold responses of riverine fish communities to land use
349 conversion across regions of the world. *Global Change Biology*, **26**, 4952–4965.
- 350 Erős, T., Comte, L., Filipe, A.F., Ruhi, A., Tedesco, P.A., Brose, U., Fortin, M.-J., ... Olden,

- 351 J.D. (2020) Effects of nonnative species on the stability of riverine fish communities.
352 *Ecography*, **43**, 1156–1166.
- 353 Fricke, R., Eschmeyer, W.N. & van der Laan, R. (eds.) (2018) Catalogue of fishes: Genera,
354 species, references.
355 <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>.
356 Electronic version accessed December 2020.
- 357 Froese, R. & Pauly, D. (eds.) (2019) FishBase. World Wide Web electronic publication.
358 Available at <http://www.fishbase.org>.
- 359 Gido, K. (2017) CFC01 Kings Creek long-term fish and crayfish community sampling at Konza
360 at Konza Prairie [Data set]. *Environmental Data Initiative*.
- 361 Goffaux, D., Grenouillet, G. & Kestemont, P. (2005) Electrofishing versus gillnet sampling for
362 the assessment of fish assemblages in large rivers. *Archiv für Hydrobiologie*, **162**, 73–90.
- 363 Heino, J. (2011) A macroecological perspective of diversity patterns in the freshwater realm.
364 *Freshwater Biology*, **56**, 1703–1722.
- 365 Hering, D., Verdonshot, P.F.M., Moog, O. & Sandin, L. (eds.) (2004) *Integrated assessment of*
366 *running waters in Europe*. Springer, Dordrecht.
- 367 Inchausti, P. & Halley, J. (2001) Investigating long-term ecological variability using the global
368 population dynamics database. *Science*, **293**, 655–657.
- 369 Kendall, B.E., Prendergast, J. & Bjørnstad, O.N. (1998) The macroecology of population
370 dynamics: taxonomic and biogeographic patterns in population cycles. *Ecology Letters*, **1**,
371 160–164.
- 372 Kuehne, L., Olden, J.D., Strecker, A., Lawler, J.J., & Theobald, D. (2017) Past, present, and
373 future of ecological integrity assessment for fresh waters. *Frontiers in Ecology and the*
374 *Environment*, **15**, 197–205.
- 375 Lehner, B., Verdin, K. & Jarvis, A. (2008) New global hydrography derived from spaceborne
376 elevation data. *Eos, Transactions American Geophysical Union*, **89**, 93.
- 377 Lévêque, C., Oberdorff, T., Paugy, D., Stiassny, M.L.J. & Tedesco, P.A. (2008) Global diversity
378 of fish (Pisces) in freshwater. *Hydrobiologia*, **595**, 545–567.
- 379 Loh, J., Green, R.E., Ricketts, T., Lamoreux, J., Jenkins, M., Kapos, V. & Randers, J. (2005) The
380 Living Planet Index: using species population time series to track trends in biodiversity.
381 *Philosophical Transactions of the Royal Society B: Biological Sciences*, **360**, 289–295.
- 382 Martin, L.J., Blossey, B. & Ellis, E. (2012) Mapping where ecologists work: biases in the global
383 distribution of terrestrial ecological observations. *Frontiers in Ecology and the*
384 *Environment*, **10**, 195–201.
- 385 Matthews, W.J. & Marsh-Matthews, E. (2017) *Stream fish community dynamics. A Critical*
386 *Synthesis*, Johns Hopkins University Press, Baltimore, Maryland.
- 387 Olden, J.D., Kennard, M.K., Leprieur, F., Tedesco, P.A., Winemiller, K.O & García-Berthou, E.
388 (2010) Conservation biogeography of freshwater fishes: past progress and future directions.
389 *Diversity and Distributions*, **16**, 496–513.
- 390 Oliveira, A.G., Gomes, L.C., Latini, J.D. & Agostinho, A.A. (2014) Implications of using a
391 variety of fishing strategies and sampling techniques across different biotopes to determine

- 392 fish species composition and diversity. *Natureza & Conservação*, **12**, 112–117.
- 393 Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V.N., Underwood,
394 E.C., ... Kassem, K.R. (2001) Terrestrial Ecoregions of the World: A New Map of Life on
395 Earth: A new global map of terrestrial ecoregions provides an innovative tool for
396 conserving biodiversity. *BioScience*, **51**, 933–938.
- 397 Paulsen, S.G., Mayo, A., Peck, D. V., Stoddard, J.L., Tarquinio, E., Holdsworth, S.M., Sickle, J.
398 Van, ... Olsen, A.R. (2008) Condition of stream ecosystems in the US: an overview of the
399 first national assessment. *Journal of the North American Benthological Society*, **27**, 812–
400 821.
- 401 Pereira, H. & Cooper, H.D. (2006) Towards the global monitoring of biodiversity change.
402 *Trends in Ecology & Evolution*, **21**, 123–129.
- 403 Portt, C.B., Coker, G.A., Ming, D.L. & Randall, R.G. (2006) A review of fish sampling methods
404 commonly used in Canadian freshwater habitats. *Canadian technical report of fisheries and
405 aquatic sciences*, **2604**, 1–58.
- 406 R Core Team (2019) R: a language and environment for statistical computing. R Foundation for
407 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- 408 Radinger, J., Britton, J.R., Carlson, S.M., Magurran, A.E., Alcaraz-Hernández, J.D., Almodóvar,
409 A., ... García-Berthou, E. (2019) Effective monitoring of freshwater fish. *Fish and
410 Fisheries*, **20**, 729–747.
- 411 Reid, A.J., Carlson, A.K., Creed, I.F., Eliason, E.J., Gell, P.A., Johnson, P.T.J., ... Cooke, S.J.
412 (2019) Emerging threats and persistent conservation challenges for freshwater biodiversity.
413 *Biological Reviews*, **94**, 849–873.
- 414 Revenga, C., Campbell, I., Abell, R., de Villiers, P. & Bryer, M. (2005) Prospects for monitoring
415 freshwater ecosystems towards the 2010 targets. *Philosophical Transactions of the Royal
416 Society B: Biological Sciences*, **360**, 397–413.
- 417 Ricker, W.E. (1975) Computation and interpretation of biological statistics of fish populations.
418 *Bulletin of the Fisheries Research Board of Canada*, **191**, 1–332.
- 419 Sibly, R.M., Barker, D., Denham, M.C., Hone, J. & Pagel, M. (2005) On the regulation of
420 populations of mammals, birds, fish, and insects. *Science*, **309**, 607–610.
- 421 Tedesco, P.A., Paradis, E., Lévêque, C. & Hugueny, B. (2017) Explaining global-scale
422 diversification patterns in actinopterygian fishes. *Journal of Biogeography*, **44**, 773–783.
- 423 Tickner, D., Opperman, J.J., Abell, R., Acreman, M., Arthington, A.H., Bunn, S.E., ... Young,
424 L. (2020) Bending the curve of global freshwater biodiversity loss: An emergency recovery
425 plan. *BioScience*, **70**, 330–342.

426

427

428 **Data Storage and Documentation**

429 We provide a temporary link to the database at <https://figshare.com/s/97c447c4f16d92baf254>

430 (confidential; for review purposes only). The database and associated DOI will be made publicly
431 available through the iDiv portal (<https://idata.idiv.de/idiv/Content/Databases>) upon publication.

432

433 **ACKNOWLEDGMENTS**

434 The data compilation is the result of a joint effort of the international working group “sYNGEO -
435 The geography of synchrony in dendritic networks” kindly supported by sDiv, the Synthesis
436 Centre of the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig,
437 funded by the German Research Foundation (FZT 118). We thank the iDiv Biodiversity
438 Informatics unit and Jitendra Gaikwad for the support received to prepare, upload and scrutinize
439 dataset integrity. We also would like to thank Angelo Agostinho, Agencia Catalana del Agua,
440 Agencia Vasca del Agua, Jammes Gammon, Iowa Department of Natural Resources, William
441 Matthews, Edie Marsh-Matthews, Minnesota Pollution Control Agency, Montana Fish, Wildlife
442 & Parks, Murray-Darling Basin Authority, North Carolina Department of Environmental
443 Quality, Okavango Research Institute, Oklahoma Water Resources Board, Office Français de la
444 Biodiversité (OFB), Parks Canada, Vincent Resh, John Rinne, Swedish University of
445 Agricultural Sciences, Toronto and Region Conservation Authority, U.K. Environmental
446 Agency, Upper Midwest Environmental Sciences Center, U.S. Environmental Protection
447 Agency, U.S. Geological Survey and Gerlinde Van Thuyne for making their monitoring data
448 available. Data collection for the SourceID #2 was supported by Fapesp - São Paulo Research
449 Foundation through the research grants to J.O.Z. (Processes Numbers: 2012/20280-5 and
450 2015/04366-5) and LC (Processes Numbers: 2001/13340-7, 2012/05983-0) and CNPq (Conselho
451 Nacional de Desenvolvimento Científico e Tecnológico) through the research grant provided to
452 LC (Process Number: 301877/2017-3). Data collection for the SourceID #3 was supported by

453 Convenio CT-2017- 001714 between University of Antioquia and Empresas Publicas de
454 Medellin. Part of the data collection for the SourceID #4 was supported by the GINOP-2.3.2.-15-
455 2016-00004 project. Data for the SourceID #6 were collected by the Queensland Department of
456 Environment and Science in collaboration with Healthy Land and Water Limited. Data collection
457 for the SourceID #10 was supported by the New Mexico Department of Game and Fish. We
458 acknowledge CNPq for research grants provided to J.R.S.V. (Process Numbers:302367/2018-7
459 and 303776/2015-3) [SourceID #13], and the Japan Society for the Promotion of Science
460 KAKENHI Grant 18K06404 to A.T. [SourceID #14]. We also acknowledge the assistance of Dr.
461 Stefano Porcellotti, Dr. Simona Piccini (UTR Grosseto) and Dr. Lorena Di Iulio Chiacchia (UTR
462 Prato), who kindly helped with original data retrieval for the SourceID #16. Data collection for
463 the SourceID #17 (Ivory Coast) was supported by the World Health Organisation (WHO),
464 Onchocerciasis Control Programme (OCP). Data collection for the SourceID #18 was supported
465 by the United States Fish and Wildlife Service and Oklahoma Department of Wildlife
466 Conservation through Endangered Species Act funding (Project E-12). Data collection for the
467 SourceID #19 (Kings Creek watershed) was supported by the NSF Long-Term Ecological
468 Research Program at Konza Prairie Biological Station. Data collection for the SourceID #20
469 (Upper Little Tennessee River Watershed) was supported by Brad Stanback, Janirve Foundation,
470 Macon County Community Foundation, Merck Family Fund, National Fish and Wildlife
471 Foundation, National Forest Foundation, NC Division of Water Quality, NC Wildlife Resources
472 Commission, Norcross Foundation, River Network, Southern Appalachian Man and the
473 Biosphere Foundation, Tennessee Valley Authority, World Wildlife Fund, and Z. Smith
474 Reynolds Foundation. Data collection for the SourceID #29 was supported by the Companhia
475 Energética de Minas Gerais (P&D ANEEL/CEMIG GT-487 and GT-599), Fundação de Amparo

476 à Pesquisa do Estado de Minas Gerais (FAPEMIG), Coordenação de Aperfeiçoamento de
477 Pessoal de Nível Superior (CAPES) and CNPq. Data collection for the SourceID #35 (Upper
478 Paraná River) was supported by the Long-Term Research Program / CNPq-Sítio 6, carried out by
479 Nupélia / UEM (Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura/Universidade
480 Estadual de Maringá). R.B.D.C. received a scholarship from CAPES and F.G.B. received a
481 student scholarship from CNPq and financial support from WWF (Programa Natureza e
482 Sociedade) for sampling done in 1999 [SourceID #37]. Data collection for the SourceID #40 was
483 supported by the New Mexico Department of Game and Fish. Data collection for the SourceID
484 #43 was supported by various town councils and the Catalan Water Agency through the
485 "Observatori de la Tordera" project (led by Dr Martí Boada).

486

487 **DATA AVAILABILITY**

488 RivFishTIME is publicly available through the iDiv Biodiversity Portal:

489 <https://doi.org/10.25829/ividiv.1873-10-4000>. We kindly ask the users to cite this paper as well as

490 the source of each primary dataset in any published material produced using these data. We

491 encourage any potential data contributor to contact LC with possible datasets to expand the

492 database. Updates of RivFishTIME will be curated through the iDiv Biodiversity Portal and also

493 released through the more specialized Freshwater Biodiversity Data Portal

494 (<https://data.freshwaterbiodiversity.eu/>).

495


496 **BIOSKETCH**

497 Lise Comte is a global change ecologist who is interested in the mechanisms shaping the

498 (re)distribution of biodiversity in an era of global environmental change, with a special emphasis


499 on freshwater fish. All co-authors share the common goal of advancing freshwater biodiversity
500 conservation and hope that RivFishTIME will generate new knowledge to help bend the curve of
501 global freshwater biodiversity loss.

502



504

505 **Figure 1.** (a) Map showing the distribution of the time-series where each time-series is
 506 represented by a dot with colors indicating the biogeographic realm and size representing fish
 507 species richness (averaged across surveys). Inset histograms display the distribution of the time-
 508 series according to latitude and longitude. Barplots show the distribution of the time-series with
 509 respect to the (b) type of abundance, and (c) primary sampling method. Note the log₁₀(x+1) y-
 510 axes in (b)-(c).



511

512 **Figure 2.** (a) Temporal distribution of the yearly surveys relative to the period covered by the
 513 database (1951-2019). Each time-series appears in rows where the background colors correspond
 514 to the biogeographical realms and white indicates sampled years. (b) Temporal distribution of the
 515 surveys with respect to the quarter of the year. Temporal characteristics of the time-series with
 516 respect to the (c) overall time span, (d) number of yearly surveys, and (e) completeness defined
 517 as the ratio between the number of yearly surveys and the overall time span (expressed in %).
 518 Note the $\log_{10}(x+1)$ y-axes in (b)-(e).

Appendix 1 - Data sources

SourceID	Citations	URLaccess
1	Agència Catalana de l'Aigua (2003); Agència Catalana de l'Aigua (2010); Agència Catalana de l'Aigua (2018)	http://aca.gencat.cat/
2	Zeni <i>et al.</i> (2017); Casatti <i>et al.</i> (2009)	–
3	Universidad de Antioquia-Empresas Publicas de Medellin (2019)	–
4	Erős <i>et al.</i> (2014)	–
5	Gammon (2013)	–
6	Ecosystem Health Monitoring Program Queensland (2019)	https://hlw.org.au/report-card/
7	Finnish electrofishing register Hertta (2019)	https://www.wp2.ymparisto.fi/koekalastus_sahko/yhteinen/Login.aspx?ReturnUrl=%2fkoekalastus_sahko
8	Sigouin (2017)	https://open.canada.ca/data/en/dataset/fe2441a6-8ae4-4884-b181-cd7ec53bd842
9	Whitney <i>et al.</i> (2016)	–
10	Gido <i>et al.</i> (2013); Gido <i>et al.</i> (2019)	–
11	Kesner and Marsh (2010)	https://www.rosemonteis.us/documents/kesner-marsh-2010
12	Griffith (2017); Griffith <i>et al.</i> (2018)	https://doi.org/10.23719/1376690
13	Occhi, V. T. & Vitule, J. R. S. (Unpublished data)	–

14	Terui <i>et al.</i> (2018)	–
15	Iowa DNR (2013)	https://data.iowa.gov/Environment/BioNet/e7yf-f5fs
16	Milardi <i>et al.</i> (2020)	–
17	Levêque <i>et al.</i> (2003)	–
18	Pyron <i>et al.</i> (1998)	–
19	Gido (2017)	https://doi.org/10.6073/pasta/150e218b069074a8ecede85a7406d43f
20	McLarney <i>et al.</i> (2013)	http://coweeta.uga.edu/dbpublic/personnel_bios.asp?id=wmclarney
21	Long Term Resource Monitoring Program (2016)	https://www.umesc.usgs.gov/data_library/fisheries/fish1_query.shtml
22	Matthews and Marsh-Matthews (2017)	https://doi.org/10.5061/dryad.2435k
23	Murray-Darling Basin Authority (2018)	https://data.gov.au/data/dataset/murray-darling-basin-fish-and-macroinvertebrate-survey
24	Minnesota Pollution Control Agency (2018)	https://cf.pca.state.mn.us/water/watershedweb/wdip/search_more.cfm?datatype=assessments
25	Montana, Fish, Wildlife & Parks (2019)	http://gis-mtftp.opendata.arcgis.com/items/8192e75218c6460ba97ba3dd0a2fb3a5
26	U.S. Geological Survey (2019)	https://aquatic.biodata.usgs.gov/clearCriteria.action
27	U.K. Environmental Agency (2019)	https://data.gov.uk/dataset/d129b21c-9e59-4913-91d2-82faef1862dd/nfpd-freshwater-fish-survey-relational-datasets

28	North Carolina Department of Environmental Quality (2018)	https://deq.nc.gov/about/divisions/water-resources/water-resources-data/water-sciences-home-page/ecosystems-branch/fish-stream-assessment-program
29	Fagundes <i>et al.</i> (2015)	–
30	Winston <i>et al.</i> (1991); Taylor (2010)	https://onlinelibrary.wiley.com/doi/full/10.1111/fwb.13211
31	Mosie and Makati (2018)	https://www.gbif.org/dataset/77929c0a-7506-4b2d-a49d-10fc3312d50d
32	Office Francais de la Biodiversite (2019)	http://www.naiades.eaufrance.fr/acces-donnees#/hydrobiologie
33	Oklahoma Water Resources Board (2019)	http://home-owrb.opendata.arcgis.com/search?tags=fish
34	Agencia Vasca del Agua (2019)	http://www.uragentzia.euskadi.eus/informazioa/ubegi/u81-0003341/eu/
35	Ortega <i>et al.</i> (2015)	–
36	Davenport, S.R. (Unpublished data)	–
37	Dala-Corte <i>et al.</i> (2017)	–
38	Bêche <i>et al.</i> (2009); The Resh Lab (2019)	https://nature.berkeley.edu/reshlab/
39	Toronto and Region Conservation Authority (2018)	https://data.trca.ca/dataset/2018-watershed-fish-community
40	U.S. Fish and Wildlife Service (2017)	–
41	Steffèrud, J. A. (Unpublished data)	–

42	Sers (2013)	https://www.slu.se/en/departments/aquatic-resources1/databases1/database-for-testfishing-in-streams/
43	Benejam <i>et al.</i> (2010); Merciai <i>et al.</i> (2017)	–
44	Miyazono and Taylor (2015)	https://bioone.org/journals/The-Southwestern-Naturalist/volume-60/issue-1/MP-02.1/Long-term-changes-in-seasonal-fish-assembly-dynamics-in-an/10.1894/MP-02.1.short
45	Rinne & Miller (2006)	–
46	Van Thuyne <i>et al.</i> (2013); Brosens <i>et al.</i> (2015)	https://ipt.inbo.be/resource?r=vis-inland-occurrences

References

- Agència Catalana de l'Aigua (2003) *Desenvolupament d'un índex d'integritat biòtica (IBICAT) basat en l'ús dels peixos com a indicadors de la qualitat ambiental dels rius a Catalunya*. Aplicació de la Directiva Marc en Política d'Aigües de la Unió Europea (2000/60/CE). 207 pages.
- Agència Catalana de l'Aigua (2010) *Ajust de l'Índex d'Integritat Biòtica (IBICAT) basat en l'ús dels peixos com a indicadors de la qualitat ambiental als rius de Catalunya*. Departament de Medi Ambient i Habitatge, Generalitat de Catalunya. 187 pages.
- Agència Catalana de l'Aigua (2018) ACORD GOV/139/2013, de 15 d'octubre, pel qual s'aprova el Programa de seguiment i control del Districte de conca fluvial de Catalunya per al període 2013-2018.
- Agencia Vasca del Agua (2019) UBEGI. Información sobre el estado de las masas de agua de la CAPV. Available at <http://www.uragentzia.euskadi.eus/informacion/ubegi/u81-0003341/es/> [Accessed 10/10/2019].
- Bêche, L.A., Connors, P.G., Resh, V.H. & Merenlender, A.M. (2009) Resilience of fishes and invertebrates to prolonged drought in two California streams. *Ecography*, **32**, 778–788.
- Benejam, L., Angermeier, P.L., Munné, A. & García-Berthou, E. (2010) Assessing effects of water abstraction on fish assemblages in Mediterranean streams. *Freshwater Biology*, **55**, 628–642.

- Brosens, D., Breine, J., Van Thuyne, G., Belpaire, C., Desmet, P. & Verreycken, H. (2015) VIS – A database on the distribution of fishes in inland and estuarine waters in Flanders, Belgium. *ZooKeys*, **475**, 119-145.
- Bunn, S.E., Abal, E.G., Smith, M.J., Choy, S.C., Fellows, C.S., Harch, B.D., ... Sheldon, F. (2010) Integration of science and monitoring of river ecosystem health to guide investments in catchment protection and rehabilitation. *Freshwater Biology*, **55**, 223–240.
- Casatti, L., Ferreira, C.P. & Carvalho, F.R. (2009) Grass-dominated stream sites exhibit low fish species diversity and dominance by guppies: an assessment of two tropical pasture river basins. *Hydrobiologia*, **632**, 273–283.
- Dala-Corte, R.B, Becker, F.G. & Melo, A.S. (2017) The importance of metacommunity processes for long-term turnover of riffle-dwelling fish assemblages depends on spatial position within a dendritic network. *Canadian Journal of Fisheries and Aquatic Sciences*, **74**, 101–115.
- Erős, T., Sály, P., Takács, P., Higgins, C.L., Bíró, P. & Schmera, D. (2014) Quantifying temporal variability in the metacommunity structure of stream fishes: the influence of non-native species and environmental drivers. *Hydrobiologia*, **722**, 31–43.
- Fagundes, D.C., Leal, C.G., Carvalho, D.R., Junqueira, N.T., Langeani, F. & Pompeu, P.S. (2015) The stream fish fauna from three regions of the Upper Paraná River basin. *Biota Neotropica*, **15**, 1–8.
- Finnish electrofishing register Hertta (2019) Available at https://www.p2.ymparisto.fi/koekalastus_sahko/yhteinen/Login.aspx?ReturnUrl=%2fkoekalastus_sahko [Accessed on 04/09/2019].
- Gammon, J. (2003) The fish communities of Big Raccoon Creek 1981-2002. Report for Heritage Environmental Services. Indianapolis, Indiana. 258 pages.
- Gido, K. (2017) CFC01 Kings Creek long-term fish and crayfish community sampling at Konza Prairie. Environmental Data Initiative. Available at <https://doi.org/10.6073/pasta/150e218b069074a8ecede85a7406d43f> [Accessed on 11/18/2019].
- Gido, K.B, Propst, D.L., Olden, J.D., & Bestgen, K.R. (2013) Multidecadal responses of native and introduced fishes to natural and altered flow regimes in the American Southwest. *Canadian Journal Fisheries and Aquatic Sciences*, **70**, 554–564.
- Gido, K.B, Propst, D.L., Whitney, J.E., Hedden, S.C., Turner, T.F. & Pilger, T.J. (2019) Pockets of resistance: Response of arid-land fish communities to climate, hydrology, and wildfire. *Freshwater Biology*, **64**, 761–777.
- Griffith, M. (2017) Combined Fish Data. U.S. EPA Office of Research and Development (ORD), doi: 10.23719/1376690.

- Griffith, M., Zheng, L. & Cormier, S. (2018) Using extirpation to evaluate ionic tolerance of freshwater fish. *Environmental Toxicology and Chemistry*, **37**, 871–883.
- Iowa DNR [Department of Natural Resources] (2013) BioNet - Iowa DNR Biological Monitoring and Assessment Database. Available at <https://programs.iowadnr.gov/bionet/> [Accessed on 02/01/2018].
- Kesner, B.R. & Marsh, P.C. (2010) Central Arizona project fish monitoring: Final report. Analysis of fish population monitoring data for selected waters of the Gila River Basin, Arizona, for the five year period 2005-2009. Contract No. R09PD32013. Submitted to U.S. Bureau of Reclamation. Tempe, Arizona: Marsh and Associates, LLC. 50 pages.
- Levêque, C., Hougard, J.M., Resh, V., Statzner, B. & Yaméogo, L. (2003) Freshwater ecology and biodiversity in the tropics: what did we learn from 30 years of onchocerciasis control and the associated biomonitoring of West African rivers? *Hydrobiologia*, **500**, 23–49.
- Long Term Resource Monitoring Program (2016) Available at https://www.umesc.usgs.gov/data_library/fisheries/fish1_query.shtml [Accessed on 06/09/2019].
- Matthews, W.J. & Marsh-Matthews, E. (2017) Data from: *Stream fish community dynamics: a critical synthesis*, Dryad, Dataset, doi: 10.5061/dryad.2435k.
- McLarney, W.O., Meador, J. & Chamblee, J. (2013) Upper Little Tennessee River biomonitoring program database. Coweeta Long Term Ecological Research Program. Available at: https://coweeta.uga.edu/dbpublic/dataset_details.asp?accession54045 [Accessed on 11/15/2019].
- Merciai, R., Molons-Sierra, C., Sabater, S. & García-Berthou, E. (2017) Water abstraction affects abundance, size-structure and growth of two threatened cyprinid fishes. *Plos One*, **12**, e0175932.
- Milardi, M., Gavioli, A., Soana, E., Lanzoni, M., Fano, E.A. & Castaldelli, G. (2020) The role of species introduction in modifying the functional diversity of native communities. *Science of the Total Environment*, **699**, 134364.
- Minnesota Pollution Control Agency (2018) Surface water data. Available at https://cf.pca.state.mn.us/water/watershedweb/wdip/search_more.cfm?datatype=assessments [Accessed on 04/03/2019].
- Miyazono, S. & Taylor, C.M. (2015) Long-term changes in seasonal fish assemblage dynamics in an adventitious desert stream. *The Southwestern Naturalist*, **60**, 72–79.
- Montana, Fish, Wildlife & Parks (2019) Fish Survey Sites. Available at <http://gis-mtftp.opendata.arcgis.com/datasets/8192e75218c6460ba97ba3dd0a2fb3a5> [Accessed on 05/21/2019].

- Mosie, I. & Makati, K. (2018) Long term time-series data on fish monitoring by Okavango Research Institute, Botswana. Version 1.2. Okavango Research Institute. Available at <https://www.gbif.org/dataset/77929c0a-7506-4b2d-a49d-10fc3312d50d> [Accessed on 05/11/2019], doi: 10.15468/4vwwzc.
- Murray-Darling Basin Authority (2018) Murray-Darling Basin Fish and Macroinvertebrate Survey. Available at <https://data.gov.au/data/dataset/7826d7c9-bcc5-48c0-832a-66aaedfe7b0f> [Accessed on 10/15/2019].
- North Carolina Department of Environmental Quality (2018) Fish Community Assessment Data. Available at <https://deq.nc.gov/about/divisions/water-resources/water-resources-data/water-sciences-home-page/biological-assessment-branch/fish-community> [Accessed on 02/06/2018].
- Office Français de la Biodiversité (2019) Suivi des éléments biologiques ‘POISSONS’ des rivières françaises. Available at <http://www.naiades.eaufrance.fr/acces-donnees#/hydrobiologie> [Accessed on 07/08/2019].
- Oklahoma Water Resources Board (2019) OWRB Open Data. Available at <http://home-owrb.opendata.arcgis.com/search?tags=fish> [Accessed on 10/16/2019].
- Ortega, J.C.G., Dias, R.M., Petry, A.C., Oliveira, E.F. & Agostinho, A.A. (2015) Spatio-temporal organization patterns in the fish assemblages of a Neotropical floodplain. *Hydrobiologia*, **745**, 31–41.
- Pyron, M., Vaughn, C.C., Winston, M.R. & Pigg J. (1998) Fish assemblage structure from 20 years of collections in the Kiamichi River, Oklahoma. *Southwestern Naturalist*, **43**, 336–343.
- Rinne, J.N., & Miller, D. (2006) Hydrology, geomorphology and management: implications for sustainability of native southwestern fishes. *Reviews in Fisheries Sciences* **14**, 91–110.
- Sers, B. (2013) Swedish Electrofishing RegiSter – SERS. Swedish University of Agricultural Sciences (SLU), Department of Aquatic Resources. Available at <http://www.slu.se/elfiskeregistret> [Accessed on 11/18/2019].
- Sigouin, D. (2017) Fish Communities – Forillon, dataset fe2441a6-8ae4-4884-b181-cd7ec53bd842. Available at <https://open.canada.ca/data/en/dataset/fe2441a6-8ae4-4884-b181-cd7ec53bd842> [Accessed on 05/07/2019].
- Taylor, C.M. (2010) Covariation among plains stream fish assemblages, flow regimes, and patterns of water use. In: *Community ecology of stream fishes: concepts, approaches, and techniques*. American Fisheries Society Symposium, 2010.
- The Resh Lab (2019) Hunting Creek and Knoxville Creek long-term data. Available at: https://nature.berkeley.edu/reshlab/data/HCKC_fish.xls [Accessed 05/19/2019].

- Terui, A., Ishiyama, N., Urabe, H., Ono, S., Finlay, J. C. & Nakamura, F. (2018) Metapopulation stability in branching river networks. *Proceedings of the National Academy of Sciences*, **115**, E5963–E5969.
- Toronto and Region Conservation Authority (TRCA) (2018) Watershed Fish Community. Available at <https://data.trca.ca/dataset/2018-watershed-fish-community> [Accessed on 09/18/2019].
- U.K. Environmental Agency (2019) National Fish Populations Database (NFPD): Freshwater fish survey relational datasets. Available at <https://data.gov.uk/dataset/d129b21c-9e59-4913-91d2-82faef1862dd/nfpd-freshwater-fish-survey-relational-datasets> [Accessed on 10/15/2019].
- Universidad de Antioquia-Empresas Publicas de Medellin. (2019) Monitoreo a la ictiofauna, su habitat y sus servicios ecosistemicos. Convenio CT 2017- 001714.
- U.S. Geological Survey (2019) BioData - Aquatic Bioassessment Data for the Nation. Available at <https://aquatic.biodata.usgs.gov/clearCriteria.action> [Accessed on 10/15/2019], doi:10.5066/F77W698B.
- U.S. Fish and Wildlife Service (2017) San Juan River basin recovery implementation program. Albuquerque, NM, 87113, USA. Data are available by contacting: <https://www.fws.gov/southwest/sjrip/>.
- Van Thuyne, G., Breine, J., Verreycken, H., De Boeck, T., Brosens, D. & Desmet, P. (2013) VIS - Fishes in inland waters in Flanders, Belgium. Research Institute for Nature and Forest (INBO). Dataset/Occurrence. Available at <https://www.gbif.org/dataset/823dc56e-f987-495c-98bf-43318719e30f> [Accessed on 04/11/2017], doi:10.15468/gzyxyd.
- Whitney, J.E., Gido, K.E., Martin, E.C. & Hase K.J. (2016) The first to arrive and the last to leave: colonisation and extinction dynamics of common and rare fishes in intermittent prairie streams. *Freshwater Biology*, **61**, 1321–1334.
- Winston, M.R., Taylor, C.M. & Pigg, J. (1991) Upstream extirpation of four minnow species due to damming of a prairie stream. *Transactions of the American Fisheries Society*, **120**, 98–105.
- Zeni, J.O., Hoinghaus, D.J. & Casatti, L. (2017) Effects of pasture conversion to sugarcane for biofuel production on stream fish assemblages in tropical agroecosystems. *Freshwater Biology*, **62**, 2026–2038.