



# A review of phototoxic plants, their phototoxic metabolites, and possible developments as photosensitizers.

Raphaelle Petit, Jonathan Izambart, Mathieu Guillou, Jackson Guedes da Silva Almeida, Raimundo Gonçalves de Oliveira Junior, Vincent Sol, Tan-Sothea Ouk, Raphael Grougnet, Lucindo Quintans-Junior, Przemyslaw Sitarek, et al.

## ► To cite this version:

Raphaelle Petit, Jonathan Izambart, Mathieu Guillou, Jackson Guedes da Silva Almeida, Raimundo Gonçalves de Oliveira Junior, et al.. A review of phototoxic plants, their phototoxic metabolites, and possible developments as photosensitizers.. Chemistry and Biodiversity, 2023, 10.1002/cbdv.202300494 . hal-04304472

HAL Id: hal-04304472

<https://hal.science/hal-04304472v1>

Submitted on 24 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Review

2 A review of phototoxic plants, their phototoxic metabolites, and possible developments as  
3 photosensitizers.

4 Raphaëlle Petit<sup>1</sup>, Jonathan Izambart<sup>1</sup>, Mathieu Guillou<sup>1</sup>, Pr.Dr. Jackson Roberto Guedes da  
5 Silva Almeida<sup>2,3</sup>, Dr. Raimundo Gonçalves de Oliveira Junior<sup>1,3</sup>, Pr.Dr. Vincent Sol<sup>3,4</sup>, Dr. Tan-  
6 Sothea Ouk<sup>3,4</sup>, Dr. Raphaël Grougnet<sup>3,5</sup>, Pr.Dr. Lucindo José Quintans-Júnior<sup>3,6</sup>, Dr.  
7 Przemyslaw Sitarek<sup>7</sup>, Pr.Dr. Valérie Thiéry<sup>1,3</sup>, Dr. Laurent Picot<sup>1,3\*</sup>

8

9 <sup>1</sup> UMR CNRS 7266 LIENSSs, La Rochelle Université, 17042 La Rochelle, France.

10 <sup>2</sup> NEPLAME, Universidade Federal do Vale do São Francisco, 56304-205, Petrolina,  
11 Pernambuco, Brazil.

12 <sup>3</sup> Franco-Brazilian Network on Natural Products, FB2NP.

13 <sup>4</sup> LABCiS, UR 22722, Université de Limoges, 87000 Limoges, France.

14 <sup>5</sup> UMR CNRS 8038 CiTCoM, Université Paris Cité, 75006 Paris, France.

15 <sup>6</sup> LANEF, Universidade Federal de Sergipe, 49100-000 São Cristóvão, Sergipe, Brazil

16 <sup>7</sup> Department of Medical Biology, Medical University of Lodz, Łódź, Poland

17

18 \*Corresponding author

19 Dr Laurent PICOT, La Rochelle Université, UMR CNRS 7266 LIENSSs, Curie B10 Faculté des  
20 Sciences et Technologies, Avenue Michel Crépeau 17042 La Rochelle, France.

21 E-mail: [laurent.picot@univ-lr.fr](mailto:laurent.picot@univ-lr.fr), homepage: <https://lienss.univ-larochelle.fr/Picot-Laurent>

22 Phone: +33 5 46 45 82 20

23


24

25



26 Raphaëlle PETIT received her Bachelor Degree in Biochemistry and her Master of Science  
27 degree in Applied Blue Biotechnology from La Rochelle University in France. She is currently  
28 doing her Doctoral Degree in Neurology at the Universitätsklinikum Jena in Germany, where  
29 she is working for the Marie Skłodowska-Curie Innovative Training Network “SmartAge” on  
30 the subject: Age-dependent gut-brain interactions in mice: modulation by physical activity.  
31 This article is based on her Master's thesis work.

32



33 Jonathan IZAMBART received his bachelor degree in Biochemistry and his Master of Science  
34 degree in Applied Blue Biotechnology in 2022 from La Rochelle University, France. He is  
35 currently working in a “France Relance” project as an assistant engineer in valorization of local  
36 seaweed biomass in Charente-Maritime. This article is based on his Master's thesis work.

37

38



43

44 Mathieu GUILLOU received his Bachelor Degree in Biochemistry and his Master Degree in  
45 Biotechnology and management in agro-industries in 2022 from La Rochelle University,  
46 France. He is currently working as a quality assistant manager in a food industry in France.  
47 This article is based on his Master's thesis work.

48



49  
50  
51 Pr.Dr. Jackson Roberto Guedes da Silva Almeida is Professor of Department of Pharmacy at  
52 the Federal University of Vale do São Francisco (Petrolina, Pernambuco, Brazil). He received  
53 his Ph.D. (Chemistry of Natural Products) degree from the Federal University of Paraíba (João  
54 Pessoa, Paraíba, Brazil) and performed his post-doctoral fellowship in Pharmaceutical Sciences  
55 at the University of São Paulo (Ribeirão Preto, São Paulo, Brazil). His research focuses on the  
56 study of chemical and pharmacological properties of medicinal plants from the Caatinga biome,  
57 used for the control of pain, inflammation and central nervous system disorders. He has  
58 experience with the extraction, isolation and characterization of the chemical structure of  
59 natural products as well as the evaluation of biological activity. He has published 296 articles  
60 in the fields of chemistry and pharmacology of natural products. He is also a co-author of 31  
61 book chapters and has directed 13 Ph.D. thesis and 27 dissertations. He has spent over 17 years  
62 working in the academic world with national and international collaborations, resulting in  
63 several scientific publications, dissertations, thesis and patents.



64  
65 Dr. Raimundo GONÇALVES DE OLIVEIRA JUNIOR received his Master of Science degree  
66 in Natural Resources (Universidade Federal do Vale do São Francisco, UNIVASF, Brazil) and  
67 PhD degree in Molecular and Cellular Biology (La Rochelle Université, France). He is  
68 currently an associate professor in Pharmacognosy at the Faculty of Pharmacy of Université  
69 Paris Cité (France). He has been the young researcher coordinator and co-founder of the  
70 Franco-Brazilian Network on Natural Products (FB2NP) since its creation in 2019, and a

71 member of the francophone association for teaching and research in pharmacognosy (AFERP,  
72 France).

73 He belongs to the Natural Products, Analysis and Synthesis (PNAS) team from the UMR  
74 CNRS 8038 CiTCoM laboratory, working on eco-responsible methods for extraction,  
75 purification and biotransformation of secondary metabolites from tropical plants (especially  
76 Brazilian native plants) with anticancer potential.

77

78



79  
80 Pr.Dr. Vincent SOL received his Master of Science and PhD degree in organic chemistry from  
81 Limoges University (France). He performed in a research position (Pr G. Mackenzie) -  
82 University of Hull, UK in 2009-2010. He is currently a full professor of Organic chemistry at  
83 the department of Chemistry of Limoges University and member of Laboratory LABCiS. He  
84 is a member of SCF “Société Chimique de France” and of SPP (Society of Porphyrins and  
85 Phthalocyanines). For 25 years, he has developed the design and the organic synthesis of  
86 different macrocycles but also the extraction of photosensitizers from natural compound. He  
87 has been working on the biological applications of this macrocycles in PhotoDynamic Therapy  
88 (PDT) against cancer cells and bacteria strains.

89



90

91

92 Dr. Tan-Sothea OUK received his Master of Science degree in Molecular and Cellular Biology  
93 (Lille University, France) and his PhD degree in Biology and Health (University of Limoges,  
94 France). He is currently an Associate professor of Biochemistry and Analytical Chemistry at  
95 University Institute of Technology in Limoges (France). He is a member of Laboratory

96 LABCiS (University of Limoges) and coordinator of the research theme “Oncogenesis,  
97 antitumoral and antimicrobial focal therapies”. He has been working on the biological  
98 evaluation of natural and synthetic compounds in antimicrobial photodynamic therapy (aPDT).  
99 He also belongs to the board of a technological center for the valorization of by-products from  
100 plants (Centre Valorization of Agroresources in Brive-la-Gaillarde, France).

101



102

103

104 Dr. Raphaël GROUGNET received his Pharmacy Master of Science and PhD degree in Natural  
105 Products chemistry from Paris Descartes University (France). He is an associate professor in  
106 Pharmacognosy at the Faculty of Pharmacy of Paris Cité University, member of the  
107 francophone association for teaching and research in pharmacognosy (AFERP) and of the  
108 Franco-Brazilian Network on Natural Products (FB2NP). He has been working on several  
109 aspects of Natural Products, mainly the development of techniques with low environmental  
110 impact for the extraction, fractionation and isolation (pressurized extraction, centrifugal  
111 partition chromatography) and the implementation of chemical libraries  
112 (polymethoxyflavones, triterpenes, alkaloids).

113

114



115

116 Pr.Dr. Lucindo Quintans-Júnior. He received a Master of Science and PhD degree in Natural  
117 Products and Bioactive Compounds from UFPB (Brazil), performing a post-doctorate  
118 internship in Neuroscience at the University of Iowa (UIOWA, USA). He is currently a Full  
119 Professor at the Department of Physiology of the Federal University of Sergipe (DFS/UFS –  
120 Brazil). He is also a member of the Brazilian Society of Pharmacognosy (SBF), of the Brazilian  
121 Society of Neuroscience and Behavior (SBNeC), The American Society of Pharmacognosy  
122 (ASP) and the American Society for Pharmacology, Experimental Therapeutics (ASPET) and

123 a member of FB2NP – Franco-Brazilian network on Natural Products  
124 (<https://fb2np.recherche.univ-lr.fr/>). He has published around 455 articles, supervised 28  
125 masters and 23 PhD students. His bibliometric indicators are h-index 48 and 8810 citations  
126 (Scopus) and h-index 47 and 8416 citations (Publons). He has been working on the  
127 pharmacological approach of essential oils; terpenes, flavonoids, and alkaloids, mainly for the  
128 management of pain and inflammation.

129



130

131

132 Dr. Przemysław SITAREK received his Master of Science degree in Biology in the specialty  
133 of biochemistry, University of Lodz and his doctoral degree in Medical Biology-genetics in  
134 2012 in Medical University of Lodz. He is an associate Professor at the Department of Medical  
135 Biology, Medical University of Lodz, where he obtained his Professorial Habilitation in  
136 Pharmaceutical Sciences in 2019. He has 10 years of teaching and research experience in the  
137 fields of plant biotechnology, medicinal plants, natural products, apoptosis induction,  
138 metabolic engineering, molecular biology. He has been working on *in vitro* transgenic roots of  
139 medicinal plants for anti-cancer drug discovery.

140

141



142

143

144 Pr.Dr. Valérie THIERY received her Master of Science and PhD degree in organic chemistry  
145 from the University of Orléans, France. After a post-doctoral fellow position in the group of  
146 Pr. Charles W. Rees at Imperial College of London in 1995-1996, she joined La Rochelle  
147 University as an assistant professor in 1996 and since 2007 she has been a Full Professor of  
148 organic chemistry (La Rochelle University, LIENSs laboratory, France). Her current research

149 interests include: medicinal chemistry applied to kinase inhibitors, heteroatomic chemistry  
150 (N,S), microwave assisted methodologies and eco-extraction processes, and valorization of  
151 marine or terrestrial alkaloids in cancerology. She is a member of “Société Chimique de  
152 France” (SCF).

153



154  
155 Dr. Laurent PICOT received his Master of Science degree in Molecular and Cellular  
156 Pharmacology in Paris 6 University and his doctoral degree in Cellular Microbiology in 2003  
157 from Rouen University, France. He is an associate Professor of Biochemistry, Immunology  
158 and Pharmacology at the department of Biotechnology of La Rochelle University, France  
159 where he obtained his Professorial Habilitation in Biochemistry and Marine Pharmacology in  
160 2023. He has 25 years of teaching and research experience in the fields of natural products,  
161 pharmacology, cancerology, marine drugs and neuroscience. He has been working on  
162 microalgae, seaweeds and medicinal plants metabolites for anti-cancer and anti-microbial drug  
163 discovery, including phototherapy, and he coordinates the Franco-Brazilian Network on  
164 Natural Products (FB2NP). He has published more than 80 scientific papers and 1 international  
165 patent in the field of Phototherapy.

166

## 167 Abstract

168 This study provides a comprehensive overview of the current knowledge regarding phototoxic  
169 terrestrial plants and their phototoxic and photosensitizing metabolites. Within the 435,000  
170 land plant species, only around 250 vascular plants have been documented as phototoxic or  
171 implicated in phototoxic occurrences in humans and animals. This work compiles a  
172 comprehensive catalog of these phototoxic plant species, organized alphabetically based on  
173 their taxonomic family. The dataset encompasses meticulous details including taxonomy,  
174 geographical distribution, vernacular names, and information on the nature and structure of  
175 their phototoxic and photosensitizing molecule(s). Subsequently, this study undertook an in-  
176 depth investigation into phototoxic molecules, resulting in the compilation of a comprehensive

177 and up-to-date list of phytochemicals exhibiting phototoxic or photosensitizing activity  
178 synthesized by terrestrial plants. For each identified molecule, an extensive review was  
179 conducted, encompassing discussions on its phototoxic activity, chemical family, occurrence  
180 in plant families or species, distribution within different plant tissues and organs, as well as the  
181 biogeographical locations of the producer species worldwide. The analysis also includes a  
182 thorough discussion on the potential use of these molecules for the development of new  
183 photosensitizers that could be used in topical or injectable formulations for antimicrobial and  
184 anticancer phototherapy as well as manufacturing of photoactive devices.

185

## 186 Abbreviations

187	GBIF	Global Biodiversity Information Facility
188	INPI	French Institute for Intellectual Property
189	PACT	Photodynamic antimicrobial therapy
190	PDT	Photodynamic Therapy
191	PS	Photosensitizer
192	ROS	Reactive Oxygen Species
193	USA	United States of America


194

## 195 Highlights

- Complete list of phototoxic and photosensitizing plant species of the world
- Complete list of plant phototoxic metabolites (Type I and Type II photosensitizations)
- Chemodiversity of phototoxic molecules in relation to botanical families
- Discussion on the possible use of plant phototoxic molecules for phototherapy

200

## 201 Graphical abstract

<p>Exploration of the biodiversity and chemodiversity of phototoxic plants</p> <p>Identification of new photosensitizers</p> <p>Innovation in tumor and antimicrobial phototherapy</p> <p>Development of photodecontaminable clothes and medical devices</p> <p>Prevention of phototoxic events</p>	  
---	--

202

## 203 1. Introduction

204 Natural products and their structural analogues play a key role in drug discovery, particularly  
205 for the treatment of cancer and infectious diseases [1]. They are more than ever a major source  
206 of inspiration for researchers to develop innovative therapies and their interest is currently  
207 revitalised in the face of chemoresistance of cancer cells and microorganisms [2]. Among the  
208 alternative strategies to chemotherapy, photodynamic therapy (PDT) and Photodynamic  
209 antimicrobial therapy (PACT) have emerged as efficient treatments to destroy cancer cells,  
210 tumour vasculature and microorganisms. PDT and PACT are based on the use of a  
211 photoactivatable molecule, the photosensitizer (PS), light, and oxygen naturally present in the  
212 biological medium. After light excitation of the PS, an energy transfer to oxygen generates a  
213 high burst of reactive oxygen species (ROS), such as singlet oxygen and superoxide radical  
214 anion. These highly oxidative species induce amino acid oxidation, proteins fragmentation,  
215 peroxidation of membrane lipids and oxidation of nucleic acids in the immediate biological  
216 environment, leading to rapid cancer cell death through necrosis, apoptosis, necroptosis or  
217 autophagy and destruction of viruses, bacteria, parasites and fungi [3]. This strategy has proven  
218 very efficient to treat solid tumors and for the eradication of pathogenic microorganisms [3–6],  
219 that represent a major public health threat in regard to the emergence of new viruses (Ebola,  
220 coronavirus, etc.), spread of resistant bacteria and fungi, and the growing public distrust of  
221 vaccines. As compared to conventional chemotherapeutic treatments, phototherapy offers

many advantages as PS usually show a low intrinsic toxicity and can be delivered easily and precisely to tumour microenvironments or infection zones. Accurate definition of the photo-irradiation area limits adverse phototoxic effects and the fast and non-specific mode of action of PS avoids the development of chemoresistance, as the life time of singlet oxygen is very short and tumor cells or microorganisms have no possibility to develop chemoresistance in response to the lethal ROS release. Additionally, the possible vectorization and targeted delivery of PS to cell and bacterial membranes, cell walls or cell organelles can further improve the selectivity of PDT.

## 2. Clinical applications of photosensitizers (PS)

The first uses of light as a therapeutic agent date back to ancient times as Egyptians, Chinese and Indians were trying to treat vitiligo, psoriasis and even skin cancer using light. However, it was not until the end of the 19th century that phototherapy became a science and was popularized by the Danish physician Niels Finsen who initiated the use of concentrated light radiation for the treatment of lupus vulgaris. During this same period, the use of PS for therapeutic purposes also began to be studied. In 1898, the concept of photosensitization emerged when Oscar Raab first observed the toxicity of acridine in the presence of light on paramecia. This finding led to extensive studies by Tappeiner, Jodl-bauer and collaborators and to the introduction of the term "photodynamic action," used by Tappeiner in 1904. In 1900, Prime, a French neurologist, observed that epileptic patients treated with oral eosin developed dermatitis in areas exposed to the sun. In 1908, Hausmann demonstrated that hematoporphyrin is a photosensitizer. He reported the destruction of paramecia and red blood cells and described in detail the symptoms of sensitized mice after exposure to light. In 1913, Friedrich Meyer-Betz performed the first human trial of hematoporphyrin, injecting himself with 200 mg of the product. As long as he remained unexposed to the sun, nothing happened, but upon exposure to sunlight, he observed a photosensitization that resulted in hyperpigmentation of the skin and generalized edema, from which he suffered for two months. Following these pioneering discoveries, PDT was progressively developed as an efficient and innovative method to treat tumors and infections, and natural or synthetic phototoxic molecules were developed as PS. The range of PS applications has expanded considerably in recent years, and they are now commonly used for the phototherapy of tumours, destruction of tumour vasculature, as well as the treatment of viral, bacterial and fungal infections of the skin, nails, wounds and mucous membranes. Some examples of routine use of PS include the treatment of acne and keratosis,

254 chronic and acute wound infections, burn infections, and solid topical and internal tumors. PS  
255 have also been more recently proposed and developed for the photo-decontamination of  
256 hospital equipment and manufacture of photodecontaminable surfaces to prepare clothing,  
257 laboratory coats and dressings for hospital and hygiene equipment. For example, a large  
258 number of surfaces and materials (cellulose, cellulose nanocrystals, nanocellulose,  
259 hemicellulose, ...) can be ionically or covalently grafted with natural or synthetic PS and  
260 developed for antimicrobial applications [7–12]. Lignin and chitosan nanoparticles grafted with  
261 PS have also been developed to obtain photoantimicrobial surfaces [13–15]. Several research  
262 groups have also carried out the functionalization of different polymers like Nylon-6,  
263 polyacrylonitrile, polyurethane, polylactic or polyolefin with different PS including cationic  
264 porphyrin, protoporphyrin IX, Bodipy, ruthenium complexes, ... [16–20]. These polymers  
265 functionalized with PS are very effective for antibacterial PDT on *S.aureus*, *E. Coli*, *P.*  
266 *aeruginosa*, for antifungal PDT and for antiviral PDT on the human coronavirus strain HCoV-  
267 229E [19]. However, despite the numerous benefits of phototherapy, only few phototoxic drugs  
268 are currently indicated for clinical use, because of pharmacokinetics limitations or preparation  
269 difficulties.

### 270 3. Searching for new phototoxic chemical scaffolds to stimulate innovation in phototherapy

271 In the seventies, Dr Thomas Dougherty, continuing and expanding upon the work of  
272 phototherapy pioneers, played a major role in the development of PS and modern  
273 photodynamic therapy (PDT) for cancer therapy [21]. The first generation of PS were porphyrins  
274 and hematoporphyrins with long clearance time, low chemical purity and short excitation  
275 wavelength [22]. Photofrin®, the first clinically approved PS for the treatment of cancer, is a  
276 complex mixture of oligomers, which makes its clinical use poorly reproducible. In addition,  
277 it has a relatively low quantum fluorescence yield and a low efficiency for ROS generation.  
278 The second generation of PS includes clinically approved or under-clinical trials drugs such as  
279 Porfimer sodium (Photofrin®), Temoporfin (Foscan®), Motexafin lutetium, Palladium  
280 bacteriopheophorbide, Purlytin®, Verteporfin (Visudyne®) and Talaporfin (Laserphyrin®). In  
281 this second generation, introduction of heavy atoms (halogens and metals) enhanced the singlet  
282 oxygen production yield and prevented PS aggregation in the bloodstream and physiological  
283 fluids. Additionally, expansion of the PS' conjugating systems increased their light absorption  
284 efficiency at wavelengths better penetrating human skin and tissues (650-700 nm).  
285 Pharmacomodulation of PS also increased their bioavailability, lipophilicity and cellular

uptake. Despite these improvements, most current PS still do not meet the requirements of optimal drugs. For example, Foscan® has an enhanced absorption in the near infrared, is very efficient, but has the disadvantage of being relatively toxic and difficult to eliminate from the body, resulting in undesirable residual phototoxicity. Visudyne® and Verteporfin®, two benzoporphyrin-derived PS, have the advantage of not causing skin photosensitivity but are relatively difficult to prepare [22,23]. Most current PS are macrocyclic tetrapyrrolic derivatives (porphyrins and their chlorine and bacteriochlorine derivatives), with high absorbency in the far-red spectral range (650-750 nm), and thus comparable fields of application for phototherapy because of their chemical homology. These gaps justify the importance of exploring new natural PS and synthetic routes, not only in order to develop drugs with improved pharmacokinetics, selectivity and reduced toxicity or systemic phototoxicity, but also to identify novel molecular scaffolds with attractive physicochemical characteristics allowing the development of innovative PS.

#### 4. Chemical ecology as a source of inspiration to identify new photosensitizers: screening phototoxic plants

A lot of terrestrial plants are known to synthesize phototoxic phytochemicals that are released and triggered in response to predation by herbivores or attack by infectious pathogens and parasites [24-30]. However, the total number of phototoxic species and metabolites has never been explored in detail although previous attempts to write comprehensive reviews on the subject are available in the literature [31,32]. Moreover, for some molecules like coumarins, hundreds have been described but many of them have never been evaluated for possible phototherapy applications [33]. Plant phototoxic molecules are synthesized as natural secondary metabolites, produced and possibly induced in response to predation, invasion or infection. Contact with the tissues of predators, invaders or pathogens, followed by sunlight photoactivation eventually kills the pathogens or burns the predators to protect the plant and exert an allelopathic effect [34-37]. This defence system is highly effective not only to repell grazers or invaders, but also to kill microorganisms that could attack the plant tissues using cell wall disrupting enzymes. The existence of this chemistry defence system has been demonstrated in diverse plants from various taxonomic groups, biomes and ecoregions of the world, supporting the hypothesis of a convergent adaptive evolutionary process that has led to a high chemodiversity of phototoxic molecules [27]. In line with these data, many wild and cultivated phototoxic plants have been reported to cause contact photodermatitis in humans. In particular,

phototoxic events are regularly reported in gardeners, farmers, children, employees of plants processing companies and consumers in contact with wild species, horticultural, agricultural or ornamental plants and essential oils. Phototoxic plants are also responsible for episodes of livestock phototoxicity, with a strong impact on the health and survival of animals, especially in countries where animals are kept in natural areas<sup>[38,39]</sup>. It is important to consider two possible mechanisms of phototoxicity<sup>[32,40]</sup>. Primary phototoxicity is induced by molecules that are directly photosensitive and will generate the production of ROS when exposed to light. The phototoxic agent contained in the plant may be phototoxic when in contact with the skin (e.g. furanocoumarins) or may reach the skin when ingested, with or without any chemical modification. The photophysical, photochemical and biochemical processes involved in the primary phototoxicity of these natural products has been described in detail in a previous review<sup>[41]</sup> and rely on the photoxidation of intracellular and membrane biomolecules<sup>[41]</sup>. In contrast, secondary phototoxicity is caused by molecules that are not directly phototoxic but can lead to the production of phototoxic compounds through liver damage when ingested. For example, phylloerythrin is produced by microbial degradation of chlorophyll in ruminant stomachs. Phylloerythrin is normally excreted in bile but accumulates in the blood and skin when hepatic damage interferes with the phylloerythrin conjugating and excreting mechanisms. In the superficial layers of the skin, phylloerythrin is phototoxic because it absorbs ultraviolet radiation of the sun. Hepatogenous sensitization can be observed for example with triterpenes and monoterpenes that are not directly phototoxic as they do not contain light-absorbing chromophores. As such, compounds considered to be photoprotective such as flavonoids or betalaines can also be responsible for secondary photosensitization after ingestion<sup>[32,42–44]</sup>.

## 5. Objective of the study

In this work, our objective was to highlight the potential of plant phototoxic molecules as possible candidates to develop new PS. For this purpose, we achieved an inventory of the biodiversity of phototoxic terrestrial plants and described the structures of phototoxic or photosensitizing molecules identified so far in terrestrial plants. First, it appeared essential to estimate the number of phototoxic plants in relation to the biodiversity of terrestrial plants, and we therefore listed all plants reported to contain phototoxic natural products or to be phototoxic in humans and livestock, detailing their taxonomy and geographical distribution worldwide. In the next section, we listed all phototoxic plant natural products, indicating, when available, their activity in various model studies, their chemical family, their physicochemical

350 characteristics, the species in which they can be found and the geographical distribution of  
351 these species. We discriminated phototoxic molecules involved in primary photosensitization  
352 from molecules involved in secondary photosensitization after ingestion by ruminants. In  
353 some cases, molecules may be phototoxic through both photosensitization processes. This  
354 analysis made it possible to highlight the chemodiversity of phototoxic molecules synthesized  
355 by terrestrial plants and to link the presence of specific phototoxic chemical families to  
356 botanical families. The described plants and phototoxic natural products reveal promising  
357 prospects for the further development of new photosensitizers, for the phototherapy of tumors  
358 and infections.

359

## 360 6. Methods

361 A literature search was conducted to include scientific publications published since 1989 in  
362 five specialised databases (ScienceDirect, Scopus, PubMed, Google Scholar, MDPI) using  
363 different combinations of keywords, the main ones being plant names but also family names  
364 of molecules, as well as other terms such as natural molecule, natural product, plant,  
365 photosensitizer, photosensitization, phototoxic and phototoxicity. A few previous studies,  
366 considered as fundamental pioneer research on phototoxicity and phototherapy were also  
367 included. A total of 84 non-redundant publications and 8 book chapters were found and kept  
368 for the review after checking that their scientific content was relevant to the scope and provided  
369 they met a sufficient scientific quality. Only publications with title, abstract and content in  
370 English were selected. Selected publications were mainly articles, but journals, book chapters  
371 or short communications were also evaluated for inclusion in the study. Articles dealing with  
372 natural photoprotective products were excluded. Information about molecules, including their  
373 chemical family, structure and molecular formula, was checked on the PubChem database and  
374 cross-checked independently by 2 chemists. The taxonomic classification of each plant and  
375 location was checked with the GBIF biodiversity database (<https://www.gbif.org/>). Wherever  
376 the presence of a plant species in a country or biome was reported in our paper, we checked  
377 that the validity of the data was supported by rigorous scientific observation or report.

## 378 7. Results

### 379 7.1. Inventory of phototoxic plants

380 To date about 435,000 land plant species, including 391,000 vascular plants, have been  
381 identified on Earth (Figure 1.) Vascular plants include angiosperms, gymnosperms, filices  
382 (ferns) and lycophytes while bryophytes (non-vascular seedless plants) include mosses,  
383 hornworts and liverworts (Figure 1.). The exact number of world plants species is for certain  
384 underestimated as botanists discover and document about 2,000 new species of flowering  
385 plants every year [45]. Among these species, no phototoxic plant was reported in the subgroup  
386 of bryophytes. Within the 391,000 vascular plant species, only around 250 species were  
387 documented as phototoxic or involved in phototoxic occurrences in humans (Figure 1.). The  
388 inaccurate number of phototoxic species is explained by the fact that in some reports, only the  
389 plant genus was described without specifying the species. Additionally, some species that  
390 contain phototoxic metabolites have not necessarily been the subject of phototoxicity reports  
391 in humans or animals.

## KINGDOM PLANTAE ~ 435 000 species



### **Subgroup VASCULAR PLANTS**

391 000 species: Angiosperms,  
gymnosperms, ficles (ferns) and  
lycophytes

392 **~250 phototoxic species**

### **Subgroup BRYOPHYTES**

#### **NON VASCULAR**

#### **SEEDLESS PLANTS**

44 000 species: Mosses,  
hornworts, liverworts

**0 phototoxic species**

393 Figure 1. Repartition of the 435,000 world plant species in taxonomic subgroups and  
394 distribution of phototoxic world plant species. Only around 250 plant species belonging to the  
395 subgroup of vascular plants have been identified as phototoxic.

396 Table 1. presents the complete inventory of phototoxic plants of the world, listed in alphabetical  
397 order within their taxonomic family. All species were catalogued with their taxonomy, location,  
398 vernacular name and, when known, the nature and structure of their phototoxic molecule(s). In  
399 several phototoxic plants, the molecule(s) responsible for the activity are still unidentified  
400 (indicated as “Unknown” in Table 1.). Phototoxic plants include species which have been  
401 known for a long time for their phototoxicity but also lesser described species that have not  
402 been studied in detail for the characterization of phototoxic molecules and represent a potential  
403 source of new molecules. The presence of phototoxic compounds was reported in a wide  
404 diversity of plant families present in different biomes of the world (Figure 2.). In particular, a  
405 large number of phototoxic species were documented in the families Apiaceae, Asteraceae,  
406 Fabaceae, Rutaceae, and Poaceae (Table 1. and Figure 2.).

Family	Plant name	Vernacular name, local name	Location	Molecule(s) responsible for phototoxicity	Chemical family	References
Acanthaceae	<i>Strobilanthes sp.</i>	Lidah Jin/Pecah biling (Malaysia)	Malaysia	Unknown	Report of flavonoids, alkaloids, triterpenoids, porphyrins and derivatives in the taxonomic family.	[46]. This paper discusses the phototoxicity of Malaysian plant extracts screened in HL60 promyelocytic leukemia cells.
Actinidiaceae	<i>Actinidia chinensis</i> Planch.	Kiwi	East Asia, Europe, New- Zealand, USA	Unknown	Unknown	[47]
Aizoaceae	<i>Tetragonia tetragonoides</i> (Pall.) Kuntze	Tetragone	Littorals, including Brazil	Chlorophylls, pheophorbide <i>a</i> , pheophorbide <i>b</i>	Porphyrins and derivatives	[48]
Amaranthaceae	<i>Alternanthera maritima</i> (Mart.) A.St.-Hil.	Seaside joyweed	East of South America, including	Unknown	Report of Flavonoids, saponins, steroids	[49]

Amaranthaceae	<i>Alternanthera philoxeroides</i> (Mart.) Griseb.	Alligator weed	Brazil, West of Africa			
			USA, South America (littoral),	Rubiadin, rubiadin-1- methyl ether	Anthraquinones	[47,50,51]
			France and East Asia			
Amaranthaceae	<i>Froelichia floridana</i> (Nutt.) Moq.	Florida snakecotton	USA, Australia	3,4-dihydroxyphenyl polyphenolic caffeate	Polyphenols	[52]
Amaranthaceae	<i>Froelichia humboldtiana</i> (Roem. & Schult.) Seub.		Brazil	Naphthodianthrone(s)	Naphthodianthrone(s)	[47,52-54]
Amaranthaceae	<i>Kochia scoparia</i> (L.) Schrad. = <i>Bassia scoparia</i> subsp. scoparia		Europe, USA Asia, Mexico	Unknown	Unknown	[55]
Amaryllidaceae	<i>Cooperia pedunculata</i>	Rain lily	Mexico, USA	Unknown	Unknown	[54]
Amaryllidaceae	<i>Zephyranthes drummondii</i> D.Don	Prairie lily	U.S.A, Mexico	Unknown	Report of Furanocoumarins	[50]
Anacardiaceae	<i>Mangifera indica</i> L.	Mango mangueira	South hemisphere, including Brazil	Unknown	Report of catechols, triterpenoids,	[56]

Anacardiaceae	<i>Pentaspadon motleyi</i> <i>Hook.fil.</i>		Asia Pacific	Pheophorbide <i>a</i> , pheophorbide <i>b</i>	Porphyrins and derivatives	[57]
Anacardiaceae	<i>Rhus succedaneum</i> = <i>Rhus</i> <i>succedanea</i> L. = <i>Toxicodendron</i> <i>succedaneum</i> (L.) Kuntze	Unknown	Asia, Brazil	Unknown	Report of catechols	[58]
Anacardiaceae	<i>Rhus verniciflua</i> Stokes = <i>Toxicodendron</i> <i>vernicifluum</i> (Stokes) F.A.Barkley		China, Korea and Japan	Unknown	Report of catechols	[56]
Anacardiaceae	<i>Schinopsis brasiliensis</i> Engl.	Baraúna	South America, including Brazil	Unknown	Report of Tannins	[59]
Anacardiaceae	<i>Schinus terebinthifolius</i> = <i>Schinus terebinthifolia</i> Raddi	Aroeira	Littorals, including Brazil	Unknown	Report of catechols	[56]
Anacardiaceae	<i>Semecarpus anacardium</i> L.fil.	Oriental cashew	India	Unknown	Report of catechols	[56]
Annonaceae	<i>Annona purpurea</i> Moç. & Sessé ex Dunal		Central America	Berberine	Isoquinoline alkaloid	[47,60]
Annonaceae	<i>Guatteria blepharophylla</i> Mart.	Envireira	South hemisphere,	Berberine, isomoschatoline, lysicamine	Oxaporphine	[61,61]

---

			including		
			Brazil		
Annonaceae	<i>Guatteria ferruginea</i> A.St.-	North	3-hydroxy-	Aporphine alkaloids	[61,62]
	Hil.	of Southern	nornuciferine,		
		America,	isomoschatoline,		
		including	liriodenine,		
		Brazil	lysicamine,		
			nornuciferine, o-		
			methylmoschatoline		
Annonaceae	<i>Guatteria latifolia</i> (Mart.)	Brazil	3-hydroxy-	Aporphine alkaloids	[61–63]
	R.E.Fr.		nornuciferine,		
			isomoschatoline,		
			liriodenine,		
			lysicamine,		
			nornuciferine, o-		
			methylmoschatoline		
Annonaceae	<i>Guatteria sellowiana</i> Schltdl.	Brazil	3-hydroxy-	Aporphine alkaloids	[61,62]
			nornuciferine,		
			isomoschatoline,		
			liriodenine,		
			lysicamine,		
			nornuciferine, o-		
			methylmoschatoline		

---

Apiaceae	<i>Ammi majus L.</i>	Bishop's weed	Southern Europe, Northern and Western Africa, Central Asia, Brazil	3,4-dihydroxyphenyl polyphenolic caffeoate, 7-Methoxycoumarin, 8-Methoxypсорален, alloimperatorin, ammirin, angelicin, archangelicin, bergapten, bergaptol, imperatorin, isobergapten, isopimpinellin, marmesin, marmesinin, methoxsalen, oxypeucedanin, sphondin, xanthotoxin	Furanocoumarins, polyphenols	[38,39,47,50,52,54,64–68]
Apiaceae	<i>Ammi visnaga (L.) Lam</i>	Khella	Argentina, Europe, Mediterranea n basin	7-Methoxycoumarin, 8-hydroxybergapten,, 8-Methoxypсорален, archangelicin, imperatorin, khellin, marmesin, methoxsalen, visnagin, xanthotoxin	Furanocoumarins	[47,50,69–71]

---

Apiaceae	<i>Anethum graveolens</i> L.	Dill	Worldwide, reported in Europe, North America, Brazil	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergapten, bergaptol, isobergapten, methoxsalen, isobergapten, sphondin	Furanocoumarins	[67]
Apiaceae	<i>Angelica archangelica</i> L.	Angelica	Europe + cold climates	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergaptol, isobergapten, isopimpinellin, oreoselone, sphondin, steroid saponins 3- spirostanol	Furanocoumarins	[47,50,54,64–67]

---

Apiaceae	<i>Angelica dahurica</i> (Fisch. ex Hoffm.) Benth. & Hook.fil. ex Franch. & Sav.		East Asia	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, imperatorin, methoxsalen, xanthotoxin	Furanocoumarins	[66]
Apiaceae	<i>Angelica lucida</i> L.	Angelica	North hemisphere	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergapten, isoimperatorin, methoxsalen, xanthotoxin	Furanocoumarins	[66]
Apiaceae	<i>Angelica sinensis</i> (Oliv.) Diels (scientific name) = <i>Angelicae tuhou radix</i> (chinese medicinal herb)	Chinese angelica	China	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, methoxsalen, xanthotoxin	Furanocoumarins	[66]

Apiaceae	<i>Angelica sylvestris L.</i>	Wild angelica	USA, Iceland, Eurasia	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, methoxsalen, xanthotoxin	Furanocoumarins	[67]
Apiaceae	<i>Anthriscus cereifolium</i> (L.) Hoffm.		Europe, USA, New Zealand	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergaptol, isobergapten, methoxsalen, sphondin	Furanocoumarins	[67]
Apiaceae	<i>Anthriscus sylvestris</i> (L.) Hoffm.	Cow parsley (EN)	Africa, New Zealand, Europe, Asia and USA	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergaptol, isobergapten, methoxsalen, sphondin	Furanocoumarins	[67]

Apiaceae	<i>Apium graveolens</i> L.	Celery	Worldwide, including Brazil	4,5',8'- trimethylpsoralen (Trioxsalen), 7- Methoxycoumarin, 8- Methoxysoralen, angelicin, archangelicin, bergapten, bergaptol, isobergapten, isoimperatorin, isopimpinellin, methoxsalen, sphondin, xanthotoxin	Furanocoumarins	[50,66,67]
Apiaceae	<i>Cnidium monnieri</i> (L.) <i>Cusson</i>	Monnier's snowparsley	East Asia	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, methoxsalen	Furanocoumarins	[66]
Apiaceae	<i>Coriandrum sativum</i> L.	Coriander	Worldwide, including Brazil	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, methoxsalen	Furanocoumarins	[66]

Apiaceae	<i>Cymopterus watsonii</i> (J.M.Coult. & Rose) = <i>Aulospermum ibapense</i> (M.E.Jones) J.M.Coult. & Rose	Spring parsley M.E.Jones	North America	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, bergapten, methoxsalen, xanthotoxin	Furanocoumarin	[47,50,54,64–67]
Apiaceae	<i>Daucus carota</i> L.	Wild carrot	North hemisphere, South Africa, Oceania, South America, some occurrences reported in Brazil	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergaptol, isobergapten, methoxsalen, sphondin	Furanocoumarins	[67]
Apiaceae	<i>Foeniculum vulgare</i> Mill.	Fennel	Worldwide, including Brazil	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergapten, bergaptol, isobergapten,	Furanocoumarins	[67]

---

			methoxsalen sphondin		
Apiaceae	<i>Glehnia littoralis</i> (A.Gray) F.Schmidt = <i>Radix glehniae</i> as chinese traditional medicine	USA and East Asia	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, isoimperatorin, methoxsalen	Furanocoumarins	[66]
Apiaceae	<i>Heracleum candidans</i> Wall. ex DC. = <i>Semenovia candidans</i> (Wall. ex DC.)	China	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, bergapten, heraclenin, methoxsalen	Furanocoumarins	[66]
Apiaceae	<i>Heracleum laciniatum</i>	Norway	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergapten, bergaptol, isobergapten, isopimpinellin, methoxsalen, sphondin	Furanocoumarins	[67]

---

---

Apiaceae	<i>Heracleum lanatum</i> Michx.		North hemisphere	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, isobergapten, methoxsalen	Furanocoumarins	[66]
Apiaceae	<i>Heracleum mantegazzianum</i> Sommier & Levier	Giant hogweed	Europe, USA	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergapten, bergaptol, isobergapten, methoxsalen, sphondin	Furanocoumarins	[54,66,67]
Apiaceae	<i>Heracleum sibiricum</i> L.		Eurasia	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, bergapten, methoxsalen	Furanocoumarins	[66]

---

Apiaceae	<i>Heracleum sphondylium</i> L.	Hogweed	North hemisphere, New Zealand	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergapten, bergaptol isobergapten, isopimpinellin, methoxsalen, sphondin	Furanocoumarins	[66,67]
Apiaceae	<i>Heracleum thomsonii</i> C.B.Clarke = <i>Semenovia thomsonii</i> (C.B.Clarke) Manden.		Asia	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, methoxsalen, sphondin	Furanocoumarins	[66]
Apiaceae	<i>Imperatoria ostruthium</i> L. = <i>Peucedanum ostruthium</i> (L.) Koch		Europe	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, isobergapten, methoxsalen, oxypeucedanin, sphondin	Furanocoumarins	[66,67]

---

Apiaceae	<i>Levisticum officinale</i> Koch	Lovage	Europe, USA, Korea, Japan, New Zealand	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergapten, bergaptol, isobergapten, methoxsalen, sphondin	Furanocoumarins	[66,67]
Apiaceae	<i>Pastinaca sativa</i> L.	Wild parsnip	North hemisphere, Australia, New Zealand	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergapten, bergaptol, imperatorin, isobergapten, isopimpinellin, methoxsalen, xanthotoxin, sphondin	Furanocoumarins	[47,50,66,67]
Apiaceae	<i>Petroselinum crispum</i> (Mill.) <i>Fuss</i>	Parsley	Europe, Americas, including	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin,	Furanocoumarins	[50]

---

---

		Brazil, South of Africa, Australia, New Zealand	methoxsalen, oxypeucedanin		
Apiaceae	<i>Petroselinum sativum</i> Hoffm. = <i>Petroselinum crispum</i> subsp. <i>crispum</i>	Europe and Central America	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, archangelicin, bergapten, bergaptol, isobergapten, isopimpinellin, methoxsalen, oxypeucedanin, sphondin, xanthotoxin	Furanocoumarins	[50,66,67]
Apiaceae	<i>Peucedanum alsaticum</i> L. = <i>Xanthoselinum alsaticum</i> (L.) Schur	Europe	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, methoxsalen	Furanocoumarins	[66]

---

Apiaceae	<i>Peucedanum cervaria</i> Linnaeus, 1753	Europe	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, methoxsalen	Furanocoumarins	[66]	
Apiaceae	<i>Peucedanum tauricum</i> M. Bieb Also known as <i>Peucedanum ruthenicum</i> M.Bieb.	Balkans (Eastern Europe)	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, bergapten, bergaptol	Furanocoumarins	[66]	
Apiaceae	<i>Pimpinella magna</i> L.	Europe	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin, bergaptol, isobergapten, methoxsalen, sphondin	Furanocoumarins	[66]	
Apiaceae	<i>Pimpinella major</i> (L.) Huds.	Europe	7-Methoxycoumarin, archangelicin, bergapten, methoxsalen	Furanocoumarins	[66]	
Apiaceae	<i>Pimpinella saxifraga</i> L.	North hemisphere	7-Methoxycoumarin, archangelicin, isopimpinellin, methoxsalen	Furanocoumarins	[66]	
Apiaceae	<i>Sphenosciadium capitellatum</i> A.Gray	Whiteheads, ranger's buttons	West USA	7-Methoxycoumarin, 8-Methoxysoralen, archangelicin,	Furanocoumarins	[54]

	also known as <i>Angelica capitellata</i> (A.Gray) Spalik, Reduron & S.R.Downie			bergapten, methoxsalen		
Araceae	<i>Aglaonema simplex</i> (Blume) Blume	Malayan sword	South East Asia	20 <sup>1-</sup> Hydroxypurpurin-7 delta-lactone ethyl methyl diester, hydroxypurpurin-7- lactone ethyl methyl diester, methyl phytol diester, pheophorbide	Porphyrins and derivatives	[47,72,73]
				<i>a</i>		
Araceae	<i>Dieffenbachia maculata</i> (Lodd.) G.Don = <i>Dieffenbachia seguine</i> (Jacq.) Schott		Tropics, no occurrence reported in	Unknown	Unknown	[74]
Arecaceae	<i>Euterpe oleracea</i> Mart.	Açaí	Brazil and Colombia	Pheophorbide <i>a</i>	Porphyrins and derivatives	[75,76]
Asparagaceae	<i>Agave lecheguilla</i> Torr.		Mexico	Unknown	Report of saponins	[50,52,54]
Asparagaceae	<i>Agave lechuguilla</i> Torr.		Mexico	Unknown	Report of Porphyrins and derivatives	[50]
Asparagaceae	<i>Nolina texana</i> S. Watson	Texas sacahuista	USA, Mexico	Unknown	Unknown	[50,52]

Asphodelaceae	<i>Aloe vera (L.) Burm. f.</i>	Aloe vera	Worldwide, including Brazil	Aloe-emodin, anthraquinone, fagopyrin, emodin, rhein	Naphthodianthrones, Anthraquinones	[47,67,77]
Asteraceae	<i>Artemisia scoparia</i>		Eurasia	Scoparone	Coumarin	[78]
Asteraceae	<i>Brachyglottis adamsii</i> (Cheeseman) B.Nord.		New Zealand	Retorsine, senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Brachyglottis hectori</i> (Buchanan) B.Nord.		New Zealand	Retorsine, senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Brachyglottis huntii</i> (F.Muell.) B.Nord.		New Zealand	Retorsine, senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Brachyglottis kirkii</i> (Hook.fil. ex Kirk) C.J.Webb = <i>Urostemon kirkii</i> (Hook.fil. ex Kirk) B.Nord.		New Zealand	Retorsine, senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Brachyglottis perdicioides</i> (Hook.fil.) Hook.fil.		New Zealand	Retorsine, senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Brachyglottis sciadophila</i> (Raoul) B.Nord.		New Zealand	Retorsine, senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]

Asteraceae	<i>Chrysanthemum segetum</i> L. = <i>Glebionis segetum</i> (L.) <i>Fourr.</i>	Europe	Unknown	Reports of Furanocoumarins see also [80–82]	[54,66] [80,81]
Asteraceae	<i>Cineraria maritima</i> L. = <i>Jacobaea maritima</i> subsp. <i>maritima</i>	Mediterranea n basin	Senecionine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Echinops latifolius</i> Tausch.	Russia, China	Thiophene	Thiophene and derivatives	[47]
Asteraceae	<i>Leuzea carthamoides</i> Willd (= <i>Rhaponticum</i> <i>carthamoides</i> )	Russia	Thiophene	Thiophene and derivatives	[83]
Asteraceae	<i>Parthenium hysterophorus</i> L.	Santa Maria feverfew	USA, Central and South Americas including Brazil, East Africa, Oceania, East Asia, India	Unknown	Unknown
Asteraceae	<i>Pittocaulon bombycophole</i> (Bullock) H.Rob. & Brettell	Mexico	Senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Pittocaulon filare</i> (Mc Vaugh) H.Rob. & Brettell	Mexico	Senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]

Asteraceae	<i>Pittocaulon hintonii</i> H.Rob. & Brettell	Mexico	Senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Pittocaulon praecox</i> (Cav.) H.Rob. & Brettell	Mexico	Senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Pittocaulon velatum</i> (Greenm.) H.Rob. & Brettell	Mexico	Senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Porophyllum obscurum</i> (Spreng.) DC.	South America, including Brazil	Thiophene	Thiophene and derivatives	[41]
Asteraceae	<i>Senecio brasiliensis</i> (Spreng.) Less.	Flor das almas	Brazil	Senecionine	Pyrrolizidine Alkaloids
Asteraceae	<i>Senecio carpathicus</i> Herbich = <i>Jacobaea abrotanifolia</i> subsp. <i>carpathica</i> (Herbich)		East Europe	Senecionine	Pyrrolizidine Alkaloids
		B.Nord. & Greuter			
Asteraceae	<i>Senecio cineraria</i> DC.	Silver ragwort	North hemisphere	Senecionine	Pyrrolizidine Alkaloids
Asteraceae	<i>Senecio douglasii</i> DC.		USA	Retorsine	Pyrrolizidine Alkaloids
Asteraceae	<i>Senecio erucifolius</i> L.		Eurasia	Senecionine	Pyrrolizidine Alkaloids
Asteraceae	<i>Senecio gibbosus</i> subsp. <i>bicolor</i> (Willd.) Peruzzi, N.G.Passal. & Soldano = <i>Jacobaea maritima</i> subsp.		Europe, Mediterranea n basin	Senecionine	Pyrrolizidine Alkaloids

---

		<i>bicolor</i> (Willd.) B.Nord. &				
<i>Greuter</i>						
Asteraceae	<i>Senecio jacalensis</i> Greenm. & Greenm.		Mexico	Senecionine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Senecio jacobaea</i> Brot. ex Willk. & Lange, 1865	Tansy ragwort	North America, Europe, Australia	Retorsine	Pyrrolizidine Alkaloids	[54]
Asteraceae	<i>Senecio othonnae</i> M.Bieb. = <i>Jacobaea othonnae</i> (M.Bieb.) C.A.Mey.		East Europe, Middle East	Senecionine, senkirkine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Senecio paludosus</i> L. = <i>Jacobaea paludosa</i> subsp. paludosa		Europe	Senecionine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Senecio riddellii</i> Torr. & A.Gray		USA	Pyrrolizidine alkaloids, riddelliine, retorsine	Pyrrolizidine Alkaloids	[79,86]
Asteraceae	<i>Senecio rupestris</i> Waldst. & Kit. = <i>Senecio squalidus</i> subsp. rupestris (Waldst. & Kit.)		East Europe	Senecionine	Pyrrolizidine Alkaloids	[79]
<i>Greuter</i>						
Asteraceae	<i>Senecio stabianus</i> Lacaita		Italy	Senecionine	Pyrrolizidine Alkaloids	[79]

---

Asteraceae	<i>Senecio subalpinus</i> Koch = <i>Jacobaea subalpina</i> (W.D.J.Koch) Pelser & Veldkamp	East Europe	Senecionine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Senecio vulgaris</i> L.	North America, Eurasia, West of South America, Australia, New Zealand	Retorsine	Pyrrolizidine Alkaloids	[54]
Asteraceae	<i>Senecio wagneri</i> Degen = <i>Tephroseris papposa</i> subsp. wagneri (Degen) B.Nord.	Bulgaria	Senecionine	Pyrrolizidine Alkaloids	[79]
Asteraceae	<i>Tagetes erecta</i> L.	Worldwide	Thiophene	Thiophene and derivatives	[41]
Bignoniaceae	<i>Oroxylum indicum</i>	Malaysia	Unknown	Report of Chrysin, baicalein, baicalin, flavonoids, iridoids in the Bignoniaceae family. Phototoxicity was demonstrated in leaf extracts.	[46]

Boraginaceae	<i>Anchusa strigosa</i> Banks & Sol.		Arabian peninsula, Middle East, Israel	Retronecine	Pyrrolizidine Alkaloids	[79]
Boraginaceae	<i>Cynoglossum officinale</i> L.	Houndstoungue	North hemisphere	Unknown	Report of Pyrrolizidine alkaloids	[54,64]
Boraginaceae	<i>Echium plantagineum</i> L.	Borage / Paterson's curse (purple viper's- bugloss)	Europe, South Africa, Oceania, South America (some occurrences in Brazil)	Unknown	Report of Pyrrolizidine alkaloids	[38,54]
Brassicaceae	<i>Brassica napus</i> L.	Rapeseed / Canola	Europe, Brazil	Unknown	Unknown	[47]
Brassicaceae	<i>Brassica rapa</i> L.	Turnip rape / mustard	Europe, Brazil	Unknown	Reports of Saponins	[47,50,54]
Bromeliaceae	<i>Ananas comosus</i> (L.) Merr.	Pineapple	Mostly Central and South America, including	Unknown	Unknown	[56]

			Brazil, also reported in Africa and Asia			
Bromeliaceae	<i>Bromelia antiacantha</i> Bertol.		Brazil	Unknown	Unknown	[87]
Celastraceae	<i>Kokoona reflexa</i>	Mata ulat	Malaysia	Unknown	Report of sesquiterpenes and triterpenoids.	[46]
					Phototoxicity was reported in the leaf extract.	
Chrysobalanaceae	<i>Maranthes corymbosa</i>	Merbatu (Malaysia)	Malaysia	Unknown	Report of flavonoids. Phototoxicity was reported in the stem extract.	[46]
Clusiaceae	<i>Calophyllum inophyllum</i>	Bintangor laut (Malaysia)	South hemisphere, including Brazil	Unknown	Report of pyranocoumarins, xanthones, triterpenes, steroids, and flavonoids in the taxonomic family. Phototoxicity was reported in the leaf extract.	[46]
Clusiaceae	<i>Garcinia atroviridis</i> Griff ex T.Anderson	Asam gelugor (Malaysia)	Malaysia	Unknown		[46]

Clusiaceae	<i>Garcinia mangostana</i> L.	Mangosteen	South hemisphere, including Brazil	Mangostin (in pericarp)	Xanthone Flavonoid	[88,89]
Commelinaceae	<i>Rhoeo spathacea</i> (Sw.) Stearn also known as <i>Tradescantia spathacea</i> Sw.		Central America	Rutin	Flavonoid	[90]
Costaceae	<i>Costus speciosus</i> (J.Koenig) Sm. = <i>Hellenia speciosa</i> (J.Koenig) S.R.Dutta	Canereed	Pacific Asia, Central America	Dioscin, diosgenin	Saponins	[91]
Euphorbiaceae	<i>Hevea brasiliensis</i> (Willd. ex A.Juss.) Müll.Arg.	Hevea	South Asia, West Africa, Central and South Americas including Brazil	Unknown	Unknown	[56]
Euphorbiaceae	<i>Ricinus communis</i> L.		Worldwide, including Brazil	<i>Ricinus communis</i> agglutinin (RCA)	Lectin	[92]
Fabaceae	<i>Arachis hypogaea</i> L.	Peanut	Worldwide, including Brazil	<i>Arachis hypogaea</i> Lectin (PNA)	Lectin, Report of Furanocoumarins,	[92]
Fabaceae	<i>Bauhinia forficata</i> Link	Cow's foot	South America,	<i>Bauhinia forficata</i> Lectin (BfL)	Lectin, Report of Furanocoumarins,	[92]

			including			
			Brazil			
Fabaceae	<i>Biserrula pelecinus</i> L.		Mediterranea	Unknown	Reports of	[47]
			n basin		Furanocoumarins	
Fabaceae	<i>Bituminaria bituminosa</i> (L.)	C.H. Stirz.	Europe	Angelicin	Furanocoumarins	[47,50,66]
Fabaceae	<i>Cassia alata</i> (L.) = <i>Senna alata</i> (L.) Roxb.	Gelenggang daun besar (Malaysia)	South Asia, West Africa, Central and South Americas including Brazil, USA	Unknown	Reports of alkaloids, flavonoids, coumarins, anthraquinones in the taxonomic family. Report of phototoxicity in the leaf extract.	[46]
Fabaceae	<i>Chamaecrista nictitans</i> (L.) Moench	Sensitive pea	Central America and South America, including Brazil	Unknown	Reports of	[93]
Fabaceae	<i>Chamaecrista serpens</i> (L.) Greene	Slender sensitive pea	Central America and South America,	Unknown	Reports of	[93]
					Furanocoumarins	

			including			
			Brazil			
Fabaceae	<i>Crotalaria retusa</i> L.	Rattleweed	Tropical belt,	Unknown	Reports of Furanocoumarins,	[36,52,85]
			including		pyrrolizidine	
			Brazil		alkaloids	
Fabaceae	<i>Cullen cinereum</i> (Lindl.)		Australia	Unknown	Reports of Furanocoumarins	[47,50]
	J.W.Grimes					
Fabaceae	<i>Dipteryx odorata</i> (Aubl.)	Cumaru / tonka	North of South America,	Nodakenetin	Furanocoumarins	[66,94]
	Forsyth f.		including			
			Brazil			
Fabaceae	<i>Entada phaseoloides</i> (L.)	Beluru/akar	South Asia,	Unknown	Unknown. Report of phototoxicity in stem extract.	[46]
	Merr.	sintok	Africa,			
		(Malaysia)	Central			
			Americ			
Fabaceae	<i>Enterolobium contortisiliquum</i> (Vell.)	Pacara earpod tree	Brazil	Unknown	Reports of porphyrin and derivatives, Furanocoumarins,	[85,93,95]
	Morong				Saponins	
Fabaceae	<i>Erythrina variegata</i> L.	Tiger's claw	India, Oceania,	Unknown	Reports of Carotenoids,	[96]
			South East		furanocoumarins	
			Asia, Pacific			
			ocean			

---

			islands, Central and South Americas including Brazil			
Fabaceae	<i>Glycine max (L.) Merr.</i>	Soybean	Americas, including Brazil, Eurasia, Africa	Lectin (glycine max)	Lectin, report of Furanocoumarins	[92]
Fabaceae	<i>Griffonia simplicifolia</i> (Vahl ex DC.) Baill.		West Africa	Griffonia simplicifolia Lectin (GSAIA4)	Lectin, Report of Furanocoumarins,	[92]
Fabaceae	<i>Holocalyx balansae</i> Micheli	Alecrim	South America, including Brazil	Unknown	Reports of Furanocoumarins, porphyrins and derivatives	[50]
Fabaceae	<i>Lotus corniculatus</i> L.	Bird's-foot trefoil	Europe, Brazil	Unknown	Reports of Furanocoumarins	[47]
Fabaceae	<i>Medicago denticulata</i> Willd. also known as <i>Medicago</i> <i>polymorpha</i> L.	Burr trefoil	Colombia, Mexico, Europe, USA and China	Unknown	Reports of Furanocoumarins	[54,64]

---

Fabaceae	<i>Medicago lupulina</i> L.	Black medic / Alfafa-lupulina	North hemisphere, Australia, New Zealand, South America including Brazil	Unknown	Reports of Furanocoumarins	[54]
Fabaceae	<i>Medicago nigra</i> L.		Worldwide, including Brazil	Unknown	Reports of Furanocoumarins	[47,54]
Fabaceae	<i>Medicago sativa</i> L.	Alfalfa	Worldwide, including Brazil	Unknown	Reports of Furanocoumarins	[47,54]
Fabaceae	<i>Melilotus officinalis</i> (L.) Lam.	Melilot	North hemisphere	Unknown	Reports of coumarins and furanocoumarins	[33,66]
Fabaceae	<i>Mimosa pteridifolia</i> Benth.	Jurema-cor-de- rosa	Brazil	Flobabenic tannins, pyrogallic tannins	Tannins	[97]
Fabaceae	<i>Mimosa tenuiflora</i> (Willd.) Poir.	Jurema preta, calumbi (Brazil)	Central America and Brazil	Flobabenic tannins, pyrogallic tannins	Tannins Reports of Furanocoumarins and saponins	[97]
Fabaceae	<i>Mimosa verrucosa</i> Benth.	Jurema-branca, jurema-de- oeiras	Brazil	Flobabenic tannins, pyrogallic tannins	Furanocoumarins	[97]

Fabaceae	<i>Peltophorum pterocarpum</i>		South hemisphere	Unknown	Report of phototoxicity in the leaf extract.	[46]
Fabaceae	<i>Psoralea corylifolia</i> L. also known as <i>Cullen corylifolium</i> (L.) Medik.		East Asia	Angelicin, bergaptol, isobergapten, sphondin	Furanocoumarins	[65–68]
Fabaceae	<i>Stryphnodendron fissuratum</i> E.M.O. Martins		Brazil	Unknown	Reports of Furanocoumarins	[98]
Fabaceae	<i>Trifolium hybridum</i> L.	Alsike clover	Europe, North America, Brazil	Unknown	Reports of Furanocoumarins	[54,64]
Fabaceae	<i>Trifolium pratense</i> L.	Red clover	Europe, Brazil	Unknown	Reports of Furanocoumarins	[47]
Fabaceae	<i>Trifolium repens</i> L.	White clover	Europe, Brazil	Unknown	Reports of Furanocoumarins	[47]
Fabaceae	<i>Vicia sativa</i> L.	Lentils	Worldwide, including Brazil	Bergapten	Furanocoumarins	[50,54]
Geraniaceae	<i>Erodium cicutarium</i> (L.) L'Hér.	Alfileria	Eurasia, North America, South of Australia, South Africa, West	Unknown	Report of Anthraquinones	[47]

---

			of South America			
Geraniaceae	<i>Erodium moschatum</i> (L.) L'Hér.	musk stork's-bill	Europe, Oceania, South Africa, Japan, Western USA, South America (not Brazil)	Unknown	Unknown	[47]
Ginkgoaceae	<i>Ginkgo biloba</i> L.	Ginkgo	North hemisphere, also found in Brazil	Unknown	Report of catechols	[56]
Haemodoraceae	<i>Haemodorum distichophyllum</i> (Hook.)		Australia, Tasmania	Phenalenone A	aromatic ketone	[99,100]
Haemodoraceae	<i>Lachnanthes tinctoria</i>	USA Atlantic coast	lachnanthocarpone (Report of tetraoxygenated derivatives in	Phenalenone derivative		[101,102]

---

---

				<i>Wachendorfia</i>		
				<i>thyrsiflora</i> )		
Heliotropiaceae	<i>Heliotropicum europeum</i> L.	European heliotrope	Eurasia, Australia, USA	Heliotrine, lasiocarpine	Pyrrolizidine Alkaloids	[54,79]
Hypericaceae	<i>Hypericum maculatum</i> Crantz		Eurasia	Hypericin	Naphtodianthrone	[50]
Hypericaceae	<i>Hypericum perforatum</i> L.	St John's wort	Worldwide, including Brazil (Mostly North hemisphere)	Anthraquinone, hyperforine, hypericin	Naphtodianthrone	[47,50,52,54,57,64,67,103–106]
Hypericaceae	<i>Hypericum pseudomaculatum</i> Bush	False spotted St. Johnswort	USA	Anthraquinone, hypericin	Anthraquinone, naphtodianthrone	[52,54,64]
Hypericaceae	<i>Hypericum punctatum</i> Lam.	Spotted St. Johnswort	USA	Anthraquinone, hypericin	Anthraquinone, naphtodianthrone	[52,54,64]
Hypericaceae	<i>Hypericum triquetrifolium</i> Turra		Mediterranea n basin	Hypericin	Naphtodianthrone	[50]
Irvingiaceae	<i>Irvingia malayana</i>	Paul kijang (Malaysia)	Malaysia, South-East Asia	Unknown		[46]

---

Lamiaceae	<i>Leonurus sibiricus L.</i>		America, Europe, Asia	Pheophorbide <i>a</i>	Porphyrins and derivatives	[73]
Lamiaceae	<i>Phlomis younghusbandii</i> Mukerjee		China	Unknown	Reports of Furanocoumarins	[66]
Lamiaceae	<i>Scutellaria barbata</i> D. Don	Barbed skullcap	Southern China	Pheophorbide <i>a</i>	Porphyrins and derivatives	[47,106]
Lamiaceae	<i>Tectona grandis</i> L.f.	Teak	Central America, Brazil, Subsaharan Africa, India and Pacific Asia	Unknown	Unknown	[107]
Lamiaceae	<i>Thymus vulgaris</i> L.	Thyme	Europa, Oceania, USA,Central and South Americas, including Brazil	Thymol	Phenol	[108]
Lauraceae	<i>Cinnamomum camphora</i> (L.) Nees & Eberm. = <i>Cinnamomum camphora</i> (L.) J.Presl	Camphor tree	Worldwide, including Brazil	Eucalyptol	Monoterpene	[108]

Lauraceae	<i>Cryptocarya griffithiana</i>	Medang (Malaysia)	Malaysia	Unknown	Report of phototoxicity in leaf extracts.	[46]
Lythraceae	<i>Lawsonia inermis</i> (L.)		South Asia, Africa, Australia, South and Central America	Unknown	Report of phototoxicity in leaf extracts.	[46]
Malvaceae	<i>Hibiscus cannabinus</i> (L.)		South hemisphere, USA, Europe	Unknown	Report of phototoxicity in leaf extracts.	[46]
Malvaceae	<i>Malachra fasciata</i> Jacq. N.V.	Roadside leafbract	Oceania, Central and South America, including Brazil	Unknown	Unknown	[36,47]
Moraceae	<i>Artocarpus integrifolia</i> L.fil. = <i>Artocarpus integer</i> (Thunb.) Merr.		Africa, Europe, South America, including Brazil	Archangelicin, bergapten, citropten, jacalin	Furanocoumarins, coumarins, lectins	[109]

Moraceae	<i>Brosimum gaudichaudii</i> Trécul	South America, Brazil	Archangelicin, bergapten, citropten, marmesin, marmesinin	Furanocoumarins	[63,66,87,110]
Moraceae	<i>Fatoua pilosa</i>	China, Indonesia, New Zealand	Rutaretin	Coumarins	[111]
Moraceae	<i>Ficus carica</i> L.	Common fig Worldwide, including Brazil	Angelisin, archangelicin, bergapten, bergaptol, citropten, isobergapten, isopimpinellin, sphondin. Report of unidentified phototoxic molecules in the Malaysian species <i>Ficus deltoidea</i> [46]	Furanocoumarins	[56,66,67]
Myrtaceae	<i>Eucalyptus globulus</i> Labill.	Eucalyptus Worldwide, including Brazil	Eucalyptol, thymol	Monoterpenes, Phenols	[108]
Myrtaceae	<i>Myrciaria cauliflora</i> (Mart.) O.Berg	Jaboticaba Americas, including Brazil	Anthocyanins, coumarins, flavonoids, phenolic acids,	Coumarins, phenols and flavonoids	[112,113]

				p-Hydroxybenzoic acid, resorcinol		
Myrtaceae	<i>Syzygium aromaticum</i> (L.) Merr. & Perry = <i>Eugenia caryophyllata</i> Thunb.	Clove	Tropical rainforest, including Brazil	Eugenol, thymol	Phenols	[108]
Nartheciaceae	<i>Narthecium ossifragum</i> (L.) Huds.	Asphodel	Europe	Episarsasapogenin, smilagenin, yamogenin	Saponins	[50,91,114]
Papaveraceae	<i>Chelidonium majus</i> L.	Celandine	Europe, Asia and South America, including Brazil	Alkaloidal protoberberine (BBR- F), berberine, chelerythrine, chelidonine, coptisine, jatorrhizine, protopine, sanguinarine, stylopine	Protoberberine Alkaloids	[47,115]
Papaveraceae	<i>Fumaria officinalis</i> L.		North hemisphere, Australia, New Zealand, South America,	Berberine		[68]

			including			
			Brazil			
Pittosporaceae	<i>Pittosporum eugenoides</i> A.Cunn.	Lemonwood	Australia, New Zealand, West of USA	Bergapten	Furanocoumarin	[47]
Pittosporaceae	<i>Pittosporum tenuifolium</i> Gaertn.	Kohuhu	Western Europe, West of USA	Bergapten	Furanocoumarin	[47]
Poaceae	<i>Avena sativa</i> L.	Oat	Europe, USA, Brazil	Unknown	Unknown	[54]
Poaceae	<i>Brachiaria brizantha</i> (A.Rich.) Stapf		Central and South Americas, including Brazil, Africa, Asia Pacific	Protodioscine	Saponins	[36,54,85,114,116,117]
Poaceae	<i>Brachiaria decumbens</i> = <i>Brachiaria eminii</i> (Mez) Robyns		Central and South Americas, including Brazil, Africa	Dichotomin, dioscin, diosgenin, protodioscine, soyasapogenin, soyasapogenol, steroid saponins 3-	Saponins	[36- 38,54,85,91,114,116,117] ]

---

				spirostanols, yamogenin	
Poaceae	<i>Brachiaria humidicola</i> (Rendle) Schweick. = <i>Brachiaria dictyoneura</i> (Fig. & De Not.) Stapf = <i>Urochloa humidicola</i> (Rendle) Morrone & Zuloaga	Central and South Americas, including Brazil, Africa	Protodioscine	Saponins	[36,54,85,114,116,117]
Poaceae	<i>Brachiaria ruziziensis</i> R.Germ & C.M.Evrard	South America, including Brazil, Africa	Unknown	Reports of Saponins	[36,85,117]
Poaceae	<i>Cynodon dactylon</i> (L.) Pers.	Bermuda grass	Worldwide	Unknown	Reports of Porphyrins and derivatives [50]
Poaceae	<i>Echinochloa crus-galli</i> (L.) P.Beauv.	Cockspur	Worldwide, including Brazil	Unknown	Reports of Anthraquinones [47]
Poaceae	<i>Echinochloa esculenta</i> (A.Braun) H.Scholz	Japanese barnyard millet	West Asia, USA, Europe, Oceania	Unknown	Reports of Anthraquinones [47]

---

Poaceae	<i>Echinochloa frumentacea</i>	Indian barnyard millet	Australia, India, Europe, USA, Africa, rare in Brazil, Asia	Unknown	Reports of Anthraquinones	[47]
Poaceae	<i>Lolium perenne L.</i>	Perennial rye grass	Europe, Brazil	Perloline	Alkaloid	[54,64]
Poaceae	<i>Panicum antidotale</i> Retz.	Blue panicum	Australia, USA, some occurrences in Europe	Diosgenin, epismilagenin, tamagenin	Saponins, steroidial sapogenins	[54]
Poaceae	<i>Panicum coloratum</i> L.	Coolah grass	Africa, U.S.A- Mexico border, one occurrence in Brazil	Diosgenin, epismilagenin, tamagenin	Saponins	[54,91]
Poaceae	<i>Panicum dichotomiflorum</i> Michx.	Millet	Americas (including Brazil), Europe, Korea, Japan, New Zealand	Dichotomin, diosgenin, epismilagenin, tamagenin	Saponins	[37,54,91]

Poaceae	<i>Panicum maximum</i> Jacq. = <i>Megathyrsus maximus</i> (Jacq.) B.K.Simon & S.W.L.Jacobs	Guinea grass	Americas, including Brazil, Africa, East Asia, Australia	Diosgenin, epismilagenin, tamagenin	Saponins, steroidial sapogenins	[54]
Poaceae	<i>Panicum miliaceum</i> L.	Millet	North hemisphere, Australia, reports in Brazil	Diosgenin, epismilagenin, tamagenin, yamogenin	Saponins, steroidial sapogenins	[54,114]
Poaceae	<i>Panicum virgatum</i> L.	Switch grass	USA, Europe, East Asia	Diosgenin, epismilagenin, tamagenin	Saponins, steroidial sapogenins	[54]
Poaceae	<i>Phyllostachys bambusoides</i> Siebold & Zucc.	Castillon bamboo	Africa, Asia, North and South America, including Brazil	20 <sup>1</sup> -hydroxypurpurin- 7 delta-lactone ethyl methyl diester, hydroxypurpurin-7- lactone ethyl methyl diester, methyl phytol diester	Porphyrins and derivatives	[47,72]
Polygonaceae	<i>Fagopyrum esculentum</i> Moench = <i>Fagopyrum sagittatum</i> Gilib.	Buckwheat	North hemisphere, Brazil, South of Africa,	Fagopyrin E, fagopyrin F, hypericin	Coumarins, naphthodianthrones	[38,50,52,54,64]

---

			Australia, New Zealand			
Polygonaceae	<i>Fagopyrum tataricum</i> (L.) Gaertn	Tartarian buckwheat	North hemisphere	Fagopyrin E, fagopyrin F, hypericin	Furanocoumarins, naphthodianthrone	[47,50,54,64]
Polygonaceae	<i>Rheum palmatum</i> L.		South-East Oceania, Europe, China, USA	Anthraquinones	Anthraquinones	[47]
Polygonaceae	<i>Rumex hymenosepalus</i> , <i>Rumex</i> sp.		North America, arid and semi-arid regions	Emodine	Anthraquinone	[118,119]
Primulaceae	<i>Myrsine coriacea</i> Merr.		Vietnam	<i>Myrsine coriacea</i> Lectin (McL)	Lectin	[109]
Ranunculaceae	<i>Helleborus niger</i> L.	Christmas rose	Europe, USA	Berberine	Isoquinoline alkaloid	[47,60]
Rhamnaceae	<i>Rhamnus frangula</i>	buckthorn or alder buckthorn	Europe, North Africa, and western Asia	Anthraquinones including phycione (=parietin=rheochrysi	Anthraquinones	[120]

---

---

				din), chrysophanol and emodine	
Rubiaceae	<i>Cinchona spp.</i>	Tropical regions of South America	Quinine	Alkaloid	[121]
Rubiaceae	<i>Galium odoratum</i> (L.) Scop.	Asperula odorata (FR)	North hemisphere	Anthraquinone, berberine, rubiadin, rubiadin-1-methyl ether,	Anthraquinones, coumarins, furanocoumarins [33]
Rubiaceae	<i>Genipa americana</i> L.	Genip	Central and South America, including Brazil	Anthraquinone, berberine, rubiadin, rubiadin-1-methyl ether,	Anthraquinones [87]
Rubiaceae	<i>Heterophyllaea lycioides</i> (Rusby) Sandwith		West of South America	5,5'-bisoranjidiol, 5-chlorosoranjidiol, 7-chlorobisoranjidiol, anthraquinone, berberine lycionine, pustuline, rubiadin,	Anthraquinones [47,50,51,122]

---

---

				rubiadin-1-methyl ether, soranjidiol		
Rubiaceae	<i>Heterophyllaea pustulata</i> Hook. f.	Cegadera	Argentina	2-hydroxy-3-methyl anthraquinone, 5,5'- bisoranjidiol, anthraquinone, berberine, damnacanthal, damnacanthol, heterophyllin, pustuline, rubiadin-1- methyl ether, rubiadin, soranjidiol, soranjidiol-1-methyl ether	Anthraquinones	[47,51,54,122-129]
Rubiaceae	<i>Rubia cordifolia</i> L.		Africa, Asia	Anthraquinone, berberine, rubiadin, rubiadin-1-methyl ether,	Anthraquinones	[68]
Rubiaceae	<i>Rubia yunnanensis</i> Diels		China	Anthraquinone, berberine rubiadin,	Anthraquinones	[130]

---

				rubiadin-1-methyl ether		
Rubiaceae	<i>Sherardia arvensis</i> L.	Blue field- madder	North America, Europe, West coast of South America, Australia, New Zealand	Anthraquinone, berberine, rubiadin, rubiadin-1-methyl ether, rubiadin, soranjidiol	Anthraquinones	[50]
Rutaceae	<i>Atalantia racemosa</i>		India	Rutaretin	Coumarins	[131]
Rutaceae	<i>Boenninghausenia</i> <i>albiflora</i>		Himalaya, China, Japan, Indonesia and South- east Asia	Suberenone	Coumarins	[132]
Rutaceae	<i>Citrus × limon</i> (L.) Burm.fil. = <i>Citrus × bergamia</i> Risso	Lemon	Worldwide, including Brazil	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, anthranilic acid, archangelicin, berberine, bergapten, bergaptol, citropten,	Alkaloids, furanocoumarins	[56,66,67,110,133]

---

				isobergapten, isopimpinellin, methoxsalen, sphondin		
Rutaceae	<i>Citrus limetta</i> Risso = <i>Citrus medica</i> L.	Mousami / sweet lemon	South America, including Brazil	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, anthranilic acid, archangelicin, berberine, bergapten, bergaptol, isobergapten,, methoxsalen, sphondin	Alkaloids, furanocoumarins	[67,133]
Rutaceae	<i>Citrus x aurantifolia</i> (Christm.) Swingle  <i>Citrus hystrix</i>	Lime	South hemisphere, including Brazil	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, anthranilic acid, archangelicin, berberine, bergapten, bergaptol, citropten, isobergapten, isopimpinellin, methoxsalen, sphondin, xanthotoxin	Alkaloids, furanocoumarins	[46,56,66,67,110,133]

---

Rutaceae	<i>Citrus maximus</i> = <i>Citrus grandis</i>		Americas, Africa, western Europe, India, China, South-east Asia, Australia and Oceania	Suberenone	Coumarin	[134]
Rutaceae	<i>Citrus x paradisi</i> Macfad.	Grapefruit / Pomelo	Americas, including Brazil, East Asia, Europe	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, anthranilic acid, archangelicin, berberine, bergapten, bergaptol, citropten, isobergapten, isopimpinellin, methoxsalen, sphondin	Alkaloids, furanocoumarins	[67,84,133]
Rutaceae	<i>Citrus x sinensis</i> (Mill.) Pers., 1806 = <i>Citrus × aurantium</i> L.	Sweet orange / blood orange	Western Africa, Western Europe, Central America and South	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, anthranilic acid, archangelicin, berberine, bergapten, bergaptol, citropten, isobergapten,	Alkaloids, furanocoumarins	[67,110,133]

Rutaceae	<i>Clausena anisata</i> (Willd.) Hook.fil., De Wild. & Staner	Horsewood	America, including Brazil	isopimpinellin, methoxsalen, sphondin		
Rutaceae	<i>Dictamnus albus</i> L.	Burning bush	Central and South of Africa, Sino-Indian coastline	7-Methoxycoumarin, anthranilic acid, archangelicin, bergapten, citropten, imperatorin, methoxsalen, oxypeucedanin	Alkaloids, furanocoumarins	[66,133]
Rutaceae	<i>Glycosmis pentaphylla</i>	orangeberry, gin berry	Europe, USA, Asia	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, anthranilic acid, archangelicin, berberine, bergapten, bergaptol, citropten, isobergapten, methoxsalen, sphondin	Alkaloids, furanocoumarins	[67,133]
Rutaceae	<i>Micromelum minutum</i>	Cemumar hitam (Malaysia)	South-east Asia, North of Australia	Glycoborinine, p-coumaroyltyramine	Carbazole alkaloids (betacarbolines)	[135]
Rutaceae			Malaysia, Indonesia, Australia	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, anthranilic acid, archangelicin,	Alkaloids, furanocoumarins	[46]

---

				berberine, bergapten, bergaptol, citropten, isobergapten, isopimpinellin, methoxsalen, sphondin, xanthotoxin		
Rutaceae	<i>Pelea anisata</i> H. Mann. = <i>Melicope anisata</i> (H.Mann) T.G.Hartley & B.C.Stone	Mokihana	Hawaii	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, anthranilic acid, archangelicin, berberine, bergapten, bergaptol, citropten, isobergapten, methoxsalen, sphondin	Alkaloids, furanocoumarins	[67,133]
Rutaceae	<i>Ruta angustifolia</i> Pers.		Europe	7-Methoxycoumarin, anthranilic acid, archangelicin, bergapten, citropten, methoxsalen, rutamarin	Alkaloids, furanocoumarins	[136]
Rutaceae	<i>Ruta chalepensis</i> L.	Fringed rue	Europe, Central America, South America	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, anthranilic acid, archangelicin,	Alkaloids, furanocoumarins, coumarins	[67,133,136]

---

---

			America, a few in Brazil	berberine, bergapten, bergaptol, citropten, isobergapten, methoxsalen, sphondin, rutacultin		
Rutaceae	<i>Ruta graveolens</i> L.	Rue, common rue, herb-of-grace	Eurasia, South America, including Brazil	7-Methoxycoumarin, 8-Methoxysoralen, angelicin, anthranilic acid, archangelicin, berberine, bergapten, bergaptol, chalepensin, citropten, isobergapten, isopimpinellin, isoimperatorin,isorutarin, marmesin, methoxsalen, rutamarin, sphondin, xanthotoxin, gravelliferone, rutacultin, suberenone, rutaretin	Alkaloids, furanocoumarins, coumarins	[50,66,67,74,133,136-139]
Rutaceae	<i>Ruta montana</i> (L.) L.		Europe, Mediterranea n basin	7-Methoxycoumarin, anthranilic acid, archangelicin,	Alkaloids, furanocoumarins	[136]

---

---

				bergapten, citropten, methoxsalen, sphondin		
Rutaceae	<i>Ruta pinnata L.fil.</i>		Spain	7-Methoxycoumarin, anthranilic acid, archangelicin, bergapten, citropten, methoxsalen	Alkaloids, furanocoumarins	[66]
Rutaceae	<i>Thamnosma montana</i> Torr. & Frem	Mojave desert rue	West of USA	7-Methoxycoumarin, 8-Methoxysoralen, anthranilic acid, archangelicin, bergapten, citropten, isoimperatorin, isopimpinellin, methoxsalen, xanthotoxin	Alkaloids, furanocoumarins	[50,133]
Rutaceae	<i>Thamnosma texana</i> (Gray) Torr.	Dutchman's breeche	USA: Arizona (Hualapai Mountains) to Texas and N. Mexico.	7-Methoxycoumarin, 8-Methoxysoralen, alloimperatorin, alloimperatorin methyl ether epoxide, anthranilic acid, archangelicin, berberine, bergapten, citropten, heraclenin,	Alkaloids, furanocoumarins	[47,50,52,54,64,66,133,1 40]

---

---

					herniarin, imperatorin, isopimpinellin, methoxsalen, osthol, oxypeucedanin, thamnosmin, xanthotoxin	
Sapindaceae	<i>Talisia esculenta</i> (A.St.-Hil.) Radlk.	Pitomba	Brazil	Unknown	Unknown	[87]
Scrophulariaceae	<i>Myoporum laetum</i> G. Forst. And <i>Myoporum tenuifolium</i>	New Zealand manatoka	American and European coastlines, Brazil, New Zealand	Ngaione	Furanosesquiterpenoi ds	[38,50]
Smilacaceae	<i>Smilax ornata</i> Lem.		Central America, brazil	Smilagenin	Saponin	[141]
Solanaceae	<i>Lycopersicon esculentum</i> Mill. = <i>Solanum lycopersicum</i> L.	Tomato	Worldwide	Bergapten	Furanocoumarin	[50]
Solanaceae	<i>Solanum betaceum</i> Cav.	Tamarillo	Worldwide	Unknown	Reports of Anthocyanes, carotenoids, chlorophylls	[142]


---

Solanaceae	<i>Solanum melongena</i> L.	Eggplant	Worldwide, including Brazil	Nasunin	Anthocyanes	[143]
Strophariaceae	<i>Agrocybe aegerita</i> (V.Brig.) Singer = <i>Cyclocybe aegerita</i> (V.Brig.) Vizzini		USA, Europe	<i>Agrocybe aegerita</i> Lectin	Lectin	[109]
Verbenaceae	<i>Lantana camara</i> L.	Common lantana	Worldwide	Lantadenes	Lantadenes triterpenoids	[36,38,50,52,54]
Verbenaceae	<i>Peronema canescens</i>	Sungkai (Malaysia)	Malaysia	Unknown	Report of phototoxic molecules in leafs.	[46]
Verbenaceae	<i>Premna odorata</i>	Bebuas (Malaysia)	South-east Asia	Unknown	Report of phototoxic molecules in leafs.	[46]
Verbenaceae	<i>Verbena officinalis</i> L.	Vervain	Worldwide	Unknown	Report of Porphyrins and derivatives	[50]
Viscaceae	<i>Viscum album</i> L.	Mistletoe	Eurasia, North America	Mistletoe ( <i>Viscum</i> <i>album</i> ) Lectin (ML)	Lectin	[109]
Ximeniaceae	<i>Ximenia americana</i> L.	Tallow wood	South hemisphere, including Brazil	Riproximin	Lectin	[109]

Zingiberaceae	<i>Curcuma longa</i> L.	Curcuma	Tropical belt, including Brazil	Curcumin	Polyphenol	[41,47,144–147]
Zygophyllaceae	<i>Tribulus terrestris</i> L.	Tribulus	Worldwide	Diosgenin, Epi- sarsasapogenin	Saponins	[50,52,91,114]

407

408 Table 1. Phototoxic plants of the world, listed in alphabetical order of their taxonomic family.  
 409 All species were catalogued with their taxonomy, location, vernacular name and, when known,  
 410 the nature and structure of their phototoxic molecule(s). In several phototoxic plants, the  
 411 molecule(s) responsible for the activity are still unidentified (Unknown).



412


413 Figure 2. Number and distribution of phototoxic plant species in the plant taxonomic families.



414

415 7.2. Inventory of phototoxic molecules


416 In a second step, a detailed search of phototoxic molecules was performed, providing the  
417 complete updated list of phototoxic phytochemicals synthesized by terrestrial plants (Table 2.).



418 For each molecule, we reviewed all papers discussing its phototoxic activity, chemical  
419 structure, chemical family, presence in different plant families or species, location in the  
420 different plant tissues and organs, biogeographical location of the producer species in the  
421 world, as well as a list of relevant references evidencing the phototoxic activity of the  
422 compound in *in vitro* or *in vivo* models. (Table 2.). For well-studied molecules, singlet-oxygen  
423 yields can be found in a previous report [148].



Chemical Family and Photosensitization type	Molecule	Chemical structure	Producer plant(s)	Biogeographical zone of the plant(s)	Plant tissue location	Biological activity and photophysical characteristics	References
Type I Photosensitization							
Alkaloid Type I	(+)-Chelidone		<i>Chelidonium majus</i> L. (Celandine) - (Papaveraceae)	Europe, Asia and South America, including Brazil	Sap, stem and bark	Pro-apoptotic in uterine cancer cells.	[115]
						$\lambda_{\text{max}} \sim 278 \text{ nm}$ depending on solvent	
						.	



Alkaloid Type I	Coptisine (C <sub>19</sub> H <sub>14</sub> NO <sub>4</sub> +)		<i>Chelidonium majus</i> L. (Celandine) - (Papaveraceae)	Europe, Asia and South America, including Brazil	Sap, stem and bark	Same comment as (+)- Chelidonine  $\lambda_{\text{max}} \sim 325$ nm depending on solvent	[115]
Alkaloid Type I	Jatrorrhizine (C <sub>20</sub> H <sub>20</sub> NO <sub>4</sub> +)		<i>Chelidonium majus</i> L. (Celandine) - (Papaveraceae)	Europe, Asia and South America, including Brazil	Sap, stem, bark	Same comment as (+)- Chelidonine.  <i>Chelidonium</i> <i>majus</i> L. is rich in numerous biologically active compounds, like protoberberine	[115]

alkaloids (e.g.,  
berberine,  
coptisine,  
protopine,  
stylopine,  
jatrorrhizine),  
flavonoids,  
phenolic acids,  
saponins and  
defense-related  
proteins

Alkaloid	Protopine		<i>Chelidonium majus L.</i> (Celandine) - (Papaveraceae)	Europe, Asia and South America, including Brazil	Sap, stem, bark	Same comment as (+)- Chelidonine	[115]
Type I		$(C_{20}H_{19}NO_5)$				$\lambda_{max} \sim 308$ nm depending on solvent	.

Alkaloid	Quinine		a) <i>Cinchona spp.</i> (Rubiaceae)	a) tropical regions of South America	Bark	$\lambda_{\text{max}} \sim 345 \text{ nm}$ depending on solvent	[121]
Type I	(C <sub>20</sub> H <sub>24</sub> N <sub>2</sub> O <sub>2</sub> )						
Alkaloid	Sanguinarine		<i>Chelidonium majus</i> L. (Celandine) - (Papaveraceae)	Europe, Asia and South America, including Brazil	Sap, stem, bark	Same comment as (+)- Chelidonine	[115]
Type I	(C <sub>20</sub> H <sub>14</sub> NO <sub>4</sub> +)					$\lambda_{\text{max}} \sim 290-310$ nm depending on solvent	

Alkaloid Type I	Stylopine (C <sub>19</sub> H <sub>17</sub> NO <sub>4</sub> )		<i>Chelidonium majus</i> L. (Celandine) - (Papaveraceae)	Europe, Asia and South America, including Brazil	Sap, stem, bark	Same comment as (+)- Chelidonine.	[115]
Anthraquinone Type I	2-Hydroxy-3 methyl anthraquinone (C <sub>15</sub> H <sub>10</sub> O <sub>4</sub> )		<i>Heterophyllaea pustulata</i> Hook. f. (Cegadera) - (Rubiaceae)	Argentina (Jujuy, Salta and Tucuman provinces)	Leaves and stems	Indicated in breast and colorectal cancer phototherapy.	[47]  $\lambda_{\text{max}} \sim 430-440$ nm depending on solvent

Anthraquinone Type I	5,5'-bisoranjidiol (C <sub>30</sub> H <sub>18</sub> O <sub>8</sub> )		a) <i>Heterophyllaea pustulata</i> Hook. f. (Cegadera) - (Rubiaceae) <sup>[122,125-127]</sup> b) <i>Heterophyllaea lycioides</i> (Rusby) Sandwith - (Rubiaceae) <sup>[122]</sup>	a) Argentina (Jujuy, Salta and Tucuman provinces) b) West of South America	Antibacterial phototoxicity on <i>S.aureus</i> [122,125-127]
Anthraquinone Type I	5-Chloro-soranjidiol (C <sub>15</sub> H <sub>9</sub> ClO <sub>4</sub> )		<i>Heterophyllaea lycioides</i> (Rusby) Sandwith - (Rubiaceae)	West of South America	Aerial parts Anthraquinone derivatives similar to those found in <i>H.pustulata</i> , but <i>H.lycioides</i> contains rare chlorinated derivatives such as 7-chlorobisoranji diol and 5-

						chloro-
						soranjidiol
Anthraquinone Type I	7-Chloro bis soranjidiol (C <sub>30</sub> H <sub>17</sub> ClO <sub>8</sub> )		<i>Heterophyllaea lycioides</i> (Rusby) Sandwith - (Rubiaceae) <sup>[122]</sup>	West of South America	Aerial parts	Same anthraquinones derivatives as found in <i>H. pustulata</i> .
Anthraquinone Type I	Aloe-emodin (C <sub>15</sub> H <sub>10</sub> O <sub>5</sub> )		<i>Aloe vera</i> (L.) Burm. f. ( <i>Aloe vera</i> ) - (Asphodelaceae) <sup>[47]</sup>	Worldwide, including Brazil	Roots and Leaves	Used as a photosensitizer  [47] Some non-PS natural products may potentiate phototoxicity including indole-3-acetic acid, carvacrol, β-glucans, and

ascorbic acid.

[106]


$\lambda_{\text{max}} \sim 430$  nm  
depending on  
solvent

Anthraquinone

Type I

Chrysophanol

(C<sub>15</sub>H<sub>10</sub>O<sub>4</sub>)





*Rhamnus frangula* [120]



(Rhamnaceae)




Europe, North  
Africa, and western  
Asia



$\lambda_{\text{max}} \sim 430-440$   
nm depending  
on solvent


[120]


Anthraquinone	Damnacanthal		<i>Heterophyllaea pustulata</i> Hook. f. (Cegadera) - (Rubiaceae)	Argentina	Phototoxic on [50,51,123,125–127] <i>S.aureus</i>
Type I					$\lambda_{\text{max}} \sim 530 \text{ nm}$ depending on solvent
Anthraquinone	Damnacanthol		<i>Heterophyllaea pustulata</i> Hook. f. (Cegadera) - (Rubiaceae)	Argentina	[50,51,123,125–127]
Type I					


Anthraquinone	Emodin = Archin		a) <i>Aloe vera</i> (L.) Burm. f. ( <i>Aloe vera</i> ) - (Asphodelaceae) <sup>[67]</sup>	a) Worldwide, including Brazil	b) Latex	$\lambda_{\text{max}} \sim 430 \text{ nm}$ depending on solvent	[67]
Type I		(C <sub>15</sub> H <sub>10</sub> O <sub>5</sub> )	b) <i>Rheum spp.</i> (Rhubarb) - (Polygonaceae) <sup>[67]</sup>	c) <i>Rhamnus spp.</i> (Buckthorns) (Rhamnaceae) <sup>[67]</sup>	d) <i>Rumex sp.</i> (Polygonaceae) <sup>[118,149]</sup>		
Anthraquinone	Heterophyllin		<i>Heterophyllaea pustulata</i> Hook. f.(Cegadera) - (Rubiaceae) <sup>[123,125,126]</sup>	Argentina			[123,125,126]
Type I		(C <sub>22</sub> H <sub>26</sub> N <sub>2</sub> O <sub>4</sub> )					

Anthraquinone Type I	Lycionine ( $C_{30}H_{18}O_9$ )		<i>Heterophyllaea lycioides</i> (Rusby) Sandwith - (Rubiaceae) [122]	West of South America	Aerial parts	See comment for 5-chloro-soranjidiol	[122]
Anthraquinone Type I	Physcion=Physcione= parietin=rheochrysidin ( $C_{11}H_{12}O_5$ )		<i>Rhamnus frangula</i> [120]	Europe, North Africa, and western Asia	$\lambda_{max}\sim 365$ nm depending on solvent		[120]
Anthraquinone Type I	Pustuline ( $C_{16}H_{12}O_4$ )		a) <i>Heterophyllaea pustulata</i> Hook. f. (Cegadera) - (Rubiaceae) [51,122,125,126]  b) <i>Heterophyllaea lycioides</i> (Rusby) Sandwith - (Rubiaceae) [51,122]	a) Argentina  b) West of South America			[51,122,125,126]

Anthraquinone	Rhein		<i>Rheum spp.</i> - (Polygonaceae) [67] (Rhubarb) - (Polygonaceae) [67]	a) Worldwide, including Brazil	a) Latex	$\lambda_{\text{max}} \sim 430-450$ nm depending on solvent	[67]
Type I			a) <i>Aloe vera</i> (L.) Burm. f. ( <i>Aloe vera</i> ) - (Asphodelaceae) [67]				
Anthraquinone	Rubiadin		(Rubiaceae) [50] a) <i>Heterophyllaea pustulata</i> Hook. f. (Cegadera) - (Rubiaceae)	a) Argentina	a) stem, leaves	Antibacterial	[47,50,51,123–129]
Type I				b) USA, South America (coastal), France and East Asia		The phototoxic anthraquinones soranjidiol (red) and rubiadin (yellow) have absorption maxima at the	
			b) <i>Alternanthera philoxeroides</i> (Mart.) Griseb. (alligator weed) - (Amaranthaceae) [47,50]	c) North America, Europe, West coast of South America,			


			Australia, New Zealand	interface of the UV-C and UV-B (~280 nm)
		<i>Galium spp.</i> - (Rubiaceae) [50]		and at 410–420 nm (visible light) [50]
		<i>Coprosma spp.</i> - (Rubiaceae) [50]		
		c) <i>Sherardia arvensis</i> - (Rubiaceae) [50]		
Anthraquinone Type I	Rubiadin-1-methyl ether (C <sub>16</sub> H <sub>12</sub> O <sub>4</sub> )		(Rubiaceae) [50] a) <i>Heterophyllaea pustulata</i> Hook. f. (Cegadera) - (Rubiaceae) [47,50,51,123–129] b) USA, South America (coastal), France and East Asia b) <i>Alternanthera philoxeroides</i> (Mart.) Griseb. (alligator weed) - (Amaranthaceae) [47,50] c) North America, Europe, West coast of South America,	a) Argentina a) Stem, leaves Phototoxicity demonstrated in human breast cancer cells [47,50]  λ <sub>max</sub> ~ 530-540 nm depending on solvent

		<i>Galium spp.</i> - (Rubiaceae) <sup>[50]</sup>	Australia, New Zealand
		<i>Coprosma spp.</i> - (Rubiaceae) <sup>[50]</sup>	
		c) <i>Sherardia arvensis</i> - (Rubiaceae)	
		[50]	
Anthraquinone Type I	Soranjidiol (C <sub>15</sub> H <sub>10</sub> O <sub>4</sub> )	 <p><i>Heterophyllaea spp.</i>- (Rubiaceae)<sup>[50]</sup></p> <p>a) <i>Heterophyllaea lycioides</i> (Rusby) Sandwith - (Rubiaceae)<sup>[50]</sup></p> <p>b) <i>Heterophyllaea pustulata</i> Hook. f. (Cegadera) - (Rubiaceae)<sup>[50,51,122–129]</sup></p> <p><i>Galium spp.</i> - (Rubiaceae)<sup>[50]</sup></p> <p><i>Coprosma spp.</i> - (Rubiaceae)<sup>[50]</sup></p> <p>c) <i>Sherardia arvensis</i> L. - (Rubiaceae)<sup>[50]</sup></p>	<p>a) West of South America</p> <p>b) Argentina</p> <p>c) North America, Europe, West coast of South America, Australia, New Zealand</p>
			Stem, leaves
			Antibacterial
			[50,51,122–129]
			Rubiadin is identified as a promising photosensitizer


Anthraquinone Type I	Soranjidiol-1-methyl ether (C <sub>16</sub> H <sub>12</sub> O <sub>4</sub> )		<i>Heterophyllaea pustulata</i> Hook. f.(Cegadera) - (Rubiaceae) [50,51,122– 129]	Argentina	High antiproliferativ e activity <i>in</i> <i>vitro</i> in human breast cancer cells	[50,51,122–129]
					Phototoxic on HSV-1 herpes virus	

Anthraquinone  
derivatives =  
naphthodianthrones  
=

Hypericin ( $C_{30}H_{16}O_8$ )  
and  
Pseudohypericin  
( $C_{30}H_{16}O_9$ )



Perylenequinones  
Type I



*Hypericum spp.* - (Hypericaceae) [67]

a) *Hypericum perforatum* L. (St John's wort) - (Hypericaceae)

[41,47,50,54,64,67,103,105,146]

b) *Hypericum triquetrifolium* Turra - (Hypericaceae) [50]

c) *Hypericum maculatum* Crantz - (Hypericaceae) [50]

d) *Hypericum pseudomaculatum* Bush (False spotted St. John's wort) - (Hypericaceae) [54,64]

e) *Hypericum punctatum* Lam. - (Hypericaceae) [54,54]

f) *Fagopyrum esculentum* Moench - (Polygonaceae) [50,54]

g) *Fagopyrum sagittatum* - (Polygonaceae) [54]

a) Pistils, fruits, stems  
Hypericin has [41,47,50,54,64,67,10  
absorption 3–106,129,146,150]  
maxima at 236  
and 592 nm [50]

[54]

causes severe  
phototoxicity in  
grazing animals  
exposed to  
visible light  
(500 – 600  
nm).

a) Worldwide,  
including Brazil  
(Mostly North  
hemisphere)

b) Mediterranean  
basin

c) Eurasia

d) USA

e) USA

f) North hemisphere,  
Brazil, South of  
Africa, Australia,  
New Zealand


g) Balkans, Japan,  
Korea, USA



h) North hemisphere




h) *Fagopyrum tataricum* (L.) Gaertn


(Tartary buckwheat) -

(Polygonaceae) [47,50,54]



Aromatic ketone (fused benzene and naphthalene ring)	Phenalenone (C <sub>13</sub> H <sub>8</sub> O)		<i>Haemodorum distichophyllum</i> (Hook.)	Australia, Tasmania	High production yield of singlet oxygen	[99,100]
Type I					λ <sub>max</sub> ~ 240-260 nm and 330-430 nm depending on solvent	




Benzo naphthyridinone alkaloid Type I	Perloline  (C <sub>20</sub> H <sub>17</sub> N <sub>2</sub> O <sub>3</sub> +) 	<i>Lolium perenne</i> L. (Perennial rye grass) - (Poaceae)	Europe, Brazil	[54]	[54,64]
Benzophenanthridin e Alkaloid Type I	Chelerythrine  (C <sub>21</sub> H <sub>18</sub> NO <sub>4</sub> +) 	<i>Chelidonium majus</i> L. (Celandine) - (Papaveraceae)	Europe, Asia and South America, including Brazil	Sap, stem and bark	Pro-apoptotic in uterine cancer cells.  $\lambda_{\text{max}} \sim 342 \text{ nm}$ depending on solvent
		Also reported in <i>Zanthoxylum clava-herculis</i> and <i>Zanthoxylum rhoifolium</i>			[115]

Carbazole alkaloid Type I	Glycoborinine (C <sub>18</sub> H <sub>17</sub> NO <sub>2</sub> )		<i>Glycosmis pentaphylla</i> (Rutaceae)	South-east Asia, North of Australia	[135]
Coumarin Type I	7-Methoxycoumarin = herniarin (C <sub>10</sub> H <sub>8</sub> O <sub>3</sub> )		(Umbelliferae) <sup>[110]</sup> (Apiaceae) <sup>[110]</sup> (Rutaceae) <sup>[110]</sup>	[110] $\lambda_{\text{max}} \sim$ 330-340 nm depending on solvent	[110,140]
			<i>Thamnosma texana</i> (Gray) Torr. (Dutchman's breeche) - (Rutaceae)	[140]	
Coumarin Type I	Ammirin		<i>Ammi majus</i> L. (Bishop's weed) - (Apiaceae) <sup>[47,50]</sup>	Southern Europe, Northern and	Highly phototoxic <sup>[47,50]</sup>



	(C <sub>14</sub> H <sub>12</sub> O <sub>3</sub> )	Western Africa, Central Asia, Brazil	λmax: 248, 260 and 302 nm depending on solvent
Coumarin Type I	Citropten = 5,7- dimethoxycoumarin = Limettin (C <sub>11</sub> H <sub>10</sub> O <sub>4</sub> )	 <p>(Rutaceae)<sup>[56]</sup></p> <p>a) <i>Citrus x sinensis</i> (Mill.) Pers., 1806 = <i>Citrus x aurantium</i> L. (Orange) - (Rutaceae)<sup>[110]</sup></p> <p>b) <i>Citrus x aurantifolia</i> (Christm.) Swingle (Lime) - (Rutaceae)<sup>[56]</sup></p> <p>c) <i>Citrus × limon</i> (L.) Burm.fil. = <i>Citrus × bergamia</i> Risso (Lemon) - (Rutaceae)<sup>[56]</sup></p> <p>d) <i>Citrus x paradisi</i> Macfad. (Grapefruit / Pomelo) - (Rutaceae)</p> <p>[56]</p> <p>(Moraceae)<sup>[56]</sup></p>	/
		<p>a) Western Africa, Western Europe, Central America and South America, including Brazil</p> <p>b) South hemisphere, including Brazil</p> <p>c) Worldwide, including Brazil</p> <p>d) Americas, including Brazil, East Asia, Europe</p>	Same comment as for 7- methoxycouma rin
			λmax~ 360-365 nm depending on solvent




e) *Ficus carica* L. (Common fig) -  
 (Moraceae)<sup>[56]</sup> e) Worldwide,  
 including Brazil

Coumarin Type I	Coumarin  (C <sub>19</sub> H <sub>18</sub> O <sub>5</sub> )		<i>Myrciaria cauliflora</i> (Mart.) O.Berg (Jaboticaba) - (Myrtaceae) <sup>[112]</sup>	Americas, including Brazil	Fruits	λ <sub>max</sub> ~ 315-340 nm depending on solvent	<sup>[112]</sup>
Coumarin	Gravelliferone (and methyl ester)  (C <sub>19</sub> H <sub>22</sub> O <sub>3</sub> )		<i>Ruta graveolens</i> L. (Rua) - (Rutaceae)	Eurasia, South America, including Brazil			<sup>[151]</sup>

Coumarin	Rutacultin (C <sub>16</sub> H <sub>18</sub> O <sub>4</sub> )		<i>Ruta graveolens</i> L. (Rua) - (Rutaceae), <i>Ruta chaleensis</i> (Rutaceae)	Eurasia, South America, including Brazil	Roots	[151]
				<i>Helietta apiculata</i> (Rutaceae)		
Coumarin Type I	Scoparone (=6,7-Dimethoxycoumarin) (C <sub>11</sub> H <sub>10</sub> O <sub>4</sub> )		<i>Artemisia scopariae</i> (virgate wormwood, capillary wormwood or redstem wormwood, yīn chén in Mandarin Chinese)	Eurasia	$\lambda_{\text{max}} \sim 300-330$ nm depending on solvent	[152]
Coumarin	Suberenone (C <sub>14</sub> H <sub>12</sub> O <sub>4</sub> )		<i>Ruta graveolens</i> L. (Rua) (Rutaceae) also present in <i>Citrus maxima</i> (Burm.) Merr. = <i>Citrus grandis</i> (L.) Osbeck	Eurasia, South America, including Brazil	Several minor related coumarins have been reported in the roots of <i>Ruta pinnata</i>	[153]
				Central and South America, including Brazil, India,		[154]. These

		Himalaya, China and South-east Asia	include
	and <i>Boenninghausenia albiflora</i> (Rutaceae)		Gravelliferone
		Himalaya, China, Japan, Indonesia and South-east Asia	methyl ether, xanthyletin, europinnarin, pinnarin, benahorin, limettin, luvangetin, bergapten and sabandinin.

Coumarin Type I	Thamnosmin (C <sub>15</sub> H <sub>14</sub> O <sub>4</sub> )		<i>Thamnosma texana</i> (Gray) Torr. (Dutchman's breeche) - (Rutaceae) [140]	Arizona (Hualapai Mountains) to Texas and N. Mexico.	λmax~ 300-330 nm depending on solvent	[140]
Coumarin Type I	Umbelliferone = 7- hydroxycoumarin (C <sub>9</sub> H <sub>6</sub> O <sub>3</sub> )		(Apiaceae) [66]		λmax~ 300-330 nm depending on solvent	[66]


Cyclic terpene ketone	Hyperforin  Type I  (C <sub>35</sub> H <sub>52</sub> O <sub>4</sub> )		<i>Hypericum perforatum</i> L. (St John's wort) - (Hypericaceae) <sup>[155]</sup>	Worldwide, including Brazil (Mostly North hemisphere)	Pistils, fruits, stems	Plant defense molecule	[155]
Furanochromone  Type I	Visnagin  (C <sub>13</sub> H <sub>10</sub> O <sub>4</sub> )		<i>Ammi visnaga</i> (L.) Lam (Khella) - (Apiaceae) <sup>[69,70]</sup>	Argentina, Europe, Mediterranean basin	Seeds	λ <sub>max</sub> ~ 315-334 nm depending on solvent	[69,70]
Furanochromone (=Furochromone)  Type I	Khellin  (C <sub>14</sub> H <sub>12</sub> O <sub>5</sub> )		<i>Ammi visnaga</i> (L.) Lam (Khella) - (Apiaceae) <sup>[69,71]</sup>	Argentina, Europe, Mediterranean basin	Seeds	Antifungal, antibacterial and antiviral activity.  Khellin could be useful as a PS against psoriasis and	[69–71]

vitiligo.  $\lambda_{\text{max}}$ ~  
250-and 338  
nm in alcohol

Furanocoumarin  
(=Furocoumarin  
= Psoralen)

4,5',8'-  
Trimethylpsoralen =  
Trioxsalen

Type I  
(C<sub>14</sub>H<sub>12</sub>O<sub>3</sub>)



*Apium graveolens L.* (celery)  
(infected by *Sclerotinia sclerotiorum*  
(Lib.) de Bary ("pink rot")) -  
(Apiaceae) [50]



Worldwide,  
including Brazil

The species  
contains other  
phototoxic  
molecules:  
Xanthotoxin,  
bergapten, 8-  
hydroxybergapt  
en,  
isopimpinellin,  
isoimperatorin,  
rutaretin

[50]

$\lambda_{\text{max}}$ ~360 nm  
depending on  
solvent

Furanocoumarin Type I	8-hydroxybergapten (C <sub>12</sub> H <sub>8</sub> O <sub>5</sub> )		a) <i>Ammi visnaga</i> (L.) Lam - (Apiaceae) <sup>[47,50]</sup>  (This molecule is also produced by b) <i>Sclerotinia sclerotiorum</i> (Lib.) de Bary ("pink rot") - (Sclerotiniaceae)	a) Argentina, Europe, Mediterranean basin  b) Europe, USA, South of Africa, East of Asia  [50]	Highly phototoxic  $\lambda_{max}$ ~ 350-360 nm depending on solvent	[47,50]
Furanocoumarin Type I	8-Methoxysoralen (8- MOP) = Methoxsalen = Xanthotoxin  (C <sub>12</sub> H <sub>8</sub> O <sub>4</sub> )		(Apiaceae) <sup>[110]</sup>  (Rutaceae) <sup>[110]</sup>		Phototoxic on the plant- pathogenic fungus <i>Colletotrichum</i> <i>acutatum</i> and the ascomycete <i>Aspergillus</i> <i>nidulans</i>	[110]


Furanocoumarin Type I	Alloimperatorin (=prangenidin)		a) <i>Ammi majus</i> L. (Bishop's weed) - (Apiaceae) <sup>[47,50]</sup>  b) <i>Thamnosma texana</i> (Gray) Torr. (Dutchman's breeche) - (Rutaceae) <sup>[140]</sup>	a) Southern Europe, Northern and Western Africa, Central Asia, Brazil  b) Texas, Mexico	Highly phototoxic	[47,50,140]
Furanocoumarin Type I	Alloimperatorin methyl ether epoxide		<i>Thamnosma texana</i> (Gray) Torr. (Dutchman's breeche, blisterweed) - (Rutaceae) <sup>[140]</sup>	Texas, Mexico	$\lambda_{\text{max}} \sim 290-310$ nm depending on solvent	[140]

Furanocoumarin

Type I

Angelinin

(C<sub>11</sub>H<sub>6</sub>O<sub>3</sub>)



Present with other furanocoumarins  
in

a) *Ammi majus* L. (Bishop's weed) -  
(Apiaceae) [66,67]

b) *Anethum graveolens* L. (Dill) -  
(Apiaceae) [67]

*Angelica spp* - (Apiaceae) [66]

c) *Angelica archangelica* L.  
(Angélique) - (Apiaceae) [67]

d) *Anthriscus sylvestris* (L.) Hoffm.  
(Cow parsley) - (Apiaceae) [67]

e) *Anthriscus cereifolium* (L.) Hoffm.  
- (Apiaceae) [67]

f) *Apium graveolens* L. (Celery) -  
(Apiaceae) [66,67]

g) *Daucus carota* L. (Wild carrot) -  
(Apiaceae) [67]

a) Southern Europe,

Northern and  
Western Africa,  
Central Asia, Brazil

b) Worldwide,  
mostly Europe and  
North America,  
found in Brazil

c) Europe + cold  
climates, very rare  
cases in Brazil

d) Africa, New  
Zealand, mostly in  
Europe, Asia and  
USA

e) Europe, USA,  
New Zealand

f) Worldwide,  
including Brazil

The content in  
furanocoumarin  
s may vary  
widely  
according to  
geographic  
location and  
climate

λ<sub>max</sub>~ 340-350  
nm depending  
on solvent

[66,67]



Americas, including

Brazil

p) *Psoralea corylifolia* L. -

(Fabaceae) [67]

n) Europe and

Central America

q) *Bituminaria bituminosa* (L.) C.H.

Stirt. - (Fabaceae) [66]

(only 1 occurrence in

Brazil, does not seem

endemic)

r) *Ficus carica* L. (Common fig) -

(Moraceae) [66,67]

o) Europe

p) Asia, 1 occurrence

s) *Citrus x aurantifolia* (Christm.) in North America, 1

Swingle (Citron vert) - (Rutaceae) [67] in Africa, 1 in Arabic

peninsula

t) *Citrus × limon* (L.) Burm.fil. =

*Citrus × bergamia* Risso (Lemon) - q) Europe

(Rutaceae) [67]

u) *Citrus limetta* Risso = *Citrus*

*medica* L. (Mousami) - (Rutaceae)

[67]

r) Worldwide

,including Brazil

- v) *Citrus x paradisi* Macfad.  
 (Grapefruit / Pomelo) - (Rutaceae) [67]
- w) *Citrus x sinensis* (Mill.) Pers.,  
 1806 = *Citrus × aurantium* L.  
 (Orange) - (Rutaceae) [67]
- x) *Dictamnus albus* L. (Burning  
 bush) - (Rutaceae) [67]
- y) *Pelea anisata* H. Mann. =  
*Melicope anisata* (H.Mann)  
 T.G.Hartley & B.C.Stone -  
 (Rutaceae) [67]
- z) *Ruta chalepensis* L. (Fringed rue) -  
 (Rutaceae) [67]
- aa) *Ruta graveolens* L. (Rue / Rua) -  
 (Rutaceae) [67]
- s) South hemisphere,  
 including Brazil
- t) Benin
- u) South America,  
 including Brazil
- v) Benin
- w) Western Africa,  
 Western Europe,  
 Central America and  
 South America,  
 including Brazil
- x) Europe, USA,  
 Asia
- y) Hawaii
- z) Europe, Central  
 America, South  
 America, a few in  
 Brazil


aa) South America,  
including Brazil,  
Eurasia

Furanocoumarin

Archangelicin

Type I

(C<sub>24</sub>H<sub>26</sub>O<sub>7</sub>)




(Apiaceae)<sup>[66]</sup>

(Rutaceae)<sup>[66]</sup>

(Fabaceae)<sup>[66]</sup>

(Moraceae)<sup>[66]</sup>

[66]

Furanocoumarin	Bergapten		a) <i>Ammi majus</i> L. (Bishop's weed) - (Apiaceae) <sup>[38,47,50,66]</sup>	a) Southern Europe, Northern and Western Africa, Central Asia, Brazil	/	Highly phototoxic in the ultraviolet A (UVA) range, 320–400 nm	[38,47,50,56,63,66,13 9,140]
Type I			b) <i>Pastinaca sativa</i> L. (Wild parsnip) - (Apiaceae) <sup>[47,50,66]</sup>	b) Eurasia, Oceania, North and South Americas, including Brazil			
(5-methoxysoralen)			c) <i>Cymopterus watsonii</i> (J.M.Coult. & Rose) M.E.Jones (Spring parsley) - (Apiaceae) <sup>[47,50]</sup>	c) North America			
(C <sub>12</sub> H <sub>8</sub> O <sub>4</sub> )			d) <i>Heracleum laciniatum</i> - (Apiaceae) <sup>[50,66]</sup>	d)			
			e) <i>Heracleum sphondylium</i> L. (Hogweed) - (Apiaceae) <sup>[50,66]</sup>	e) North hemisphere, New Zealand			
			f) <i>Foeniculum vulgare</i> Mill. (Fennel) - (Apiaceae) <sup>[50]</sup>	f) Worldwide, including Brazil			
			g) <i>Anethum graveolens</i> L. (Dill) - (Apiaceae) <sup>[50]</sup>	g) Worldwide, mostly Europe and North America, found in Brazil			
			Angelica spp - (Apiaceae) <sup>[50,66]</sup>				

- |   |   |
|---|---|
| <p>h) <i>Petroselinum sativum</i> Hoffm.=<br/> <i>Petroselinum crispum</i> subsp. <i>crispum</i><br/>           - (Apiaceae) [50,66]</p> <p>i) <i>Pimpinella major</i> (L.) Huds. -<br/>           (Apiaceae) [66]</p> <p>j) <i>Levisticum officinale</i> Koch<br/>           (Lovage) - (Apiaceae) [66]</p> <p>k) <i>Apium graveolens</i> L. (Celery) -<br/>           (Apiaceae) [66]</p> <p>l) <i>Peucedanum tauricum</i> M. Bieb<br/>           (Also known as <i>Peucedanum ruthenicum</i> M.Bieb.) - (Apiaceae) [66]<br/>           (Rutaceae) [56]</p> <p>m) <i>Ruta graveolens</i> L. (Rue / Rua) -<br/>           (Rutaceae) [47,56,138]</p> | <p>h) Europe and<br/>           Central America<br/>           (only 1 occurrence in<br/>           Brazil, does not seem<br/>           endemic)</p> <p>i) Europe</p> <p>j) Europe, USA,<br/>           Korea, Japan, New<br/>           Zealand</p> <p>k) Worldwide,<br/>           including Brazil</p> <p>l) Balkans (Eastern<br/>           Europe)</p> <p>m) Eurasia, South<br/>           America, including<br/>           Brazil</p> <p>n) Texas, Mexico</p> |
|---|---|

- n) *Thamnosma texana* (Gray) Torr. (Dutchman's breeche) - (Rutaceae) [47,50,140]
- o) *Citrus spp.* (e.g. limes, bergamot orange, grapefruit, Earl Grey tea) - (Rutaceae) [50,66]
- p) *Citrus × limon* (L.) Burm.fil. = *Citrus × bergamia* Risso (Lemon) - (Rutaceae) [50,56,66]
- q) *Citrus x paradisi* Macfad. - (Grapefruit / Pomelo) (Rutaceae) [50,56,66]
- r) *Thamnosma montana* Torr. & Frem - (Mohave desertrue) (Rutaceae) [50]
- s) *Citrus x aurantifolia* (Christm.) Swingle (Lime) - (Rutaceae) [56]
- Various *Pittosporum* spp. - (Pittosporaceae) [50]
- o) Worldwide, including Brazil
- p) Worldwide, including Brazil
- q) Americas, including Brazil, East
- Asia, Europe
- r) West of USA
- s) South hemisphere, including Brazil
- t) Australia, New Zealand, West of USA
- u) Western Europe, West of USA
- v) Worldwide, including Brazil

t) *Pittosporum eugenoides* A.Cunn. - w) Worldwide,  
(Pittosporaceae) [50] including Brazil

u) *Pittosporum tenuifolium* Gaertn. - x) Europe, USA,  
(Pittosporaceae) [50] South of Africa, East  
of Asia

y) Brazil  
v) *Vicia sativa* L. - (Fabaceae) [50]  
w) The cultivated tomato : z) Worldwide  
*Lycopersicon esculentum* Mill - ,including Brazil  
(Solanaceae) [50]

x) *Sclerotinia sclerotiorum* (Lib.) de  
Bary (“pink rot”) - (Sclerotiniaceae)  
[50]

(Moraceae) [56]

y) *Brosimum gaudichaudii* Trécul  
(mamacadela or inharé) - (Moraceae)  
[63]

z) *Ficus carica* L. (Common fig) -  
(Moraceae) [56]

Furanocoumarin	Bergaptol		Reports of various furanocoumarins that may include bergaptol in:	a) Southern Europe, Northern and Western Africa, Central Asia, Brazil	Involved in photodermatitis cases in humans.	[66,67]
Type I						
	(C <sub>11</sub> H <sub>6</sub> O <sub>4</sub> )					
			a) <i>Ammi majus</i> L. (Bishop's weed) - (Apiaceae) <sup>[67]</sup>	b) Worldwide, mostly Europe and North America, found in Brazil		
			b) <i>Anethum graveolens</i> L. (Dill) - (Apiaceae) <sup>[67]</sup>			
			c) <i>Angelica archangelica</i> L. (Angélique) - (Apiaceae) <sup>[67]</sup>	c) Europe + cold climates, very rare cases in Brazil		
			d) <i>Anthriscus sylvestris</i> (L.) Hoffm. (Cow parsley) - (Apiaceae) <sup>[67]</sup>	d) Africa, New Zealand, mostly in		
			e) <i>Anthriscus cereifolium</i> (L.) Hoffm. - (Apiaceae) <sup>[67]</sup>	e) Europe, Asia and USA		
			f) <i>Apium graveolens</i> L. (Celery) - (Apiaceae) <sup>[67]</sup>	e) Europe, USA, New Zealand		
			g) <i>Daucus carota</i> L. (Wild carrot) - (Apiaceae) <sup>[67]</sup>	f) Worldwide, including Brazil		

- h) *Foeniculum vulgare* Mill. (Fennel) g) North hemisphere,  
- (Apiaceae) [67] South Africa,  
Oceania, South America, some  
occurrences found in

i) *Heracleum laciniatum* auct. scand.,  
non Hornem.- (Apiaceae) [67] Brazil

j) *Heracleum mantegazzianum*

Sommier & Levier (Giant hogweed)  
- (Apiaceae) [67] h) Worldwide,  
including Brazil

k) *Heracleum sphondylium* L.  
(Hogweed) - (Apiaceae) [67] i) Norway

l) *Levisticum officinale* Koch  
(Lovage) - (Apiaceae) [67] j) Europe, North  
America, New Zealand

m) *Pastinaca sativa* L. (Wild  
parsnip) - (Apiaceae) [67] k) North hemisphere,  
New Zealand

n) *Petroselinum sativum* Hoffm.=  
*Petroselinum crispum* subsp. *crispum*  
- (Apiaceae) [67] l) Europe, USA,  
Korea, Japan, New Zealand

o) *Peucedanum ostruthium* Linnaeus,  
1753 (Masterwort) - (Apiaceae) [67] m) Eurasia, Oceania,  
North and South

- p) *Peucedanum tauricum* M. Bieb Americas, including  
     Also known as *Peucedanum ruthenicum* M.Bieb. - (Apiaceae) [67] Brazil
- q) *Psoralea corylifolia* L. - Central America  
     (Fabaceae) [67] n) Europe and  
     o) Europe
- r) *Ficus carica* L. (Common fig) - p) Balkans (Eastern  
     (Moraceae) [67] Europe)
- s) *Citrus x aurantifolia* (Christm.)  
     Swingle (Lime) - (Rutaceae) [67]
- t) *Citrus × limon* (L.) Burm.fil. = q) East Asia  
     *Citrus × bergamia* Risso (Lemon) -  
     (Rutaceae) [67]
- u) *Citrus limetta* Risso = *Citrus medica* L. (Mousami) - (Rutaceae) r) Worldwide,  
     including Brazil
- v) *Citrus x paradisi* Macfad. s) South hemisphere,  
     (Grapefruit / Pomelo) - (Rutaceae) including Brazil
- [67]

- w) *Citrus x sinensis* (Mill.) Pers.,  
1806 = *Citrus × aurantium* L.  
(Orange) - (Rutaceae) [67]

x) *Dictamnus albus* L. (Burning  
bush) - (Rutaceae) [67]

y) *Pelea anisata* H. Mann. =  
*Melicope anisata* (H.Mann)  
T.G.Hartley & B.C.Stone -  
(Rutaceae) [67]

z) *Ruta chalepensis* L. (Fringed rue) - Central America and  
(Rutaceae) [67]

aa) *Ruta graveolens* L. (Rue / Rua) -  
(Rutaceae) [67]

t) Worldwide,  
including Brazil

u) South America,  
including Brazil

v) Americas,

including Brazil, East  
Asia, Europe

w) Western Africa,  
Western Europe,

x) Europe, USA,  
Asia

y) Hawaii


z) Europe, Central  
America, South  
America, a few in  
Brazil

Furanocoumarin

Type I

Chalepensin

(C<sub>16</sub>H<sub>14</sub>O<sub>3</sub>)



*Ruta graveolens* L. (Rue / Rua)

(Rutaceae) and *Clausena sp.*,  
(Rutaceae) [47,50]

Eurasia, South

America, including  
Brazil

Highly  
phototoxic

[47,50,156]

λmax~ 325-360  
nm depending  
on solvent


aa) Eurasia, South  
America, including  
Brazil

Furanocoumarin

Heraclenin

Type I

(C<sub>16</sub>H<sub>14</sub>O<sub>5</sub>)



a) *Thamnosma texana* (Gray) Torr.

(Dutchman's breeche) - (Rutaceae)

[140]

a) Texas, Mexico

b) China

λmax~ 260 nm

depending on  
solvent

[66,140]


Furanocoumarin

Herniarin

Type I

= (7-

Methoxycoumarin)



*Thamnosma texana* (Gray) Torr.

(Dutchman's breeche) - (Rutaceae)

[140]

Texas, Mexico


λmax~ 358 nm

depending on  
solvent

[140]

Same comment  
as Heraclenin

(C<sub>10</sub>H<sub>8</sub>O<sub>3</sub>)


Furanocoumarin	Imperatorin		a) <i>Ammi majus</i> L. - (Apiaceae) [38,47,50]	a) Southern Europe, Northern and Western Africa, Central Asia, Brazil	$\lambda_{\text{max}} \sim 255, [38,50,66,67,140,157]$
Type I			b) <i>Ammi visnaga</i> (L.) Lam - (Apiaceae) [50,157]		268 and 286 nm depending on solvent
	(C <sub>16</sub> H <sub>14</sub> O <sub>4</sub> )		c) <i>Pastinaca sativa</i> L. - (Apiaceae) [50,157]	b) Argentina, Europe, Mediterranean basin	Highly phototoxic
			d) <i>Thamnosma texana</i> (Gray) Torr. - (Apiaceae) [140]	c) Eurasia, Oceania, North and South Americas, including Brazil	Same comment as Heraclenin.
			e) <i>Angelica dahurica</i> (Fisch. ex Hoffm.) Benth. & Hook.fil. ex Franch. & Sav. - (Apiaceae) [66]	d) Arizona (Hualapai Mountains) to Texas and N. Mexico.	
			f) <i>Clausena anisata</i> (Willd.) Hook.fil., De Wild. & Staner - (Rutaceae) [66]	e) East Asia	

Furanocoumarin

Isobergapten

Type I

(C<sub>12</sub>H<sub>8</sub>O<sub>4</sub>)



Reports of various furanocoumarins  
that may include isobergapten in :

a) *Ammi majus* L. (Bishop's weed) -  
(Apiaceae) [67]

b) *Anethum graveolens* L. (Dill) -  
(Apiaceae) [67]

c) *Angelica archangelica* L.  
(Angélique) - (Apiaceae) [67]

d) *Anthriscus sylvestris* (L.) Hoffm.  
(Cow parsley) - (Apiaceae) [67]

e) *Anthriscus cereifolium* (L.) Hoffm.  
- (Apiaceae) [67]

f) Central and South  
of Africa, Sino-  
Indian coastline

a) Southern Europe,  
Northern and  
Western Africa,  
Central Asia, Brazil

b) Worldwide,  
mostly Europe and  
North America,  
found in Brazil

c) Europe + cold  
climates, very rare  
cases in Brazil

d) Africa, New  
Zealand, mostly in

USA

λmax~ 340-350  
nm depending  
on solvent

[66,67]

Responsible for  
numerous cases  
of  
photodermatitis  
in humans.




- n) *Petroselinum sativum* Hoffm.= l) Europe, USA,  
*Petroselinum crispum* subsp. *crispum* Korea, Japan, New  
- (Apiaceae) [67] Zealand
- o) *Peucedanum ostruthium* Linnaeus, m) Eurasia, Oceania,  
1753 (Masterwort) - (Apiaceae) [67] North and South  
Americas, including
- p) *Pimpinella magna* L. - (Apiaceae) Brazil  
[67]
- q) *Heracleum lanatum* - (Apiaceae) n) Europe and  
[66] Central America
- o) Europe
- p) Europe
- q) North hemisphere
- r) *Psoralea corylifolia* L. -  
(Fabaceae) [67]
- r) East Asia
- s) *Ficus carica* L. (Common fig) -  
(Moraceae) [66,67]  
s) Worldwide,  
including Brazil

- t) *Citrus x aurantifolia* (Christm.)  
 Swingle (Lime) - (Rutaceae) [67]  
 t) South hemisphere,  
 including Brazil
- u) *Citrus × limon* (L.) Burm.fil. =  
*Citrus × bergamia* Risso (Lemon) -  
 (Rutaceae) [67]  
 u) Worldwide,  
 including Brazil
- v) *Citrus limetta* Risso = *Citrus medica* L. (Mousami) - (Rutaceae)  
 [67]  
 v) South America,  
 including Brazil
- w) *Citrus x paradisi* Macfad.  
 (Grapefruit / Pomelo) - (Rutaceae)  
 [67]  
 w) Americas,  
 including Brazil, East  
 Asia, Europe
- x) *Citrus x sinensis* (Mill.) Pers.,  
 1806 = *Citrus × aurantium* L.  
 (Orange) - (Rutaceae) [67]  
 x) Western Africa,  
 Western Europe,  
 Central America and  
 South America,  
 including Brazil
- y) *Dictamnus albus* L. (Burning  
 bush) - (Rutaceae) [67]  
 y) Europe, USA,
- e) *Pelea anisata* H. Mann. =  
*Melicope anisata* (H.Mann)  
 z) Hawaii

T.G.Hartley & B.C.Stone - aa) Europe, Central  
(Rutaceae) [67] America, South  
America, a few in  
aa) *Ruta chalepensis* L. (Fringed rue) Brazil  
- (Rutaceae) [67]  
ab) *Ruta graveolens* L. (Rue / Rua) - ab) Eurasia, South  
(Rutaceae) [67] America, including  
Brazil

Furanocoumarin	Isoimperatorin Type I (C <sub>16</sub> H <sub>14</sub> O <sub>4</sub> )		a) <i>Thamnosma montana</i> Torr. & Frem (Mohave desertree) - (Rutaceae) <sup>[50]</sup>  b) <i>Ruta graveolens</i> L. (Rue / Rua) - (Rutaceae) <sup>[50,157]</sup>  c) <i>Apium graveolens</i> L. (celery) (infected by <i>Sclerotinia sclerotiorum</i> (Lib.) de Bary ("pink rot")) - (Apiaceae) <sup>[50]</sup>  d) <i>Glehnia littoralis</i> (A.Gray) F.Schmidt = <i>Radix glehniae</i> - (Apiaceae) <sup>[66]</sup>  e) <i>Angelica lucida</i> L. (Angelica) - (Apiaceae) <sup>[66]</sup>	a) West of USA  b) Eurasia, South America, including Brazil  c) Worldwide, including Brazil  d) USA and East of Asia  e) North hemisphere	Highly phototoxic  λmax~ 330-340 nm depending on solvent	[50,66,157]
----------------	---	--	--	---	--	-------------

Furanocoumarin	Isopimpinellin		(Rutaceae) <sup>[56]</sup>	a) Eurasia, South America, including Brazil	Highly phototoxic, responsible for cattle phototoxicity, particularly in sheeps.
Type I			a) <i>Ruta graveolens</i> L. (Rue / Rua) - (Rutaceae) <sup>[47,50]</sup>	b) Arizona (Hualapai Mountains) to Texas and N. Mexico.	$\lambda_{\text{max}} \sim 330-340$ nm depending on solvent
	(C <sub>13</sub> H <sub>10</sub> O <sub>5</sub> )		b) <i>Thamnosma texana</i> (Gray) Torr. (Dutchman's breeche) - (Rutaceae) <sup>[47,50,140]</sup>	c) West of USA	
			c) <i>Thamnosma montana</i> Torr. & Frem (Mohave desertrue) - (Rutaceae) <sup>[50]</sup>	d) Worldwide, including Brazil	
			d) <i>Citrus × limon</i> (L.) Burm.fil. = <i>Citrus × bergamia</i> Risso (Lemon) - (Rutaceae) <sup>[56,66]</sup>	e) Western Africa, Western Europe, Central America and South America, including Brazil	Phototoxic on the plant-pathogenic fungus
			e) <i>Citrus x sinensis</i> (Mill.) Pers., 1806 = <i>Citrus × aurantium</i> L. (Orange) - (Rutaceae) <sup>[110]</sup>	f) South hemisphere, including Brazil	<i>Colletotrichum acutatum</i> and the ascomycete
			f) <i>Citrus x aurantifolia</i> (Christm.) Swingle (Lime) - (Rutaceae) <sup>[56]</sup>		

- g) *Citrus x paradisi* Macfad.  
 (Grapefruit / Pomelo) - (Rutaceae) including Brazil, East  
 [56] Americas,  
 Asia, Europe
- Aspergillus  
*nidulans* <sup>[110]</sup>
- h) *Pastinaca sativa* L. (Wild parsnip)  
 - (Apiaceae) <sup>[50,66,157]</sup>  
 h) Eurasia, Oceania,
- i) *Apium graveolens* L. (Celery) -  
 (Apiaceae) <sup>[50,66]</sup> North and South  
 Americas, including  
 Brazil
- j) *Petroselinum sativum* Hoffm. =  
*Petroselinum crispum* subsp. *crispum* i) Worldwide,  
 - (Apiaceae) <sup>[66]</sup> including Brazil
- k) *Heracleum sphondylium* L. j) Europe and Central  
 (Hogweed) - (Apiaceae) <sup>[66]</sup> America
- l) *Ammi majus* L. (Bishop's weed) - k) North hemisphere,  
 (Apiaceae) <sup>[38,66]</sup> New Zealand
- m) *Angelica archangelica* L. l) Southern Europe,  
 (Angelica) - (Apiaceae) <sup>[66]</sup> Northern and  
 Western Africa,
- n) *Heracleum laciniatum* - Central Asia, Brazil  
 (Apiaceae) <sup>[66]</sup>

		o) <i>Pimpinella saxifraga</i> L. - (Apiaceae) [66]	m) Europe + cold climates	
		p) <i>Sclerotinia sclerotiorum</i> (Lib.) de Bary (Pink rot) - (Sclerotiniaceae) [50]	n) Norway o) North hemisphere	
		(Moraceae) [56]		
		q) <i>Ficus carica</i> L. (Common fig) - (Moraceae) (de Menezes <i>et al.</i> , 2014)	p) Europe, USA, South of Africa, East of Asia	
			q) Worldwide, including Brazil	
Furanocoumarin Type I	Marmesin = nodakenetin (C <sub>14</sub> H <sub>14</sub> O <sub>4</sub> )		a) <i>Ammi majus</i> L. (Bishop's weed) - (Apiaceae) [38,47,50]  b) <i>Ammi visnaga</i> (L.) Lam (Khella) - (Apiaceae) [47,71]	Highly phototoxic, responsible for phototoxicity in cattle
			a) Southern Europe, Northern and Western Africa, Central Asia, Brazil  b) Argentina, Europe, Mediterranean basin	[38,47,50,71,110]

c) *Ruta graveolens* L. (Rue / Rua) -

(Rutaceae)<sup>[47,50]</sup>

c) Eurasia, South America, including

Brazil

d) *Brosimum gaudichaudii* Trécul -

(Moraceae)<sup>[110]</sup>

d) South America,

Brazil

$\lambda_{\text{max}}$ ~ 300-320


nm depending

on solvent

Furanocoumarin Marmesinin = Ammijin

Type I

(C<sub>20</sub>H<sub>24</sub>O<sub>9</sub>)



a) *Ammi majus* L. (Bishop's weed) -  
(Apiaceae)<sup>[38,47,50]</sup>

a) Southern Europe,  
Northern and  
Western Africa,  
Central Asia, Brazil

Highly [38,47,50,110]

phototoxic,  
responsible for

phototoxic  
events in cattle

d) *Brosimum gaudichaudii* Trécul -




(Moraceae)<sup>[110]</sup>

b) South America,  
Brazil

$\lambda_{\text{max}}$ ~ 330-340


nm depending

on solvent

Furanocoumarin Type I	Nodakenetin (C <sub>14</sub> H <sub>14</sub> O <sub>4</sub> )		<i>Dipteryx odorata</i> Willd, ( <i>Coumarouna odorata</i> Aubl.) - (Fabaceae) [66]	North of South America, including Brazil	λmax~ 330-340 nm depending on solvent	[66]
Furanocoumarin Type I	Osthol (C <sub>15</sub> H <sub>16</sub> O <sub>3</sub> )		<i>Thamnosma texana</i> (Gray) Torr. (Dutchman's breeches) - (Rutaceae) [140]	Arizona (Hualapai Mountains) to Texas and N. Mexico.	λmax~ 310-320 nm depending on solvent	[140]
Furanocoumarin Type I	Oxypeucedanin = (4-[(3,3-Dimethyloxiran-2-		a) <i>Ammi majus</i> L. (Bishop's weed) - (Apiaceae) [47,50]  b) <i>Petroselinum crispum</i> (Mill.) Fuss (Parsley) - (Apiaceae) [50]	a) Southern Europe, Northern and Western Africa, Central Asia, Brazil  a) Seeds, fruits	Highly phototoxic [140]  λmax~ 310-320 nm depending on solvent	[47,50,66,140]

yl)methoxy]furo[3,2-g]chromen-7-one)

(C<sub>16</sub>H<sub>14</sub>O<sub>5</sub>)



(-)-Oxypeucedanin

c) *Peucedanum ostruthium* Linnaeus, 1753 (Masterwort) - (Apiaceae)<sup>[66]</sup>

b) Worldwide, including Brazil

d) *Thamnosma texana* (Gray) Torr. (Dutchman's breeche) - (Rutaceae)

c) Europe

[140]

e) *Clausena anisata* (Willd.)

Hook.fil., De Wild. & Staner (Horsewood) - (Rutaceae)<sup>[66]</sup>

d) Arizona (Hualapai Mountains) to Texas and N. Mexico.


e) Central and South of Africa, Sino-Indian coastline

Furanocoumarin

Type I

Rutamarin

(C<sub>21</sub>H<sub>24</sub>O<sub>5</sub>)



a) *Ruta graveolens* L. (Rue / Rua) - (Rutaceae)

a) Eurasia, South America, including Brazil

b) *Ruta angustifolia* Pers. - (Rutaceae)

b) Europe

$\lambda_{\text{max}}$  ~ 310-320 nm depending on solvent

Phototoxic,  
antibacterial,  
antifungal,  
antiviral,  
antiparasitic,


<sup>[136]</sup>

insecticidal and  
larvicidal

Furanocoumarin

Rutaretin

C<sub>14</sub>H<sub>14</sub>O<sub>5</sub>



(a) *Ruta graveolens* L. (Rua)  
(Rutaceae)

(a) Eurasia, South  
America, including  
Brazil

[151]

also present in (b) *Atalantia  
racemosa* (Rutaceae),  
(c) *Fatoua pilosa* (Moraceae)

(b) India

(c) China, Indonesia,  
New-Zealand

[131]


[111]

Furanocoumarin

Sphondin

Type I

(C<sub>12</sub>H<sub>8</sub>O<sub>4</sub>)



Present with other furanocoumarins

in:

a) Southern Europe,  
Northern and  
Western Africa,  
Central Asia, Brazil

a) *Ammi majus* L. (Bishop's weed) -  
(Apiaceae)<sup>[67]</sup>

b) *Anethum graveolens* L. (Dill) -  
(Apiaceae)<sup>[67]</sup>

c) *Angelica archangelica* L.  
(Angélique) - (Apiaceae)<sup>[67]</sup>

d) *Anthriscus sylvestris* (L.) Hoffm.  
(Cow parsley) - (Apiaceae)<sup>[67]</sup>

e) *Anthriscus cereifolium* (L.) Hoffm.  
- (Apiaceae)<sup>[67]</sup>

f) *Apium graveolens* L. (Celery) -  
(Apiaceae)<sup>[67]</sup>

g) *Daucus carota* L. (Wild carrot) -  
(Apiaceae)<sup>[67]</sup>

a) Southern Europe,

Northern and  
Western Africa,

Central Asia, Brazil

b) Worldwide,  
mostly Europe and

North America,  
found in Brazil

c) Europe + cold  
climates, very rare  
cases in Brazil

d) Africa, New  
Zealand, mostly in

Europe, Asia and  
USA

e) Europe, USA,  
New Zealand

f) Worldwide,  
including Brazil

$\lambda_{\text{max}} \sim 310-320$

nm depending  
on solvent

<sup>[66,67]</sup>

- h) *Foeniculum vulgare* Mill. (Fennel) g) North hemisphere,  
- (Apiaceae) [67] South Africa,  
Oceania, South America, some  
occurrences found in

i) *Heracleum laciniatum* auct. scand.,  
non Hornem.- (Apiaceae) [67] Brazil

j) *Heracleum mantegazzianum*

Sommier & Levier (Giant hogweed)  
- (Apiaceae) [67] h) Worldwide,  
including Brazil

k) *Heracleum sphondylium* L.  
(Hogweed) - (Apiaceae) [67] i) Norway

l) *Levisticum officinale* Koch  
(Lovage) - (Apiaceae) [67] j) Europe, North  
America, New Zealand

m) *Pastinaca sativa* L. (Wild  
parsnip) - (Apiaceae) [66,67] k) North hemisphere,  
New Zealand

n) *Petroselinum sativum* Hoffm.=  
*Petroselinum crispum* subsp. *crispum*  
- (Apiaceae) [67] l) Europe, USA,  
Korea, Japan, New Zealand

o) *Peucedanum ostruthium* Linnaeus,  
1753 (Masterwort) - (Apiaceae) [67] m) Eurasia, Oceania,  
North and South

p) *Pimpinella magna* L. - (Apiaceae) Americas, including

[67]

Brazil

q) *Heracleum thomsoni* - (Apiaceae) n) Europe and

[67]

Central America

r) *Psoralea corylifolia* L. -

(Fabaceae)

o) Europe

p) Europe

q) Asia

s) *Ficus carica* L. (Common fig) -

(Moraceae)

[66,67]

r) East Asia

t) *Citrus x aurantifolia* (Christm.)

Swingle (Lime) - (Rutaceae)

[67]

s) Worldwide,

u) *Citrus × limon* (L.) Burm.fil. =

*Citrus × bergamia* Risso (Lemon) -

(Rutaceae)

[67]

t) South hemisphere,

v) *Citrus limetta* Risso = *Citrus medica* L. (Mousami) - (Rutaceae)


including Brazil

[67]

- w) *Citrus x paradisi* Macfad.  
 (Grapefruit / Pomelo) - (Rutaceae) [67] u) Worldwide,  
 including Brazil
- x) *Citrus x sinensis* (Mill.) Pers.,  
 1806 = *Citrus × aurantium* L.  
 (Orange) - (Rutaceae) [67] v) South America,  
 including Brazil
- y) *Dictamnus albus* L. (Burning  
 bush) - (Rutaceae) [67] w) Americas,  
 including Brazil, East
- e) *Pelea anisata* H. Mann. =  
*Melicope anisata* (H.Mann)  
 T.G.Hartley & B.C.Stone -  
 (Rutaceae) [67] x) Western Africa,  
 Western Europe,  
 Central America and  
 South America,  
 including Brazil
- aa) *Ruta chalepensis* L. (Fringed rue)  
 - (Rutaceae) [67] y) Europe, USA,  
 Asia
- ab) *Ruta graveolens* L. (Rue / Rua) -  
 (Rutaceae) [67] z) Hawaii  
 aa) Europe, Central  
 America, South  
 America, a few in  
 Brazil

Furanocoumarin  
glycoside  
Type I  
(C<sub>20</sub>H<sub>24</sub>O<sub>10</sub>)

Isorutarin




*Ruta graveolens* L. (Rua) -  
(Rutaceae)

ab) Eurasia, South  
America, including  
Brazil

Numerous  
reports of  
phototoxic  
events with this  
species.

[47,50]

$\lambda_{\text{max}} \sim 310-320$   
nm depending  
on solvent

Isoquinoline alkaloid  Type I	Berberine  (C <sub>20</sub> H <sub>18</sub> NO <sub>4</sub> +)		(Annonaceae) [60,157]  a) <i>Annona purpurea</i> Moç. & Sessé ex Dunal (Sncoya) - (Annonaceae) [47]	a) Central America  b) Europe, USA	a) leaves  b) leaves	Berberin is used in Traditional Chinese Medicine  because of its multiple biological activities including antidiabetic, anti- inflammatory, antimalarial, antimicrobial, anticancer and
			(Ranunculaceae) [157]  b) <i>Helleborus niger</i> L. (Rose de Noël) - (Ranunculaceae) [157]	c) Europe, Asia and South America, including Brazil		
			(Berberidaceae) [157]			
			(Juglandaceae) [47]			
			(Magnoliaceae) [157]			
			(Menispermaceae) [157]			

(Rubiaceae) [157]

antioxidant activities.


(Rutaceae) [157]


(Papaveraceae) [157]

c) *Chelidonium majus* L. (Celandine)  
- (Papaveraceae) [115]

Berberin is fluorescent and capable of producing  $^1\text{O}_2$  and radical species when exposed to UVA irradiation. Its phototoxicity is associated to apoptosis after ROS generation and, autophagy in cancer cell lines (« photokilling » activity).

$\lambda_{\text{max}} \sim 345\text{nm}$   
depending on  
solvent

Naphthodianthrones	Fagopyrin :			[38,50,54,64,67,157]
Type I	E : ( $C_{39}H_{32}N_2O_8$ )		<i>Fagopyrum spp.</i> - (Polygonaceae) [64,67] a) North hemisphere Leaves and flowers	Highly Phototoxic [54]
	F : ( $C_{38}H_{30}N_2O_8$ )		b) North hemisphere, a) <i>Fagopyrum tataricum</i> (L.) Gaertn Brazil, South of (Tartarian buckwheat) - Africa, Australia, (Polygonaceae) [54,64,157] New Zealand	$\lambda_{\text{max}} \sim 550\text{-}590$ nm depending on solvent
		Fagopyrin E	b) <i>Fagopyrum esculentum</i> Moench - c) Balkans, Japan, (Polygonaceae) [38,50,54,64] Korea, USA	




c) *Fagopyrum sagittatum* -  
(Polygonaceae) [54]

Protofagopyrin :

Fagopyrin F

(C<sub>40</sub>H<sub>36</sub>N<sub>2</sub>O<sub>8</sub>)




Protofagopyrin

Oxoaporphine  
alkaloid

Type I

Lysicamine  
 $(C_{18}H_{13}NO_3)$



*Guatteria spp.* - (Annonaceae)<sup>[62]</sup>

a) *Guatteria blepharophylla* Mart.-

(Annonaceae)<sup>[61]</sup>

a) South hemisphere,  
including Brazil

b) *Guatteria ferruginea* A.St.-Hil. -

(Annonaceae)<sup>[62]</sup>

b) North of Southern  
America, including  
Brazil

c) *Guatteria latifolia* (Mart.) R.E.Fr.

- (Annonaceae)<sup>[62]</sup>

c) Brazil

d) Brazil

d) *Guatteria sellowiana* Schltld. -



(Annonaceae)<sup>[62]</sup>

Identification  
of Isoquinoline  
alkaloids in  
aerial parts of 3



*Guatteria*  
species found  
in Brazil.

[158]

$\lambda_{max}$ ~282nm  
depending on  
solvent

Perylenequinone Type I	Hypocrellin B (C <sub>30</sub> H <sub>24</sub> O <sub>9</sub> )		<i>Hypocrella bambusae</i> (Berk. & Broome) Sacc. - (Parasitic fungi Clavicipitaceae) <sup>[105]</sup>	China	$\lambda_{\text{max}} \sim 502-510$ nm in polar solvents like ethanol or DMSO	[105,106,129]
Phenalenone derivative	Lachnanthocarpone (C <sub>19</sub> H <sub>12</sub> O <sub>3</sub> )		<i>Lachnantes tinctoria</i> – (Haemodoraceae)	USA Atlantic coast	Phototoxic on <i>Staphylococcus epidermidis</i>  Report of lachnanthocarpone tetraoxxygenated derivatives (tetraoxigenate d phenylphenalen ones) in <i>Wachendorfia thyrsiflora</i>	[101,102,159]

(Haemodoracea  
e)

Phenylpropanoid	Hydroxycinnamic acid		<i>Myrciaria cauliflora</i> (Mart.) O.Berg	Americas, including	Fruits	[112]	[112]
Type I	=		(Jaboticaba) - (Myrtaceae)	Brazil			
	(E)- <i>p</i> -coumaric acid					$\lambda_{\text{max}} \sim 310-320$	
	(C <sub>9</sub> H <sub>8</sub> O <sub>3</sub> )					nm depending	
						on solvent	
Polyphenol	3,4-Dihydroxyphenyl		a) <i>Froelichia floridana</i> (Nutt.) Moq.	a) USA, Australia		$\lambda_{\text{max}} \sim 325-330$	[52]
Type I	polyphenolic caffeate		- (Amaranthaceae) <sup>[52]</sup>			nm depending	
	=3,4-					on solvent	
	Dihydroxycinnamate		b) <i>Ammi majus</i> L. (Bishop's weed) -	Northern and			
	=caffein acid		(Apiaceae) <sup>[52]</sup>	Western Africa,			
				Central Asia, Brazil			
		(C <sub>14</sub> H <sub>16</sub> O <sub>4</sub> )					

Polyphenol Type I	Curcumin  (C <sub>21</sub> H <sub>20</sub> O <sub>6</sub> )		<i>Curcuma longa</i> L. (Curcuma) - (Zingiberaceae) [41,145–147,157]	Tropical belt, including Brazil	Rhizomes	Phototoxicity demonstrated in several cancer cell lines and bacteria.	[41,67,106,129,145– 147,150,157]
Protoberberine Alkaloid Type I	Protoberberine  (C <sub>17</sub> H <sub>14</sub> N <sup>+</sup> )		<i>Chelidonium majus</i> L. (Celandine) - (Papaveraceae) [115,157]	Europe, Asia and South America, including Brazil	Leaves and stems during flowering	Phototoxic and Pro-apoptotic in uterine cancer cells	[115,157]


Pyrrolizidine

Dehydromonocrotaline

Alkaloid

Type I

(C<sub>16</sub>H<sub>21</sub>NO<sub>6</sub>)



[79]


Pyrrolizidine

Heliotrine

Alkaloid

Type I

(C<sub>16</sub>H<sub>27</sub>NO<sub>5</sub>)



*Heliotropicum europeum* (European heliotrope) - (Heliotropiaceae) [54]



USA

λ<sub>max</sub>~323 nm

depending on  
solvent. See


Dehydromonoc  
rotaline  
comment

[54,79]

Pyrrolizidine	Lasiocarpine		<i>Heliotropicum europeum</i> (European heliotrope) - (Heliotriaceae) <sup>[54]</sup>	Eurasia, Australia, USA	See Dehydromonocrotaline comment	[54,79]
Alkaloid						
Type I						
(C <sub>21</sub> H <sub>33</sub> NO <sub>7</sub> )					λmax~265 nm depending on solvent	
Pyrrolizidine	Retronecine		<i>Anchusa strigosa</i> Banks & Sol. - (Boraginaceae) <sup>[79]</sup>	Arabian peninsula, Middle East, Israel	Roots	See Dehydromonocrotaline comment
Alkaloid						
Type I						
(C <sub>8</sub> H <sub>13</sub> NO <sub>2</sub> )					λmax~330 nm depending on solvent	

Pyrrolizidine  
Alkaloid  
Type I

Retrorsine  
(C<sub>18</sub>H<sub>25</sub>NO<sub>6</sub>)



- a) *Brachyglottis hectori* (Buchanan)  
B.Nord. - (Asteraceae) a) New Zealand
- b) *Brachyglottis adamsii*  
(Cheeseman) B.Nord. - (Asteraceae) b) New Zealand
- c) *Brachyglottis huntii* (F.Muell.)  
B.Nord. - (Asteraceae) c) New Zealand
- d) *Brachyglottis perdicioides*  
(Hook.fil.) Hook.fil. - (Asteraceae) d) New Zealand
- e) *Brachyglottis sciadophila* (Raoul)  
B.Nord. - (Asteraceae) e) New Zealand
- f) *Brachyglottis kirkii* (Hook.fil. ex  
Kirk) C.J.Webb= *Urostemon kirkii* f) New Zealand
- g) *Senecio jacobaea* Brot. ex Willk.  
& Lange, 1865 (Tansy ragwort) - g) North America,  
Europe, Australia
- h) Etats-Unis
- i) USA
- j) North America,  
Eurasia, West of  
South America,  
Australia, New  
Zealand

See  
Dehydromonoc  
rotaline  
comment.

[54,79]

λ<sub>max</sub>~ 250 nm  
depending on  
solvent

h) *Senecio riddellii* Torr. & A.Gray -

(Asteraceae)

i) *Senecio douglasii* DC. -

(Asteraceae)

j) *Senecio vulgaris* L. - (Asteraceae)


Pyrrolizidine

Riddelline

Alkaloid

Type I

(C<sub>18</sub>H<sub>23</sub>NO<sub>6</sub>)



*Senecio riddellii* Torr. & A.Gray -

(Asteraceae)<sup>[79]</sup>


USA

See

[79]

Dehydromonoc  
rotaline  
comment

λ<sub>max</sub>~ 335 nm  
depending on  
solvent

Pyrrolizidine														
Alkaloid														
Type I														
	Senecionine (C <sub>18</sub> H <sub>25</sub> NO <sub>5</sub> )		<i>Brachyglottis</i> spp - (Asteraceae)	a) New Zealand										
			a) <i>Brachyglottis hectori</i> (Buchanan) B.Nord. - (Asteraceae)	b) New Zealand										
			b) <i>Brachyglottis adamsii</i> (Cheeseman) B.Nord. - (Asteraceae)	c) New Zealand										
			c) <i>Brachyglottis huntii</i> (F.Muell.) B.Nord. - (Asteraceae)	d) New Zealand										
			d) <i>Brachyglottis perdicioides</i> (Hook.fil.) Hook.fil. - (Asteraceae)	e) New Zealand										
			e) <i>Brachyglottis sciadophila</i> (Raoul) B.Nord. - (Asteraceae)	f) New Zealand										
			f) <i>Brachyglottis kirkii</i> (Hook.fil. ex Kirk) C.J.Webb = <i>Urostemon kirkii</i> (Hook.fil. ex Kirk) B.Nord. - (Asteraceae)	g) Mexico										
				h) Mexico										
				i) Mexico										
				j) Mexico										
				k) Mexico										
				l) Europe, Mediterranean basin										
			g) <i>Pittocaulon praecox</i> (Cav.) H.Rob. & Brettell - (Asteraceae)	m) North hemisphere										

See  
Dehydromonoc  
rotaline  
comment.

[79]

λ<sub>max</sub>~ 245-250  
nm depending  
on solvent

- h) *Pittocaulon bombycophole*  
 (Bullock) H.Rob. & Brettell -  
 (Asteraceae)
- n) Mediterranean  
 basin
- i) *Pittocaulon velatum* (Greenm.)  
 H.Rob. & Brettell - (Asteraceae)
- p) Eurasia
- j) *Pittocaulon hintonii* H.Rob. &  
 Brettell - (Asteraceae)
- q) East Europe,  
 Middle East
- k) *Pittocaulon filare* (Mc Vaugh)  
 H.Rob. & Brettell - (Asteraceae)
- r) Bulgaria
- Senecio spp. - (Asteraceae)
- s) East Europe
- t) East Europe
- l) *Senecio gibbosus* subsp. *bicolor* -  
 (Asteraceae)
- u) Europe
- m) *Senecio cineraria* DC. -  
 (Asteraceae)
- v) East Europe
- n) *Cineraria maritima* L.= *Jacobaea*  
*maritima* subsp. *maritima* -  
 (Asteraceae)

o) *Senecio stadianus* Lacaita -

(Asteraceae)

p) *Senecio erucifolius* L. -

(Asteraceae)

q) *Senecio othonnae* M.Bieb.=

*Jacobaea othonnae* (M.Bieb.)

C.A.Mey. - (Asteraceae)

r) *Senecio wagneri* Degen =

*Tephroseris papposa* subsp. *wagneri*

(Degen) B.Nord. - (Asteraceae)

s) *Senecio subalpinus* Koch =

*Jacobaea subalpina* (W.D.J.Koch)

Pelser & Veldkamp - (Asteraceae)

t) *Senecio carpathicus* Herbich =

*Jacobaea abrotanifolia* subsp.


*carpathica* (Herbich) B.Nord. &

Greuter - (Asteraceae)

u) *Senecio paludosus* L. = *Jacobsaea*  
*paludosa* subsp. *paludosa* -  
(Asteraceae)

v) *Senecio rupestris* Waldst. & Kit.=  
*Senecio squalidus* subsp. *rupestris*  
(Waldst. & Kit.) Greuter-  
(Asteraceae)

Pyrrolizidine  
Alkaloid  
Type I  
(C<sub>19</sub>H<sub>27</sub>NO<sub>6</sub>)




- |  |                |
|--|----------------|
| a) <i>Brachyglottis hectori</i> (Buchanan)<br>B.Nord. - (Asteraceae)         | a) New Zealand |
| b) <i>Brachyglottis adamsii</i><br>(Cheeseman) B.Nord. - (Asteraceae)        | b) New Zealand |
| c) <i>Brachyglottis huntii</i> (F.Muell.)<br>B.Nord. - (Asteraceae)          | c) New Zealand |
| d) <i>Brachyglottis perdicioides</i><br>(Hook.fil.) Hook.fil. - (Asteraceae) | d) New Zealand |
|  | e) New Zealand |
|  | f) New Zealand |

See  
Dehydromonoc  
rotaline  
comment.

$\lambda_{\text{max}}$  ~ 275-280  
nm depending  
on solvent

- e) *Brachyglottis sciadophila* (Raoul)  
B.Nord. - (Asteraceae)
- f) *Brachyglottis kirkii* (Hook.fil. ex Kirk) C.J.Webb = *Urostemon kirkii*  
(Hook.fil. ex Kirk) B.Nord. -  
(Asteraceae)
- g) Mexico
- h) Mexico
- i) Mexico
- j) Mexico
- k) Mexico
- l) East Europe,  
Middle East
- g) *Pittocaulon praecox* (Cav.)  
H.Rob. & Brettell - (Asteraceae)
- h) *Pittocaulon bombycophole*  
(Bullock) H.Rob. & Brettell -  
(Asteraceae)
- i) *Pittocaulon velatum* (Greenm.)  
H.Rob. & Brettell - (Asteraceae)
- j) *Pittocaulon hintonii* H.Rob. &  
Brettell - (Asteraceae)
- k) *Pittocaulon filare* (Mc Vaugh)  
H.Rob. & Brettell - (Asteraceae)

		1) <i>Senecio othonnae</i> M.Bieb.= <i>Jacobsaea othonnae</i> (M.Bieb.) C.A.Mey. - (Asteraceae)			
Thiophene Type I	Thiophene and derivatives (C <sub>4</sub> H <sub>4</sub> S)		(Asteraceae) <sup>[47]</sup> <i>Echinops</i> spp. - (Asteraceae) <sup>[47]</sup> a) <i>Echinops latifolius</i> Tausch. - (Asteraceae) <sup>[47,106]</sup> <i>Tagetes</i> spp. - (Asteraceae) <sup>[41,157]</sup> b) <i>Tagetes erecta</i> L. - (Asteraceae) [41] <i>Artemisia</i> spp. - (Asteraceae) <sup>[157]</sup> <i>Balsamorhiza</i> spp. - (Asteraceae) <sup>[157]</sup> <i>Blumea</i> spp. - (Asteraceae) <sup>[47]</sup> <i>Pluchea</i> spp. - (Asteraceae) <sup>[47]</sup>	a) Russia, China b) Worldwide, including Brazil c) Russia d) South America, including Brazil	c) Root d) Aerial part ROS production after UV irradiation <sup>[157]</sup> Phototoxic in skin and uterine cancer cell lines <sup>[106]</sup> $\lambda_{\text{max}} \sim 210-225$ nm depending on solvent

*Porophyllum* spp. - (Asteraceae) [47]

*Eclipta* spp. - (Asteraceae) [47]

*Heliopsis* spp. - (Asteraceae) [41]

*Rudbeckia* spp. - (Asteraceae) [41]

c) *Leuzea carthamoides* Willd. -  
(Asteraceae) [41]

d) *Porophyllum obscurum* (Spreng.)  
DC. - (Asteraceae) [41]

Genus *Berkheya* - (Asteraceae) [41]

Genus *Dyssodia* - (Asteraceae) [41]

Genus *Echinops* - (Asteraceae) [41]

Genus *Bidens* - (Asteraceae) [41,160]

Xanthone	Mangostin		<i>Garcinia mangostana</i> L. (Mangosteen) - (Clusiaceae) <sup>[88]</sup>	South hemisphere, including Brazil	Pericarp	$\lambda_{\text{max}} \sim 243,$ $317$ and $352$ nm depending on solvent	[41,89,96]
Type I							

(C<sub>24</sub>H<sub>26</sub>O<sub>6</sub>)

Type II  
Photosensitization

Anthocyanins or anthocyanins	Anthocyanins (flavonoids)		a) <i>Myrciaria cauliflora</i> (Mart.) O.Berg (Jaboticaba) - (Myrtaceae) [112]	a) Americas, including Brazil	Fruits	Reports in superior plants	[96,112]
Type II							


(Anthocyanins are  
usually considered  
as photoprotective  
molecules whose

(C<sub>15</sub>H<sub>11</sub>O+)

b) Red fruits, red-purple dark  
vegetables, black rice<sup>[96]</sup>

biosynthesis is induced by high solar irradiation.

However, their ingestion by cattle can lead to type II photosensitization.)

Furanosesquiterpene Type II	Ngaione (=ipomeamarone) (C <sub>15</sub> H <sub>22</sub> O <sub>3</sub> )		<i>Myoporum laetum</i> G. Forst. (New Zealand manatoka) - (Scrophulariaceae)	American and European coastlines, including Brazil	Oils	Hepato-induced photosensitization	[38,54]
Lectin Type II	Lectin <i>Arachis hypogaea</i> : PNA	Protein	<i>Arachis hypogaea</i> L. (Peanut) - (Fabaceae) <sup>[109]</sup>	Worldwide, including Brazil		Studied as PS in epidermoid carcinoma, colorectal carcinoma,	[109]

hepato-  
carcinoma cell  
lines.

Lectin Type II	Lectin <i>Artocarpus integrifolia</i> : Jacalin	Protein	<i>Artocarpus integrifolia</i> L.fil. = <i>Artocarpus integer</i> (Thunb.) Merr. - (Moraceae) <sup>[109]</sup>	Africa, Europe, South America, including Brazil	Studied as PS in epidermoid carcinoma, colorectal carcinoma, hepato- carcinoma cell lines.	[109]	
Lectin Type II	Lectin <i>Bauhinia forficata</i> : BfL	Protein	<i>Bauhinia forficata</i> Link (Cow's foot) - (Fabaceae) <sup>[109]</sup>	South America, including Brazil	/	Studied as PS in the NCI-60 tumor cell lines panel.  Inhibits the adhesion of	[109]



cancer cells on  
laminin,  
collagen I and  
fibronectin and  
induces  
necrosis of  
cancer cell and  
DNA  
fragmentation.

Lectin	Lectin <i>Glycine max</i>	Protein	<i>Glycine max</i> (L.) Merr. (Soybean) (P) - (Fabaceae) <sup>[109]</sup>	Americas, including Brazil, Eurasia, Africa	Studied as PS in various cancer cell lines including hepato- carcinoma, melanoma, choriocarcinom a cells.	<sup>[109]</sup>
Type II					ROS-dependent induction of autophagy,	

					apoptosis and DNA damage in cells	
Lectin Type II	Lectin <i>Griffonia simplicifolia</i> : GSAIA4	Protein	<i>Griffonia simplicifolia</i> (Vahl ex DC.) Baill. - (Fabaceae) <sup>[109]</sup>	West Africa	Studied as PS in various cancer cell lines including Melanoma and colorectal cells.	<sup>[109]</sup>
Lectin Type II	Lectin <i>Myrsine coriacea</i> : McL	Protein Unknown structure	<i>Myrsine coriacea</i> Merr. - (Primulaceae) <sup>[109]</sup>	Vietnam	Studied as PS in colon cancer cells.  ROS-induced apoptosis evidenced by DNA fragmentation and Cytochrome c	<sup>[109]</sup>

					release in the cytosol	
Lectin Type II	Lectin <i>Ricinus communis</i> : RCA	Protein Unknown structure	<i>Ricinus communis</i> L. - (Euphorbiaceae) [109]	Worldwide, including Brazil	/	Studied as PS in melanoma and hepatoma cell lines.
					ROS-induced apoptosis evidenced by chromatin condensation and nuclear fragmentation	[109]
Lectin Type II	Lectin <i>Viscum album</i> (Mistletoe) : ML	Protein	<i>Viscum album</i> L. (Gui) - (Viscaceae) [109]	Eurasia, North America	Studied as PS in breast, hepatocarcinoma and colorectal cancer cells.	

					ROS-induced apoptosis in colorectal cancer cells, regulation of $\beta$ -catenin expression	
Lectin Type II	Lectin <i>Ximenia americana</i> : Riproximin	Protein Unknown structure	<i>Ximenia americana</i> L. (Tallow wood) - (Ximeniaceae) <sup>[109]</sup>	South hemisphere, including Brazil	Studied as PS in breast carcinoma, brain, prostate, colon, leukemia cell lines.	<sup>[109]</sup>




Monoterpene Type II	(+)-Eucalyptol = 1,8-cinéole (C <sub>10</sub> H <sub>18</sub> O)		a) <i>Eucalyptus globulus</i> Labill. (Eucalyptus / Gommier bleu) - (Myrtaceae) <sup>[108]</sup>  b) <i>Cinnamomum camphora</i> (L.) Nees & Eberm. = <i>Cinnamomum camphora</i> (L.) J.Presl (Camphrier / Laurier de Chine) - (Lauraceae) <sup>[108]</sup>	a) Worldwide, including Brazil  b) Worldwide, including Brazil	Widely distributed in several plant species	Widely distributed in plant essential oils.	First published study evaluating the use of essential oils as PS, in addition to their well-known antibacterial activities.	Eucalyptol, eugenol and thymol have been studied as potential PS.	[108]
Pentacyclic triterpenes Type II	Lantadenes A = (C <sub>35</sub> H <sub>54</sub> O <sub>5</sub> )		<i>Lantana camara</i> L. (Common lantana) - (Verbenaceae)	Worldwide			Reports of photosensitization after hepatic metabolism (hepato-induced)		[54]

B = (C<sub>35</sub>H<sub>52</sub>O<sub>5</sub>)

photosensitizati  
on)

C = (C<sub>35</sub>H<sub>54</sub>O<sub>5</sub>)

D = (C<sub>34</sub>H<sub>52</sub>O<sub>5</sub>)

Phenolic compound	Eugenol		<i>Syzygium aromaticum</i> (L.) Merr. & Perry = <i>Eugenia caryophyllata</i> Thunb. (Clove) - (Myrtaceae) <sup>[108]</sup>	Tropical rainforest, including Brazil	Clove (flower bud)	See comment for Eucalyptol	[108]
Type II							
Phenolic compound <i>p</i> -hydroxy-benzoic acid			<i>Myrciaria cauliflora</i> (Mart.) O.Berg (Jaboticaba = Brazilian fruit) - (Myrtaceae)	Americas, including Brazil	Fruits		[112]
Type II							
Phenolic compound	Resorcinol		<i>Myrciaria cauliflora</i> (Mart.) O.Berg (Jaboticaba) - (Myrtaceae)	This plant is native to Brazil, and is distributed	Fruits		[56,112]
Type II							


(C<sub>6</sub>H<sub>6</sub>O<sub>2</sub>)

Phenolic compound

Type II

(C<sub>10</sub>H<sub>14</sub>O)

Thymol



a) *Thymus vulgaris* L. (Thyme)

b) *Eucalyptus globulus* Labill.  
(Eucalyptus / Gommier bleu) -  
(Myrtaceae)

c) *Syzygium aromaticum* (L.) Merr.  
& Perry = *Eugenia caryophyllata*  
Thunb. (Clove) - (Myrtaceae)

throughout the  
Atlantic Forest  
biome.

a) Europa, Oceania, Clove, thyme, oregano, Phototoxic in  
USA, Central and cinnamon and bacteria,  
South Americas, peppermint oils, (as viruses, fungi,  
including Brazil well as other volatile protozoans.

b) Worldwide,  
including Brazil

c) Tropical  
rainforest, including  
Brazil

[108]


Widely used as  
conservative in  
pharmaceutical  
products.

See Eucalyptol  
comment.

Saponin

Dichotomin

Type II  
(C<sub>57</sub>H<sub>94</sub>O<sub>26</sub>)



a) *Brachiaria decumbens* Stapf =  
*Brachiaria eminii* (Mez) Robyns -  
(Poaceae) [114]

a) Central and South  
Americas, including  
Brazil, Africa

[114]

b) *Panicum dichotomiflorum* Michx.

(Millet) - (Poaceae) [114]

b) Americas


(including Brazil)

Europe, Korea and  
Japan, New Zealand

Saponin

Dioscin

Type II  
(C<sub>45</sub>H<sub>72</sub>O<sub>16</sub>)



a) *Brachiaria decumbens* Stapf =  
*Brachiaria eminii* (Mez) Robyns -  
(Poaceae) [114]

a) Central and South  
Americas, including  
Brazil, Africa

[116,161]

b) *Costus speciosus* (J.Koenig) Sm.=

*Hellenia speciosa* (J.Koenig)

S.R.Dutta (Canereed) - (Costaceae)

b) Pacific Asia,

Central America

[91]

Saponin	Diosgenin		a) <i>Costus speciosus</i> (J.Koenig) Sm.= <i>Hellenia speciosa</i> (J.Koenig) S.R.Dutta (Canereed) - (Costaceae) [91]	a) Pacific Asia, Central America	a) Rhizomes	[54,91,114]
Type II			b) <i>Panicum dichotomiflorum</i> Michx. (Smooth witchgrass) - (Poaceae)	Europe, Korea and Japan, New Zealand [54,91]	b) Americas (including Brazil)	Reported in biliary crystals associated to hepatogenous photosensitizati
	(C <sub>27</sub> H <sub>42</sub> O <sub>3</sub> )		c) <i>Brachiaria decumbens</i> Stapf = <i>Brachiaria eminii</i> (Mez) Robyns (Poaceae) (Meagher, Smith and Wilkins, 2001; Muniandy <i>et al.</i> , 2020)	Central and South Americas, including Brazil, Africa	d) Australia, USA, some occurences in Europe	on of sheep, goats and cattle
			d) <i>Panicum antidotale</i> Retz. (Blue panicum) - (Poaceae) <sup>[54]</sup>	e) Africa, U.S.A- Mexico border, one occurence in Brazil	f) Americas, including Brazil,	
			e) <i>Panicum coloratum</i> L. (Coolah grass) - (Poaceae) <sup>[54]</sup>			
			f) <i>Panicum maximum</i> Jacq. = <i>Megathyrsus maximus</i> (Jacq.)			

B.K.Simon & S.W.L.Jacobs (Guinea Africa, East Asia,  
grass) - (Poaceae) [54] Australia

g) *Panicum miliaceum* L. (Millet) - g) North hemisphere,  
(Poaceae) [54,114] Australia, rares cases  
in Brazil

h) *Panicum virgatum* L. (Switch  
grass) - (Poaceae) [54] h) USA, Europe, East  
Asia

i) *Tribulus terrestris* L. (Tribule  
terrestre) - (Zygophyllaceae) [91,114] i) Worldwide,  
including Brazil


Saponin	Episarsasapogenin		<i>Panicum spp.</i> - (Poaceae)	a) Worldwide, including Brazil	[91]
Type II	(C <sub>27</sub> H <sub>44</sub> O <sub>3</sub> )		a) <i>Tribulus terrestris</i> L. (Tribule terrestre) - (Zygophyllaceae)	b) Europe	Involved in the hepatogenous photosensitizati on of cattle. Identified in biliary crystals.



b) *Narthecium ossifragum* (L.) Huds.




(Nathrécie des marais) -

(Nartheciaceae)

Saponin	Epismilagenin		<i>Panicum spp.</i> - (Poaceae) <sup>[91]</sup>	a) South Africa	Involved in the hepatogenous photosensitization of cattle. <sup>[54,91,162]</sup>
Type II			a) <i>Tribulus terrestris</i> - <sup>[162]</sup>		
	(C <sub>27</sub> H <sub>44</sub> O <sub>3</sub> )		b) <i>Panicum antidotale</i> Retz. (Blue panicum) - (Poaceae) <sup>[54]</sup>	b) Australia, USA, some occurrences in Europe	
	= (3 $\alpha$ ,5 $\beta$ ,25R)-Spirostan-3-ol		c) <i>Panicum coloratum</i> L. (Coolah grass) - (Poaceae) <sup>[54]</sup>	c) Africa, U.S.A-Mexico border, one occurrence in Brazil	
			d) <i>Panicum dichotomiflorum</i> Michx. (Millet) - (Poaceae) <sup>[54]</sup>	d) Americas	
			e) <i>Panicum maximum</i> Jacq. = <i>Megathyrsus maximus</i> (Jacq.) B.K.Simon & S.W.L.Jacobs (Guinea grass) - (Poaceae) <sup>[54]</sup>	(including Brazil) Europe, Korea and Japan, New Zealand	
			f) <i>Panicum miliaceum</i> L. (Millet) - (Poaceae) <sup>[54]</sup>	e) Americas, including Brazil,	

		g) <i>Panicum virgatum</i> L. (Switch grass) - (Poaceae) <sup>[54]</sup>	Africa, East Asia, Australia
		f) North hemisphere, Australia, rares cases in Brazil	
		g) USA, Europe, East Asia	
Saponin	Protodioscine		a) <i>Brachiaria decumbens</i> Stapf = <i>Brachiaria eminii</i> (Mez) Robyns - (Poaceae)
Type II	(C <sub>51</sub> H <sub>84</sub> O <sub>22</sub> )		a) Central and South Americas, including Brazil, Africa
		b) <i>Brachiaria brizantha</i> (A.Rich.) Stapf - (Poaceae)	b) Central and South Americas, including Brazil, Africa, Asia
		c) <i>Brachiaria humidicola</i> (Rendle) Schweick.= <i>Brachiaria dictyoneura</i> (Fig. & De Not.) Stapf - (Poaceae)	c) Central and South Americas, including Brazil, Africa
			Grasses of the genus <i>Brachiaria</i> and leaves Reports of Phototoxicity in cattle (buffalos). <sup>[116]</sup>



Saponin	Smilagenin		<i>Narthecium ossifragum</i> (L.) Huds. (Swamp natherica) - (Nartheciaceae) [91,114]	Europe	Involved in the hepatogenous photosensitizati on of cattle. Identified in biliary crystals.	[91,114]
Type II						
Saponin	Soyasapogenol		<i>Brachiaria decumbens</i> Stapf = <i>Brachiaria eminii</i> (Mez) Robyns - (Poaceae)	Central and South Americas, including Brazil, Africa	Saponins often found in forages include dioscin/protodi oscin, soyasaponin, diogenin, soyasapogenol, and yamogenin	[114]
Type II	A: R=OH on C <sub>21</sub>  B: R=H on C <sub>21</sub>					
	(C <sub>30</sub> H <sub>50</sub> O <sub>4</sub> or C <sub>30</sub> H <sub>50</sub> O <sub>3</sub> )					

Saponin	Soyasaponin I		<i>Brachiaria decumbens</i> Stapf = <i>Brachiaria eminii</i> (Mez) Robyns - (Poaceae) [114]	Central and South Americas, including Brazil, Africa	See the soyasapogenol comment	[114]
Type II	(C <sub>48</sub> H <sub>78</sub> O <sub>18</sub> )				Glycine max (soybean)	
Saponin	Spirostanol		<i>Brachiaria decumbens</i> Stapf = <i>Brachiaria eminii</i> (Mez) Robyns - (Poaceae) [38]	Central and South Americas, including Brazil, Africa		[38]
Type II	C <sub>27</sub> H <sub>44</sub> O <sub>3</sub>					
Saponin	Yamogenin		a) <i>Brachiaria decumbens</i> Stapf = <i>Brachiaria eminii</i> (Mez) Robyns - (Poaceae) [91,114]  b) <i>Panicum miliaceum</i> L. (Millet) - (Poaceae) [114]	a) Central and South Americas, including Brazil, Africa  b) North hemisphere, Australia, rares cases in Brazil		[91,114]
Type II	(C <sub>27</sub> H <sub>42</sub> O <sub>3</sub> )				c) Europe	
						[91,114]

c) *Narthecium ossifragum* (L.) Huds.

(Nathrécie des marais) -

(Nartheciaceae)<sup>[114]</sup>

Tannin Type II	Pyrogallic and flobabenic tannins		a) <i>Mimosa pteridifolia</i> Benth. - (Fabaceae)  b) <i>Mimosa verrucosa</i> Benth. (Jurema-branca, Jurema-de-oeiras) - (Fabaceae)  c) <i>Mimosa tenuiflora</i> Willd. Poir. (Jurema preta, calumbi) - (Fabaceae)	a) Brazil  b) Brazil  c) Central America and Brazil	[97]
Triterpene Type II	Ursolic acid  (C <sub>30</sub> H <sub>48</sub> O <sub>3</sub> )		Abundant triterpene present in several species <sup>[163]</sup> including  <i>Malus pumila</i> (apple tree)  <i>Ocimum basilicum</i> (basil)	Nanovectorizati on strategy using indocyanine green and ursolic acid to treat cancers by	[144,163]

*Vaccinium spp.* (bilberries,  
blueberries, cranberries)

laser  
phototherapy

*Olea europaea* (olive)

*Origanum vulgare* (oregano)

*Rosmarinus officinalis* (rosemary)

*Salvia officinalis* (sage, salvia)

*Thymus vulgaris* L. (common thyme)

---

Type I and II Mixed


Photosensitization

Oxoaporphine alkaloid	Isomoschatoline (3-Hydroxy-1,2-dimethoxy-7H-dibenzo[de,g]quinoline-7-one)		<i>Guatteria spp.</i> - (Annonaceae) <sup>[62]</sup>	a) South hemisphere, including Brazil	Bark, crude extract	Antibacterial	[62,62]
Type I and II			a) <i>Guatteria blepharophylla</i> Mart.- (Annonaceae) <sup>[61]</sup>	b) North of Southern America, including Brazil		phototoxicity in Gram + and Gram - bacteria and <i>Candida</i> yeasts.	
			b) <i>Guatteria ferruginea</i> A.St.-Hil. - (Annonaceae) <sup>[62]</sup>	c) Brazil			Induction of type I and II photochemical

(C<sub>18</sub>H<sub>13</sub>NO<sub>4</sub>)

Porphyrins and  
derivatives  
Type I and II

20<sup>1</sup>-Hydroxypurpurin-7  
delta-lactone diesters  
and derivatives  
(dimethyl esters, ethyl  
methyl diester, methyl  
phytetyl diester)  
  
(Hydroxy purpurin 7-  
lactone dimethyl ester :  
 $C_{36}H_{38}N_4O_7$ )



Hydroxy purpurin 7-  
lactone dimethyl ester

c) *Guatteria latifolia* (Mart.) R.E.Fr.  
- (Annonaceae) [62]

d) Brazil

d) *Guatteria sellowiana* Schleidl. -  
(Annonaceae) [62]

mechanisms.  $\lambda$   
max~600-700  
nm depending  
on solvent

a) *Aglaonema simplex* (Blume)  
(Malayan sword) - (Araceae)

b) *Phyllostachys bambusoides*  
Siebold & Zucc (Japanese bamboo) -  
(Poaceae)

a) South East Asia  
b) Africa, Asia,  
North and South  
America, including

Brazil

a) Leaves and stems  
b) Leaves

Highly  
phototoxic and  
pro-apoptotic  
against human  
promyelocytic  
leukemia  
(HL60) and  
human  
leukemia cells  
(CMK-7).


[47,72,106]

Porphyrins and derivatives

Type I and II (phylloerythrin derivative, see paragraph 4.)

**Chlorophyll *a***

(C<sub>55</sub>H<sub>72</sub>MgN<sub>4</sub>O<sub>5</sub>)



Present in all photosynthetic organisms- Chlorophyll *a* ecophysiological function is to ensure photon collection for oxygenic photosynthesis in photosystems so its primary function is not to act as a phototoxic compound. However, chlorophylls and derivatives (e.g. pheophorbides and derivatives) show high phototoxicity when purified.

λ<sub>max</sub> chlorophyll *a*: 430, 660 nm depending on solvent

λ<sub>max</sub>~ 430-440 nm depending on solvent

[106]

**Chlorophyll *a*** can be found at

a massic concentration of up to 2% in leaves and is therefore an accessible, functionalizable pigment and an excellent starting material for the development of new PS.

[48,76,96]


Several chlorophyll *a* derivatives have a high phototoxic activity, for example chlorin, which shows a characteristic absorption spectrum at 400 nm and 680 nm.

Chlorophyll *a* is also the precursor of Pheophorbide *a*, its dephytylated and demetallized metabolite, that

exhibits potent phototoxic activity and can be obtained through chemical or enzymatic treatment. [48]

Porphyrins and derivatives  
Type I and II

Chlorophyllin *a*  
(C<sub>34</sub>H<sub>34</sub>MgN<sub>4</sub>O<sub>6</sub>)



a) *Spinacia oleracea* L. (Spinach) -  
(Amaranthaceae) [157]

b) Many plant species

a) Eurasia, North and South Americas,  
including Brazil

a) Leaves

Phototoxic at [106,129,146,157]  
405 nm on  
*Bacillus cereus*,  
*Listeria monocytogenes*,  
*Salmonella enterica*,  
*Escherichia coli*,  
*Pseudomonas aeruginosa*,  
*Enterococcus*


*faecalis* and  
fungi. [146]

Efficient for the  
photodecontam  
ination of  
plums (*Prunus*  
*domestica*),  
tomatoes  
(*Solanum*  
*lycopersicum*),  
cauliflowers  
(*Brassica*  
*oleracea*) and  
red peppers  
(*Capsicum*  
*annum*),  
naturally  
contaminated  
with mesophilic  
bacteria.

Chlorophyllin  
is activated in  
the wavelength  
range of 600–  
670 nm [106,157]

Porphyrins and  
derivatives  
Type I and II

Pheophorbide *a*  
(C<sub>35</sub>H<sub>36</sub>N<sub>4</sub>O<sub>5</sub>)



a) *Leonurus sibiricus* L. -  
(Lamiaceae) [73]  
  
b) *Scutellaria barbata* D. Don  
(Barbed skullcap) - (Lamiaceae)  
(Mansoori *et al.*, 2019; Foresto *et al.*,  
2021b)  
  
c) *Pentaspadon motleyi* -  
(Anacardiaceae) [73]

a) Americas,  
including Brazil,  
Europe, Asia  
  
b) Southern China  
  
c) Asia Pacific  
  
d) South East Asia


λ<sub>max</sub> [48,50,73,76,106,129,  
pheophorbide 157]  
*a*: 410, 666 nm

depending on  
solvent

Pheophorbide *a*  
and derivatives  
have also been  
reported as the  
molecules  
responsible for  
the

Porphyrins and  
derivatives  
Type I and II

Pheophorbide *b*  
(C<sub>35</sub>H<sub>34</sub>N<sub>4</sub>O<sub>6</sub>)



d) *Aglaonema simplex* Blume  
(Malayan sword) - (Araceae) [73]

e) *Euterpe oleracea* Mart. (Pinot) -  
(Araceae) [76]

j) *Tetragonia tetragonoides* (Pall.)  
Kuntze (Tetragone) - (Aizoaceae) [48]

e) Brazil and  
Colombia

f) Coastlines,  
including Brazil

a) Asia Pacific

a) *Pentaspadon motleyi* Hook.fil. -  
(Anacardiaceae) [73]

b) *Tetragonia tetragonoides* (Pall.)  
Kuntze (Tetragone) - (Aizoaceae) [48]

phototoxicity of  
microalgae  
extracts such as  
*Skeletonema*  
*sp.* ethanolic  
extract by some  
authors of this  
work (French  
patent  
FR20/03712)


Phototoxic on [48,73]

human  
colorectal  
carcinoma  
cells. [106]

$\lambda_{\text{max}}$   
pheophorbide  
*b*: 435, 654 nm  
depending on  
solvent

424 Table 2. List of phototoxic plants natural products, classified in alphabetic order. For each molecule are presented the raw formula, molecular  
425 structure, chemical family, producing species, geographical localization of producing species, plant tissue(s) or organ(s) containing the molecule(s)  
426 and biological activity in phototoxicity models. Phototoxicity events in humans and animals are also indicated when available.

427 The distribution of plant phototoxic molecules in different chemical families is presented in  
428 Figure 3. A total of 125 phototoxic molecules was listed, including 35 coumarins and  
429 furanocoumarins, 23 quinones/anthraquinones/naphtodianthrone, 22 alkaloids, 11 saponins, 9  
430 phenolics and polyphenols, 9 lectins, 8 terpenes, 5 porphyrins and derivatives, 2  
431 furanochromones and 1 thiophene.



432  
433 Figure 3. Repartition of the 125 plants phototoxic natural products in chemical families. The  
434 top five chemical families including the highest number of phototoxic compounds are  
435 coumarins and furanocoumarins (29), quinones, anthraquinones and naphtodianthrone (22),  
436 alkaloids (20), saponins (11) and lectins (9).

437  
438 Table 3. summarizes the distribution of phototoxic metabolites by botanical families. These  
439 data can be useful to select a botanical family for the identification and/or purification of  
440 phototoxic metabolites that belong to a targeted chemical family.

441

442

443

444

445

446

	Amaranthaceae	Anacardiaceae	Annonaceae	Apiaceae	Asteraceae	Fabaceae	Hypericaceae	Lamiaceae	Poaceae	Rubiaceae	Rutaceae
Alkaloids											
Catechols											
Coumarins and Furanocoumarins											
Lectins											
Polyphenols											
Porphyrins and derivatives											
Quinones/Anthraquinones/Naphtodianthrone											
Saponins											
Tannins											
Thiophenes											

447

448 Table 3. Distribution of phototoxic metabolites in the 11 major botanical families containing phototoxic species. A dark gray rectangle indicates  
 449 the discriminant presence of these metabolites in the botanical family, a light gray rectangle indicates the non-specific biosynthesis of these  
 450 metabolites (porphyrins and derivatives are photon collectors for oxygenic photosynthesis, but can be highly phototoxic once purified).

451

452

453 8. Conclusion

454 In this study, we have inventoried phototoxic plants and their phototoxic metabolites as a  
455 reservoir of chemodiversity for the possible development of photosensitizers for antimicrobial  
456 and antitumor phototherapy. Based on our results, we show that most biomes of the world  
457 contain plant species of major interest for the isolation of natural phototoxic molecules and that  
458 this high biodiversity is also related to a high chemodiversity. This implies that the biosynthesis  
459 of phototoxic phytochemicals is probably a convergent evolutionary trait that has evolved in a  
460 large taxonomic diversity of terrestrial plants, in reaction to a wide diversity of aggressions by  
461 predators and microorganisms. However, only around 250 vascular plant species (less than  
462 0.09% of all species), have been identified as phototoxic, containing a total of 125 phototoxic  
463 compounds, some of which are occurring together in a given species. These data suggest that  
464 this evolutionary strategy remains relatively rare for plants, although it can be observed in most  
465 ecoregions of the world and in a wide variety of taxonomic families. Based on our data, we  
466 could demonstrate that the botanical families Amaranthaceae, Anacardiaceae, Annonaceae,  
467 Apiaceae, Asteraceae, Fabaceae, Hypericaceae, Lamiaceae, Poaceae and Rutaceae have an  
468 important metabolic potential for the biosynthesis of phototoxic secondary metabolites,  
469 inducing primary photosensitizers (coumarins and furanocoumarins,  
470 quinones/anthraquinones/naphtdianthrones, alkaloids, porphyrins and derivatives,  
471 furanochromones and thiophenes) or secondary photosensitizers (saponins, phenolics and  
472 polyphenols, lectins, terpenes, chlorophylls and derivatives). Additional metabolites present in  
473 terrestrial plants, such as folates and pterins derivatives can also be considered as  
474 photosensitizing molecules with potential interest for the development of pharmacomodulated  
475 phototoxic derivatives<sup>[164]</sup>. Our study also demonstrates that several phototoxic species contain  
476 unidentified molecules and it would be interesting to study them to obtain new chemical  
477 scaffolds with improved ROS production yields, selective photoactivation parameters,  
478 interesting potential for pharmacomodulation or possible bioavailability improvements.

479 This review brought together scattered information from the literature about the world's plants  
480 ability to produce natural phototoxic compounds with potential for phototherapy. Our  
481 bibliographic research also revealed that a few species not belonging to the plant group also  
482 synthesize phototoxic molecules. These included

483 (i) *Teloschistes flavicans* (Sw.) Norman, a golden hair lichen (*Teloschistaceae*) present in  
484 Europe, South Africa, Oceania, Central and South Americas, including Brazil (Caatinga,

485 Amazônia, Mata Atlântica and Cerrado) synthesizing parietin, an anthraquinone pigment, [129].  
486 There are very few scientific articles demonstrating the presence of phototoxic molecules in  
487 lichens, but the finding in this species suggests that this hypothesis should be investigated  
488 further (on the other hand, several publications attest to the presence of photoprotective  
489 molecules in lichens [165]).

490 (ii) *Sclerotium rolfsii* Sacc. *Athelia rolfsii* (Curzi) C.C.Tu & Kimbr. (Atheliaceae), a soil fungus  
491 particularly prevalent in warm tropical and subtropical regions, which causes rotting of  
492 agronomic plants, present in Europe, USA, South of Africa, East of Asia and Brazil (Pampa  
493 and Mata Atlântica), whose phototoxicity was related to the synthesis of furanocoumarins  
494 (4,5',8-Trimethylpsoralen, 8-hydroxybergapten, bergapten, isoimperatorin, isopimpinellin,  
495 xanthotoxin) [50,66].

496 (iii) *Alternaria alternata* (Fr.) Keissl. (Pleosporaceae), a worldwide plant pathogenic fungus  
497 (often simply designated as "Alternaria"), agent of leaf spot disease, causing black spots, rot  
498 and rust on a wide range of plant species, present in Brazil in Caatinga, Pampa, Mata Atlântica,  
499 Cerrado and producing Altertoxin I, a phototoxic perylene dione [105].

500 (iv) *Cercospora kikuchii* (Tak.Matsumoto & Tomoy.) M.W.Gardner (Mycosphaerellaceae), a  
501 worldwide phytopathogenic fungus, responsible for the *Cercospora* leaf blight and purple seed  
502 stain diseases and producing the toxin cercosporin, a perylenequinone. Related  
503 perylenequinone fungal toxins with potential phototoxicity include cercosporin (*Cercospora*  
504 sp.), hypocrellin A (*Hypocrella bambusae*), Phleochrome (*Cladosporium* spp) and  
505 Elsinochrome A (*Elsinoe* spp)) [105].

506 (v) Basidiomycetes and sordariomycetes [166,167]: *Agaricus bisporus* (J.E.Lange) Imbach  
507 (Agaricaceae), an edible basidiomycete mushroom present in Europe, USA, Australia, Asia  
508 and America), contains a phototoxic lectin [109]. *Cortinarius croceus* acetone extract was  
509 characterized by an EC<sub>50</sub>, 9.3 J cm<sup>-2</sup> of 1 µg L<sup>-1</sup> against cells of a lung cancer cell-line (A549) [166].  
510 *Cortinarius austrovenetus* was also reported to contain austroventin, a precursor of hypericin  
511 that is oxidized when insects attack the fruiting bodies of the basidiomycete [166]. *Inonotus*  
512 *obliquus*, *Tapinella atrotomentosa*, *Roesleria subterranea* and *Metarrhizium brunneum*  
513 (sordariomycete) were also reported as containing phototoxic molecules inducing singlet  
514 oxygen production [166]. Other phototoxic natural products have been reported from  
515 Basidiomycetes, including questin, funalenone and 7,7'-emodinphysscion [166]. The small

516 number of articles published in the field suggests that these organisms represent an interesting  
517 but under-explored source of photosensitizers.

518 (vi) Microalgae species containing phototoxic porphyrin derivatives (e.g. ethanolic extract of  
519 diatoms from the genus *Skeletonema*, French patent FR20/03712 published by our group on  
520 the 15th of October 2021 by the French national institute of intellectual property INPI).

521 (vii) Bacterial, archaeal or cyanobacterial species containing phototoxic pigments such as  
522 prodigiosin, violacein or carotenoids.

523 The low number of phototoxic molecules identified in microorganisms suggests that these  
524 taxonomic groups have not been extensively screened yet. Additionally, no phototoxic  
525 molecules have been reported from bryophytes, filices and lycophytes so far, but it cannot be  
526 ruled out that these species have a potential to synthesize phototoxic molecules.

527 9. Future perspectives

528 Plant phototoxic molecules have a high potential for the development of new PS that could be  
529 used in topical or injectable formulations or grafted on biomaterials. Applications of these new  
530 PS could include the photodecontamination of abiotic surfaces (e.g decontamination of labs  
531 and hospitals workplans), treatment of topical infections caused by bacteria, viruses and fungi  
532 (e.g. chronic diabetic wounds, acne, nail mycosis), phototherapy of topical and internal tumors  
533 and manufacturing of photoactive devices (e.g. tissues, fibers, clothes, bandages). The chemical  
534 diversity of plant molecules inducing primary photosensitization implies that they have  
535 different bioavailabilities and different maximum absorption wavelengths, which makes it  
536 possible to envisage their differential use for topical or internal photoactivations using LASER.  
537 They could also be used as chemical leads for pharmacomodulation to selectively target  
538 microorganisms or cells to be destroyed (e.g. grafting of antibiotics or oligosaccharides to  
539 target bacterial membranes, grafting of polyamines to target tumor cell membranes, coupling  
540 to antibodies for vectorization against tumor antigens). Hardly any phototoxic molecules have  
541 been the subject of pharmaceutical or clinical studies so far, and some of them are found in  
542 common plants of agronomic, cosmetic, perfumery or medicinal interest, suggesting a possible  
543 secondary valorization of these species. Indeed, the vast majority are common species that are  
544 easy to cultivate, and it can be hypothesized that some valuable economic chains could be  
545 developed to produce PS (some species being already cultivated for the production of fruits,  
546 oils, essential oils, perfumes, spices, etc.).

547 The sustainable industrial production of natural photosensitisers from phototoxic plants will  
548 undoubtedly benefit from advances in other areas of medicinal chemistry and pharmacognosy  
549 research, such as the development of eco-friendly based extraction and purification processes  
550 [168,169], hemisynthesis or biotechnological production of bioactive metabolites in plant  
551 bioreactor cell cultures [170]. Finally, this study re-emphasises the importance of preserving  
552 plants as a pharmaceutical resource for humanity, as some species at risk of extinction could  
553 contain molecules allowing major innovations in the field of clinical phototherapy.

554

555 Declaration of competing interest

556 The authors declare that they have no known competing financial interests or personal  
557 relationships that could have appeared to influence the work reported in this paper.

558 Authors' contributions

559 Conceptualization: LP, VT, JGDSA, RGDOJ, VS; Data collection, analysis and curation: RP,  
560 JI, MG, RGDOJ, VT, LP; Writing, review and editing: All authors; Funding acquisition: LP  
561 and VT; Coordination and supervision: LP.

562 Acknowledgements

563 We are grateful to the French Cancer League (Ligue Nationale contre le Cancer, comité 17) for  
564 financial support, and the French western canceropole for scientific support. This work was  
565 carried out in the framework of the Franco-Brazilian Network on Natural Products (FB2NP).

566

567 Keywords

568 Natural Product, Plant, Photodynamic therapy, Photosensitizer, Phototoxicity.

569

570 References

- 571 [1] G. M. Cragg, D. J. Newman, K. M. Snader, *J. Nat. Prod.* **1997**, *60*, 52–60.  
572 [2] A. G. Atanasov, S. B. Zotchev, V. M. Dirsch, C. T. Supuran, *Nat. Rev. Drug Discov.*  
573 **2021**, *20*, 200–216.

- 574 [3] T. J. Dougherty, C. J. Gomer, B. W. Henderson, G. Jori, D. Kessel, M. Korbelik, J.  
575 Moan, Q. Peng, *J. Natl. Cancer Inst.* **1998**, *90*, 889–905.  
576 [4] D. E. J. G. J. Dolmans, D. Fukumura, R. K. Jain, *Nat. Rev. Cancer* **2003**, *3*, 380–387.  
577 [5] M. R. Hamblin, T. Hasan, *Photochem. Photobiol. Sci.* **2004**, *3*, 436–450.  
578 [6] T. G. St. Denis, T. Dai, L. Izikson, C. Astrakas, R. R. Anderson, M. R. Hamblin, G. P.  
579 Tegos, *Virulence* **2011**, *2*, 509–520.  
580 [7] E. Feese, H. Sadeghifar, H. S. Gracz, D. S. Argyropoulos, R. A. Ghiladi,  
581 *Biomacromolecules* **2011**, *12*, 3528–3539.  
582 [8] B. L. Carpenter, E. Feese, H. Sadeghifar, D. S. Argyropoulos, R. A. Ghiladi,  
583 *Photochem. Photobiol.* **2012**, *88*, 527–536.  
584 [9] C. Ringot, V. Sol, M. Barrière, N. Saad, P. Bressollier, R. Granet, P. Couleaud, C.  
585 Frochot, P. Krausz, *Biomacromolecules* **2011**, *12*, 1716–1723.  
586 [10] C. Ringot, N. Saad, F. Brégier, P. Bressollier, E. Poli, V. Chaleix, T. S. Ouk, V. Sol,  
587 *Photochem. Photobiol. Sci.* **2018**, *17*, 1780–1786.  
588 [11] J. Dong, R. A. Ghiladi, Q. Wang, Y. Cai, Q. Wei, *Nanotechnology* **2018**, *29*, 265601.  
589 [12] A. Elkhalil, C. Christie, C. Vernisse, T.-S. Ouk, R. Lucas, V. Chaleix, V. Sol, *ACS  
590 Appl. Bio Mater.* **2021**, *4*, 7204–7212.  
591 [13] T. Tsai, H.-F. Chien, T.-H. Wang, C.-T. Huang, Y.-B. Ker, C.-T. Chen, *Antimicrob.  
592 Agents Chemother.* **2011**, DOI 10.1128/AAC.00550-10.  
593 [14] A. Shrestha, M. R. Hamblin, A. Kishen, *Antimicrob. Agents Chemother.* **2012**.  
594 [15] N. Maldonado-Carmona, G. Marchand, N. Villandier, T.-S. Ouk, M. M. Pereira, M. J.  
595 F. Calvete, C. A. Calliste, A. Źak, M. Piksa, K. J. Pawlik, K. Matczyszyn, S. Leroy-  
596 Lhez, *Front. Microbiol.* **2020**, *11*, 2846.  
597 [16] M. Merchán, T. S. Ouk, P. Kubát, K. Lang, C. Coelho, V. Verney, S. Commereuc, F.  
598 Leroux, V. Sol, C. Taviot-Guého, *J. Mater. Chem. B* **2013**, *1*, 2139–2146.  
599 [17] Q. Wang, W. Chen, Q. Zhang, R. A. Ghiladi, Q. Wei, *Appl. Surf. Sci.* **2018**, *457*, 247–  
600 255.  
601 [18] N. Soliman, V. Sol, T.-S. Ouk, C. M. Thomas, G. Gasser, *Pharmaceutics* **2020**, *12*,  
602 961.  
603 [19] C. R. Ghareeb, B. S. T. Peddinti, S. C. Kisthardt, F. Scholle, R. J. Spontak, R. A.  
604 Ghiladi, *Front. Med.* **2021**, *8*, 1092.  
605 [20] K. R. Stoll, F. Scholle, J. Zhu, X. Zhang, R. A. Ghiladi, *Photochem. Photobiol. Sci.*  
606 **2019**, *18*, 1923–1932.  
607 [21] M. R. Hamblin, *Photochem. Photobiol.* **2020**, *96*, 506–516.  
608 [22] J. Kou, D. Dou, L. Yang, *Oncotarget* **2017**, *8*, 81591–81603.  
609 [23] R. Bonnett, *Chem. Soc. Rev.* **1995**, *24*, 19–33.  
610 [24] G. S. Fraenkel, *Science* **1959**, DOI 10.1126/science.129.3361.1466.  
611 [25] M. Berenbaum, *Science* **1978**, DOI 10.1126/science.201.4355.532.  
612 [26] T. Arnason, P. Morand, J. Salvador, I. Reyes, J. Lambert, N. Towers, *Phytochemistry*  
613 **1983**, *22*, 594–595.  
614 [27] E. Kourany, J. T. Arnason, E. Schneider, *Physiol. Mol. Plant Pathol.* **1988**, *33*, 287–  
615 297.  
616 [28] J. T. A. T. Arnason, M. A. B. A. Bernards, *Can. J. Zool.* **2010**, DOI 10.1139/Z10-038.  
617 [29] V. M. S. dos Reis, *An. Bras. Dermatol.* **2010**, *85*, 479–489.  
618 [30] E. F. F. Matias, K. K. A. Santos, J. G. M. Costa, H. D. M. Coutinho, *Indian J. Med.  
619 Res.* **2010**, *132*, 520–522.  
620 [31] K. R. Downum, *New Phytol.* **1992**, *122*, 401–420.  
621 [32] G. H. N. Towers, in *Prog. Phytochem.*, **1980**, pp. 183–200.  
622 [33] A. Garrard, in *Encycl. Toxicol. Third Ed.* (Ed.: P. Wexler), Academic Press, Oxford,  
623 **2014**, pp. 1052–1054.

- 624 [34] G. Guillet, J. Harmatha, T. G. Waddell, B. J. Philogène, J. T. Arnason, *Photochem.*  
625 *Photobiol.* **2000**, *71*, 111–115.
- 626 [35] J. C. Quinn, A. Kessell, L. A. Weston, *Int. J. Mol. Sci.* **2014**, *15*, 1441–1465.
- 627 [36] V. O. de Araújo, T. S. Oliveira Neto, S. V. D. Simões, T. K. F. da Silva, F. Riet-  
628 Correa, R. B. Lucena, *Toxicon* **2017**, *138*, 184–187.
- 629 [37] D. L. Lelis, L. N. Rennó, M. L. Chizzotti, C. E. R. Pereira, J. C. P. Silva, L. G. T.  
630 Moreira, F. B. P. Carvalho, F. H. M. Chizzotti, *Small Rumin. Res.* **2018**, *169*, 24–28.
- 631 [38] C. H. Tokarnia, J. Döbereiner, P. V. Peixoto, *Toxicon* **2002**, *40*, 1635–1660.
- 632 [39] A. L. Schild, A. C. Motta, F. Riet-Correa, F. C. Karam, F. B. Grecco, in *Poisonous*  
633 *Plants Relat. Toxins* (Eds.: T. Acamovic, C.S. Stewart, T.W. Pennycott), CABI,  
634 Wallingford, **2004**, pp. 162–166.
- 635 [40] A. E. Johnson, *J. Natl. Cancer Inst.* **1982**, *69*, 253–258.
- 636 [41] B. Siewert, H. Stuppner, *Phytomedicine* **2019**, *60*, 152985.
- 637 [42] P. Ferreira da Silva, L. Paulo, A. Barbaflava, F. Elisei, F. H. Quina, A. L. Maçanita,  
638 *Chem. – Eur. J.* **2012**, *18*, 3736–3744.
- 639 [43] M. A. Guerrero-Rubio, J. Escribano, F. García-Carmona, F. Gandía-Herrero, *Trends*  
640 *Plant Sci.* **2020**, *25*, 159–175.
- 641 [44] S. J. Galitzer, F. W. Oehme, *Vet. Sci. Commun.* **1978**, *2*, 217–230.
- 642 [45] M. J. M. Christenhusz, J. W. Byng, *Phytotaxa* **2016**, *261*, 201–217.
- 643 [46] C. Y. Ong, S. K. Ling, R. M. Ali, C. F. Chee, Z. A. Samah, A. S. H. Ho, S. H. Teo, H.  
644 B. Lee, *J. Photochem. Photobiol. B* **2009**, *96*, 216–222.
- 645 [47] E. Foresto, P. Gilardi, L. E. Ibarra, I. S. Cogno, *Phytomedicine Plus* **2021**, *1*, 100044.
- 646 [48] K. da Silva Souza Campanholi, J. M. Jaski, R. C. da Silva Junior, A. B. Zanqui, D.  
647 Lazarin-Bidóia, C. M. da Silva, E. A. da Silva, N. Hioka, C. V. Nakamura, L. Cardozo-  
648 Filho, W. Caetano, *J. Photochem. Photobiol. B* **2020**, *203*, 111763.
- 649 [49] A. Gasparetto, T. F. Lapinski, S. R. Zamuner, S. Khouri, L. P. Alves, E. Munin, M. J.  
650 Salvador, *J. Photochem. Photobiol. B* **2010**, *99*, 15–20.
- 651 [50] M. G. Collett, *Toxicon X* **2019**, *3*, 100012.
- 652 [51] M. G. Collett, S. M. Taylor, *Toxicon* **2019**, *167*, 172–173.
- 653 [52] D. S. Santos, C. C. B. Silva, V. O. Araújo, M. de Fátima Souza, P. B. Lacerda-Lucena,  
654 S. V. D. Simões, F. Riet-Correa, R. B. Lucena, *Toxicon* **2017**, *125*, 65–69.
- 655 [53] S. N. R. Knupp, C. C. Borburema, T. S. Oliveira Neto, R. de Medeiros, L. S. Knupp, F.  
656 Riet-Correa, R. B. Lucena, *Pesqui. Veterinária Bras.* **2014**, *34*, 1191–1195.
- 657 [54] A. Rashmir-Raven, R. S. McConnico, in *Robinsons Curr. Ther. Equine Med. Seventh*  
658 *Ed.* (Eds.: K.A. Sprayberry, N.E. Robinson), W.B. Saunders, St. Louis, **2015**, pp. 536–  
659 542.
- 660 [55] A. Zink, J. Ring, *N. Engl. J. Med.* **2014**, *371*, 559.
- 661 [56] J. P. Niemeyer-Corbellini, O. Lupi, L. Klotz, L. Montelo, D. M. Elston, V. Haddad, S.  
662 K. Tyring, in *Trop. Dermatol. Second Ed.* (Eds.: S.K. Tyring, O. Lupi, U.R. Hengge),  
663 Elsevier, **2017**, pp. 443–470.
- 664 [57] D. E. J. G. J. Dolmans, D. Fukumura, R. K. Jain, *Nat. Rev. Cancer* **2003**, *3*, 380–387.
- 665 [58] J. G. B. Derraik, *N. Z. Med. J.* **2007**, *120*.
- 666 [59] A. L. N. Formiga Filho, V. S. M. Carneiro, E. A. Souza, R. L. Santos, M. H. C. V.  
667 Catão, A. C. D. Medeiros, *Photomed. Laser Surg.* **2015**, *33*, 240–245.
- 668 [60] N. Luiza Andreazza, C. Vevert-Bizet, G. Bourg-Heckly, F. Sureau, M. José Salvador,  
669 S. Bonneau, *Int. J. Pharm.* **2016**, *510*, 240–249.
- 670 [61] N. L. Andreazza, C. Caramano de Lourenço, Á. J. Hernandez-Tasco, M. L. B.  
671 Pinheiro, M. É. Alves Stefanello, E. Vilaça Costa, M. J. Salvador, *J. Photochem.*  
672 *Photobiol. B* **2016**, *160*, 154–162.

- 673 [62] A. R. dos Santos, C. Pires, F. A. Marques, A. Q. Lobão, B. H. L. N. S. Maia, *Biochem.*  
674 *Syst. Ecol.* **2017**, *73*, 1–2.
- 675 [63] R. D. Machado, M. C. de Moraes, E. C. da Conceição, B. G. Vaz, A. R. Chaves, K. R.  
676 Rezende, *J. Pharm. Biomed. Anal.* **2020**, *191*, 113593.
- 677 [64] B. L. Stegelmeier, *Clin. Tech. Equine Pract.* **2002**, *1*, 81–88.
- 678 [65] E. M. Nagy, L. Dalla Via, L. Ronconi, D. Fregona, *Curr. Pharm. Des.* **2010**, *16*, 1863–  
679 1876.
- 680 [66] J. A. D. Río, L. Díaz, D. García-Bernal, M. Blanquer, A. Ortúño, E. Correal, J. M.  
681 Moraleda, in *Stud. Nat. Prod. Chem.* (Ed.: Atta-ur-Rahman), Elsevier, **2014**, pp. 145–  
682 195.
- 683 [67] L. P. Christensen, in *Polyphenols Prev. Treat. Hum. Dis. Second Ed.* (Eds.: R.R.  
684 Watson, V.R. Preedy, S. Zibadi), Academic Press, **2018**, pp. 349–384.
- 685 [68] A. Miron, L. Nahar, E. Gille, S. D. Sarker, in *Annu. Rep. Med. Chem.* (Eds.: S.D.  
686 Sarker, L. Nahar), Academic Press, **2020**, pp. 297–325.
- 687 [69] A. A. Abu-Hashem, M. El-Shazly, *Eur. J. Med. Chem.* **2015**, *90*, 633–665.
- 688 [70] A. S. Bhagavathula, A. J. M. Al-Khatib, A. A. Elnour, N. M. AlKalbani, A. Shehab,  
689 *Pharmacogn. Res.* **2015**, *7*, 397.
- 690 [71] N. Khalil, M. Bishr, S. Desouky, O. Salama, *Molecules* **2020**, *25*, 301.
- 691 [72] K. Kim, Y. Kawano, Y. Yamazaki, *Anticancer Res.* **2003**, *23*, 2355–61.
- 692 [73] W. W. Jong, P. J. Tan, F. A. Kamarulzaman, M. Mejin, D. Lim, I. Ang, M. Naming, T.  
693 C. Yeo, A. S. H. Ho, S. H. Teo, H. B. Lee, *Chem. Biodivers.* **2013**, *10*, 1475–1486.
- 694 [74] S. K. Tyring, O. Lupi, U. R. Hengge, Eds. , in *Trop. Dermatol.*, Churchill Livingstone,  
695 Edinburgh, **2006**, pp. 439–467.
- 696 [75] V. Monge-Fuentes, L. A. Muehlmann, J. P. F. Longo, J. R. Silva, M. L. Fascineli, P. de  
697 Souza, F. Faria, I. A. Degterev, A. Rodriguez, F. P. Carneiro, C. M. Lucci, P. Escobar,  
698 R. F. B. Amorim, R. B. Azevedo, *J. Photochem. Photobiol. B* **2017**, *166*, 301–310.
- 699 [76] S. K. Kandagatla, R. T. Uhl, T. N. Graf, N. H. Oberlies, G. M. Raner, *Nat. Prod.*  
700 *Commun.* **2019**, *14*, DOI 10.1177/1934578x19852443.
- 701 [77] M. Akhdar, R. Abedini, S. Tavakolpour, Z. Gholibeigian, A. Azizpour, *J. Herb. Med.*  
702 **2021**, *28*, 100442.
- 703 [78] W. Ma, Y. Lee, K. H. Row, *Sep. Sci. Technol. Phila.* **2017**, *52*, 2547–2556.
- 704 [79] J. Tamariz, E. Burgueño-Tapia, M. A. Vázquez, F. Delgado, in *Alkaloids Chem. Biol.*  
705 (Ed.: H.-J. Knölker), Academic Press, **2018**, pp. 1–314.
- 706 [80] E. L. Camm, G. H. N. Towers, J. C. Mitchell, *Phytochemistry* **1975**, *14*, 2007–2011.
- 707 [81] B. R. Scott, M. A. Pathak, G. R. Mohn, *Mutat. Res.* **1976**, *39*, 29–74.
- 708 [82] G. N. Towers, T. Arnason, C. K. Wat, E. A. Graham, J. Lam, J. L. Mitchell, *Contact*  
709 *Dermatitis* **1979**, *5*, 140–144.
- 710 [83] *Fitoterapia* **2006**, *77*, 194–198.
- 711 [84] J. P. Niemeyer-Corbellini, O. Lupi, L. Klotz, L. Montelo, D. M. Elston, V. Haddad, S.  
712 K. Tyring, in *Trop. Dermatol. Second Ed.* (Eds.: S.K. Tyring, O. Lupi, U.R. Hengge),  
713 Elsevier, **2017**, pp. 443–470.
- 714 [85] M. Machado, M. P. Miguel, J. P. Terra, J. A. Ferreira, F. Riet-Correa, M. B. de Castro,  
715 *Toxicon* **2019**, *168*, 122–125.
- 716 [86] Z. Zhou, L. Zhang, Z. Zhang, Z. Liu, *Asian J. Pharm. Sci.* **2021**, DOI  
717 10.1016/j.ajps.2020.12.003.
- 718 [87] A. T. de Carvalho, M. M. Paes, M. S. Cunha, G. C. Brandão, A. M. Mapeli, V. C.  
719 Rescia, S. A. Oesterreich, G. R. Villas-Boas, *Molecules* **2020**, *25*, DOI  
720 10.3390/molecules25173879.
- 721 [88] T. R. Govindachari, P. S. Kalyanaraman, N. Muthukumaraswamy, B. R. Pai,  
722 *Tetrahedron* **1971**, *27*, 3919–3926.

- 723 [89] B. Ovalle-Magallanes, D. Eugenio-Pérez, J. Pedraza-Chaverri, *Food Chem. Toxicol.*  
724 *Int. J. Publ. Br. Ind. Biol. Res. Assoc.* **2017**, *109*, 102–122.
- 725 [90] H. A. Maddah, V. Berry, S. K. Behura, *Renew. Sustain. Energy Rev.* **2020**, *121*,  
726 109678.
- 727 [91] L. P. Meagher, B. L. Smith, A. L. Wilkins, *Anim. Feed Sci. Technol.* **2001**, *91*, 157–  
728 170.
- 729 [92] P. Zarrintaj, Z. Ahmadi, M. Hosseinnezhad, M. R. Saeb, P. Laheurte, M. Mozafari,  
730 *Mater. Today Proc.* **2018**, *5*, 15836–15844.
- 731 [93] M. F. F. Mendonça, L. A. Pimentel, P. V. Leal, J. C. Oliveira Filho, L. G. Caymmi, A.  
732 W. O. Silva, R. S. Jesus, T. C. Peixoto, *Toxicon* **2021**, *193*, 13–20.
- 733 [94] S. S. Garg, J. Gupta, S. Sharma, D. Sahu, *Eur. J. Pharm. Sci.* **2020**, *152*, 105424.
- 734 [95] T. Acamovic, C. S. Stewart, T. W. Pennycott, Eds., *Poisonous Plants and Related*  
735 *Toxins*, CABI Pub, Wallingford, Oxon, UK ; Cambridge, MA, USA, **2004**.
- 736 [96] H. A. Maddah, V. Berry, S. K. Behura, *Renew. Sustain. Energy Rev.* **2020**, *121*,  
737 109678.
- 738 [97] S. A. de N. M. Silva, A. B. Barros, J. M. T. Souza, A. F. Moura, A. R. de Araújo, M.  
739 G. A. Mendes, T. C. Daboit, D. A. da Silva, A. J. Araújo, J. D. B. Marinho Filho,  
740 *Phytochem. Lett.* **2020**, *39*, 173–181.
- 741 [98] C. C. Guizelini, R. C. Pupin, J. A. C. Arredondo, L. L. Robalinho, R. N. Bernardo, W.  
742 J. Silva, G. L. Franco, R. A. A. Lemos, *Toxicon* **2021**, *194*, 11–16.
- 743 [99] I. R. C. Bick, A. J. Blackman, *Aust. J. Chem.* **1973**, *26*, 1377–1380.
- 744 [100] R. Thomas, *Pure Appl. Chem.* **1973**, *34*, 515–528.
- 745 [101] J. M. Edwards, U. Weiss, *Phytochemistry* **1970**, *9*, 1653–1657.
- 746 [102] J. M. Kornfeld, J. M. Edwards, *BBA - Gen. Subj.* **1972**, *286*, 88–90.
- 747 [103] A. Kamuhabwa, P. Agostinis, B. Ahmed, W. Landuyt, B. van Cleynenbreugel, H. van  
748 Poppel, P. de Witte, *Photochem. Photobiol. Sci. Off. J. Eur. Photochem. Assoc. Eur. Soc. Photobiol.* **2004**, *3*, 772–780.
- 749 [104] D. Kacerovská, K. Pizinger, F. Majer, F. Smíd, *Photochem. Photobiol.* **2008**, *84*, 779–  
750 785.
- 751 [105] M. E. Daub, S. Herrero, K.-R. Chung, *Antioxid. Redox Signal.* **2013**, *19*, 970–989.
- 752 [106] B. Mansoori, A. Mohammadi, M. Amin Doustvandi, F. Mohammadnejad, F. Kamari,  
753 M. F. Gjerstorff, B. Baradaran, M. R. Hamblin, *Photodiagnosis Photodyn. Ther.* **2019**,  
754 26, 395–404.
- 755 [107] C. de Menezes Furtado, F. S. E. D. V. de Faria, R. B. Azevedo, K. Py-Daniel, A. L.  
756 dos Santos Camara, J. R. da Silva, E. de Holanda Oliveira, A. F. R. Rodriguez, I. A.  
757 Degterev, *J. Photochem. Photobiol. B* **2017**, *167*, 242–248.
- 758 [108] M. S. Marqués-Calvo, F. Codony, G. Agustí, C. Lahera, *Photodiagnosis Photodyn.*  
759 *Ther.* **2017**, *17*, 180–184.
- 760 [109] P. Zarrintaj, Z. Ahmadi, M. Hosseinnezhad, M. R. Saeb, P. Laheurte, M. Mozafari,  
761 *Mater. Today Proc.* **2018**, *5*, 15836–15844.
- 762 [110] H. D. de Menezes, A. C. Pereira, G. T. P. Brancini, H. C. de Leão, N. S. Massola  
763 Júnior, L. Bachmann, M. Wainwright, J. K. Bastos, G. U. L. Braga, *J. Photochem.*  
764 *Photobiol. B* **2014**, *131*, 74–83.
- 765 [111] C.-C. Chiang, M.-J. Cheng, C.-F. Peng, H.-Y. Huang, I.-S. Chen, *Chem. Biodivers.*  
766 **2010**, *7*, 1728–1736.
- 767 [112] D. P. dos Santos, D. P. S. Lopes, S. P. de Melo Calado, C. V. Gonçalves, I. P. R.  
768 Muniz, I. S. Ribeiro, M. P. L. Galantini, R. A. A. da Silva, *J. Photochem. Photobiol. B*  
769 **2019**, *191*, 107–115.
- 770 [113] M. Schulz, S. K. T. Seraglio, P. Brugnerotto, L. V. Gonzaga, A. C. O. Costa, R. Fett,  
771 *Food Res. Int.* **2020**, *137*, 109744.

- 773 [114] K. V. Muniandy, E. L. T. Chung, M. S. Jaapar, M. H. M. Hamdan, A. Salleh, F. F. A.  
774 Jesse, *Toxicon* **2020**, *174*, 26–31.
- 775 [115] A. Warowicka, Ł. Popenda, G. Bartkowiak, O. Musidlak, J. Litowczenko-Cybulska, D.  
776 Kuźma, R. Nawrot, S. Jurga, A. Goździcka-Józefiak, *Phytomedicine* **2019**, *64*, 152919.
- 777 [116] C. H. S. De Oliveira, J. D. Barbosa, C. M. C. Oliveira, E. Bastianetto, M. M. Melo, M.  
778 Haraguchi, L. G. L. Freitas, M. X. Silva, R. C. Leite, *Toxicon* **2013**, *73*, 121–129.
- 779 [117] G. M. L. Diamantino, F. Pierezan, M. I. C. Ferreira, W. S. D. Rocha, V. M. O. Veiga,  
780 C. E. Martins, M. O. Veiga, B. Soto-Blanco, *Toxicon* **2020**, *185*, 1–4.
- 781 [118] J.-J. Li, Y.-X. Li, N. Li, H.-T. Zhu, D. Wang, Y.-J. Zhang, *Nat. Prod. Bioprospecting*  
782 **2022**, *12*, DOI 10.1007/s13659-022-00346-z.
- 783 [119] L. Buchalter, *J. Pharm. Sci.* **1969**, *58*, 904–904.
- 784 [120] R. S. Gonçalves, E. L. Silva, N. Hioka, C. V. Nakamura, M. L. Bruschi, W. Caetano,  
785 *Nat. Prod. Res.* **2018**, *32*, 366–369.
- 786 [121] J. D. Spikes, *J. Photochem. Photobiol. B* **1998**, *42*, 1–11.
- 787 [122] J. A. Dimmer, S. C. Núñez Montoya, C. S. Mendoza, J. L. Cabrera, *Phytochemistry*  
788 **2017**, *137*, 94–100.
- 789 [123] S. C. Núñez Montoya, L. R. Comini, M. Sarmiento, C. Becerra, I. Albesa, G. A.  
790 Argüello, J. L. Cabrera, *J. Photochem. Photobiol. B* **2005**, *78*, 77–83.
- 791 [124] S. C. Núñez Montoya, L. R. Comini, B. Rumie Vittar, I. M. Fernández, V. A. Rivarola,  
792 J. L. Cabrera, *Toxicon* **2008**, *51*, 1409–1415.
- 793 [125] L. R. Comini, S. C. Núñez Montoya, M. Sarmiento, J. L. Cabrera, G. A. Argüello, *J.*  
794 *Photochem. Photobiol. Chem.* **2007**, *188*, 185–191.
- 795 [126] L. R. Comini, I. M. Fernandez, N. B. R. Vittar, S. C. Núñez Montoya, J. L. Cabrera, V.  
796 A. Rivarola, *Phytomedicine* **2011**, *18*, 1093–1095.
- 797 [127] L. R. Comini, S. C. Núñez Montoya, P. L. Páez, G. A. Argüello, I. Albesa, J. L.  
798 Cabrera, *J. Photochem. Photobiol. B* **2011**, *102*, 108–114.
- 799 [128] N. B. Rumie Vittar, L. Comini, I. M. Fernandez, E. Agostini, S. Nuñez-Montoya, J. L.  
800 Cabrera, V. A. Rivarola, *Photodiagnosis Photodyn. Ther.* **2014**, *11*, 182–192.
- 801 [129] I. S. Cogno, P. Gilardi, L. Comini, S. C. Núñez-Montoya, J. L. Cabrera, V. A.  
802 Rivarola, *Photodiagnosis Photodyn. Ther.* **2020**, *31*, 101852.
- 803 [130] M.-J. Liou, P.-L. Wu, T.-S. Wu, *Chem. Pharm. Bull. (Tokyo)* **2002**, *50*, 276–279.
- 804 [131] D. L. Luthria, V. Ramakrishnan, G. S. Verma, B. R. Prabhu, A. Banerji, *J. Agric. Food  
805 Chem.* **1989**, *37*, 1435–1437.
- 806 [132] P. Basnet, S. Kadota, K. Manandhar, M. D. Manandhar, T. Namba, *Planta Med.* **1993**,  
807 *59*, 384–386.
- 808 [133] W. A. Kukula-Koch, J. Widelski, in *Pharmacognosy* (Eds.: S. Badal, R. Delgoda),  
809 Academic Press, Boston, **2017**, pp. 163–198.
- 810 [134] M. H. Shahrajabian, *Curr. Rev. Clin. Exp. Pharmacol.* **2022**, DOI  
811 10.2174/2772432817666220510143902.
- 812 [135] Y. Chen, B. Yang, J. Xu, T. Zheng, H. Fan, G.-Z. Yang, *Yao Xue Xue Bao* **2012**, *47*,  
813 1646–1652.
- 814 [136] A. T. Coimbra, S. Ferreira, A. P. Duarte, *J. Ethnopharmacol.* **2020**, *260*, 113076.
- 815 [137] O. Schimmer, I. Kühne, *Mutat. Res.* **1990**, *243*, 57–62.
- 816 [138] D. Wessner, H. Hofmann, J. Ring, *Contact Dermatitis* **1999**, *41*, 232–232.
- 817 [139] S. Arias-Santiago, M. Fernández-Pugnaire, F. Almazán-Fernández, C. Serrano-Falcón,  
818 S. Serrano-Ortega, *Rheumatol. Oxf. Engl.* **2009**, *48*, 1401.
- 819 [140] Ernest H. Oertli, Ross C. beier, G.Wayne Ivie and Loyd D.Rowe, *Phytochemistry*  
820 **1984**, *23*, 439–441.
- 821 [141] P. K. Mukherjee, in *Qual. Control Eval. Herb. Drugs* (Ed.: P.K. Mukherjee), Elsevier,  
822 **2019**, pp. 237–328.

- 823 [142] S. Wang, F. Zhu, *Trends Food Sci. Technol.* **2020**, *95*, 45–58.  
824 [143] G. Calogero, G. D. Marco, *Sol. Energy Mater. Sol. Cells* **2008**, *92*, 1341–1346.  
825 [144] R. Zhao, G. Zheng, L. Fan, Z. Shen, K. Jiang, Y. Guo, J.-W. Shao, *Acta Biomater.*  
826 **2018**, *70*, 197–210.  
827 [145] L. D. Dias, K. C. Blanco, I. S. Mfouo-Tynga, N. M. Inada, V. S. Bagnato, *J.  
828 Photochem. Photobiol. C Photochem. Rev.* **2020**, *45*, 100384.  
829 [146] Z. Luksiene, in *Innov. Food Process. Technol.* (Eds.: K. Knoerzer, K.  
830 Muthukumarappan), Elsevier, Oxford, **2021**, pp. 368–384.  
831 [147] C. Wei, F. Zhang, L. Song, X. Chen, X. Meng, *Food Control* **2021**, *123*, 107683.  
832 [148] R. W. Redmond, J. N. Gamlin, *Photochem. Photobiol.* **1999**, *70*, 391–475.  
833 [149] A. Vasas, O. Orbán-Gyapai, J. Hohmann, *J. Ethnopharmacol.* **2015**, *175*, 198–228.  
834 [150] M. L. Bruschi, J. B. da Silva, H. C. Rosseto, *Curr. Pharm. Des.* **2019**, *25*, 2279–2291.  
835 [151] J. Reisch, K. Szendrei, E. Minker, I. Novák, *Experientia* **1968**, *24*, 992.  
836 [152] J. R. S. Hoult, M. Payá, *Gen. Pharmacol. Vasc. Syst.* **1996**, *27*, 713–722.  
837 [153] P. Ainiwaer, M. Nueraihemaiti, Z. Li, D. Zang, L. Jiang, Y. Li, H. A. Aisa, *Fitoterapia*  
838 **2022**, *156*, DOI 10.1016/j.fitote.2021.105094.  
839 [154] A. G. González, R. Estévez, I. Jaraiz, *Phytochemistry* **1971**, *10*, 1621–1626.  
840 [155] B. I. Pia Schiavone, L. Verotta, A. Rosato, M. Marilena, S. Gibbons, E. Bombardelli,  
841 C. Franchini, F. Corbo, *Anticancer Agents Med. Chem.* **2014**, *14*, 1397–1401.  
842 [156] L. Nahar, S. Al-Majmaie, A. Al-Groshi, A. Rasul, S. D. Sarker, *Molecules* **2021**, *26*,  
843 1609.  
844 [157] E. Foresto, P. Gilardi, L. E. Ibarra, I. S. Cogno, *Phytomedicine Plus* **2021**, *1*, 100044.  
845 [158] N. L. Andreazza, C. Caramano de Lourenço, Á. J. Hernandez-Tasco, M. L. B.  
846 Pinheiro, M. É. Alves Stefanello, E. Vilaça Costa, M. J. Salvador, *J. Photochem.  
847 Photobiol. B* **2016**, *160*, 154–162.  
848 [159] T. Munde, S. Brand, W. Hidalgo, R. K. Maddula, A. Svatoš, B. Schneider,  
849 *Phytochemistry* **2013**, *91*, 165–176.  
850 [160] T. Arnason, C.-K. Wat, K. Downum, E. Yamamoto, E. Graham, G. H. N. Towers,  
851 *Can. J. Microbiol.* **1980**, *26*, 698–705.  
852 [161] L. P. Meagher, B. L. Smith, A. L. Wilkins, *Anim. Feed Sci. Technol.* **2001**, *91*, 157–  
853 170.  
854 [162] G. F. Bath, E. M. van Tonder, P. A. Basson, *J. S. Afr. Vet. Assoc.* **1978**, *49*, 23–25.  
855 [163] S. T. Cargnin, S. B. Gnoatto, *Food Chem.* **2017**, *220*, 477–489.  
856 [164] L. F. Stakhov, V. G. Ladygin, L. N. Stakhova, *Biol. Bull.* **2005**, *32*, 121–127.  
857 [165] B. Legouin, F. Lohézic-Le Dévéhat, S. Ferron, I. Rouaud, P. Le Pogam, L. Cornevin,  
858 M. Bertrand, J. Boustie, *Mol. Basel Switz.* **2017**, *22*, DOI 10.3390/molecules22071162.  
859 [166] B. Siewert, P. Vrabl, F. Hammerle, I. Bingger, H. Stuppner, *RSC Adv.* **2019**, *9*, 4545–  
860 4552.  
861 [167] B. Siewert, *Photochem. Photobiol. Sci.* **2021**, *20*, 475–488.  
862 [168] F. Chemat, M. A. Vian, G. Cravotto, *Int. J. Mol. Sci.* **2012**, *13*, 8615–8627.  
863 [169] R. Gonçalves de Oliveira-Júnior, R. Grougnet, P.-E. Bodet, A. Bonnet, E. Nicolau, A.  
864 Jebali, J. Rumin, L. Picot, *Algal Res.* **2020**, *51*, DOI 10.1016/j.algal.2020.102035.  
865 [170] T. Kowalczyk, A. Merecz-Sadowska, L. Picot, I. Brčić Karačonji, J. Wieczfinska, T.  
866 Śliwiński, P. Sitarek, *Molecules* **2022**, *27*, 795.

867