



HAL
open science

Better together? A mediation analysis of general practitioners' performance in multi-professional group practice

Anna Zaytseva, Pierre Verger, Bruno Ventelou

► **To cite this version:**

Anna Zaytseva, Pierre Verger, Bruno Ventelou. Better together? A mediation analysis of general practitioners' performance in multi-professional group practice. 2023. hal-04305092

HAL Id: hal-04305092

<https://hal.science/hal-04305092>

Preprint submitted on 24 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Better together?
**A mediation analysis of general practitioners' performance
in multi-professional group practice**

**Anna Zaytseva
Pierre Verger
Bruno Ventelou**

WP 2023 - Nr 25

Better together? A mediation analysis of general practitioners' performance in multi-professional group practice

Anna Zaytseva^{1,2}, Pierre Verger³, and Bruno Ventelou²

¹SBS-EM, DULBEA, Université libre de Bruxelles, Brussels Belgium

²Aix-Marseille University, CNRS, AMSE, Marseille France

³Observatoire Régional de la Santé Provence-Alpes-Côte d'Azur, Southeastern Regional Health Observatory, Marseille France

November 17, 2023

Abstract

Objectives: To analyse how general practitioners (GPs) respond to insufficient GP supply in their practice area in terms of quantity and quality of care, and how this response can be mediated by enrolment in integrated primary care teams (multi-professional group practices (MGP)).

Methods: We used three representative cross-sectional surveys (2019-2020) of 1,209 French GPs. Using structural equations, we assumed that low GP density influences GPs' work-related stress (mediator 1) as well as their use of e-health tools (mediator 2) and ultimately quantity and quality of care. Quantity (respectively quality) of care were approximated by demand absorption capacities (respectively frequencies of vaccine recommendations). We estimated an additional specification where enrolment in an MGP was a mediator between GP density and the two mediators defined above.

Results: GP density was significantly and positively associated with work-related stress, which was consecutively associated with deteriorated demand absorption capacity. Higher use of e-health tools was associated with greater involvement in vaccine recommendations. Lastly, GPs in MGP tend to use more e-health tools than those practicing outside MGP, with a favourable effect on quality of care.

Discussion: This study demonstrates that a lower level of work-related stress is the key mediator in handling patients' requests. Correcting for the self-selection into MGP, we amend some unstable results contained in the literature: there is no significant mediation effect of enrolment in integrated primary care teams on the quantity of care, but rather an effect on the quality of care. Although probably disappointing for the quantity of care provided, our results pinpoint a novel added value of enrolment in an integrated practice as a response to decreasing GP density.

Keywords: general practitioners, medically underserved area, integrated care, France

JEL Keywords: I11, I18

We thank our panel of GPs as well as the bureau de professions de santé of DREES, French Ministry of Health for

1 Introduction

Primary care remains an important concern for the policymakers, as emphasized by the 2018 Astana declaration following the Global Conference on Primary Health Care, which celebrated the fortieth anniversary of the Declaration of Alma-Ata (WHO, 2018). Resulting from this meeting, the 2019 Global Monitoring Report on progress on universal health coverage (WHO, 2019) indicates that reorganization of care provision, and especially the improvement of primary care coordination is one of the key strategies to tackle the inequalities in access to primary care and achieve universal health coverage.

As other developed countries, France is currently facing a shortage of primary care workforce, due to mass retirement of self-employed general practitioners (GPs) that are ensuring more than 90% of primary care associated with population growth and population aging as well as lack of anticipation of this demographic evolution (Anguis et al., 2021). According to the French Ministry of Health, the number of GPs is supposed to decline at least up to 2026. Once adjusted for healthcare demand, the 2021 GP density level should be attained once again by 2036.

In the meantime, a clear trend towards integration of primary care providers is observed: in a 2010 survey, 54% of GPs indicated practicing in a group practice; there were already 61% in 2019 (Chaput et al., 2019). However, most of these group practices only share premises, and/or back-office functions.

Up to now, the most advanced form of integrated healthcare teams in France is the so-called multi-professional primary care group practice (MGP) [in French: “maison de santé pluridisciplinaire”]. The MGPs are created on a voluntary basis and are required to have at least two full-time equivalent GPs and one paramedic. This policy was launched in 2007 and has proven to be extremely popular among physicians: in 2020, more than 1,300 MGPs were actively operating compared to less than 20 in 2008 (Chevillard and Mousquès, 2021).

Current literature mostly demonstrates improvements in the quantity of care provided in integrated healthcare teams, e.g., longer patient lists and more home and office visits (Cassou et al., 2020; Kerrissey et al., 2017; Lalani et al., 2020; Mousquès and Daniel, 2015; Singer et al., 2020). However, evidence is mixed and scarce regarding the gains related to integrated practices in terms of quality of care.

This paper contributes to this literature exploring the impact of existing variability in GPs’ accessibility both on quantity and quality of care using data from a representative panel of 1,209 French private practice GPs. In addition, we sought to provide evidence on the efficiency of the MGP policy by estimating the mediation effect of MGP enrolment on quantity and quality of primary care provided. To our knowledge,

their continuous and valuable support throughout the data collection process. We warmly thank Alain Paraponaris, Anne-Laure Samson, Anthony Scott, Paul Frappé, Julien Mousquès, as well as the participants of the 21st *Journées Louis-André Gérard-Varet* (LAGV) conference for their helpful comments.

The project leading to this publication has received funding from the French government under the "France 2030" investment plan managed by the French National Research Agency (reference: ANR-17- EURE-0020) and from Excellence Initiative of Aix-Marseille University - A*MIDEX as well as from French Institute for Public Health Research grant (reference IReSP-LIAAP18-HSR-008).

A.Z. has benefitted a PhD grant from the (non-profit) Sud Provence-Alpes-Côte d’Azur regional council during the research for this article.

this is the first study to analyse the effects of GP density on quantity and quality of primary care using mediation analysis, which allows to disentangle indirect and direct effects of GP density and the real effect of the MGP policy. To preface our results, we provide limited evidence on the efficiency of the MGP policy – at least the way it has been implemented in France.

2 Conceptual framework

We start by arguing that low GP density might hinder both the quality and the quantity of care provided. This section presents the conceptual framework (based on relevant literature) that sheds light on the potential mechanisms, both direct and indirect, that might explain this relationship.

Extensive literature documents higher workload for GPs practicing in rural or remote areas (Groenewegen et al., 2020; Hassel et al., 2019; Lurquin et al., 2021; Steinhäuser et al., 2011). In addition, higher workload might result in shorter consultation length (Deveugele et al., 2002; Irving et al., 2017). Growing literature points out work-related stress as one of the important mediators of the GP productivity (Soler et al., 2008; Wallace et al., 2009). According to a recent French study in the Upper Normandy region, GP density below 7.75 per 10,000 inhabitants is positively correlated with all of three dimensions (emotional exhaustion, depersonalization, and personal accomplishment) of Maslach Burnout Inventory (Picquendar et al., 2019; Maslach et al., 1997). Wallace et al. (2009) also argue that lack of GPs is one of the reasons for their professional burnout. Overall, GPs have been found to have relatively high levels of burnout: two recent literature reviews indicate that the burnout incidence is around a third among the GPs (Karuna et al., 2021; Shen et al., 2022). Higher levels of burnout are in turn associated with suboptimal productivity (Shen et al., 2022; Wallace et al., 2009). Recent studies that took place during the Covid-19 pandemic indicate that work-related stress among healthcare professionals was entirely due to subjective factors, i.e., the perceptions of their working conditions (Spányik et al., 2022).

Empirically, practice in underserved areas is frequently associated with worse patients' health outcomes obtained from the use of primary care services, especially regarding time-consuming activities such as prevention (Lane et al., 2021; Silhol et al., 2020; Zyzanski et al., 1998). This is precisely the case for vaccination activity, in which GPs may play a key role in many countries.

In France, vaccination coverage remains suboptimal, especially against seasonal influenza (Robert et al., 2020) and human papillomavirus (Eliès et al., 2022), while almost half of the population is vaccine-hesitant (Larson et al., 2016; Rey et al., 2018). Moreover, a quarter of GPs declare that some of the recommended vaccines are not useful (Verger et al., 2015).

While the evidence is mixed, expanding literature suggests that e-health solutions, e.g., use of electronic health records combined with automated reminders, might be a convenient tool to improve the vaccination rates from 5 up to 20 percentage points (Derrough et al., 2017; Dumit et al., 2018; Groom et al., 2015; Jacobson Vann et al., 2018). For example, an US study indicates that introduction of automated reminders within the electronic health record significantly improves the pertussis postpartum immunization rate

(Morgan et al., 2015). Two recent literature reviews reveal greater levels of HPV vaccine initiation and completion using communication technologies, including electronic health records (Francis et al., 2017; Niccolai and Hansen, 2015). Another review (Okoli et al., 2021) pinpoints an important increase (19%) in seasonal influenza in adult patients and a smaller, but still significant, increase in paediatric patients associated with the use of e-health tools.


Greater use of e-health tools is also promoted to bridge inequalities in access to primary care (Petrie et al., 2021; Wakerman and Humphreys, 2011). The implementation of e-health technologies, and especially, telemedicine has been even more encouraged since the beginning of Covid-19 pandemic, both by the patients and the health authorities (Davin-Casalena et al., 2021; Woodall et al., 2021). In addition, use of e-health tools appears cost-saving due to better administrative efficiency (Patel et al., 2015).

Finally, interprofessional collaboration is supposed to deal with both the quantity and the quality of care provided. On the one hand, it is beneficial to the GPs wellbeing and is supposed to reduce their workload due to better practice organization (Kerrissey et al., 2017; Singer et al., 2020). On the other hand, better coordination and proximity with specialists and other health professionals allows to improve the quality of care, especially for the patients with complex needs (Colla et al., 2020; De Sutter et al., 2019; Singer et al., 2020; Smith et al., 2021). These points suggest that the practice in an integrated care setting is an 'upstream' (opening) determinant of both the quantity and quality of care. Nonetheless, GPs might have diverse attitudes towards cooperation and prescriptions management with specialists and other health professionals (Carrier et al., 2022).

Following this theoretical framework, we were inclined to test the following chain of pathways (Figure 1): low density leads to (1) increase in work-related stress that influences GPs' productivity; and (2) higher use of e-health tools that in turn affect the quality of care provided. We then add (3) enrollment in MGP as an 'opening' mediator that influences both the other two mediators and the outcome variables.

This is the theoretical framework which will be tested by using an appropriate statistical analysis (see next section).

Figure 1: The direct and indirect effects between GP density and quantity and quality of care



Latent variables (factors) are represented by circles, and measured variables (items) by rectangles; arrows point in the hypothesized direction of influence. The total effect of low GP density on quantity (DV1) and quality of care (DV2) is represented by dashed arrows. The effect of low GP density on each potential mediator is represented by paths δ_1 , γ_1 and η_1 . The effect of potential mediators on quantity and quality of care is represented by paths α_2 , β_2 , α_3 , and β_3 . The direct effect, i.e., the effect not explained by the use of mediators, of low GP density on quantity and quality of care is represented by paths α_1 and β_1 .

3 Materials and Methods

3.1 Study population

We used data from the fourth round (2018-2022) of a national panel survey of French private practice (non salaried) GPs, designed to collect information about GPs' medical behaviors, working conditions and opinions on public health policies. Participating GPs were randomly selected from a French National Registry of Health Care Workers as of January 1st, 2018.

Sampling was stratified for gender, age (<50, 50-59, 60+), workload (annual number of office and home visits in 2017; in terciles) and GP density of practice area (low, moderate, high). The survey is representative of GPs practicing in France (except Mayotte). GPs planning to retire or to move before the end of data collection, those exclusively practicing alternative medicine or those having less than 200 patients as a gatekeeper were excluded. Of the 12,022 eligible GPs contacted, 3,304 (27%) completed the inclusion questionnaire and undertook to answer six future cross-sectional surveys (one every nine months). The sample benefits from the French "public statistics" label of the National Authority for Statistical Information (*Conseil National de l'Information Statistique*) that confirms the quality of the processes used to produce the data and their compliance with the best practices, e.g., the burden that the survey implies for the subjects and the degree of consultation with users.

3.2 Procedure and questionnaire

We exploit three waves of the survey with data collected respectively between October 2018 and April 2019 (first questionnaire), December 2019 and March 2020 (dummy variable described below) and October 2020 and November 2020 (second questionnaire). All the questionnaires were pilot-tested for clarity and face validity; they were collected either online or by professional interviewers using computer-assisted telephone interviews. GPs received compensation equivalent to one consultation fee (25 euros) for their participation in each survey.

First questionnaire dealt with GPs' perceptions regarding the evolution of availability of medical services in their practice area. More precisely, it collected information about (1) their perception of the GP density of their practice area (from 1=very insufficient to 4=completely sufficient); (2) perceived difficulties when dealing with patients' requests (3 items): whether they werethese difficulties resulted in refusing new patients (both enrolled or casual) or increasing waiting times; (3) work-related stress (3 items): whether the working hours adapted well to their personal commitments, they worked longer than desired or cut on training time and (4) use of e-health tools (3 items): electronic patient records, e-prescribing software and secured email service. We constructed a dummy variable to isolate the GPs reporting a practice in an area with 'very insufficient' GP density.

Second questionnaire addressed GPs' self-reported vaccine recommendation frequency (1 = never to 4 = always) in six specific vaccine situations, chosen because their current vaccination coverage rates in France do not meet official public health objectives: (1) measles, mumps, and rubella for non-immune adolescents

(catch-up), (2) pertussis in the immediate postpartum period for new mothers who did not receive a dose prior to their pregnancy, (3) meningococcal C for 12-month-old infants, (4) human papillomavirus for children aged 11–14, (5) hepatitis B for adolescents (catch-up), and (6) seasonal influenza for adults under 65 with chronic condition.

We use a dummy variable indicating GPs reporting enrollment in MGP. GPs were also asked about their (part-time) practice of alternative medicine, e.g., homeopathy and acupuncture.

We also enriched the survey data by the French Deprivation index (FDEP) calculated at the municipality level in 2015 (Rey et al., 2009). This score was constructed by the French National Institute of Health and Medical Research (Inserm) from National Institute of Statistics and Economic Studies (Insee) data on four dimensions: (1) the unemployment rate in the active population aged 15-64; (2) the percentage of blue-collar workers in the active population aged 15-64; (3) the percentage of high school graduates in the population aged over 15 years old and (4) the median income per consumption unit. The FDEP index varies between -4 and 4: the higher the score, the greater the deprivation. We used a dummy variable to isolate the most deprived municipalities (with positive FDEP index).

3.3 Statistical analysis

For the descriptive analyses, data were weighted to match the nationwide GP population in terms of stratification variables to correct for possible systematic non-response bias.

We used confirmatory factor analysis (CFA) to construct four latent variables of interest: (1) difficulties when dealing with patients’ requests (demand absorption capacity, a proxy of quantity of care, DV1), (2) self-reported vaccine recommendation frequency (a proxy for quality of care, DV2), (3) work-related stress (mediator 1, M1), and (4) use of e-health tools (mediator 2, M2). Full list of variables used to construct the latent variables is presented in Tables 1 and 2.

We draw on the conceptual framework described above to motivate our mediation analysis. We assumed that low GP density influences GPs’ work-related stress (M1) as well as their use of e-health tools (M2) and ultimately the quantity of care (DV1) and the quality of care (DV2) (Model 1). We therefore estimate the following reduced-form base model (Model 1) for quantity of care (DV1) and the quality of care (DV2):

$$Quantity_i = \alpha_0 + \alpha_1 Density_i + \alpha_2 Stress_i + \alpha_3 X_i + \epsilon_i \tag{1}$$

$$Quality_i = \beta_0 + \beta_1 Density_i + \beta_2 Ehealth_i + \beta_3 X_i + \zeta_i \tag{2}$$

with $Quantity_i$ as the quantity of care provided by the GP i , explained by the GPs density $Density_i$, her work-related stress $Stress_i$ and a vector of covariates X_i (age, gender, deprivation index and occasional practice of complementary medicine). The specification is similar for the quality of care $Quality_i$. We assume that the error terms ϵ_i and ζ_i are uncorrelated.

To estimate the indirect effects, we modify equations (1) and (2) as follows:

$$Quantity_i = \alpha_0 + \alpha_1 Density_i + \alpha_2 Stress_i(Density_i) + \alpha_3 X_i + \epsilon_i \quad (3)$$

$$Quality_i = \beta_0 + \beta_1 Density_i + \beta_2 Ehealth_i(Density_i) + \beta_3 X_i + \zeta_i \quad (4)$$

We assume that the indirect effect of GP density on quantity of care (DV1) is mediated through GPs' work-related stress (M1), while the indirect effect on the quality of care (DV2) is mediated through the use of e-health tools (M2). We disentangle the direct effect from the indirect effect by total differentiation with respect to $Density_i$:

$$\frac{dQuantity_i}{dDensity_i} = \alpha_1 + \left(\frac{\partial Quantity_i}{\partial Stress_i} \times \frac{\partial Stress_i}{\partial Density_i} \right) \quad (5)$$

$$\frac{dQuality_i}{dDensity_i} = \beta_1 + \left(\frac{\partial Quality_i}{\partial Ehealth_i} \times \frac{\partial Ehealth_i}{\partial Density_i} \right) \quad (6)$$

While the left-hand side displays the total effect of equation (5), the right-hand side decomposes the total effect into (1) the direct effect α_1 of GP density on quantity of care, and (2) the indirect effect (in parentheses) as cross-derivatives. Equation (6) is the equivalent of (5) for the of GP density on quality of care.

The mediating factors are estimated in two additional equations below:

$$Stress_i = \delta_0 + \delta_1 Density_i + \delta_2 X_i + \mu_i \quad (7)$$

$$Ehealth_i = \gamma_0 + \gamma_1 Density_i + \gamma_2 X_i + \nu_i \quad (8)$$

Moreover, we estimate a second model where we consider enrollment in MGP as an "opening" mediator (M3) between GP density and the two mediators defined above (Model 2). Consequently, the reduced-form model is adjusted as follows:

$$Quantity_i = \alpha_0 + \alpha_1 Density_i + \alpha_2 Stress_i + \alpha_3 MGP_i + \alpha_4 X_i + \epsilon_i \quad (9)$$

$$Quality_i = \beta_0 + \beta_1 Density_i + \beta_2 Ehealth_i + \beta_3 MGP_i + \beta_4 X_i + \zeta_i \quad (10)$$

with MGP_i acting as a mediator for the two dependent variables. In addition, MGP_i affects both the work-related stress and the use of e-health tools, so that (3) and (4) become

$$Quantity_i = \alpha_0 + \alpha_1 Density_i + \alpha_2 Stress_i(MGP_i(Density_i), Density_i) + \alpha_3 MGP_i(Density_i) + \alpha_4 X_i + \epsilon_i \quad (11)$$

$$Quality_i = \beta_0 + \beta_1 Density_i + \beta_2 Ehealth_i(MGP_i(Density_i), Density_i) + \beta_3 MGP_i(Density_i) + \beta_4 X_i + \zeta_i \quad (12)$$

so that the decomposition of the total effect is

$$\frac{dQuantity_i}{dDensity_i} = \alpha_1 + \frac{\partial Quantity_i}{\partial Stress_i} \left[\frac{\partial Stress_i}{\partial MGP_i} \times \frac{\partial MGP_i}{\partial Density_i} + \frac{\partial Stress_i}{\partial Density_i} \right] + \left(\frac{\partial Quantity_i}{\partial MGP_i} \times \frac{\partial MGP_i}{\partial Density_i} \right) \quad (13)$$

$$\frac{dQuality_i}{dDensity_i} = \beta_1 + \frac{\partial Quality_i}{\partial Ehealth_i} \left[\frac{\partial Ehealth_i}{\partial MGP_i} \times \frac{\partial MGP_i}{\partial Density_i} + \frac{\partial Ehealth_i}{\partial Density_i} \right] + \left(\frac{\partial Quality_i}{\partial MGP_i} \times \frac{\partial MGP_i}{\partial Density_i} \right) \quad (14)$$

Equations (13) and (14) could be also written as

$$\frac{dQuantity_i}{dDensity_i} = \alpha_1 + \alpha_2 [\lambda \times \eta_1 + \delta_1] + (\alpha_3 \times \eta_1) \quad (15)$$

$$\frac{dQuality_i}{dDensity_i} = \beta_1 + \beta_2 [\theta \times \eta_1 + \gamma_1] + (\beta_3 \times \eta_1) \quad (16)$$

The mediating factors are estimated in three additional equations below:

$$Stress_i = \delta_0 + \delta_1 Density_i + \delta_2 X_i + \lambda MGP_i + \mu_i \quad (17)$$

$$Ehealth_i = \gamma_0 + \gamma_1 Density_i + \gamma_2 X_i + \theta MGP_i + \nu_i \quad (18)$$

$$MGP_i = \eta_0 + \eta_1 Density_i + \eta_2 X_i + \omega_i \quad (19)$$

The issue of the potential endogeneity of variables (MGP in particular) due to omitted variables is addressed through the inclusion of adequate controls in the multiple-stage least squares analysis (since instrumental variables method is not recommended to deal with omitted variables, as it would require a specific instrument for each link assessed in the SEM). The risk of reverse causality is handled by the outline of the structural equation modelling (Collier, 2020; Pearl, 2009; VanderWeele and Shpitser, 2011). For example, note that in our case (Model 2) the endogeneity of practice in MGP to GP density is included in the structural modelling, with the reverse causality excluded (that practice in MGP causes GP density to decline, which is not a plausible assumption). However, we do provide the results that include covariance between practice in MGP and GP density (the causality flows both ways) as a robustness check. We also test a model that includes the effects of work-related stress on quality of care and that of use of e-health tools on quantity of care.

Model fit was assessed by the root mean square error of approximation (RMSEA), the comparative fit index (CFI), the Tucker-Lewis index (TLI), and the standardized root mean square residual (SRMR). Models with CFI>0.90, TLI>0.90, RMSEA<0.08, and SRMR<0.08 were considered to fit reasonably or well (Hooper et al., 2008).

All analyses were based on two-sided P values and conducted with Stata 14 (StataCorp, College Station, Texas).

4 Results

4.1 Summary statistics

Of 1,209 French GPs still enrolled in 2020, 40% of GPs were female, about a third were younger than 50 years old (Table 1). Twelve percent were enrolled in an MGP and 15% occasionally practiced complementary medicine. Twenty-one percent reported that they practiced in an area with very insufficient GP density compared to 10% that were objectively practicing in an underserved area, i.e., an area where a patient could have had access to less than 2.8 consultations per year (Ministère des solidarités et de la santé, 2021). Among the 10% of GPs that were objectively practicing in an area with very insufficient GP supply, only 38% had the matching perception. The proportion of those reporting to practice in an area with very insufficient GP density was significantly higher (29%) for those practicing in an MGP. More than a half (56%) were practicing in deprived areas; this proportion was significantly higher (70%) for those enrolled in MGP.

More than four GPs out of ten reported that their working hours did not adapt well to their personal commitments or that they had to cut on training time. Seventy percent reported that they had to work longer than desired.

The vast majority of GPs declared using diverse e-health tools: 92% used electronic patient record and secure email services, and 85% used e-prescribing software. These proportions were significantly higher for GPs enrolled in MGP (99%, 99% and 94% respectively).

Around half of GPs were able to accept new patients: 44% for consultations with enrolled patients and 57% for casual consultations (Table 2). Forty-six percent managed to maintain their waiting times compared to 32% of GPs in MGP.

Almost all GPs (92%) claimed to always or often recommend meningococcal C vaccine for 12-month-old infants (97% in MGP) and seasonal influenza for adults with chronic conditions. More than 80% recommended measles, mumps, and rubella catch-up for adolescents, pertussis for new mothers and human papillomavirus vaccines for children aged 11-14. Two-third of GPs declared recommending hepatitis B catch-up for adolescents. Some vaccines, e.g., hepatitis B catch-up, human papillomavirus, a seasonal influenza, were less systematically recommended by the GPs.

Table 1: Sample characteristics, national panel of French GPs (n=1,209)

%	Total	MGP
Thinks that GP density in the area is very insufficient	21.59	29.04**
MGP	11.69	-
Age		
<50	33.68	48.97
50-59	40.67	34.72
≥60	25.65	16.31
Female	39.13	40.62
Occasional practice of complementary medicine	15.83	15.62
French DEPrivation index > 0	56.44	69.87***
Work-related stress		
Working hours do not adapt well to their personal commitments	40.96	42.76
Works longer than desired	70.70	76.01
Cuts on training time	43.10	49.03
Use of e-health tools		
Uses electronic patient records	91.59	99.66***
Uses e-prescribing software	84.53	93.77***
Uses secured email service	91.49	98.87***

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

GPs: general practitioners; MGP: multi-professional group practices

Source: DREES, ORS and URPS Provence-Alpes-Côte d'Azur and Pays de la Loire, 4^{ème} Panel d'observation des pratiques et des conditions d'exercice en médecine générale de ville.

Note: weighted data.

Table 2: Dependent variables, national panel of French GPs (n=1,209)

Demand absorption capacity	%	No		Yes	
<u>Does not</u> refuse new patients (enrolled)	Total in MGP	55.62	49.88	44.38	50.12
<u>Does not</u> refuse new patients (casual)	Total in MGP	42.97	48.6	57.03	51.40
<u>Does not</u> increase waiting times	Total in MGP	53.66	67.61***	46.34	32.39***
Vaccine recommendations	%	Never	Sometimes	Often	Always
MMR for non-immune adolescents (catch-up)	Total in MGP	1.42	13.80	24.90	59.89
Pertussis postpartum†	Total in MGP	0.18	14.81	21.30	63.71
Meningococcal C for 12-month-old infants	Total in MGP	7.73	10.76	20.05	61.46
HPV for children aged 11–14	Total in MGP	5.01	10.81	22.27	61.90
Hepatitis B for adolescents (catch-up)	Total in MGP	1.19	6.80	12.17	79.84
Seasonal influenza for adults under 65 with chronic condition	Total in MGP	0.37***	2.33***	6.03***	91.27***
HPV for children aged 11–14	Total in MGP	3.31	14.85	42.26	39.58
Hepatitis B for adolescents (catch-up)	Total in MGP	3.15	7.82	41.10	47.93
Seasonal influenza for adults under 65 with chronic condition	Total in MGP	1.97	23.25	37.10	37.68
HPV for children aged 11–14	Total in MGP	2.20	21.71	39.54	36.56
Seasonal influenza for adults under 65 with chronic condition	Total in MGP	0.09	6.71	37.88	55.31
HPV for children aged 11–14	Total in MGP	0.18	5.75	40.29	53.78

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

GPs: general practitioners; MGP: multi-professional group practices; MMR: Measles, mumps, and rubella; HPV: Human papillomavirus

Source: DREES, ORS and URPS Provence-Alpes-Côte d'Azur and Pays de la Loire, 4^{ème} Panel d'observation des pratiques et des conditions d'exercice en médecine générale de ville.

Note: weighted data.

† Pertussis in the immediate postpartum period for new mothers who did not receive a dose prior to their pregnancy.

4.2 Mediation analysis results

The results of the mediation analysis are presented in Table 3. Total standardized effects are displayed in order to compare across models.

Overall, the quantity of care delivered was negatively affected by perceived GP density as well as work-related stress. The quality of care was not significantly affected by the GP density, while higher levels of quality were linked to greater use of e-health tools. Female GPs were likely to have lower demand absorption capacity and to recommend vaccines more often. Older GPs were more likely to absorb demand more efficiently. Practice of complementary medicine was associated with lower frequency of vaccine recommendations, while greater workload had a positive impact on quality. Adding MGP as a mediator (Model 2) had no significant impact on the outcome variables.

As for the mediating factors, low perceived GP density was associated with higher levels of work-related stress and more enrollment in MGP. Older GPs were less likely to use e-health tools and enroll in MGP. Finally, there was a positive association between practice in a deprived area and enrollment in a MGP.

When disentangling between direct and indirect effects, in the absence of potential mediators, our results suggest that the relation between GP density and quantity of care delivered was negative and significant ($\alpha_1 = -0.220$); there was no significant effect found on the quality of care (Figure 2). In the presence of mediators (work-related stress (M1) and use of e-health tools (M2)), GP density was significantly and positively associated with work-related stress ($\delta_1 = 0.122$), which was consecutively associated with a degraded demand absorption capacity ($\alpha_2 = -0.191$). This sequence of correlations shows that the initial correlation between GP density and demand absorption capacity is mediated by work-related stress and this mediating effect can be quantified by the product of these coefficients ($\delta_1 \times \alpha_2 = -0.023$). Moreover, higher use of e-health tools (M2) was associated with greater vaccine recommendations frequency ($\beta_2 = 0.317$). Lastly, after the introduction of enrollment in MGP as a “mediator of mediators” (Figure 3), we found that GPs in MGP tended to use more e-health tools, with a favorable effect on our measure of quality of care, vaccine recommendations. However, we failed to find any significant improvements of practice in MGP on work-related stress. These results hold when accounting for covariance between GP density and practice in MGP as well as when we include the effects of work-related stress on quality of care and that of use of e-health tools on quantity of care.

Table 3: Mediation analysis results, SEM estimation, standardized total effects

Standardized coefficients, total effects		Outcome variables				Mediators				
		Quantity of care		Quality of care		Work-related stress		E-health		MGP
		Model 1	Model 2	Model 1	Model 2	Model 1	Model 2	Model 1	Model 2	Model 2
Mediators	Work-related stress	-0,187*** (0,061)	-0,192*** (0,041)	-	-	-	-	-	-	-
	E-health	-	-	0,166*** (0,047)	0,168*** (0,036)	-	-	-	-	-
	MGP	-	0,043 (0,062)	-	0,022 (0,049)	-	-0,018 (0,043)	-	0,096** (0,036)	-
Explanatory variable	Perceived GP density	-0,371** (0,040)	-0,371*** (0,043)	0,034 (0,035)	0,053 (0,039)	0,210*** (0,042)	0,211*** (0,047)	0,037 (0,035)	0,039 (0,039)	0,078** (0,032)
Control variables	Female	-0,179*** (0,042)	-0,171*** (0,044)	0,142*** (0,038)	0,134*** (0,040)	-0,081 (0,046)	-0,062 (0,038)	-0,047 (0,047)	-0,053 (0,039)	-0,055 (0,032)
	Age (ref. <50 years)									
	50-59	0,062 (0,042)	0,070 (0,046)	-0,024 (0,038)	-0,023 (0,045)	0,005 (0,046)	-0,003 (0,037)	-0,166*** (0,050)	-0,155*** (0,039)	-0,179*** (0,033)
	≥ 60	0,132*** (0,044)	0,159*** (0,041)	-0,070 (0,044)	-0,064 (0,036)	-0,054 (0,049)	-0,067 (0,037)	-0,394*** (0,043)	-0,370*** (0,036)	-0,191*** (0,030)
	Complementary medicine (ref.No)	0,056 (0,039)	0,054 (0,041)	-0,087** (0,035)	-0,086** (0,037)	0,029 (0,043)	0,012 (0,035)	-0,049 (0,044)	-0,050 (0,037)	0,022 (0,030)
	Deprivation index>0	-0,024 (0,039)	-0,038 (0,048)	-0,051 (0,035)	-0,061 (0,043)	-0,030 (0,043)	-0,016 (0,035)	0,078** (0,051)	0,066 (0,042)	0,150*** (0,035)
	Workload (ref.Q1)									
	Q2-Q3	-0,038 (0,046)	-0,037 (0,051)	0,061 (0,041)	0,064 (0,043)	0,085 (0,050)	0,095 (0,041)	0,073 (0,051)	0,077 (0,044)	0,054 (0,036)
	Q4	-0,001*** (0,049)	0,016 (0,031)	0,146*** (0,042)	0,145*** (0,037)	0,216*** (0,051)	0,230*** (0,042)	-0,012*** (0,042)	-0,030 (0,045)	-0,025 (0,085)


* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

Notes: non-weighted data

Goodness of fit, Model 1: RMSEA=0,028; CFI=0,933; TLI=0,915; SRMR=0,030; $R^2=0,427$


Goodness of fit, Model 2: RMSEA=0,029; CFI=0,926; TLI=0,902; SRMR=0,031; $R^2=0,462$

Figure 2: Unstandardized direct effects of paths of the multiple mediation model on demand absorption capacity and vaccine recommendation frequency (Model 1)



* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Non-weighted data. Coefficients are estimated from linear regression equations. Model was adjusted for age, gender, deprivation index and occasional practice of complementary medicine. Latent variables (factors) are represented by circles, and measured variables (items) by rectangles; arrows point in the hypothesized direction of influence. Goodness of fit: RMSEA=0,028; CFI=0,933; TLI=0,915; SRMR=0,030; $R^2=0,427$

Figure 3: Unstandardized direct effects of paths on demand absorption capacity and vaccine recommendation frequency (Model 2)



* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$. Non-weighted data. Coefficients are estimated from linear regression equations. Model was adjusted for age, gender, deprivation index and occasional practice of complementary medicine. Latent variables (factors) are represented by circles, and measured variables (items) by rectangles; arrows point in the hypothesized direction of influence. Goodness of fit: RMSEA=0,029; CFI=0,926; TLI=0,902; SRMR=0,031; $R^2=0,462$

5 Discussion & conclusion

To the best of our knowledge, this is the first study to analyse the underlying mediation effects behind the relationship between GP density and quantity and quality of primary care that also analyses integrated primary care teams' option as a possible response to the decreasing density.

We found that lower density is associated with lower demand absorption capacity as a proxy for quantity of care. We confirmed, at the country level, the existence of a negative relationship between GP density and work-related stress, in line with Picquendar et al. (2019). We show that lower level of work-related stress is a key mediator in handling patients' requests (Shen et al., 2022; Wallace et al., 2009).

As the statistical framework (using structural equations) allows correcting for the self-selection into MGP, we also amend some unstable results contained in the related literature: contrary to Cassou et al. (2020) we do not find a significant impact of enrollment in MGP on the demand absorption capacity. A difference between their work and ours might stem from the sample of GPs that are included in both research papers. The study population of Cassou et al. (2020) is restricted to MGPs accredited by Social Security that represent about half of French MGPs, that are subject to stricter requirements from the health authorities to receive an add-on payment. For instance, the accredited MGPs are supposed to have extended hours to ensure the continuity of care, which might explain the increase in the number of patients seen by the GP that they observe.

Even outside the French context, the finding that there is no significant effect of an enrolment in integrated practices on absorption capacity is unusual (but not without precedent, see Nolte and McKee (2009)) The heterogeneity in configuration of the integrated practices that we have in our study might be an explanation: while MGPs require two full-time equivalent GPs and one paramedic to be recognized as such by health authorities, the size and list of professionals available at the practice might vary across the country. Therefore, the size of the integrated practices might be insufficient to fully absorb the demand and/or its composition (e.g., in paramedics) might not be adapted to the characteristics of the practice area, e.g., does not consider the part of the aging population that generates more care demand. Another plausible explanation might be that once enrolled in an MGP, GPs adjust their working hours to achieve better work-life balance. Finally, it is possible that the efficiency gains generated by enrolment in an MGP are not reflected in quantity, but rather in quality (Singer et al., 2020; Zaytseva et al., 2022).

GP density indirectly affects the quality of care. As expected, more MGPs are located in underserved areas (Chevallard et al., 2019). While we do not find a direct effect of practicing in MGP on the quality of care, we found that practicing in MGP is indirectly able to generate better quality of care. Enrolment in MGP was associated with higher use of e-health tools and, consecutively, with higher frequency of vaccine recommendations: e-health tools can be used to systematically identify unvaccinated patients in a practice and send an automated reminder to the GP when the patient visits that a vaccination is due (Derrough et al., 2017; Dumit et al., 2018; Groom et al., 2015; Jacobson Vann et al., 2018; Morgan et al., 2015; Niccolai and Hansen, 2015; Okoli et al., 2021; Rey et al., 2018).

A limitation of our research is the use of survey data, so social desirability or conformity biases cannot

be ruled out. Another limitation is that we consider only a set of three mediators (work-related stress, use of e-health tools and enrolment in an MGP), while other factors, such as consultation length or trust in institutions, may be of importance both in prevention activities and vaccination. While the choice of mediators is justified by our conceptual framework, existing literature and data availability, further studies are needed to consider a broader set of mediators and/or outcome variables related to the quality and quantity of care.

To conclude, this study shows that an enrolment in MGP as a response to a decreasing medical density, although disappointing in terms of quantity of care supplied, has the potential to improve the quality of care provided to the patients. While a previous study that investigated the performance of GPs in MGP during Covid-19 pandemics found that GPs in MGP ensured better follow-up of patients with chronic conditions during the lockdown than those practicing outside MGP (Zaytseva et al., 2022), our results strengthen the evidence of better qualitative performance of GPs when considering their regular practice styles, such as their vaccination recommendation practices. Thus, further development of integrated primary care appears beneficial to the patients located in the underserved areas and even more so since one of the requirements to establish an MGP is the participation in public health, prevention, health education and social activities of their choice in their catchment area.

References

- Anguis, Marie, Maxime Bergeat, Jacques Pisarik, Noémie Vergier, and H el ene Chaput,** “Quelle d emographie r ecente et   venir pour les professions m edicales et pharmaceutique ? Constat et projections d emographiques [What is the recent and future demography of the medical and pharmaceutical professions? Findings and demographic projections.],” *Les Dossiers de la DREES*, March 2021, (76), 74.
- Carrier, H el ene, Anna Zaytseva, Aur elie Bocquier, Patrick Villani, Martin Fortin, and Pierre Verger,** “General practitioners’ attitude towards cooperation with other health professionals in managing patients with multimorbidity and polypharmacy: A cross-sectional study,” *European Journal of General Practice*, 2022, *28* (1), 109–117.
- Cassou, Matthieu, Julien Mousqu es, and Carine Franc,** “General practitioners’ income and activity: the impact of multi-professional group practice in France,” *The European Journal of Health Economics*, December 2020, *21* (9), 1295–1315.
- Chaput, H el ene, Martin Monziols, Lisa Fressard, Pierre Verger, Bruno Ventelou, and Anna Zaytseva,** “Plus de 80 % des m edecins g en eralistes lib eraux de moins de 50 ans exercent en groupe [More than 80% of private GPs under 50 years old are enrolled in a group practice],” *Etudes et r esultats*, 2019, (1114).
- Chevillard, Guillaume and Julien Mousqu es,** “Medically underserved areas: are primary care teams efficient at attracting and retaining general practitioners?,” *Social Science & Medicine*, October 2021, *287*, 114358.
- , – , **V eronique Lucas-Gabrielli, and St ephane Rican,** “Has the diffusion of primary care teams in France improved attraction and retention of general practitioners in rural areas?,” *Health Policy*, May 2019, *123* (5), 508–515.
- Colla, Carrie, Wendy Yang, Alexander J. Mainor, Ellen Meara, Marietou H. Ouayogode, Valerie A. Lewis, Stephen Shortell, and Elliott Fisher,** “Organizational integration, practice capabilities, and outcomes in clinically complex medicare beneficiaries,” *Health Services Research*, December 2020, *55* (S3), 1085–1097.
- Collier, Joel E.,** *Applied Structural Equation Modeling Using AMOS: Basic to Advanced Techniques*, New York: Routledge, June 2020.
- Davin-Casalena, B., M. Jardin, H. Guerrera, J. Mabile, H. Tr ehard, D. Lapalus, C. M enager, S. Nauleau, V. Cassaro, P. Verger, and V. Guagliardo,** “The impact of the COVID-19 epidemic on primary care in South-eastern France: implementation of a real-time monitoring system based on regional health insurance system data,” *Revue d’ pid emiologie et de Sant e Publique*, October 2021, *69* (5), 255–264.

- Derrough, Tarik, Kate Olsson, Vincenza Gianfredi, Francois Simondon, Harald Heijbel, Niklas Danielsson, Piotr Kramarz, and Lucia Pastore-Celentano**, “Immunisation Information Systems – useful tools for monitoring vaccination programmes in EU/EEA countries, 2016,” *Eurosurveillance*, April 2017, *22* (17), 30519. Publisher: European Centre for Disease Prevention and Control.
- Deveugele, Myriam, Anselm Derese, Atie van den Brink-Muinen, Jozien Bensing, and Jan De Maeseneer**, “Consultation length in general practice: cross sectional study in six European countries,” *BMJ*, August 2002, *325* (7362), 472.
- Dumit, Elsy Maria, David Novillo-Ortiz, Marcela Contreras, Martha Velandia, and M. Carolina Danovaro-Holliday**, “The use of eHealth with immunizations: An overview of systematic reviews,” *Vaccine*, December 2018, *36* (52), 7923–7928.
- Eliès, Antoine, Claire Bonneau, Sophie Houzard, Roman Rouzier, and Delphine Héquet**, “Impact of catch-up human papillomavirus vaccination on cervical conization rate in a real-life population in France,” *PLOS ONE*, March 2022, *17* (3), e0264821.
- Francis, Diane B., Joan R. Cates, Kyla P. Garrett Wagner, Tracey Zola, Jenny E. Fitter, and Tamera Coyne-Beasley**, “Communication technologies to improve HPV vaccination initiation and completion: A systematic review,” *Patient Education and Counseling*, July 2017, *100* (7), 1280–1286.
- Groenewegen, Peter P, Mark W G Bosmans, Wienke G W Boerma, and Peter Spreeuwenberg**, “The primary care workforce in Europe: a cross-sectional international comparison of rural and urban areas and changes between 1993 and 2011,” *European Journal of Public Health*, September 2020, *30* (Supplement_4), iv12–iv17.
- Groom, Holly, David P. Hopkins, Laura J. Pabst, Jennifer Murphy Morgan, Mona Patel, Ned Calonge, Rebecca Coyle, Kevin Dombkowski, Amy V. Groom, Mary Beth Kurilo, Bobby Rasulnia, Abigail Shefer, Cecile Town, Pascale M. Wortley, Jane Zucker, and the Community Preventive Services Task Force**, “Immunization Information Systems to Increase Vaccination Rates: A Community Guide Systematic Review,” *Journal of Public Health Management and Practice*, 2015, *21* (3), 227–248. Publisher: Lippincott Williams & Wilkins.
- Hooper, Daire, Joseph Coughlan, and Michael R. Mullen**, “Structural Equation Modelling: Guidelines for Determining Model Fit.,” *Electronic Journal of Business Research Methods*, 2008, *6* (1), 53–60.
- Irving, Greg, Ana Luisa Neves, Hajira Dambha-Miller, Ai Oishi, Hiroko Tagashira, Anistasiya Verho, and John Holden**, “International variations in primary care physician consultation time: a systematic review of 67 countries,” *BMJ Open*, October 2017, *7* (10), e017902.
- Karuna, Christo, Victoria Palmer, Anthony Scott, and Jane Gunn**, “Prevalence of burnout among GPs: a systematic review and meta-analysis,” *British Journal of General Practice*, December 2021.

- Kerrissey, Michaela J., Jonathan R. Clark, Mark W. Friedberg, Wei Jiang, Ashley K. Fryer, Molly Frean, Stephen M. Shortell, Patricia P. Ramsay, Lawrence P. Casalino, and Sara J. Singer**, “Medical Group Structural Integration May Not Ensure That Care Is Integrated, From The Patient’s Perspective,” *Health Affairs*, May 2017, *36* (5), 885–892.
- Lalani, Mirza, Sonia Bussu, and Martin Marshall**, “Understanding integrated care at the frontline using organisational learning theory: A participatory evaluation of multi-professional teams in East London,” *Social Science & Medicine*, October 2020, *262*, 113254.
- Lane, Natasha E., Vicki Ling, Richard H. Glazier, and Thérèse A. Stukel**, “Primary care physician volume and quality of care for older adults with dementia: a retrospective cohort study,” *BMC Family Practice*, December 2021, *22* (1), 51.
- Larson, Heidi J., Alexandre de Figueiredo, Zhao Xiahong, William S. Schulz, Pierre Verger, Iain G. Johnston, Alex R. Cook, and Nick S. Jones**, “The State of Vaccine Confidence 2016: Global Insights Through a 67-Country Survey,” *EBioMedicine*, September 2016.
- Lurquin, Kellou, Colin, and Letrilliart**, “Comparison of rural and urban French GPs’ activity: a cross-sectional study,” *Rural and Remote Health*, September 2021.
- Maslach, Christina, Susan E. Jackson, and Michael P. Leiter**, “Maslach Burnout Inventory: Third edition,” in “Evaluating stress: A book of resources,” Lanham, MD, US: Scarecrow Education, 1997, pp. 191–218.
- Ministère des solidarités et de la santé**, “Arrêté du 1er octobre 2021 modifiant l’arrêté du 13 novembre 2017 relatif à la méthodologie applicable à la profession de médecin pour la détermination des zones prévues au 1^ode l’article L. 1434-4 du code de la santé publique [Decree of 1 October 2021 amending the decree of 13 November 2017 relating to the methodology applicable to the physician profession for determining the zones provided in 1^oof Article L. 1434-4 of the Public Health Code],” 2021.
- Morgan, Jamie L., Sangameshwar R. Baggari, Wendy Chung, Julia Ritch, Donald D. McIntire, and Jeanne S. Sheffield**, “Association of a Best-Practice Alert and Prenatal Administration With Tetanus Toxoid, Reduced Diphtheria Toxoid, and Acellular Pertussis Vaccination Rates,” *Obstetrics & Gynecology*, August 2015, *126* (2), 333–337.
- Mousquès, Julien and Fabien Daniel**, “Multiprofessional Group Practices Generate Gains in Terms of Productivity and Expenditures. Results of the Evaluation of Multidisciplinary Group Practices (MGP), Health Care Networks (HCN) and Health Care Centers (HCC) Participating in Experiments with New Modes of Remuneration (ENMR),” *Questions d’économie de la santé*, June 2015, (210), 6.
- Niccolai, Linda M. and Caitlin E. Hansen**, “Practice- and Community-Based Interventions to Increase Human Papillomavirus Vaccine Coverage: A Systematic Review,” *JAMA pediatrics*, July 2015, *169* (7), 686–692.

- Nolte, Ellen and Martin McKee**, *Integration and chronic care: a review. In: Caring for people with chronic conditions: a health system perspective. European Observatory on Health Systems and Policies Series*, Open Univ. Press, 2009. OCLC: 613115886.
- Okoli, George N., Viraj K. Reddy, Otto L. T. Lam, Tiba Abdulwahid, Nicole Askin, Edward Thommes, Ayman Chit, Ahmed M. Abou-Setta, and Salaheddin M. Mahmud**, “Interventions on health care providers to improve seasonal influenza vaccination rates among patients: a systematic review and meta-analysis of the evidence since 2000,” *Family Practice*, July 2021, *38* (4), 524–536.
- Patel, Mona, Laura Pabst, Sajal Chattopadhyay, David Hopkins, Holly Groom, Stuart Myerburg, and Jennifer Murphy Morgan**, “Economic Review of Immunization Information Systems to Increase Vaccination Rates: A Community Guide Systematic Review,” *Journal of Public Health Management and Practice*, May 2015, *21* (3), 253–262.
- Pearl, Judea**, *Causality: Models, Reasoning, and Inference*, 2 ed., Cambridge: Cambridge University Press, 2009.
- Petrie, Samuel, Dean Carson, Paul Peters, Anna-Karin Hurtig, Michele LeBlanc, Holly Simpson, Jaymie Barnabe, Mikayla Young, Mara Ostafichuk, Heidi Hodge, Justin Gladman, Matilda Smale, and Manuel Gonzalez Garcia**, “What a Pandemic Has Taught Us About the Potential for Innovation in Rural Health: Commencing an Ethnography in Canada, the United States, Sweden, and Australia,” *Frontiers in Public Health*, December 2021, *9*, 768624.
- Picquendar, Guillaume, Antoine Guedon, Fanny Moulinet, and Matthieu Schuers**, “Influence of medical shortage on GP burnout: a cross-sectional study,” *Family Practice*, May 2019, *36* (3), 291–296.
- Rey, Dominique, Lisa Fressard, Sébastien Cortaredona, Aurélie Bocquier, Arnaud Gautier, Patrick Peretti-Watel, Pierre Verger, and on behalf of the Baromètre santé 2016 Group**, “Vaccine hesitancy in the French population in 2016, and its association with vaccine uptake and perceived vaccine risk–benefit balance,” *Eurosurveillance*, April 2018, *23* (17), 17–00816.
- Rey, Grégoire, Eric Jouglu, Anne Fouillet, and Denis Hémon**, “Ecological association between a deprivation index and mortality in France over the period 1997 - 2001: variations with spatial scale, degree of urbanicity, age, gender and cause of death,” *BMC public health*, January 2009, *9*, 33.
- Robert, J., B. Detournay, M.C. Levant, M. Uhart, J. Gourmelen, and J.M. Cohen**, “Flu vaccine coverage for recommended populations in France,” *Médecine et Maladies Infectieuses*, November 2020, *50* (8), 670–675.
- Shen, Xin, Hongbin Xu, Jing Feng, Jun Ye, Zuxun Lu, and Yong Gan**, “The global prevalence of burnout among general practitioners: a systematic review and meta-analysis,” *Family Practice*, January 2022, p. cmab180.

- Silhol, Julien, Blandine Legendre, and Martin Monziols**, “General Practitioners’ medical practices in territories that have become subject to high-priority intervention (“Zones d’intervention prioritaire (ZIP)”),” *Insee Analyses*, May 2020, (51).
- Singer, Sara J., Anna D. Sinaiko, Maike V. Tietschert, Michaela Kerrissey, Russell S. Phillips, Veronique Martin, Grace Joseph, Hassina Bahadurzada, and Denis Agniel**, “Care integration within and outside health system boundaries,” *Health Services Research*, December 2020, *55* (S3), 1033–1048.
- Smith, Susan M, Emma Wallace, Tom O’Dowd, and Martin Fortin**, “Interventions for improving outcomes in patients with multimorbidity in primary care and community settings,” *Cochrane Database of Systematic Reviews*, January 2021, *2021* (5).
- Soler, J. K., H. Yaman, M. Esteva, F. Dobbs, R. S. Asenova, M. Katic, Z. Ozvacic, J. P. Desgranges, A. Moreau, C. Lionis, P. Kotanyi, F. Carelli, P. R. Nowak, Z. de Aguiar Sa Azeredo, E. Marklund, D. Churchill, M. Ungan, and (European General Practice Research Network Burnout Study Group)**, “Burnout in European family doctors: the EGPRN study,” *Family Practice*, August 2008, *25* (4), 245–265.
- Spányik, András, Dávid Simon, Adrien Rigó, Mark D. Griffiths, and Zsolt Demetrovics**, “Subjective COVID-19-related work factors predict stress, burnout, and depression among healthcare workers during the COVID-19 pandemic but not objective factors,” *PLOS ONE*, August 2022, *17* (8), e0270156.
- Steinhaeuser, Jost, Stefanie Joos, Joachim Szecsenyi, and Antje Miksch**, “A comparison of the workload of rural and urban primary care physicians in Germany: analysis of a questionnaire survey,” *BMC Family Practice*, October 2011, *12* (1), 112.
- Sutter, Marieke De, An De Sutter, Nora Sundahl, Tom Declercq, and Peter Decat**, “Inter-professional collaboration reduces the burden of caring for patients with mental illnesses in primary healthcare. A realist evaluation study,” *European Journal of General Practice*, October 2019, *25* (4), 236–242.
- van Hassel, Daniël, Robert Verheij, and Ronald Batenburg**, “Assessing the variation in workload among general practitioners in urban and rural areas: An analysis based on SMS time sampling data,” *The International Journal of Health Planning and Management*, 2019, *34* (1), e474–e486.
- VanderWeele, Tyler J. and Ilya Shpitser**, “A new criterion for confounder selection,” *Biometrics*, December 2011, *67* (4), 1406–1413.
- Vann, Julie C. Jacobson, Robert M. Jacobson, Tamera Coyne-Beasley, Josephine K. Asafu-Adjei, and Peter G. Szilagyi**, “Patient reminder and recall interventions to improve immunization rates,” *The Cochrane Database of Systematic Reviews*, January 2018, *1*, CD003941.

- Verger, Pierre, Lisa Fressard, Fanny Collange, Arnaud Gautier, Christine Jestin, Odile Lounay, Jocelyn Raude, Céline Pulcini, and Patrick Peretti-Watel**, “Vaccine Hesitancy Among General Practitioners and Its Determinants During Controversies: A National Cross-sectional Survey in France,” *EBioMedicine*, August 2015, *2* (8), 891–897.
- Wakerman, John and John S. Humphreys**, “Sustainable primary health care services in rural and remote areas: Innovation and evidence,” *Australian Journal of Rural Health*, June 2011, *19* (3), 118–124.
- Wallace, Jean E, Jane B Lemaire, and William A Ghali**, “Physician wellness: a missing quality indicator,” *The Lancet*, November 2009, *374* (9702), 1714–1721.
- WHO**, “Declaration of Astana. Global conference on primary health care,” Technical Report, World Health Organization and the United Nations Children’s Fund October 2018.
- , “Primary Health Care on the Road to Universal Health Coverage. 2019 Monitoring Report,” Technical Report, World Health Organization 2019.
- Woodall, Tasha, Melinda Ramage, John T. LaBruyere, William McLean, and Casey R. Tak**, “Telemedicine Services During COVID-19: Considerations for Medically Underserved Populations,” *The Journal of Rural Health*, January 2021, *37* (1), 231–234.
- Zaytseva, Anna, Pierre Verger, and Bruno Ventelou**, “United, can we be stronger? Did French general practitioners in multi-professional groups provide more chronic care follow-up during lockdown?,” *BMC Health Services Research*, December 2022, *22* (1), 519.
- Zyzanski, S. J., K. C. Stange, D. Langa, and S. A. Flocke**, “Trade-offs in high-volume primary care practice,” *The Journal of Family Practice*, May 1998, *46* (5), 397–402.