



HAL
open science

Non-motor symptoms and quality of life in patients with PRRT2-related paroxysmal kinesigenic dyskinesia

Asya Ekmen, Mohamed Doulazmi, Aurélie Méneret, Prasanthi Jegatheesan, Anais Hervé, Philippe Damier, Domitille Gras, Agathe Roubertie, Juliette Piard, Eugenie Mutez, et al.

► To cite this version:

Asya Ekmen, Mohamed Doulazmi, Aurélie Méneret, Prasanthi Jegatheesan, Anais Hervé, et al.. Non-motor symptoms and quality of life in patients with PRRT2-related paroxysmal kinesigenic dyskinesia. *Movement Disorders Clinical Practice*, 2023, 10 (7), 10.1002/mdc3.13795 . hal-04306310

HAL Id: hal-04306310

<https://hal.science/hal-04306310>

Submitted on 24 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 *Title: Non-motor symptoms and quality of life in patients with PRRT2-related paroxysmal*
2 *kinesigenic dyskinesia*

3 Authors: Asya Ekmen^{1,2} MD, PhD, Mohamed Doulazmi³ PhD, Aurélie Méneret^{1,2} MD, PhD,
4 Prasanthi Jegatheesan^{1,2} PhD, Anais Hervé¹ RN, Philippe Damier⁴ MD, PhD, Domitille Gras¹
5 MD, PhD, Agathe Roubertie⁵ MD, PhD, Juliette Piard⁶ MD, PhD, Eugenie Mutez⁷ MD, PhD,
6 Clément Tarrano^{1,2} MD, Quentin Welniarz^{1,2} PhD, Marie Vidailhet^{1,2} MD, Yulia Worbe^{1,2} MD,
7 PhD*, Cécile Gallea^{1*} PhD, Emmanuel Roze^{1,2} MD, PhD*†.

8 † Corresponding author

9 * Equal contribution as senior authors

10 Affiliations :

11 ¹Sorbonne Université, INSERM, CNRS, Paris Brain Institute;

12 ²APHP Hôpital de La Pitié Salpêtrière et Saint-Antoine

13 ³Sorbonne University, Adaptation Biologique et Vieillesse (UMR8256), Institut de
14 Biologie Paris Seine, CNRS, Paris France

15 ⁴University of Nantes, CHU Nantes, CIC 1314;

16 ⁵INM, Université de Montpellier, INSERM, CHU Montpellier, Département Neuropédiatrie

17 ⁶Centre de Génétique Humaine, CHU, Besançon, France ; INSERM UMR1231, Génétique des
18 Anomalies du Développement, Université de Bourgogne, Dijon, France

19 ⁷ Univ. Lille, Inserm, CHU Lille, U1172 – LilNCog - Lille Neuroscience and Cognition.

20 **Word count:** text 2766 words, abstract 229 words

21 **Corresponding authors contact information:**

22 Address: ICM, 47-83 boulevard de l'hôpital, 75651 Paris Cedex 13, France;

23 Phone: +33 (0)157 27 46 27; Email: flamand.roze.75012@gmail.com

24 **Running title:** PRRT2 mutations and non-motor manifestations

25 **Key words:** PRRT2, paroxysmal disorders, non-motor symptoms, quality of life, stigma

26 **ABSTRACT**

27 **Background:** Monoallelic pathogenic variants of *PRRT2* often result in paroxysmal kinesigenic
28 dyskinesia (PKD). Little is known about health-related quality of life (HrQoL), non-motor
29 manifestations, self-esteem, and stigma in patients with PKD.

30 **Objectives:** We investigated non-motor symptoms and how they related to HrQoL in a
31 genetically homogeneous group of *PRRT2*-PKD patients. We paid special attention to
32 perceived stigmatisation and self-esteem.

33 **Methods:** We prospectively enrolled 21 consecutive PKD patients with a pathogenic variant of
34 *PRRT2*, and 21 healthy controls matched for age and sex. They were evaluated with dedicated
35 standardized tests for non-motor symptoms, HrQoL, anxiety, depression, stigma, self-esteem,
36 sleep, fatigue, pain, and psychological well-being.

37 **Results:** Patients reported an alteration of the physical aspects of HrQoL, regardless of the
38 presence of residual paroxysmal episodes. Non-motor manifestations were frequent, and were
39 an important determinant of the alteration of HrQoL. In addition, patients perceived a higher
40 level of stigmatization which positively correlated with a delay in diagnosis ($p=0.615$, $p=0.003$)
41 and the fear of being judged ($p=0.452$, $p=0.04$), but not with the presence of paroxysmal
42 episodes ($p=0.203$, $p=0.379$).

43 **Conclusions:** Our findings have important implications for care givers concerning patient
44 management and medical education about paroxysmal dyskinesia. *PRRT2*-PKD patients should
45 be screened for non-motor disorders in routine care. A long history of misdiagnosis may play a
46 role in the high level of perceived stigmatization. Improving knowledge about diagnostic clues
47 suggestive of PKD is mandatory.

48 Monoallelic pathogenic variants of the Proline-rich transmembrane protein 2 (*PRRT2*) gene
49 cause synaptopathy and can result in a large variety of paroxysmal neurological disorders (1–
50 3). The most frequent disorder is Paroxysmal Kinesigenic Dyskinesia (PKD) which can be the
51 sole manifestation of the disease in a large number of patients. PKD is characterized by attacks
52 of dystonia and/or chorea, typically triggered by the sudden initiation of movements or brisk
53 modification of an ongoing movement, that lasts less than a minute (4). They usually start in
54 childhood or adolescence and peak during puberty, often with a high frequency of abnormal
55 movement episodes – up to 100 per day – and complete remission between episodes. *PRRT2*-
56 PKD can be treated with low doses of anti-epileptic drugs that modulate the activity of neuronal
57 sodium channels and usually suppress the attacks (5).

58 *PRRT2* is the main culprit gene for PKD (6,7) and *PRRT2*-PKD likely reflects a primary
59 dysfunction of the cerebellum (8,9). The cerebellum has been involved in the pathogenesis of
60 several dystonic syndromes (10). It also plays a role in non-motor processes such as cognition,
61 sleep, and emotional regulation (11,12). Growing evidence indicates that non-motor
62 manifestations are part of the clinical spectrum in various dystonic disorders, including
63 idiopathic adult-onset focal dystonia and genetic forms of dystonia (13–18). Non-motor
64 manifestations can be an important determinant of health-related quality of life (HrQoL),
65 beyond the motor symptoms (15,16,19–23). Perceived stigmatization and diminished self-
66 esteem can also influence quality of life in dystonic disorders (24–26). However, HrQoL, non-
67 motor manifestations, self-esteem, and stigma have been poorly studied to date in patients with
68 paroxysmal dystonic disorders, with only a few pilot findings (27).

69 Here, we investigated non-motor symptoms and how they relate to quality of life in a genetically
70 homogeneous group of consecutive *PRRT2*-PKD patients and their age- and sex-matched
71 controls. We paid special attention to perceived stigmatization and self-esteem.

72 **METHODS**

73 *Study Population*

74 In this exploratory study, we enrolled 21 consecutive PKD patients with a pathogenic variant
75 of *PRRT2* manifesting exclusively as PKD, and healthy controls (HC) matched for age and sex.
76 HCs were recruited from the database of the Pitié-Salpêtrière Clinical Investigation Center
77 (Paris, France). Prior to data collection, the full protocol was approved by the Ethics Committee
78 “Comité de Protection des Personnes Sud-Méditerranée” (reference: 17.060), and was
79 registered on clinicaltrial.gov (NCT03481491). In accordance with the Declaration of Helsinki,
80 all participants provided written informed consent.

81 *Study Instruments*

82 We collected the patients’ history, demographic, genetic, and clinical information including
83 initial misdiagnoses. Disease duration was calculated as the time in years between symptom
84 onset and the day of the inclusion visit. Age at disease onset was taken as the age at which
85 symptoms first occurred. Delay to diagnosis was calculated as the time in years between the
86 patient’s first medical visit for their symptoms and their PKD diagnosis. The residual presence
87 of paroxysmal events was defined as having at least one episode during the month prior to
88 inclusion in the study. Patients were considered treated or non-treated depending on whether
89 they were receiving a treatment for PKD. Non-motor aspects were investigated using
90 standardized questionnaires: i) the Dystonia Non-motor Symptoms (DNMS) questionnaire,
91 which can detect non-motor problems comprising sleep disorders, autonomic dysfunction,
92 fatigue, psychiatric manifestations, stigma, and sensory symptoms (20); ii) the Stigma Scale for
93 Chronic Illness which measures perceived stigmatization (SSCI-8)(28); iii) the Hospital
94 Anxiety and Depression Scale (HADS) to investigate anxiety and depression with two specific
95 sub-scales (29); iv) the Brief Fear of Negative Evaluation Scale (BFNE)(30), the Rosenberg

96 Self Esteem Scale (RSES)(31), and the Warwick-Edinburgh Mental Wellbeing Scales
97 (WEMWBS)(32) which respectively quantify perceived stigmatization, fear of being judged by
98 others, self-esteem, and mental well-being; v) the Medical Outcomes Study 36-Item Short-
99 Form Health Survey (SF-36)(33) to evaluate HrQoL; vi) the Visual Analogue Scale of Pain
100 (34), the Fatigue Severity Scale (FSS-9)(35) and the Pittsburgh Sleep Quality Index (PSQI)(36)
101 to measure pain, fatigue and sleep disturbances, respectively.

102 Component analyses showed that SF-36 measures two distinct concepts, namely a physical
103 dimension, represented by the Physical Component Summary (PCS), and a mental dimension,
104 represented by the Mental Component Summary (MCS). PCS is composed of four primary
105 domains (physical functioning, physical role limitations, bodily pain, and general health). MCS
106 is also composed of four primary domains (social functioning, emotional role limitations,
107 mental health, and vitality). The steps for calculating the PCS and MCS have been previously
108 described (37). The SF-36 scales are standardized using a linear z-score transformation. Z-
109 scores are calculated by subtracting subscale means for a general US population sample from
110 each individual's subscale scores and dividing the difference by the standard deviation of the
111 US sample. Each of the eight z-scores is then multiplied by the corresponding factor scoring
112 coefficient for the scale. There are two different sets of factor scoring coefficients, one for the
113 PCS and another for the MCS. The products of the z-scores and factor scoring coefficients for
114 the PCS are then summed together, and a similar calculation is performed for the MCS. Each
115 resulting sum is multiplied by 10 and added to 50 to linearly transform the PCS or MCS to the
116 T-score metric, which has a mean of 50 and a standard deviation of 10 for the US general
117 population.

118 ***Statistical analysis***

119 Statistical analysis was conducted with the R software (4.0.2) system. Normality in the variable
120 distributions was assessed by the Shapiro-Wilk test. Furthermore, the Levene test was
121 performed to probe homogeneity of variances across groups.

122 We performed group analyses comparing patients to controls. Variables that failed the Shapiro-
123 Wilk or the Levene test were analyzed with nonparametric statistics. Categorical variables were
124 compared using Pearson's χ^2 test or Fisher's exact test. Quantitative variables were analyzed
125 using the Student's t-test or the Mann-Whitney's test.

126 To further explore HrQoL alteration in the patient group, we first used Spearman correlation
127 (continuous variables) and rank-biserial correlation (binary variables: sex and treatment)
128 coefficients to examine the associations between variables which were different between the
129 patients and controls, and between those variables and important characteristics (age, sex, age
130 at onset, disease duration, delay to diagnosis, treatment, or the persistence of PKD attacks) that
131 could influence HrQoL.

132 Second, we performed a multivariable linear regression model. Five variables were selected to
133 be included in the model: three variables were selected based on the statistical significance on
134 between-group comparisons and three variables were selected because we thought they might
135 be important to explain HrQoL (age at onset, treatment - being treated or not - and persistence
136 of paroxysmal events). We only considered the PCS score as the dependent variable (the MCS
137 was not different between the patient and control groups), and the above-mentioned variables
138 as independent variables. The multivariable regression model was based on the stepwise
139 method. Assumptions of using the linear regression method included normal distribution of
140 PCS and lack of outlier data.

141 The data are expressed as mean \pm -SD, and $p < 0.05$ was assumed to be statistically significant.

142 **RESULTS**

143 We analyzed data from 21 (mean age 33.7y) consecutive PKD patients with a pathogenic
144 variant of *PRRT2*, and 21 matched healthy controls (HC). The patients' characteristics and
145 group comparisons are summarized in **Table 1**.

146 *Quality of life evaluation*

147 Compared to the HC, the patients had an altered HrQoL in four domains of the SF-36, namely
148 physical functioning, bodily pain, energy/fatigue, and general health perceptions (**Figure**
149 **1**). Accordingly, the PCS score was lower showing a poorer physical component of HrQoL for
150 the patients ($p < 0.001$). There was no difference between the groups for MCS.

151 *Global non-motor evaluation*

152 There was a group difference for non-motor symptoms (**Table 2**): 100% of the patients and
153 29% of the HCs ($p < 0.001$) reported non-motor symptoms causing impairment of activities of
154 daily living on the DNMS questionnaire. 52% of patients and 9.5% of HCs reported at least
155 four non-motor symptoms. In the patient group, the most prevalent non-motor symptoms were
156 loss of self-confidence due to stigma (100%), sleep disorders (71%) (particularly difficulty to
157 fall asleep), anxiety (48%), fatigue (43%), and depression (29%). The non-motor symptoms
158 burden (i.e., total DNMS score) was higher in the patients (3.76 ± 3.33) than in the HCs (0.57
159 ± 1.52 ; $p < 0.001$).

160 *Psychological evaluation*

161 HADS scores evaluating anxiety and depression related symptoms were similar between the
162 groups ($p = 0.91$), and no subject exceeded cut-off scores. The patients had lower self-esteem
163 (RSES score) ($p = 0.01$) and their fear of being judged by others (BNFE score) tended to be

164 higher ($p=0.067$). Global well-being (WEMWBS) was similar in the patients and HCs
165 ($p=0.152$). The patients had a higher perceived stigmatization with a mean SSCI-8 score of
166 18.8 ($p<0.001$).

167 *Sleep, fatigue, and pain*

168 The patients reported a poorer quality of sleep (PSQI-total score) although not significant
169 ($p=0.116$). They needed more time to fall asleep (30 minutes for patients and 14 minutes for
170 HCs, $p=0.032$). The patients also reported more fatigue (FSS) although the difference was not
171 significant ($p=0.133$). They did not report more pain (VAS) ($p=1$) compared to the HCs.

172 *Misdiagnosis*

173 62 % of the patients were initially misdiagnosed. Wrong diagnoses included functional disorder
174 (6 patients), epilepsy (4 patients), panic attack (2 patients), Tourette syndrome (2 patients), and
175 dystonia (1 patient). Two of them sequentially received two wrong diagnoses (epilepsy and
176 functional disorder for one; Tourette syndrome and functional disorder for the other).

177 *Correlation analysis*

178 When investigating the possible determinants of PCS alteration in the patients, we found that
179 PCS was not related to the presence of residual PKD attacks but negatively correlated with the
180 stigma score SSCI8 ($\rho=-0.480$, $p=0.028$) and time needed to fall asleep ($\rho=-0.544$, $p=0.011$).
181 Higher stigma scores correlated with a higher NMS burden ($\rho=0.554$, $p=0.009$), a greater delay
182 to PKD diagnosis ($\rho=0.615$, $p=0.003$), and lower physical components of HrQoL ($\rho=-0.480$,
183 $p=0.028$).

184 HrQoL was not correlated with the demographical and clinical variables of age, sex, age at
185 onset, disease duration, treatment, or persistence of PKD attacks. Detailed information can be
186 found in **Figure 2** and **Supplementary Material**.

187

188 ***Regression analysis***

189 The independence of PCS predictors in the patients was assessed by stepwise linear regression,
190 with PCS as a response variable and age at onset, DNMS, SSCI, RSES scores and treatment
191 (being treated or not for PKD), and persistence of paroxysmal events as predictive variables
192 (Table 2). From this model (Adjusted $R^2= 0.245$), only the total DNMS was found to predict
193 PCS in the patients ($\beta = -2.06$, 95% CI $-3.628 - -0.487$, $p = 0.013$). Finally, the global non-
194 motor burden was inversely related to the PCS.

195

196 **DISCUSSION**

197 Patients with PKD caused by pathogenic variants of *PRRT2* reported an alteration of the
198 physical aspects of HrQoL, regardless of the presence of residual paroxysmal episodes or use
199 of treatments. Non-motor manifestations observed in these patients were an important
200 determinant of the alteration of HrQoL. In addition, patients perceived a high level of
201 stigmatization positively correlated with delay in diagnosis but not with the presence of
202 paroxysmal episodes. As observed in other dystonic syndromes, non-motor symptoms may be
203 an entire part of the phenotype of patients with paroxysmal dystonic manifestations and should
204 be investigated and recognized in clinical practice. Our findings have important implications
205 for the management of these patients and for medical education about PKD.

206 The main strengths of our work are the clinical (typical PKD) and genetic (*PRRT2* pathogenic
207 variant) homogeneity of the patient group, the presence of a matched comparison group, and
208 the prospective and standardized data collection. Our study was innovative in that we focused
209 on non-motor aspects which represents an emerging concern in the field of dystonic disorders.
210 A few limitations deserve to be mentioned. First, some patients were on chronic treatment,

211 mainly comprising low dose carbamazepine. Although we observed no relevant difference
212 between medicated and non-medicated patients, chronic treatments might have influenced our
213 results. In addition, carbamazepine is commonly used in *PRRT2*-related PKD and thus our
214 results remain overall relevant for the *PRRT2* population. Of note, patients with a milder form
215 of the disease are less likely to receive medication and this should be taken into account when
216 interpreting the lack of difference between medicated and non-medicated patients. Secondly,
217 we binarized the presence or absence of residual PKD attacks in the past month instead of
218 considering their exact reported frequency. The main reason for this was that frequency values
219 are unstable and fluctuant in this disorder, being highly dependent on exposure to triggering
220 factors in activities of daily living. Thirdly, our study only involved a small sample of patients
221 and is thus exploratory, warranting confirmation from larger studies.

222

223 HrQoL is altered in patients with dystonia, including various idiopathic adult-onset focal
224 dystonia (38) and genetic dystonia (14,39,40). A seminal study found an alteration of HrQoL
225 in a large but genetically heterogeneous sample of patients with PKD (27). In this study, the
226 patients' data were compared with normative data in the absence of a control group. In keeping
227 with the genetic heterogeneity of the population, the majority of these heterogeneous PKD
228 patients had an incomplete response to treatment, which is unusual in *PRRT2*-PKD.

229 We also found an alteration of HrQoL in our group of *PRRT2*-PKD patients, as compared with
230 controls. Interestingly, this altered HrQoL was not linked to the presence of residual paroxysmal
231 events or to treatment, suggesting a contribution of non-motor manifestations. Indeed, the
232 global non-motor burden was the only determinant of altered physical components of HrQoL
233 retained by the stepwise regression analysis in our patients.

234 When screened with the DNMS, our *PRRT2*-PKD patients more frequently reported non-motor
235 symptoms as compared to the controls, especially loss of self-confidence due to stigma and
236 sleep problems. Investigation with semi-quantitative dedicated scales similarly identified loss
237 of self-esteem and stigma in the patients, as well as a non-significant trend toward more fatigue
238 and poorer sleep quality compared to the controls. The single questions related to depression
239 and anxiety of the DNMS yielded different responses for patients and controls, which was not
240 confirmed by the between-groups comparison on the HADS subscores. This discrepancy is not
241 surprising as the accuracy of the specific anxiety and depression DNMS questions has not been
242 validated against the anxiety/depression severity scales. Overall, non-motor symptoms have
243 been identified as part of the phenotype in various dystonic disorders and they are an important
244 determinant of quality of life (13,19). In particular, the prevalence of psychiatric disorders, pain,
245 sleep problems and fatigue is often greater in patients with idiopathic adult-onset focal dystonia
246 (20,21,38,41–43). For example, a controlled study using the same evaluation tool (DNMS) in
247 patients with idiopathic adult-onset cervical dystonia revealed that pain, insomnia, and stigma
248 were highly prevalent and that non-motor symptoms were major determinants of their
249 HrQoL(21). A higher prevalence of psychiatric disorders, sleep problems, and fatigue was also
250 found in a mixed group of patients with dystonia – mostly genetic dystonia (dopa-responsive
251 dystonia, myoclonus-dystonia, and idiopathic adult-onset cervical dystonia) – as compared with
252 controls (22). Likewise, non-motor symptoms were also linked to HrQoL in this study.

253 Stigmatization was perceived as important in our *PRRT2*-PKD patients. For example, perceived
254 stigmatization evaluated with the same measurement tool was higher in our patients than in
255 patients with multiple sclerosis or epilepsy, which can be considered to be more visible and
256 debilitating disorders (28,44,45). The young age of our patients may have influenced this result
257 since young patients with neurological disorders seem to be more prone to feelings of
258 stigmatization, as demonstrated in the context of cervical dystonia and other neurological

259 disorders (21,28). As the perceived stigmatization was tightly linked to delay to diagnosis, we
260 suggest that a long history of misdiagnosis – often involving a wrong diagnosis of functional
261 disorders (29% of patients) – may play a role in the high level of perceived stigmatization (44–
262 47). This emphasizes the need to increase knowledge about the diagnostic clues suggestive of
263 PKD.

264 Interestingly, the global non-motor burden did not correlate with the presence of residual
265 paroxysmal episodes. It is therefore possible that the additional non-motor phenotype is not a
266 consequence of motor problems but rather reflects *PRRT2*-related brain dysfunction. If so, then
267 non-motor manifestations could be related to the well-known *PRRT2*-related cerebellar
268 dysfunction (8,48,49). For example, sleep disorders, especially insomnia, have been linked to
269 structural and functional abnormalities of the cerebellum (49,50,51). Consistently, the
270 cerebellum has sleep-related activity patterns, and its output neurons, which express clock
271 genes, are connected with the wake-sleep brain network (49, 51). Alternatively, non-motor
272 manifestations could be the consequence of other brain dysfunctions considering the wide
273 expression of *PRRT2* in the brain (52).

274 Paroxysmal disorders, such as PKD, are often seen as more “benign” compared with continuous
275 movement disorder. Our findings indicate that early diagnosis is also critical in PKD, and
276 suggest that more attention should be paid to the hidden non-motor part of the disease. For this
277 reason, some patients, at least, would benefit from long-term follow-up by a neurologist even
278 after optimal control of the paroxysmal episodes.

279

280 **AUTHOR ROLES**

281 (1) Research project: A. Conception, B. Organization, C. Execution;

282 (2) Statistical Analysis: A. Design, B. Execution, C. Review and Critique;

283 (3) Manuscript: A. Writing of the first draft, B. Review and Critique.

284 A.E.: 1A, 1B, 1C, 2A, 2B, 2C, 3A

285 M.D.: 2A, 2B, 2C, 3A

286 P.J.: 1B, 1C

287 A.H.: 1B, 1C

288 A.M., P.D., J.P., A.R., E.M., C.T., Q.W., M.V: 1C, 3B

289 C.G., Y.W, 1A, 3B

290 E. R. 1A, 1B, 1C, 2C, 3A

291 **FUNDING SOURCES AND CONFLICT OF INTEREST:** This work received financial
292 support from « Association des malades atteints de dystonie (AMADYS) » from « Fondation
293 pour la Recherche Médicale (FRM) » (prix Line-Pomaret, grant attributed to A.E.) and from
294 MERZ-Pharma and Everpharma. This project was also supported by « Agence Nationale de la
295 Recherche (ANR) » under the frame of the European Joint Programme on Rare Diseases (EJP
296 RD, ANR-16-CE37-0003-03). In addition, this project has received funding from the European
297 Union's Horizon 2020 research and innovation programme under the EJP RD COFUND-EJP
298 N° 825575 - EurDyscover. The authors declare that there are no conflicts of interest relevant to
299 this work.

300 **FINANCIAL DISCLOSURES FOR THE PREVIOUS 12 MONTHS:**

301 A.E. has no additional disclosures to report

302 M.D. has no additional disclosures to report

303 A.M. received speaker honoraria from AbbVie and travel support from Merz.

304 P.J. has no additional disclosures to report.

305 A.H. has no additional disclosures to report

306 C.T. received a PhD research grant from the “Fondation pour la Recherche Médicale”

307 Q.W. has no additional disclosures to report

308 P.D. Received honorarium for conferences from Novartis, Roche and Sun Pharma, and for
309 participation to advisory board from Kyowa Kirin.

310 A.R. received honorarium for advisory board from PTC Therapeutics.

311 E.M. received honoraria for speech from Merz-Pharma.

312 J.P. has no additional disclosures to report

313 M.V. has no additional disclosures to report and no conflict of interest

314 Y.W. has no additional disclosures to report and no conflict of interest

315 C.G. has no additional disclosures to report

316 E.R. received honorarium for speech from Orkyn, Aguetant, Elivie and for participating in an
317 advisory board from Merz-Pharma. He received research support from Merz-Pharma, Orkyn,

318 Aguetant, Elivie, Ipsen, Everpharma, Fondation Desmarest, AMADYS, ADCY5.org, Fonds

319 de dotation Patrick Brou de Laurière, Agence Nationale de la Recherche, Société Française de
320 Médecine Esthétique, Dystonia Medical Research Foundation.

321 **ETHICAL COMPLIANCE STATEMENT**

322 This study was approved by the Regional Ethics Committee “Comité de Protection des
323 Personnes Sud-Méditerranée” (reference 17.060) and was registered on ClinicalTrials.gov
324 (NCT03481491). Informed consent was obtained from all study participants. We confirm that
325 we have read the Journal’s position on issues involved in ethical publication and affirm that this
326 work is consistent with those guidelines.

327 **REFERENCES**

- 328 1. Ebrahimi-Fakhari D, Saffari A, Westenberger A, Klein C. The evolving spectrum of
329 *PRRT2* -associated paroxysmal diseases. *Brain*. 2015 Dec;138(12):3476–95.
- 330 2. Landolfi A, Barone P, Erro R. The Spectrum of *PRRT2*-Associated Disorders: Update
331 on Clinical Features and Pathophysiology. *Front Neurol*. 2021 Mar 4;12:629747.
- 332 3. Méneret A, Gaudebout C, Riant F, Vidailhet M, Depienne C, Roze E. *PRRT2*
333 mutations and paroxysmal disorders. *Eur J Neurol*. 2013 Jun;20(6):872–8.
- 334 4. Méneret A, Roze E. Paroxysmal movement disorders: An update. *Revue*
335 *Neurologique*. 2016 Aug;172(8–9):433–45.
- 336 5. Méneret A, Garcin B, Frismand S, Lannuzel A, Mariani LL, Roze E. Treatable
337 Hyperkinetic Movement Disorders Not to Be Missed. *Front Neurol*. 2021 Dec 1;12:659805.
- 338 6. Chen YP, Song W, Yang J, Zheng ZZ, Huang R, Chen K, et al. *PRRT2* mutation
339 screening in patients with paroxysmal kinesigenic dyskinesia from Southwest China. *Eur J*
340 *Neurol*. 2014 Jan;21(1):174–6.
- 341 7. Meneret A, Grabli D, Depienne C, Gaudebout C, Picard F, Durr A, et al. *PRRT2*
342 mutations: A major cause of paroxysmal kinesigenic dyskinesia in the European population.

- 343 Neurology. 2012 Jul 10;79(2):170–4.
- 344 8. Ekmen A, Meneret A, Valabregue R, Beranger B, Worbe Y, Lamy JC, et al.
345 Cerebellum Dysfunction in Patients With *PRRT2* -Related Paroxysmal Dyskinesia.
346 Neurology. 2022 Mar 8;98(10):e1077–89.
- 347 9. Lu B, Lou SS, Xu RS, Kong DL, Wu RJ, Zhang J, et al. Cerebellar spreading
348 depolarization mediates paroxysmal movement disorder. Cell Reports. 2021
349 Sep;36(12):109743.
- 350 10. Bologna M, Berardelli A. The cerebellum and dystonia. In: Handbook of Clinical
351 Neurology [Internet]. Elsevier; 2018 [cited 2022 Oct 8]. p. 259–72. Available from:
352 <https://linkinghub.elsevier.com/retrieve/pii/B9780444641892000172>
- 353 11. Schmahmann JD. The cerebellum and cognition. Neuroscience Letters. 2019
354 Jan;688:62–75.
- 355 12. Canto CB, Onuki Y, Bruinsma B, van der Werf YD, De Zeeuw CI. The Sleeping
356 Cerebellum. Trends in Neurosciences. 2017 May;40(5):309–23.
- 357 13. Junker J, Berman BD, Hall J, Wahba DW, Brandt V, Perlmutter JS, et al. Quality of
358 life in isolated dystonia: non-motor manifestations matter. J Neurol Neurosurg Psychiatry.
359 2021 Jun;92(6):622–8.
- 360 14. Peall KJ. Non-motor symptoms in genetically defined dystonia: Homogenous groups
361 require systematic assessment. 2015;10.
- 362 15. Rafee S, Ndukwe I, O’Riordan S, Hutchinson M. Reliability of DNMSQUEST as a
363 Screening Tool for Mood Disorders in Cervical Dystonia. Movement Disord Clin Pract. 2021
364 Aug;8(6):925–31.
- 365 16. Smit M, Kuiper A, Han V, Jiawan VCR, Douma G, van Harten B, et al. Psychiatric
366 co-morbidity is highly prevalent in idiopathic cervical dystonia and significantly influences
367 health-related quality of life: Results of a controlled study. Parkinsonism & Related

- 368 Disorders. 2016 Sep;30:7–12.
- 369 17. Wadon ME, Bailey GA, Yilmaz Z, Hubbard E, AlSaeed M, Robinson A, et al.
370 Non-motor phenotypic subgroups in adult-onset idiopathic, isolated, focal cervical dystonia.
371 Brain Behav [Internet]. 2021 Aug [cited 2022 Aug 26];11(8). Available from:
372 <https://onlinelibrary.wiley.com/doi/10.1002/brb3.2292>
- 373 18. Bailey GA, Martin E, Peall KJ. Cognitive and Neuropsychiatric Impairment in
374 Dystonia. Curr Neurol Neurosci Rep [Internet]. 2022 Oct 6 [cited 2022 Oct 11]; Available
375 from: <https://link.springer.com/10.1007/s11910-022-01233-3>
- 376 19. Smit M, Albanese A, Benson M, Edwards MJ, Graessner H, Hutchinson M, et al.
377 Dystonia Management: What to Expect From the Future? The Perspectives of Patients and
378 Clinicians Within DystoniaNet Europe. Front Neurol. 2021 Jun 3;12:646841.
- 379 20. Klingelhofer L, Chaudhuri KR, Kamm C, Martinez-Martin P, Bhatia K, Sauerbier A,
380 et al. Validation of a self-completed Dystonia Non-Motor Symptoms Questionnaire. Annals
381 of Clinical and Translational Neurology. 2019;6(10):2054–65.
- 382 21. Klingelhofer L, Kaiser M, Sauerbier A, Untucht R, Wienecke M, Mammadova K, et
383 al. Emotional well-being and pain could be a greater determinant of quality of life compared
384 to motor severity in cervical dystonia. J Neural Transm. 2021 Mar;128(3):305–14.
- 385 22. Timmers ER, Smit M, Kuiper A, Bartels AL, van der Veen S, van der Stouwe AMM,
386 et al. Myoclonus-dystonia: Distinctive motor and non-motor phenotype from other dystonia
387 syndromes. Parkinsonism & Related Disorders. 2019 Dec;69:85–90.
- 388 23. Wagle Shukla A, Brown R, Heese K, Jones J, Rodriguez RL, Malaty IM, et al. High
389 rates of fatigue and sleep disturbances in dystonia. International Journal of Neuroscience.
390 2016 Oct 2;126(10):928–35.
- 391 24. Ben-Shlomo Y. What are the determinants of quality of life in people with cervical
392 dystonia? Journal of Neurology, Neurosurgery & Psychiatry. 2002 May 1;72(5):608–14.

- 393 25. Gharzai L, Harney J, Higgins S, High R, Xu L, Inciarte D, et al. Alarming levels of
394 stigma toward generalized dystonia: A cross-cultural comparison. *Clinical Parkinsonism &*
395 *Related Disorders*. 2020;3:100059.
- 396 26. Rinnerthaler M, Mueller J, Weichbold V, Wenning GK, Poewe W. Social
397 stigmatization in patients with cranial and cervical dystonia. *Mov Disord*. 2006
398 Oct;21(10):1636–40.
- 399 27. Tian WT, Huang XJ, Liu XL, Shen JY, Liang GL, Zhu CX, et al. Depression, anxiety,
400 and quality of life in paroxysmal kinesigenic dyskinesia patients. *Chinese Medical Journal*.
401 2017;130(17):2088–94.
- 402 28. Molina Y, Choi SW, Cella D, Rao D. The Stigma Scale for Chronic Illnesses 8-Item
403 Version (SSCI-8): Development, Validation and Use Across Neurological Conditions. *IntJ*
404 *Behav Med*. 2013 Sep;20(3):450–60.
- 405 29. Bocéréan C, Dupret E. A validation study of the Hospital Anxiety and Depression
406 Scale (HADS) in a large sample of French employees. *BMC Psychiatry*. 2014 Dec;14(1):354.
- 407 30. Leary MR. A brief version of the Fear of Negative Evaluation Scale. *Personality and*
408 *Social Psychology Bulletin*. 1983. 371–375 p.
- 409 31. Sinclair SJ, Blais MA, Gansler DA, Sandberg E, Bistis K, LoCicero A. Psychometric
410 properties of the Rosenberg Self-Esteem Scale: overall and across demographic groups living
411 within the United States. *Eval Health Prof*. 2010 Mar;33(1):56–80.
- 412 32. Trousselard M, Steiler D, Dutheil F, Claverie D, Canini F, Fenouillet F, et al.
413 Validation of the Warwick-Edinburgh Mental Well-Being Scale (WEMWBS) in French
414 psychiatric and general populations. *Psychiatry Res*. 2016 Nov 30;245:282–90.
- 415 33. Ware JEJ, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I.
416 Conceptual framework and item selection. *Med Care*. 1992 Jun;30(6):473–83.
- 417 34. Boonstra AM, Schiphorst Preuper HR, Reneman MF, Posthumus JB, Stewart RE.

- 418 Reliability and validity of the visual analogue scale for disability in patients with chronic
419 musculoskeletal pain. *Int J Rehabil Res.* 2008 Jun;31(2):165–9.
- 420 35. Krupp LB, LaRocca NG, Muir-Nash J, Steinberg AD. The Fatigue Severity Scale:
421 Application to Patients With Multiple Sclerosis and Systemic Lupus Erythematosus. *Archives*
422 *of Neurology.* 1989 Oct 1;46(10):1121–3.
- 423 36. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep
424 quality index: A new instrument for psychiatric practice and research. *Psychiatry Research.*
425 1989 May 1;28(2):193–213.
- 426 37. Ware JEJ, Kosinski M, Bayliss MS, McHorney CA, Rogers WH, Raczek A.
427 Comparison of methods for the scoring and statistical analysis of SF-36 health profile and
428 summary measures: summary of results from the Medical Outcomes Study. *Med Care.* 1995
429 Apr;33(4 Suppl):AS264-279.
- 430 38. Yang J, Shao N, Song W, Wei Q, Ou R, Wu Y, et al. Nonmotor symptoms in primary
431 adult-onset cervical dystonia and blepharospasm. *Brain and Behavior.* 2017;7(2):1–6.
- 432 39. Coenen MA, Eggink H, Tijssen MA, Spikman JM. Cognition in childhood dystonia: a
433 systematic review. *Dev Med Child Neurol.* 2018 Mar;60(3):244–55.
- 434 40. Eggink H, Coenen MA, de Jong R, Toonen RF, Eissens MH, Veenstra WS, et al.
435 Motor and non-motor determinants of health-related quality of life in young dystonia patients.
436 *Parkinsonism and Related Disorders.* 2019;58(August):50–5.
- 437 41. Novaretti N, Cunha ALN, Bezerra TC, Pereira MAP, De Oliveira DS, Macruz Brito
438 MMC, et al. The prevalence and correlation of non-motor symptoms in adult patients with
439 idiopathic focal or segmental dystonia. *Tremor and Other Hyperkinetic Movements.*
440 2019;9:1–7.
- 441 42. Ferrazzano G, Berardelli I, Conte A, Baione V, Concolato C, Belvisi D, et al. Motor
442 and non-motor symptoms in blepharospasm: clinical and pathophysiological implications.

- 443 Journal of Neurology. 2019;266(11):2780–5.
- 444 43. Berman BD, Junker J, Shelton E, Sillau SH, Jinnah HA, Perlmutter JS, et al.
445 Psychiatric associations of adult-onset focal dystonia phenotypes. *J Neurol Neurosurg*
446 *Psychiatry*. 2017 Jul;88(7):595–602.
- 447 44. Dayapoglu N, Ayyıldız Nİ, Şeker D. Fear of negative evaluation and the concealment
448 of their disease by epilepsy patients. *Epilepsy & Behavior*. 2020 Dec;113:107539.
- 449 45. Kuramochi I, Iwayama T, Horikawa N, Shimotsu S, Watanabe S, Yamanouchi H, et
450 al. Development and validation of the Epilepsy Self-Stigma Scale. *Epilepsia Open*. 2021
451 Dec;6(4):748–56.
- 452 46. Klinke ME, Hjartardóttir TE, Hauksdóttir A, Jónsdóttir H, Hjaltason H, Andrésdóttir
453 GT. Moving from stigmatization toward competent interdisciplinary care of patients with
454 functional neurological disorders: focus group interviews. *Disability and Rehabilitation*. 2021
455 Apr 24;43(9):1237–46.
- 456 47. Ahern L, Stone J, Sharpe MC. Attitudes of neuroscience nurses toward patients with
457 conversion symptoms. *Psychosomatics*. 2009 Aug;50(4):336–9.
- 458 48. Argyropoulos GPD, van Dun K, Adamaszek M, Leggio M, Manto M, Masciullo M, et
459 al. The Cerebellar Cognitive Affective/Schmahmann Syndrome: a Task Force Paper.
460 *Cerebellum*. 2020 Feb;19(1):102–25.
- 461 49. Salazar Leon LE, Sillitoe RV. Potential Interactions Between Cerebellar Dysfunction
462 and Sleep Disturbances in Dystonia. *Dyst*. 2022 Oct 4;1:10691.
- 463 50. Holub F, Petri R, Schiel J, Feige B, Rutter MK, Tamm S, et al. Association between
464 insomnia symptoms and functional connectivity in the UK Biobank cohort (n = 29,423). *J*
465 *Sleep Res*. 2022 Dec 18; e13790
- 466 51 Canto CB, Onuki Y, Bruinsma B, Van der Werf YD, De Zeeuw CI. The sleeping
467 cerebellum. *Trends Neurosci* 2017 May; 40(5):309-23

468 52. Chen WJ, Lin Y, Xiong ZQ, Wei W, Ni W, Tan GH, et al. Exome sequencing
469 identifies truncating mutations in PRRT2 that cause paroxysmal kinesigenic dyskinesia. Nat
470 Genet. 2011 Nov 20;43(12):1252–5.

471

472 **FIGURE LEGENDS**

473

474 **Figure 1:** Radar chart with the eight domains of the Quality of life (Short Form-36) for PRRT2
475 patients and Healthy Controls (red line: PRRT2 patients, blue line: Healthy Controls). Each
476 spoke on the radar chart represents an SF-36 domain on a scale of 0-100, with higher scores
477 representing better health-related quality of life. The four green domains (Physical functioning,
478 Role physical, Bodily pain and General health) contribute to the scoring of PCS measure. The
479 four orange domains (Vitality, Social functioning, Role emotional and Mental Health)
480 contribute to the scoring of MCS measure. *** $p < 0.001$, ** $p < 0.01$, * < 0.05 , using *Student's t-*
481 *test or the Mann-Whitney's test* as appropriate.

482 **Figure 2:** Correlation matrix across Age, Sex, Age at onset, Disease duration, Diagnostic delay,
483 PCS, DNMS, SSC18, RSES, Paroxysmal events and Treatment. Color of each box indicates the
484 significance of the correlation (Spearman's ρ or rank-biserial correlation coefficient), while size
485 of the circles indicates its magnitude.

486

487

488 **TABLES**

Table 1: Characteristics of subjects and results of the non-motor evaluation

Characteristics	PRRT2 Patients (n = 21)	Healthy Controls (n = 21)	P value
Age; Age range (years)	33.66 (\pm 12.37); 17 – 51	31.7 (\pm 12.1); 21 – 56	0.55 [‡]
Sex (female/male)	7/14	7/14	1 [†]
Age at onset (years)	9.3 (\pm 3.0)	–	
Disease duration (years)	22.4 (\pm 12.5)	–	
Diagnostic delay (years)	9.33 (\pm 10.78)	–	
Treated patients n=11) CBZ/LTG/LEV)	9/1/1	–	
Residual Paroxysmal events (yes/no)	6/15		
PCS	52.53 (\pm 6.34)	58.52 (\pm 4.02)	0.001[¶]
MCS	41.57 (\pm 10.80)	42.59 (\pm 9.36)	0.743 [¶]
DNMS	3.76 (\pm 1.64)	0.57 (\pm 1.21)	0.000[‡]
PSQI	6.19 (\pm 3.71)	4.43 (\pm 3.38)	0.116 [¶]
HA	2.05 (\pm 2.06)	2.38 (\pm 2.80)	0.612
HD	4.10 (\pm 2.59)	4.19 (\pm 3.87)	0.969 [‡]
SSCI-8	18.38 (\pm 4.86)	10.00 (\pm 2.06)	0.000[‡]
RSES	21.05 (\pm 5.71)	25.52 (\pm 3.59)	0.003[‡]
BFNE	39.04 (\pm 9.74)	34.76 (\pm 8.32)	0.067 [¶]
WEMWBS	50.57 (\pm 9.58)	53.71 (\pm 9.97)	0.152 [¶]
VAS	5%	0%	1 [†]
FSS	25.19 (\pm 17.32)	17.71 (\pm 10.32)	0.133 [‡]

Legend: **CBZ** = carbamazepine; **LTG** = lamotrigine; **LEV** = levetiracetam; **PCS** = Physical Component Summary; **MCS** = Mental Component Summary; **DNMS** = Dystonia Non-Motor Symptoms questionnaire; **PSQI** = Pittsburgh Sleep Quality Index; **HA** = Hospital Anxiety scale; **HD** = Hospital Depression scale; **SSCI-8** = Stigma Scale for Chronic Illnesses 8-item; **RSES** = Rosenberg Self-Esteem Scale; **BFNE** = Brief Fear of Negative Evaluation Scale; **WEMWBS** = Warwick-Edinburgh Mental Wellbeing Scale; **VAS** = Visual Analogue Scale for pain; **FSS** = Fatigue Severity Scale.

Quantitative data are expressed as mean (\pm SD).

[†] Student's t-test; [‡] Pearson's χ^2 test; [‡] Mann-Whitney test; [†] Fisher exact test; Significant results indicated in bold: $p < 0.05$.

490
491

Table 2: Non-motor symptoms

Symptoms	Patients♦	Controls♦	p-value
Sleep disorders	61.90%	9.52%	< 0.001 †
Difficulties to fall asleep	61.90%	9.52%	< 0.001 †
Dizziness	0%	0%	—
Fatigue	42.86%	14.29%	0.040 *†
Anxiety	47.62%	19.05%	0.050*†
Depression	28.57%	0%	0.008 *§
Loss of self-confidence due to stigma	100%	0%	< 0.001 †
Flat moods	28.57%	4.76%	0.038 *§
Chewing/swallowing	0%	0%	—
Unpleasant sensations	0%	0%	—
Speech	0%	0%	—
Vision	0%	0%	—
Pain	4.76%	0%	0.311§
Balance/walking	0%	0%	—

Legend: ♦ Percentage of positive response of the subjects to the 14 items of the dystonia non motor symptoms questionnaire † Pearson’s χ^2 test; § Fisher exact. Significant results indicated in bold: p<0.05.

492

493

494 **SUPPLEMENTARY LEGENDS**

495

496 ***Supplementary Table 1:*** Data of the eight domains of the Short Form (36) health survey for

497 PRRT2 patients and Healthy Controls (HC). PCS= Physical Component Summary,

498 MCS=Mental Component Summary, PF=Physical functioning, RP=Role physical, RE=Role

499 emotional, VT=Vitality, MH=Mental Health, SF=Social functioning, BP=Bodily pain and

500 GH=General health, ¶ Student's t-test; ‡ Mann-Whitney test.

501

502 ***Supplementary Table 2:*** Correlations across Age, Sex, Age at onset, Disease

503 duration, Diagnostic delay, Physical Component Summary (PCS) of the Short Form (36)

504 health survey, Dystonia non-motor symptoms scale (DNMS), Stigma Scale for Chronic

505 Illness (SSCI8), Rosenberg Self Esteem Scale (RSES), Paroxysmal Events and Treatment.

506

507




Figure 1. Radar chart with the eight domains of the Quality of life (Short Form-36) for PRRT2 patients and Healthy Controls (red line: PRRT2 patients, blue line: Healthy Controls). Each spoke on the radar chart represents an SF-36 domain on a scale of 0-100, with higher scores representing better health-related quality of life. The four green domains (Physical functioning, Role physical, Bodily pain and General health) contribute to the scoring of PCS measure. The four orange domains (Vitality, Social functioning, Role emotional and Mental Health) contribute to the scoring of MCS measure. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, using Student's t-test or the Mann-Whitney's test as appropriate.

1075x642mm (118 x 118 DPI)




Figure 2. Correlation matrix across Age, Sex, Age at onset, Disease duration, Diagnostic delay, PCS, DNMS, SSCI8, RSES, Paroxysmal events and Treatment. Color of each box indicates the significance of the correlation (Spearman's ρ), while size of the circles indicates its magnitude.

904x462mm (236 x 236 DPI)

1 **Supplementary Results**

2 **Quality of life**

3 Supplementary Table 1: Quality of life (Short Form-36).

4 PCS= Physical Component Summary, MCS=Mental Component Summary, PF=Physical
 5 functioning, RP=Role physical, RE=Role emotional, VT=Vitality, MH=Mental Health,
 6 SF=Social functioning, BP=Bodily pain and GH=General health, ¶ Student's t-test; ‡ Mann-
 7 Whitney test;

8

9

	Healthy controls (mean±sd)	PRRT2 patients (mean±sd)	Mean difference (95% Confidence Interval)	p- value
PF	97.62±6.82	92.86±7.68	4.76 (0.23-9.29)	0.006‡
RP	80.95±30.52	67.86±35.48	13.1 (-7.56-33.75)	0.189‡
RE	73±41.69	69.86±43.35	3.14 (-23.38-29.67)	0.768‡
VT	59.05±12.81	48.57±16.89	10.48 (1.11-19.85)	0.029¶
MH	65.71±14.56	59.24±13.42	6.48 (-2.26-15.21)	0.142¶
SF	72.76±21.83	78.14±21.91	-5.38 (-19.02-8.26)	0.417‡
BP	97.9±4.45	85±21.49	12.9 (2.96-22.85)	0.023‡

GH	84.29±10.28	61.43±22.92	22.86 (11.62-34.09)	< 0.001 [¶]
PCS	58.52±4.02	52.53±6.34	6 (2.67-9.32)	0.001 [¶]
MCS	42.59±9.36	41.57±10.8	1.03 (-5.28-7.33)	0.96 [‡]

10

11 Supplementary Table 2: Correlations across Age, Sex, Age at onset, Disease12 duration, Diagnostic delay, PCS, DNMS, SSCI8, RSES, Paroxysmal Events and Treatment.

	Age	Sex	Age at onset	Disease duration	Diagnostic delay	PCS	DNMS	SSCI-8	RSES	Paroxysmal events	Treatment
Age	1	-0.008	0.122	0.944**	0.435*	0.02	0.405	0.282	0.734**	-0.044	-0.075
Sex		1	0.623**	-0.167	-0.176	0.1	0.246	-0.295	-0.336	-0.224	-0.143
Age at onset			1	-0.155	-0.169	-0.182	0.217	-0.199	-0.133	0.105	-0.488*
Disease duration				1	0.504*	0.048	0.355	0.378	0.739**	-0.061	0.05
Diagnostic delay					1	-0.016	0.442*	0.615**	0.439*	-0.009	0.328
PCS						1	-0.405	-0.480*	0.068	0.035	0.15
DNMS							1	0.554**	-0.044	-0.292	0.323
SSCI-8								1	0.174	0.203	0.211
RSES									1	0.053	-0.076
Paroxysmal events										1	-0.447*
Treatment											1

13

14