



HAL
open science

Large population limit of the spectrum of killed birth-and-death processes

J.-R Chazottes, P Collet, S Méléard

► **To cite this version:**

J.-R Chazottes, P Collet, S Méléard. Large population limit of the spectrum of killed birth-and-death processes. *Journal of Functional Analysis*, 2023, 285 (9), pp.110092. 10.1016/j.jfa.2023.110092 . hal-04306715

HAL Id: hal-04306715

<https://hal.science/hal-04306715>

Submitted on 25 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Large population limit of the spectrum of killed birth-and-death processes

J.-R. Chazottes^(a), P. Collet^(a), S. Méléard^(b)

^(a) CPHT, CNRS, Ecole polytechnique, IP Paris, Palaiseau, France

^(b) CMAP, CNRS, Ecole polytechnique, IP Paris, Palaiseau, France

March 22, 2023

Abstract

We consider a general class of birth-and-death processes with state space $\{0, 1, 2, 3, \dots\}$ which describes the size of a population going eventually to extinction with probability one. We obtain the complete spectrum of the generator of the process killed at 0 in the large population limit, that is, we scale the process by a parameter K , and take the limit $K \rightarrow +\infty$. We assume that the differential equation $dx/dt = b(x) - d(x)$ describing the infinite population limit (in any finite-time interval) has a repulsive fixed point at 0, and an attractive fixed point $x_* > 0$. We prove that, asymptotically, the spectrum is the superposition of two spectra. One is the spectrum of the generator of an Ornstein-Uhlenbeck process, which is $n(b'(x_*) - d'(x_*))$, $n \geq 0$. The other one is the spectrum of a continuous-time binary branching process conditioned on non-extinction, and is given by $n(d'(0) - b'(0))$, $n \geq 1$. A major difficulty is that different scales and function spaces are involved. We work at the level of the eigenfunctions that we split over different regions, and study their asymptotic dependence on K in each region. In particular, we prove that the spectral gap goes to $\min\{b'(0) - d'(0), d'(x_*) - b'(x_*)\}$. This work complements a previous work of ours in which we studied the approximation of the quasi-stationary distribution and of the mean time to extinction.

Key-words: Ornstein-Uhlenbeck process, quantum harmonic oscillator, binary branching process, Jacobi operators, Dirichlet form, spectral gap, Fréchet-Kolmogorov-Riesz compactness criterion, discrete orthogonal polynomials, quasi-eigenvectors, quasi-stationary distribution.

Contents

1	Introduction, heuristics, and main result	3
1.1	The context	3
1.2	Notations for basic function spaces	4
1.3	Heuristics	4
1.4	Main result	6
1.5	Consequences on relaxation times	9
1.6	Organization of the paper	10
2	Standing assumptions	10
3	Proof of Theorem 1.2	11
3.1	Some useful operators	11
3.2	Steps of the proof of Theorem 1.2	16
4	Properties of the eigenvectors	19
5	Proofs of Theorems 3.5 and 3.7	26
5.1	Proof of Theorem 3.5	26
5.2	Proof of Theorem 3.7	27
6	Fréchet-Kolmogorov-Riesz compactness criterion and Dirichlet form	30
6.1	Fréchet-Kolmogorov-Riesz compactness criterion	30
6.2	Dirichlet form for the operator \mathcal{L}_K	36
7	Spectral theory of \mathcal{M}_0	38
A	Local maximum principle and consequences thereof	40
B	Quasi-eigenvalues and quasi-eigenvectors of self-adjoint operators	41

1 Introduction, heuristics, and main result

1.1 The context

We consider a class of birth-and-death processes $(X_t^K)_{t \geq 0}$ with state space $\mathbb{Z}_{\geq 0}$ ¹ which describes how the size of a single population evolves according to birth and death rates of the form

$$\lambda_n^{(K)} = Kb \left(\frac{n}{K} \right) \quad \text{and} \quad \mu_n^{(K)} = Kd \left(\frac{n}{K} \right) \quad (1.1)$$

where $n \geq 1$, and $K \in \mathbb{Z}_{>0}$ is a scaling parameter, which can be for instance interpreted as the ‘carrying capacity’ in the logistic model. We suppose that $b(0) = d(0) = 0$, implying that 0 is an absorbing state for the process, modeling extinction, and our assumptions are such that the probability to reach this state is equal to one. The unique stationary distribution is the Dirac measure at 0, so a relevant distribution to look for is a quasi-stationary distribution. A probability measure $\nu^{(K)}$ on the positive integers is a quasi-stationary distribution if, for all $t > 0$ and for all subsets $A \subset \mathbb{Z}_{>0}$, one has $\mathbb{P}_{\nu^{(K)}}(X_t^K \in A | T_0^{(K)} > t) = \nu^{(K)}(A)$, where $T_0^{(K)}$ is the extinction time, that is, the smallest $t > 0$ such that $X_t^{(K)} = 0$. In other words, a quasi-stationary distribution plays the role of a stationary distribution when conditioning upon non-extinction. We refer to [6, 11] for more informations about quasi-stationary distributions.

When $K \rightarrow +\infty$, the trajectories of the rescaled process $(K^{-1}X_t^{(K)})_{t > 0}$ converge in probability, in any fixed time-window, to the solutions of the differential equation

$$\frac{dx}{dt} = b(x) - d(x) \quad (1.2)$$

if the initial condition state is for instance of the form $\lfloor Kx_0 \rfloor$ for a given $x_0 > 0$. We assume that the functions b and d only vanish at 0, and that

$$d'(0) - b'(0) < 0$$

meaning that the fixed point 0 is repulsive. We also assume that there is a unique attractive fixed point $x_* > 0$, that is

$$b(x_*) = d(x_*) \quad \text{and} \quad b'(x_*) - d'(x_*) < 0.$$

We will give the complete set of assumptions on the functions b and d later on (see Section 2).

A famous example is the so-called logistic process for which $b(x) = \lambda x$, $d(x) = x(\mu + x)$, where λ and μ are positive real numbers. We assume that $\lambda > \mu$ and we have $x_* = \lambda - \mu$.

¹We denote by $\mathbb{Z}_{\geq 0}$ the set of non-negative integers, and by $\mathbb{Z}_{>0}$ the set of positive integers.

In [5] we obtained the precise asymptotic behavior of the first eigenvalue of the generator L_K of the process killed at 0, and also of the law of the extinction time starting from the quasi-stationary distribution (among other results). Here we go further and obtain the complete spectrum of the generator of the killed process, in the limit $K \rightarrow +\infty$. In particular, the knowledge of the spectral gap allows us to obtain the time of relaxation for the process conditioned on non-extinction to obey the quasi-stationary distribution.

1.2 Notations for basic function spaces

We denote by

- \mathcal{D} the space of C^∞ \mathbb{C} -valued functions with compact support on \mathbb{R} ;
- c_{00} the space of \mathbb{C} -valued sequences with finitely many nonzero values;
- ℓ^2 the space of square-summable \mathbb{C} -valued sequences equipped with the standard scalar product.
- L^2 the space of square-integrable \mathbb{C} -valued functions with respect to Lebesgue measure on \mathbb{R} .

We will define several operators on c_{00} and will consider their closure on ℓ^2 . For simplicity, we will use the same notation for an operator and its closure. As we will see later, there is no ambiguity on the extensions.

1.3 Heuristics

The fundamental object in this paper is the spectrum of the following operator that we momentarily define on c_{00} :

$$(L_K v)(n) = \lambda_n^{(K)}(v(n+1) - v(n)) + \mu_n^{(K)}(v(n-1)\mathbb{1}_{\{n>1\}} - v(n)) \quad (1.3)$$

for $n \in \mathbb{Z}_{>0}$. The idea is to ‘localize’ this operator either around $n = \lfloor Kx_* \rfloor$ or $n = 1$, which corresponds in the dynamical system to the fixed point x_* or the fixed point 0. A natural idea would be to ‘cut’ the operators in order to differentiate these two dynamics. However the main difficulty is that the two different pieces involve different scales and different function spaces. Since we don’t know how to cope with this problem at the level of operators, we work at the level of the eigenfunctions that we will split on different regions, and study their asymptotic dependence on K in each region.

To have an idea of the different scales involved in the problem, let us first study the asymptotic behavior of the birth and death rates. By Taylor expansion around Kx_* we have

$$\lambda_n^{(K)} = K b(x_*) + (n - Kx_*) b'(x_*) + \mathcal{O}\left(\frac{(n - Kx_*)^2}{K}\right)$$

and

$$\mu_n^{(K)} = K d(x_*) + (n - Kx_*) d'(x_*) + \mathcal{O}\left(\frac{(n - Kx_*)^2}{K}\right).$$

Taking $v^{(K)}(n) = u((n - Kx_*)/\sqrt{K})$ with $u \in \mathcal{D}$, we get

$$(L_K v^{(K)})(n) = \mathcal{O}u_* u\left(\frac{n - Kx_*}{\sqrt{K}}\right) + \mathcal{O}\left(\frac{1}{\sqrt{K}}\right)$$

where

$$\mathcal{O}u_* f(x) = b(x_*) f''(x) + (b'(x_*) - d'(x_*)) x f'(x) \quad (1.4)$$

is the generator of the Ornstein-Uhlenbeck process on \mathbb{R} which satisfies the stochastic differential equation $dX_t = (b'(x_*) - d'(x_*)) X_t dt + \sqrt{2b(x_*)} dB_t$, where $(B_t)_{t \geq 0}$ is a one-dimensional Brownian motion. It is well known (see Remark 3.1) that the spectrum of $\mathcal{O}u_*$ is $-S_1$, where

$$S_1 = (d'(x_*) - b'(x_*)) \mathbb{Z}_{\geq 0} \quad (1.5)$$

in the space $L^2\left(\sqrt{\frac{d'(x_*) - b'(x_*)}{2\pi b(x_*)}} e^{-\frac{(d'(x_*) - b'(x_*)) x^2}{2b(x_*)}} dx\right)$.

Now we look at n near 1. By Taylor expansion, we have

$$\lambda_n^{(K)} = n \left(b'(0) + \mathcal{O}\left(\frac{n}{K}\right) \right) \quad \text{and} \quad \mu_n^{(K)} = n \left(b'(0) + \mathcal{O}\left(\frac{n}{K}\right) \right).$$

If $v \in c_{00}$, then we get

$$\|L_K v - \mathcal{Q}_0 v\|_{\ell^2} \leq \frac{\mathcal{O}(1)}{K}$$

where

$$(\mathcal{Q}_0 v)(n) = b'(0) n (v(n+1) - v(n)) + d'(0) n (v(n-1) \mathbb{1}_{\{n>1\}} - v(n))$$

which is the generator of a (continuous-time) binary branching process killed at 0. We shall prove later on (see Theorem 7.1 in Section 7) that, in a suitable 'weighted' ℓ^2 space defined below, the spectrum of \mathcal{Q}_0 is $-S_2$ where

$$S_2 = (b'(0) - d'(0)) \mathbb{Z}_{>0}. \quad (1.6)$$

The previous observations suggest that the limit of the spectrum of the generator of the birth-and-death process $(X_t^{(K)})_{t \geq 0}$, in an appropriate space, is

$$(d'(0) - b'(0)) \mathbb{Z}_{>0} \cup (b'(x_*) - d'(x_*)) \mathbb{Z}_{\geq 0}.$$

Notice that all the elements of this set are negative and this is not a disjoint union in general. The logistic model is an example illustrating this, since $d'(0) - b'(0) = b'(x_*) - d'(x_*) = \mu - \lambda$, so we will have asymptotic double eigenvalues in this case.

We will prove that the limit of the spectrum of L_K is obtained from the explicit spectra of the above two operators. Notice that one limit is a differential operator whereas the other one is still a finite-difference operator. We look for the asymptotics of (a part of) the spectrum of a discrete operator in terms of the spectrum of a differential operator. This is a reverse approach with respect to numerical analysis where the spectra of limiting differential operators are obtained from the knowledge (computation) of the spectrum of finite-difference operators. See for instance [4].

1.4 Main result

For each $K \in \mathbb{Z}_{>0}$, the sequence of numbers

$$\pi_n^{(K)} := \frac{\lambda_1^{(K)} \cdots \lambda_{n-1}^{(K)}}{\mu_1^{(K)} \cdots \mu_n^{(K)}}, \quad n \geq 2 \quad \text{and} \quad \pi_1^{(K)} := \frac{1}{\mu_1^{(K)}} \quad (1.7)$$

naturally shows up in the study of birth-and-death processes. We will give below a set of assumptions on the functions b and d , defining the differential equation (1.2), ensuring that the process reaches 0 in finite time with probability one, that the mean-time to extinction is finite, and that the quasi-stationary distribution exists and is unique.

Let $\ell^2(\pi^{(K)})$ be the space of \mathbb{C} -valued sequences $(v_n)_{n \geq 1}$ such that

$$\sum_{n \geq 1} |v(n)|^2 \pi_n^{(K)} < +\infty.$$

This is a Hilbert space when endowed with the scalar product

$$\langle v, w \rangle_{\pi^{(K)}} := \sum_{n \geq 1} \bar{v}(n) w(n) \pi_n^{(K)}.$$

From [5, Theorem 3.1], we have the following properties. The operator L_K (defined in (1.3)) is closable in $\ell^2(\pi^{(K)})$. The closure (which we denote by the same symbol) is self-adjoint and has a compact resolvent, hence its spectrum is discrete, composed of simple eigenvalues which are negative real numbers, and the corresponding eigenvectors are orthogonal. (More precisely, Theorem 3.1 in [5] states that the maximal eigenvalue is simple, but in the proof we actually show that all eigenvalues are indeed simple.) We normalize these eigenvectors and we can assume that they are real (since they are defined by a second-order real recurrence relation whose solution is determined by choosing the first element). We write

$$L_K \psi_j^{(K)} = -\rho_j^{(K)} \psi_j^{(K)} \quad (1.8)$$

where we order the eigenvalues $-\rho_j^{(K)}$ in decreasing order as j increases. To emphasize that all operators considered in this paper are negative, we have decided to write their eigenvalues under the form $-\rho$, with $\rho > 0$.

As shown in [5], the quasi-stationary distribution exists, is unique, and given by

$$\nu^{(K)}(\{n\}) = \frac{\pi_n^{(K)} \psi_0^{(K)}(n)}{\langle \psi_0^{(K)}, \mathbf{1} \rangle_{\pi^{(K)}}}, \quad n \in \mathbb{Z}_{>0} \quad (1.9)$$

where $\mathbf{1} = (1, 1, \dots)$. Note that it also follows from Theorem 3.2 and Lemma 9.3 in [5] that there exists $D > 1$ such that for all $K \in \mathbb{Z}_{>0}$, we have $D^{-1} \leq \psi_0^{(K)} \leq D$. Therefore, the Hilbert spaces $\ell^2(\nu^{(K)})$ and $\ell^2(\pi^{(K)})$ are isomorphic.

Let $S_t^{(K)} f(n) := \mathbb{E}_n \left(f(X_t^{(K)}) \mathbf{1}_{\{T_0^{(K)} > t\}} \right)$ ($t \geq 0$) be the semigroup of the killed process, where $n \in \mathbb{Z}_{>0}$ and $f \in \ell^\infty$. The following result justifies that we look for the spectrum of L_K in $\ell^2(\pi^{(K)})$.

PROPOSITION 1.1. *The semigroup $(S_t^{(K)})_{t \geq 0}$, defined on c_{00} , extends to a C_0 -contraction semigroup on $\ell^2(\nu^{(K)})$.*

We refer to [18] for definitions and properties of C_0 -contraction semigroups.

PROOF. We follow the argument of Proposition 8.1.8 p. 162 in [3]. Since $\nu^{(K)}$ is a quasi-stationary distribution, for $f \in c_{00}$ and $t \geq 0$, we have

$$\begin{aligned} \int |S_t^{(K)} f|^2 d\nu^{(K)} &\leq \int d\nu^{(K)}(n) \mathbb{E}_n \left(|f(X_t^{(K)})|^2 \mathbf{1}_{\{T_0^{(K)} > t\}} \right) \\ &= e^{-\rho_0^{(K)} t} \int |f|^2 d\nu^{(K)}. \end{aligned}$$

Therefore

$$\|S_t^{(K)} f\|_{\ell^2(\nu^{(K)})} \leq e^{-\frac{\rho_0^{(K)} t}{2}} \|f\|_{\ell^2(\nu^{(K)})}, \quad t \geq 0.$$

Since c_{00} is dense in $\ell^2(\nu^{(K)})$, we get

$$\|S_t^{(K)}\|_{\ell^2(\nu^{(K)})} \leq e^{-\frac{\rho_0^{(K)} t}{2}}, \quad t \geq 0.$$

This implies that $(S_t^{(K)})_{t \geq 0}$ extends to a contraction semigroup in $\ell^2(\nu^{(K)})$. Now, since $\mathbb{P}_n(X_t^{(K)} = m, T_0^{(K)} > t) \rightarrow \delta_{n,m}$, as $t \rightarrow 0$, then for any $f \in c_{00}$, $S_t^{(K)} f \rightarrow f$ pointwise, hence by dominated convergence we obtain $S_t^{(K)} f \rightarrow f$ in $\ell^2(\nu^{(K)})$ as $t \rightarrow 0$. The proposition follows from the contraction property obtained above and the fact that c_{00} is dense in $\ell^2(\nu^{(K)})$. ■

Observe that the same result holds in $\ell^p(\nu^{(K)})$ for any $1 \leq p < +\infty$ (with a similar proof). In the case $p = 2$, see also [7], Theorem 5.2.1 for the C_0 property.

Recall from [5] that we also have

$$\mathbb{P}_{\nu^{(K)}}(T_0^{(K)} > t) = e^{-\rho_0^{(K)} t}, \quad t > 0$$

and the mean-time to extinction starting from $\nu^{(K)}$ is

$$\begin{aligned} \mathbb{E}_{\nu^{(K)}} [T_0^{(K)}] &= \frac{1}{\rho_0^{(K)}} = \frac{\sqrt{2\pi} \exp\left(K \int_0^{x_*} \log \frac{b(x)}{d(x)} dx\right)}{b(x_*) \left(\sqrt{\frac{b(1/K)}{d(1/K)}} - \sqrt{\frac{d(1/K)}{b(1/K)}}\right) \sqrt{KH''(x_*)}} \left(1 + \mathcal{O}\left(\frac{(\log K)^3}{\sqrt{K}}\right)\right). \end{aligned}$$

(See the next section for the definition of H .) In the logistic model this gives (recall that $x_* = \lambda - \mu$)

$$\mathbb{E}_{\nu^{(K)}} [T_0^{(K)}] = \frac{\sqrt{2\pi} \mu \exp\left(K(\lambda - \mu + \mu \log \frac{\mu}{\lambda})\right)}{(\lambda - \mu)^2 \sqrt{K}} \left(1 + \mathcal{O}\left(\frac{(\log K)^3}{\sqrt{K}}\right)\right).$$

Thus $\rho_0^{(K)}$ is exponentially small in K , and we also proved in [5] that the ‘spectral gap’ satisfies (see Theorem 3.3 in [5])

$$\rho_1^{(K)} - \rho_0^{(K)} \geq \frac{\mathcal{O}(1)}{\log K}, \quad K \in \mathbb{Z}_{>1}.$$

This lower bound goes to 0 as $K \rightarrow +\infty$, so it suggests that the spectral gap could vanish in the limit. Actually, this is not the case, as Corollary 1.3 shows. As a matter of fact, we can compute explicitly the spectral gap in the limit.

In fact we will fully describe the asymptotics of all eigenvalues, which is the content of our main theorem. To state it, we need to order $S_1 \cup S_2$ to take care of possible multiplicities. Recall that S_1 and S_2 have been defined in (1.5) and (1.6) and that they are positive sequences. This is done through the definition of a non-decreasing infinite (positive) sequence $(\eta_n)_{n \geq 0}$. Let $\eta_0 = 0$. We construct this sequence recursively as follows.

- ◇ If $\eta_n \in S_1 \Delta S_2$, then $\eta_{n+1} = \min \{\eta : \eta \in S_1 \cup S_2 : \eta > \eta_n\}$.
- ◇ If $\eta_n \in S_1 \cap S_2$, then
 - If $\eta_{n-1} = \eta_n$, then $\eta_{n+1} = \min \{\eta : \eta \in S_1 \cup S_2 : \eta > \eta_n\}$.
 - If $\eta_{n-1} < \eta_n$, then $\eta_{n+1} = \eta_n$.

The main result of this paper, whose assumptions will be stated in Section 2, is the following.

THEOREM 1.2 (Convergence of the spectrum).

The spectrum of L_K in $\ell^2(\pi^{(K)})$ converges pointwise to $(-\eta_n)_{n \geq 0}$ when K tends to infinity. In other words

$$\lim_{K \rightarrow +\infty} \rho_j^{(K)} = \eta_j, \quad \forall j \in \mathbb{Z}_{\geq 0}.$$

COROLLARY 1.3. *When $K \rightarrow +\infty$, the spectral gap $\rho_1^{(K)} - \rho_0^{(K)}$ converges to*

$$\min \{b'(0) - d'(0), d'(x_*) - b'(x_*)\}.$$

EXAMPLES. *In the logistic model, we have $d'(0) - b'(0) = b'(x_*) - d'(x_*) = \mu - \lambda$ (asymptotic double eigenvalues), and the spectral gap is equal to $\lambda - \mu$.*

Another example is the Ayala-Gilpin-Ehrenfeld model [1] defined by $b(x) = \lambda x$, $d(x) = x(\mu + x^\theta)$ where $\theta \in (0, 1)$ is a parameter, and $\lambda > \mu$. In this case, $d'(0) - b'(0) = \mu - \lambda$, $x_ = (\lambda - \mu)^{\frac{1}{\theta}}$, and $b'(x_*) - d'(x_*) = \theta(\mu - \lambda)$, so the spectral gap is $\theta(\lambda - \mu)$.*

Yet another example is Smith's model [17] defined by $b(x) = \lambda x/(1+x)$, $d(x) = (x(\mu+x))/(1+x)$, where $\lambda > \mu$. One easily finds $d'(0) - b'(0) = \mu - \lambda$, $x_ = \lambda - \mu$, and $b'(x_*) - d'(x_*) = (\mu - \lambda)/(1 + \lambda - \mu)$, so the spectral gap is $(\lambda - \mu)/(1 + \lambda - \mu)$. A last example is Nisbet-Gurney model [12] defined by $b(x) = \lambda x e^{1-x/c}$, $d(x) = \mu x$, where $\lambda, \mu, c > 0$ and $\mu < e\lambda$. We have $d'(0) - b'(0) = \mu - e\lambda$, $x_* = c(1 - \log(\mu/\lambda))$, and $b'(x_*) - d'(x_*) = \mu(\log(\mu/\lambda) - 1)$. Hence the spectral gap is equal to $\mu(1 - \log(\mu/\lambda))$.*

1.5 Consequences on relaxation times

Recall that the spectral gap is the inverse of the relaxation time to the quasi-stationary distribution, namely for $\psi \in \ell^2(\pi^{(K)})$, $t > 0$ and $K \in \mathbb{Z}_{>0}$ we have

$$\left\| e^{\rho_0^{(K)} t} S_t^{(K)} \psi - \psi_0^{(K)} \langle \psi_0^{(K)}, \mathbf{1} \rangle_{\pi^{(K)}} \nu^{(K)}(\psi) \right\|_{\ell^2(\pi^{(K)})} \leq \|\psi\|_{\ell^2(\pi^{(K)})} e^{-(\rho_1^{(K)} - \rho_0^{(K)})t}.$$

From Corollary 1.3, it turns out that the relaxation time converges to a finite limit as K tends to infinity.

We can also characterize the decay of correlations for the so-called Q -process, namely the birth-and-death process conditioned on survival. Recall that the Q -process is the irreducible Markov process with state space $\mathbb{Z}_{>0}$, defined by the semigroup

$$R_t^{(K)} g = e^{\rho_0^{(K)} t} \frac{1}{\psi_0^{(K)}} S_t^{(K)} (g \psi_0^{(K)}).$$

It satisfies $R_t^{(K)} \mathbf{1}_{\{n \geq 1\}} = \mathbf{1}_{\{n \geq 1\}}$. Denoting by $(R_t^{(K)})^\dagger$ the adjoint operator to $R_t^{(K)}$, the unique invariant distribution $\mathfrak{m}^{(K)}$ of the Q -process, defined by $(R_t^{(K)})^\dagger \mathfrak{m}^{(K)} = \mathfrak{m}^{(K)}$, is related to $\nu^{(K)}$ by $\mathfrak{m}^{(K)}(g) = \nu^{(K)}(\psi_0^{(K)} g)$ where $\nu^{(K)}$ has been defined in (1.9). Indeed we have $g \in \ell^2(\mathfrak{m}^{(K)})$ if and only if $\psi_0^{(K)} g \in \ell^2(\pi^{(K)})$, and

$$\|g\|_{\ell^2(\mathfrak{m}^{(K)})}^2 = \frac{\|\psi_0^{(K)} g\|_{\ell^2(\pi^{(K)})}^2}{\langle \psi_0^{(K)}, \mathbf{1} \rangle_{\pi^{(K)}}}.$$

Hence, we get the following result.

PROPOSITION 1.4. *Let $g \in \ell^2(\mathfrak{m}^{(K)})$ and $K \in \mathbb{Z}_{>0}$. Then for all $t > 0$*

$$\|R_t^{(K)} g - \langle \psi_0^{(K)}, \mathbf{1} \rangle_{\pi^{(K)}} \mathfrak{m}^{(K)}(g)\|_{\ell^2(\mathfrak{m}^{(K)})} \leq \|g\|_{\ell^2(\mathfrak{m}^{(K)})} e^{-(\rho_1^{(K)} - \rho_0^{(K)})t}.$$

Furthermore, for $g_1, g_2 \in \ell^2(\mathfrak{m}^{(K)})$ and for all $t > 0$, we have

$$\begin{aligned} & \left| \int R_t^{(K)} g_1 \cdot g_2 \, d\mathfrak{m}^{(K)} - \int g_1 \, d\mathfrak{m}^{(K)} \int g_2 \, d\mathfrak{m}^{(K)} \right| \\ & \leq \|g_1\|_{\ell^2(\mathfrak{m}^{(K)})} \|g_2\|_{\ell^2(\mathfrak{m}^{(K)})} e^{-(\rho_1^{(K)} - \rho_0^{(K)})t}. \end{aligned}$$

As before, it follows from Corollary 1.3 that the rate of decay of correlations converges when K goes to infinity.

1.6 Organization of the paper

The proof of Theorem 1.2 relies on two results stated in Section 3. Theorem 3.5 ensures that the set $S_1 \cup S_2$ is contained in the set of accumulation points of the eigenvalues of L_K when K tends to infinity. More precisely, we will not work with L_K in $\ell^2(\pi^{(K)})$, but with the conjugated operator \mathcal{L}_K on the ‘flat’ Hilbert space ℓ^2 . The proof is based on the construction of quasi-eigenvectors and is given in Section 5. The second result is Theorem 3.7 which ensures that all the previous accumulation points are contained in $S_1 \cup S_2$ taking care of eventual multiplicities. Its proof, given in section 5, relies on two propositions. The first one (Proposition 4.1) is the splitting of the eigenvectors of \mathcal{L}_K into two dominant parts, one localized near the origin, the other one near $\lfloor Kx_* \rfloor$. The second (Proposition 4.5) relies on compactness arguments of each piece of the previous splitting. Section 6 collects various auxiliary results (some of more general nature).

Let us emphasize that one of the main difficulties of the proof is that the two pieces of the spectrum correspond to limiting operators which are obtained at different scales and live in different function spaces.

2 Standing assumptions

We work under the assumptions of [5] which we recall for convenience. Part of them were already stated in the introduction.

The functions $b, d : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ defining the differential equation (1.2) are supposed to be such that

$$b(0) = d(0) = 0$$

and the functions $x \mapsto b(x)/x$ and $x \mapsto d(x)/x$ are defined on \mathbb{R}_+ and assumed to be positive, twice differentiable and increasing (in particular the sequences $(\lambda_n^{(K)})_{n \geq 1}$ and $(\mu_n^{(K)})_{n \geq 1}$ defined in (1.1) are increasing for each K).

We start by the biologically relevant assumptions:

- $d'(0) > 0$ and $b'(0) > d'(0)$ (i.e., births prevail at low density).
- There is a unique $x_* > 0$ such that $b(x_*) = d(x_*)$, so x_* is the only positive fixed point of (1.2).
- $\lim_{x \rightarrow +\infty} \frac{b(x)}{d(x)} = 0$ (i.e., deaths prevail over births at large densities).

We assume that $b'(x_*) \neq d'(x_*)$ (genericity condition). The remaining (technical) assumptions are the following:

- $\int_{\frac{x_*}{2}}^{+\infty} \frac{dx}{d(x)} < +\infty$ and $\sup_{x>0} \left(\frac{d'(x)}{d(x)} - \frac{1}{x} \right) < +\infty$.
- The function $x \mapsto \log \frac{d(x)}{b(x)}$ is increasing on \mathbb{R}_+ .
- The function $H : \mathbb{R}_+ \rightarrow \mathbb{R}$ defined by $H(x) = \int_{x_*}^x \log \frac{d(s)}{b(s)} ds$ is three times differentiable, and $\sup_{x \in \mathbb{R}_+} (1 + x^2) |H'''(x)| < +\infty$.

The assumptions imply that 0 is a repulsive (or unstable) fixed point of (1.2), whereas x_* is an attractive (or stable) one, that is, $b'(x_*) < d'(x_*)$. It also follows that $H''(x_*) > 0$. These assumptions are satisfied for many, if not all, classical examples. We gave four examples above (see end of Subsection 1.4).

As explained in [5], the above conditions imply (for each $K \in \mathbb{Z}_{>0}$) the following properties (where $\pi_n^{(K)}$ is defined in (1.7)):

- $\sum_{n \geq 1} (\lambda_n^{(K)} \pi_n^{(K)})^{-1} = +\infty$, which is a necessary and sufficient condition for the process to reach 0 in finite time with probability one.
- $\sum_{n \geq 1} \pi_n^{(K)} < +\infty$, which implies finiteness of the mean time to extinction.
- $\sum_{n \geq 1} (\lambda_n^{(K)} \pi_n^{(K)})^{-1} \sum_{i \geq n+1} \pi_i^{(K)} < +\infty$, which is a necessary and sufficient condition for existence and uniqueness of the quasi-stationary distribution.

We add a last condition to the previous ones, namely

$$\lim_{x \rightarrow +\infty} \frac{\log b(x)}{x} = \lim_{x \rightarrow +\infty} \frac{\log d(x)}{x} = 0. \quad (2.1)$$

We could avoid it, but it makes our life easier and we don't have any natural example which does not satisfy it. As the above examples show (see right after Corollary 1.3), b and d are usually polynomial or rational functions, so (2.1) is amply satisfied.

3 Proof of Theorem 1.2

3.1 Some useful operators

Instead of working with the operator L_K (defined in (1.3)) on the weighted Hilbert space $\ell^2(\pi^{(K)})$, we find more convenient to work on the ‘flat’ Hilbert space ℓ^2 . We thus introduce the conjugated operator

$$\mathcal{L}_K = (\Pi^{(K)})^{\frac{1}{2}} L_K (\Pi^{(K)})^{-\frac{1}{2}}$$

where $\Pi^{(K)}$ denotes the multiplication operator

$$\Pi^{(K)}v(n) = \pi_n^{(K)}v(n)$$

for $v \in c_{00}$ and $n \in \mathbb{Z}_{>0}$. One can check that

$$\begin{aligned} (\mathcal{L}_K v)(n) &= \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} v(n+1) + \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}} v(n-1) \mathbb{1}_{\{n>1\}} - (\lambda_n^{(K)} + \mu_n^{(K)}) v(n) \end{aligned}$$

for $n \in \mathbb{Z}_{>0}$. We denote also by \mathcal{L}_K its closure in ℓ^2 and by $\text{Dom}(\mathcal{L}_K)$ its domain, and we have

$$\mathcal{L}_K \phi_j^{(K)} = -\rho_j^{(K)} \phi_j^{(K)} \quad (3.1)$$

where the eigenvalues $-\rho_j^{(K)}$ are that of L_K (see (1.8)), and $\phi^{(K)} = (\Pi^{(K)})^{\frac{1}{2}} \psi^{(K)}$. To capture the behavior of the eigenvectors of \mathcal{L}_K near $\lfloor Kx_* \rfloor$ at scale \sqrt{K} , we are going to embed ℓ^2 into L^2 . For this purpose we define for each $K \in \mathbb{Z}_{>0}$ the functions

$$e_n^{(K)}(x) = K^{\frac{1}{4}} \mathbb{1}_{I_n^{(K)}}(x), \quad x \in \mathbb{R}, \quad n \in \mathbb{Z}_{>0}$$

where

$$I_n^{(K)} = \left[\frac{n-0.5}{\sqrt{K}} - x_*\sqrt{K}, \frac{n+0.5}{\sqrt{K}} - x_*\sqrt{K} \right].$$

The functions $e_n^{(K)}$ are orthogonal and of norm one in L^2 . They form a basis of a sub-Hilbert space \mathcal{H}_K of piecewise constant functions in L^2 . We define two maps denoted by Q_K and P_K as follows:

$$Q_K : \ell^2 \rightarrow L^2, \quad Q_K u(x) = \sum_{n \geq 1} u(n) e_n^{(K)}(x)$$

and

$$P_K : L^2 \rightarrow \ell^2, \quad P_K f(n) = \int f(x) e_n^{(K)}(x) dx, \quad n \in \mathbb{Z}_{>0}. \quad (3.2)$$

We will use the following properties of P_K and Q_K stated as two lemmas.

LEMMA 3.1. *The two maps P_K and Q_K are continuous. For each $K \in \mathbb{Z}_{>0}$, the map Q_K is a surjective isometry between ℓ^2 and \mathcal{H}_K . The map P_K is the adjoint of Q_K . Moreover*

$$P_K Q_K = \text{Id}_{\ell^2} \quad \text{and} \quad Q_K P_K = \mathcal{P}_K$$

where \mathcal{P}_K is the orthogonal projection on \mathcal{H}_K in L^2 .

The only non trivial point is the continuity of P_K . It follows from the identity

$$\int f^2(x) dx = \sum_{n \geq 1} \int_{I_n^{(K)}} \left(f(x) - \frac{1}{|I_n^{(K)}|} \int_{I_n^{(K)}} f(y) dy \right)^2 dx + \sum_{n \geq 1} \frac{1}{|I_n^{(K)}|^2} \left(\int_{I_n^{(K)}} f(y) dy \right)^2.$$

The proof of the other statements is left to the reader.

LEMMA 3.2. *Let $f \in C^1(\mathbb{R})$ and assume that there exists $a > 0$ and $A > 0$ such that*

$$|f(x)| + |f'(x)| \leq A e^{-a|x|}, \quad x \in \mathbb{R}.$$

Then

- (i) $\lim_{K \rightarrow +\infty} \|f - Q_K P_K f\|_{L^2} = 0.$
- (ii) $\lim_{K \rightarrow +\infty} \|P_K f\|_{\ell^2} = \|f\|_{L^2}.$

PROOF. Let us prove (i). We have $Q_K P_K f(x) = e_{n(x)}^{(K)}(x) \int e_{n(x)}^{(K)}(y) f(y) dy$ for $(n(x) - 0.5)/\sqrt{K} - x_* \leq x \leq (n(x) + 0.5)/\sqrt{K} - x_*$. We get from our hypothesis

$$Q_K P_K f(x) = f(x) + \mathcal{O}(K^{-\frac{1}{2}}) e^{-a|x|}$$

and the result follows.

We now prove (ii). From the isometric property of Q_K on the space of piecewise functions \mathcal{H}_K , we get

$$\|P_K f\|_{\ell^2} = \|Q_K P_K f\|_{L^2}$$

and the result follows from (i). ■

We now introduce the operator

$$\mathcal{L}_K = Q_K \mathcal{L}_K P_K$$

with domain $Q_K \text{Dom}(\mathcal{L}_K) \oplus \mathcal{H}_K^\perp$. It is left to the reader to prove that this operator is closed (using the relations of Lemma 3.1). Since \mathcal{L}_K when acting on \mathcal{H}_K is conjugated to \mathcal{L}_K , we have

$$\mathcal{L}_K \varphi_j^{(K)} = -\rho_j^{(K)} \varphi_j^{(K)}$$

where $\varphi_j^{(K)} = Q_K \phi_j^{(K)}$, and the $\phi_j^{(K)}$'s are defined in (3.1). We will prove in the next proposition that the operator \mathcal{L}_K converges pointwise in L^2 on a subset of functions, when $K \rightarrow +\infty$, to the operator

$$\begin{aligned} \mathcal{H}_* f(x) = & \tag{3.3} \\ & b(x_*) \frac{d^2 f(x)}{dx^2} - \frac{(d'(x_*) - b'(x_*))^2}{4b(x_*)} x^2 f(x) + \frac{d'(x_*) - b'(x_*)}{2} f(x). \end{aligned}$$

PROPOSITION 3.3. *Let $f \in C^3(\mathbb{R})$ and assume that there exist $a > 0$ and $A > 0$ such that*

$$\sum_{j=0}^3 |f^{(j)}(x)| \leq A e^{-a|x|}, \quad x \in \mathbb{R}.$$

Then

$$\lim_{K \rightarrow +\infty} \|\mathcal{L}_K f - \mathcal{H}_* f\|_{L^2} = 0.$$

PROOF. By the assumption made on f , it follows easily that

$$\lim_{K \rightarrow +\infty} \|\mathbb{1}_{\{|\cdot| > (\log K)^2\}} \mathcal{H}_* f\|_{L^2} = 0.$$

We have

$$\begin{aligned} \mathcal{L}_K f(x) = & \sum_{n \geq 1} e_n^{(K)}(x) \left(\sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} \int e_{n+1}^{(K)}(y) f(y) dy \tag{3.4} \right. \\ & \left. + \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}} \mathbb{1}_{\{n > 1\}} \int e_{n-1}^{(K)}(y) f(y) dy - (\lambda_n^{(K)} + \mu_n^{(K)}) \int e_n^{(K)}(y) f(y) dy \right). \end{aligned}$$

It follows easily from the assumption made on f and assumption (2.1) that

$$\lim_{K \rightarrow +\infty} \|\mathbb{1}_{\{|\cdot| > (\log K)^2\}} \mathcal{L}_K f\|_{L^2} = 0.$$

Therefore we only have to consider $|x| \leq (\log K)^2$. Note also that for such an x , the sum in (3.4) reduces to one element for K large enough, namely $n = n(x) = \lfloor Kx_* + \sqrt{K}x + 0.5 \rfloor$. For $x \in \mathbb{R}$, we have

$$\begin{aligned} \mathcal{L}_K f(x) = & e_{n(x)}^{(K)}(x) \left(\sqrt{\lambda_{n(x)}^{(K)} \mu_{n(x)+1}^{(K)}} \int e_{n(x)+1}^{(K)}(y) f(y) dy \right. \\ & + \sqrt{\lambda_{n(x)-1}^{(K)} \mu_{n(x)}^{(K)}} \int e_{n(x)-1}^{(K)}(y) f(y) dy \\ & \left. - (\lambda_{n(x)}^{(K)} + \mu_{n(x)}^{(K)}) \int e_{n(x)}^{(K)}(y) f(y) dy \right) \\ = & e_{n(x)}^{(K)}(x) \left(\sqrt{\lambda_{n(x)}^{(K)} \mu_{n(x)+1}^{(K)}} \int e_{n(x)}^{(K)}(y) f\left(y - \frac{1}{\sqrt{K}}\right) dy \right. \\ & \left. + \sqrt{\lambda_{n(x)-1}^{(K)} \mu_{n(x)}^{(K)}} \int e_{n(x)}^{(K)}(y) f\left(y + \frac{1}{\sqrt{K}}\right) dy \right) \end{aligned}$$

$$-(\lambda_{n(x)}^{(K)} + \mu_{n(x)}^{(K)}) \int e_{n(x)}^{(K)}(y) f(y) dy \Big).$$

Now we have

$$\begin{aligned} & \int e_{n(x)}^{(K)}(y) f(y) dy \\ &= K^{\frac{1}{4}} \int_{-\frac{1}{2\sqrt{K}}}^{\frac{1}{2\sqrt{K}}} f\left(\frac{n(x) - Kx_*}{\sqrt{K}} - s\right) ds \\ &= K^{-\frac{1}{4}} f\left(\frac{n(x) - Kx_*}{\sqrt{K}}\right) + \frac{K^{-\frac{5}{4}}}{24} f''\left(\frac{n(x) - Kx_*}{\sqrt{K}}\right) + \mathcal{O}\left(K^{-\frac{7}{4}}\right) \end{aligned}$$

where the error term is uniform in x . Similarly

$$\begin{aligned} & \int e_{n(x)}^{(K)}(y) f\left(y \pm \frac{1}{\sqrt{K}}\right) dy \\ &= \int e_{n(x)}^{(K)}(y) f(y) dy \pm \frac{1}{\sqrt{K}} \int e_{n(x)}^{(K)}(y) f'(y) dy \\ &\quad + \frac{1}{2K} \int e_{n(x)}^{(K)}(y) f''(y) dy + \mathcal{O}\left(K^{-\frac{7}{4}}\right) \\ &= K^{-\frac{1}{4}} f\left(\frac{n(x) - Kx_*}{\sqrt{K}}\right) \pm K^{-\frac{3}{4}} f'\left(\frac{n(x) - Kx_*}{\sqrt{K}}\right) \\ &\quad + \frac{13 K^{-\frac{5}{4}}}{24} f''\left(\frac{n(x) - Kx_*}{\sqrt{K}}\right) + \mathcal{O}\left(K^{-\frac{7}{4}}\right). \end{aligned}$$

Recall that

$$\lambda_n^{(K)} = Kb\left(\frac{n}{K}\right) \quad \text{and} \quad \mu_n^{(K)} = Kd\left(\frac{n}{K}\right)$$

hence

$$\lambda_n^{(K)} = Kb(x_*) + (n - Kx_*) b'(x_*) + \mathcal{O}\left(\frac{(n - Kx_*)^2}{K}\right)$$

and

$$\mu_n^{(K)} = Kd(x_*) + (n - Kx_*) d'(x_*) + \mathcal{O}\left(\frac{(n - Kx_*)^2}{K}\right).$$

After a tedious but straightforward computation, we obtain that

$$\begin{aligned} \mathcal{L}_K f(x) &= \mathcal{H}_* f\left(\frac{n(x) - Kx_*}{\sqrt{K}}\right) + \mathcal{O}\left(K^{-\frac{1}{4}}\right) \\ &= \mathcal{H}_* f(x) + \mathcal{O}\left(K^{-\frac{1}{4}}(\log K)^4\right) \end{aligned}$$

and the error term is uniform in $|x| \leq (\log K)^2$. We get

$$\|\mathbb{1}_{\{|\cdot| \leq (\log K)^2\}} (\mathcal{L}_K f - \mathcal{H}_* f)\|_{L^2} = \mathcal{O}\left(K^{-\frac{1}{4}}(\log K)^6\right)$$

and the result follows. \blacksquare

REMARK 3.1. *Let us recall the relationship between the generator of the Ornstein-Uhlenbeck process (1.4) and the operator (3.3) which is, up to a minus sign and a shift, the Schrödinger operator for the quantum harmonic oscillator. We refer to e.g. [2, Chapter 3] or [13, Sections 4.4 and 4.9]. In L^2 , the eigenvalues of \mathcal{H}_* are $-(d'(x_*) - b'(x_*))n$, $n \in \mathbb{Z}_{\geq 0}$, and the corresponding eigenfunctions are*

$$\psi_n(x) = \frac{1}{\sqrt{2^n n!}} \left(\frac{(d'(x_*) - b'(x_*))^2}{2\pi b(x_*)} \right)^{\frac{1}{4}} e^{-\frac{d'(x_*) - b'(x_*)}{4b(x_*)} x^2} H_n \left(\sqrt{\frac{d'(x_*) - b'(x_*)}{2b(x_*)}} x \right)$$

where $(H_n)_n$ is the family of the physicists' Hermite polynomials defined by

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2}.$$

One can check that \mathcal{H}_* is conjugated to the generator of the Ornstein-Uhlenbeck process (1.4) acting on $L^2 \left(\sqrt{\frac{d'(x_*) - b'(x_*)}{2\pi b(x_*)}} e^{-\frac{d'(x_*) - b'(x_*)}{2b(x_*)} x^2} dx \right)$ in the following way: $\psi_0^{-1} \mathcal{H}_* (\psi_0 f) = \mathcal{O} \mathcal{U}_* f$.

In Proposition 3.4 we prove that the operator \mathcal{L}_K converges pointwise in ℓ^2 on c_{00} , when K tends to infinity, to the operator \mathcal{M}_0 defined for $v \in c_{00}$ by

$$\begin{aligned} (\mathcal{M}_0 v)(n) = & \quad (3.5) \\ & \sqrt{b'(0) d'(0) n(n+1)} v(n+1) + \sqrt{b'(0) d'(0) n(n-1)} v(n-1) \mathbb{1}_{\{n>1\}} \\ & - n(b'(0) + d'(0))v(n). \end{aligned}$$

Here again we denote the operator on c_{00} and its closure by the same letter.

PROPOSITION 3.4. *Let $u \in c_{00}$. Then*

$$\lim_{K \rightarrow +\infty} \mathcal{L}_K u = \mathcal{M}_0 u$$

where \mathcal{M}_0 is defined in (3.5).

PROOF. Follows from the fact that for each fixed n

$$\lim_{K \rightarrow +\infty} \lambda_n^{(K)} = b'(0) n \quad \text{and} \quad \lim_{K \rightarrow +\infty} \mu_n^{(K)} = d'(0) n.$$

■

3.2 Steps of the proof of Theorem 1.2

The proof of Theorem 1.2 relies on the following two theorems whose proofs are postponed to Section 5. Recall that for any fixed K , the spectrum $\text{Sp}(\mathcal{L}_K)$ is discrete, and let

$$G = \bigcup_{j=0}^{+\infty} (\rho_j^{(K)})^{\text{acc}}$$

where $(\rho_j^{(K)})^{\text{acc}}$ is the set of accumulation points of $(\rho_j^{(K)})$ when $K \rightarrow +\infty$.

THEOREM 3.5. *We have*

$$S_1 \cup S_2 \subset G$$

where S_1 and S_2 are defined in (1.5) and (1.6).

This theorem is proved in Section 5.1.

COROLLARY 3.6. *For every fixed j we have*

$$\limsup_{K \rightarrow +\infty} \rho_j^{(K)} < +\infty.$$

PROOF. We proceed by contradiction. Assume that there exists j_0 such that

$$\limsup_{K \rightarrow +\infty} \rho_{j_0}^{(K)} = +\infty.$$

Let $j_c = \min\{0 < \ell \leq j_0 : \limsup_{K \rightarrow +\infty} \rho_\ell^{(K)} = +\infty\}$. Hence there exists $\alpha < +\infty$ such that $\limsup_{K \rightarrow +\infty} \rho_{j_c-1}^{(K)} = \alpha$. By definition of j_c , there exists a diverging sequence $(K_p)_{p \geq 1}$ such that $\lim_{p \rightarrow +\infty} \rho_{j_c}^{(K_p)} = +\infty$. Let $\rho \in S_1 \cup S_2$ such that $\rho > \alpha$.

If $j \leq j_c - 1$, we have $\limsup_{p \rightarrow +\infty} \rho_j^{(K_p)} \leq \limsup_{p \rightarrow +\infty} \rho_{j_c-1}^{(K_p)} \leq \alpha < \rho$. For all $j \geq j_c$, we have $\liminf_{p \rightarrow +\infty} \rho_j^{(K_p)} \geq \liminf_{p \rightarrow +\infty} \rho_{j_c}^{(K_p)} = +\infty$. This implies $\rho \notin G$, contradicting Theorem 3.5. ■

THEOREM 3.7. *We have*

$$S_1 \cup S_2 \supset G.$$

Moreover, for each $j \in \mathbb{Z}_{\geq 0}$, let $(K_p)_{p \geq 1}$ be a diverging sequence such that

$$\lim_{p \rightarrow +\infty} \rho_j^{(K_p)} = \rho_*$$

where ρ_* is finite by Corollary 3.6. Then

1) If $\rho_* \in S_1 \Delta S_2$ then

$$\liminf_{p \rightarrow +\infty} \min \left\{ \left| \rho_{j+1}^{(K_p)} - \rho_* \right|, \left| \rho_{j-1}^{(K_p)} - \rho_* \right| \right\} > 0. \quad (3.6)$$

Moreover there are only two cases:

- (a) If $\rho_* \in S_1$ then there exists a diverging sequence of integers $(p_\ell)_{\ell \geq 1}$ such that $Q_{K_{p_\ell}} \phi_j^{(K_{p_\ell})} \xrightarrow{L^2} \varphi_*$, where ρ_* and φ_* are such that $\mathcal{H}_* \varphi_* = -\rho_* \varphi_*$.
- (b) If $\rho_* \in S_2$ then $\phi_j^{(K_p)} \xrightarrow{\ell^2} \phi_*$, where ρ_* and ϕ_* are such that $\mathcal{M}_0 \phi_* = -\rho_* \phi_*$.

2) If $\rho_* \in S_1 \cap S_2$ then we have the following two assertions:

- (a) There exists a diverging sequence of integers $(p_\ell)_{\ell \geq 1}$ such that:

$$\text{either } \lim_{\ell \rightarrow +\infty} \rho_{j+1}^{(K_{p_\ell})} = \rho_* \quad \text{or} \quad \lim_{\ell \rightarrow +\infty} \rho_{j-1}^{(K_{p_\ell})} = \rho_*.$$

- (b) We have

$$\liminf_{p \rightarrow +\infty} \min \left\{ \left| \rho_{j+1}^{(K_p)} - \rho_* \right|, \left| \rho_{j-1}^{(K_p)} - \rho_* \right| \right\} = 0 \quad (3.7)$$

and

$$\liminf_{p \rightarrow +\infty} \max \left\{ \left| \rho_{j+1}^{(K_p)} - \rho_* \right|, \left| \rho_{j-1}^{(K_p)} - \rho_* \right| \right\} > 0. \quad (3.8)$$

Note that (3.6) means that if $\rho_* \in S_1 \Delta S_2$ then $-\rho_*$ is a simple asymptotic eigenvalue, and either $-\rho_*$ is an eigenvalue of \mathcal{H}_* if $\rho_* \in S_1$, or of \mathcal{M}_0 if $\rho_* \in S_2$. In addition, (3.7) and (3.8) mean that if $\rho_* \in S_1 \cap S_2$, then $-\rho_*$ is a double asymptotic eigenvalue which is an eigenvalue of both \mathcal{H}_* and \mathcal{M}_0 .

PROOF OF THEOREM 1.2. The proof is recursive. For $j = 0$ it follows from [5] that $\lim_{K \rightarrow +\infty} \rho_0^{(K)} = 0$. Let $j \geq 0$ and assume that for $\ell \leq j$ (if any) $\lim_{K \rightarrow +\infty} \rho_\ell^{(K)} = \eta_\ell$. We now prove that $\lim_{K \rightarrow +\infty} \rho_{j+1}^{(K)} = \eta_{j+1}$. There are several cases to consider.

- If $\eta_j \in S_1 \Delta S_2$, we claim that $\liminf_{K \rightarrow +\infty} \rho_{j+1}^{(K)} \geq \eta_{j+1}$. Otherwise, by Theorem 3.7 and the recursive hypothesis, there would exist $K_p \rightarrow +\infty$ such that $\rho_{j+1}^{(K_p)} \rightarrow \eta_* < \eta_{j+1}$. Since by the recursive hypothesis $\lim_{K \rightarrow +\infty} \rho_j^{(K_p)} = \eta_j$, we have $\eta_* \geq \eta_j$. From the first statement of Theorem 3.7 it follows that $\eta_* = \eta_j$. This contradicts 1) of Theorem 3.7. We now claim that $\limsup_{K \rightarrow +\infty} \rho_{j+1}^{(K)} \leq \eta_{j+1}$. Otherwise, by Theorem 3.7 and the recursive hypothesis, there would exist $K_p \rightarrow +\infty$ such that $\rho_{j+1}^{(K_p)} \rightarrow \eta_* > \eta_{j+1}$. This implies that $\eta_{j+1} \notin G$, contradicting Theorem 3.5.

Hence, in this case, $\lim_{K \rightarrow +\infty} \rho_{j+1}^{(K)} = \eta_{j+1}$.

- If $\eta_j \in S_1 \cap S_2$ (which implies $j > 0$), we have two cases:

- If $\eta_{j-1} = \eta_j$, then we claim that $\liminf_{K \rightarrow +\infty} \rho_{j+1}^{(K)} \geq \eta_{j+1}$. Otherwise, by the same argument as before, there would exist $K_p \rightarrow +\infty$ such that $\rho_{j+1}^{(K_p)} \rightarrow \eta_j$, contradicting (3.8) in Theorem 3.7.

We now claim that $\limsup_{K \rightarrow +\infty} \rho_{j+1}^{(K)} \leq \eta_{j+1}$. Otherwise, as before, this would contradict that $\eta_{j+1} \in G$.

Hence, in this case, $\lim_{K \rightarrow +\infty} \rho_{j+1}^{(K)} = \eta_{j+1}$.

- If $\eta_{j-1} < \eta_j$, then we obviously have $\liminf_{K \rightarrow +\infty} \rho_{j+1}^{(K)} \geq \eta_j$. If $\limsup_{K \rightarrow +\infty} \rho_{j+1}^{(K)} > \eta_j$, then there exists $K_p \rightarrow +\infty$ such that $\rho_{j+1}^{(K_p)} \rightarrow \eta_* > \eta_j$, contradicting (3.7) in Theorem 3.7. Hence, in this case, $\lim_{K \rightarrow +\infty} \rho_{j+1}^{(K)} = \eta_{j+1}$.

Therefore we $\lim_{K \rightarrow +\infty} \rho_{j+1}^{(K)} = \eta_{j+1}$. As announced, the proof of Theorem 1.2 follows recursively.

4 Properties of the eigenvectors

In this section, we prove that, for K large enough, the eigenvectors $\phi_j^{(K)}$ of \mathcal{L}_K (see (3.1)) are functions whose representation is sketched in the figure. An eigenvector is ‘negligible’ outside the union of a neighborhood of 1, and a neighborhood of Kx_* . It is ‘non-negligible’ in at least one of these neighborhoods.




Figure: Schematic representation of one of the three possible ‘shapes’ of the eigenvectors $\phi_j^{(K)}$ (scales are not respected).

To separate the different behaviors, we introduce a ‘potential’ defined by

$$V_n(K) = \lambda_n^{(K)} + \mu_n^{(K)} - \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} - \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}} \mathbb{1}_{\{n>1\}}. \quad (4.1)$$

For $\eta > 0$, let $n_g(K, \eta)$ and $n_d(K, \eta)$ be integers such that $\llbracket n_g(K, \eta), n_d(K, \eta) \rrbracket$ is the maximal interval containing $Kx_*/2$ such that

$$\inf_{n \in \llbracket n_g(K, \eta), n_d(K, \eta) \rrbracket} (V_n(K) - \eta) > 0.$$

Let $n_\ell(K) = \lfloor (\log K)^2 \rfloor$ and $n_r(K) = \lfloor Kx_* - K^{\frac{2}{3}} \log K \rfloor$. It follows from our assumptions that for K large enough

$$1 < n_g(K, \eta) < n_\ell(K) \ll \frac{Kx_*}{2} < n_r(K) < n_d(K, \eta).$$

PROPOSITION 4.1. *For any $\eta > 0$, there exists $a_\eta > 0$, and $K_\eta > 0$ such that, if $K > K_\eta$, and if ϕ is of norm one in ℓ^2 and satisfies*

$$\mathcal{L}_K \phi = -\rho \phi \quad \text{where } \rho < \eta$$

then

$$\sup_{n_\ell(K) \leq n \leq n_r(K)} |\phi(n)| \leq e^{-a_\eta (\log K)^2}.$$

PROOF. Let us consider an eigenvector ϕ of norm one in ℓ^2 satisfying $\mathcal{L}_K \phi = -\rho \phi$.

If $\phi(n_g(K, \eta)) \neq 0$, define $\tilde{n}_g(K, \eta) = n_g(K, \eta)$ and if needed, change the sign of ϕ such that $\phi(\tilde{n}_g(K, \eta)) > 0$. If $\phi(n_g(K, \eta)) = 0$, define $\tilde{n}_g(K, \eta) = n_g(K, \eta) + 1$ and if needed, change the sign of ϕ such that $\phi(\tilde{n}_g(K, \eta)) > 0$ (note that $\phi(n_g(K, \eta)) = \phi(n_g(K, \eta) + 1) = 0$ contradicts the normalization since ϕ solves a second-order recurrence relation). Then, changing the definition of $n_g(K, \eta)$ if necessary, we can assume that $\phi(n_g(K, \eta)) > 0$.

Thanks to the local maximum-minimum principle (see Proposition A.2) we only have four cases.

- 1) $\phi(n_g(K, \eta) + 1) \geq \phi(n_g(K, \eta))$ and $\phi(n)$ is increasing on $\llbracket n_g(K, \eta), n_d(K, \eta) \rrbracket$.
- 2) $\phi(n_g(K, \eta) + 1) < \phi(n_g(K, \eta))$ and $\phi(n)$ is decreasing and stays nonnegative on $\llbracket n_g(K, \eta), n_d(K, \eta) \rrbracket$.
- 3) $\phi(n_g(K, \eta) + 1) < \phi(n_g(K, \eta))$ and $\phi(n)$ has a minimum in the interval $n_{\min}(K)$ in $\llbracket n_g(K, \eta) - 1, n_d(K, \eta) - 1 \rrbracket$ and $\phi(n_{\min}(K)) \geq 0$. Note that $\phi(n)$ is decreasing on $\llbracket n_g(K, \eta), n_{\min}(K) \rrbracket$ and increasing on $\llbracket n_{\min}(K), n_d(K, \eta) \rrbracket$.
- 4) $\phi(n)$ is decreasing on $\llbracket n_g(K, \eta), n_d(K, \eta) \rrbracket$ and $\phi(n_d(K, \eta)) < 0$.

We first observe that since $b'(0) > d'(0)$ we have

$$\lim_{K \rightarrow +\infty} \sup_{n \in \llbracket \frac{n_\ell(K)}{2}, n_\ell(K) \rrbracket} \frac{\sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}}}{\lambda_n^{(K)} + \mu_n^{(K)} - \eta - \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}}}$$

$$= \frac{\sqrt{b'(0) d'(0)}}{b'(0) + d'(0) - \sqrt{b'(0) d'(0)}} < 1.$$

We also observe that there exists $c > 0$ such that for K large enough, and any $n \in \llbracket n_r(K), n_d(K, \eta) \rrbracket$, we have

$$\frac{\sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}}}{\lambda_n^{(K)} + \mu_n^{(K)} - \eta - \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}}} \leq \frac{1}{1 + \frac{c(n-Kx_*)^2}{K^2}}.$$

The result follows by inspecting the monotonicity in the different cases and using the last part of Proposition (A.1). ■

PROPOSITION 4.2. *Let $\phi \in \text{Dom}(\mathcal{L}_K) \subset \ell^2$ of norm 1, satisfying*

$$\mathcal{L}_K \phi = -\rho \phi$$

for some real ρ . Then there exist $C(\phi) > 0$ and an integer $r(\phi)$ such that for all $n \geq r(\phi)$ we have

$$|\phi(n)| \leq C(\phi) 2^{-n}.$$

Moreover, $(\phi(n))_{n \geq r(\phi)}$ does not vanish and is of constant sign.

PROOF. It follows from our hypothesis that for any $K > 1$ there exists an integer $r_0(K)$ such that, for all $n \geq r_0(K)$, we have

$$0 < \frac{\sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}}}{\lambda_n^{(K)} + \mu_{n+1}^{(K)} + \rho - \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}}} \leq \frac{1}{2}.$$

We can assume that $(\phi(n))_n$ is a sequence of real numbers and $\phi(r_0(K)) > 0$. We start by proving that $(\phi(n))_n$ is positive and decreasing for $n \geq r_0(K)$. There are only the following four possibilities.

1. $\phi(r_0(K) + 1) \geq \phi(r_0(K))$. It follows from Proposition A.3 that ϕ is increasing for $n \geq r_0(K)$, contradicting that ϕ has norm 1.
2. $\phi(r_0(K) + 1) < \phi(r_0(K))$, and there exists $r' > r_0(K)$ such that $\phi(r') < 0$ and ϕ decreases on $\llbracket r_0(K), r' \rrbracket$, and $\phi \geq 0$ on $\llbracket r_0(K), r' - 1 \rrbracket$. Then by Proposition A.3, ϕ is decreasing for $n \geq r'$, contradicting that ϕ is normalized.
3. $\phi(r_0(K) + 1) < \phi(r_0(K))$, and there exists $r' > r_0(K)$ such that $\phi(r') < 0$ and ϕ is not monotonous on $\llbracket r_0(K), r' \rrbracket$, and $\phi \geq 0$ on $\llbracket r_0(K), r' - 1 \rrbracket$. Then there exists $r'' < r'$ such that ϕ is decreasing on $\llbracket r_0(K), r'' \rrbracket$, and such that $\phi(r'' + 1) \geq \phi(r'')$. If $\phi(r'') > 0$, then we are in case 1. If $\phi(r'' + 1) > 0$, it follows from Proposition A.3 that ϕ is increasing for $n \geq r'' + 1$, contradicting that ϕ has norm 1. If $\phi(r'' + 1) = \phi(r''') = 0$ then ϕ is the null sequence as solution of a second order equation, which leads to a contradiction.

4. $\phi(r_0(K) + 1) < \phi(r_0(K))$, $\phi \geq 0$. Suppose that there exists a local minimum at r''' (finite). Then if $\phi(r''' + 1) \geq \phi(r''') > 0$, we are in case 1. If $\phi(r''' + 1) > \phi(r''') = 0$, then it follows from Proposition A.3 that ϕ is increasing for $n \geq r''' + 1$, contradicting that ϕ has norm 1. If $\phi(r''' + 1) = \phi(r''') = 0$ then ϕ is the null sequence, which leads to a contradiction.

Therefore ϕ is strictly positive and monotone decreasing. The result then follows by using the last part of Proposition A.1.

■

Let us now prove two key lemmas.

LEMMA 4.3. *Let $\phi^K \in \text{Dom}(\mathcal{L}_K)$ be a normalized sequence such that*

$$\mathcal{L}_K \phi^{(K)} = -\rho^{(K)} \phi^{(K)}.$$

Assume that there exists a diverging sequence (K_p) such that

$$0 \leq \lim_{p \rightarrow +\infty} \rho^{(K_p)} = \rho_* < +\infty$$

and

$$\limsup_{p \rightarrow +\infty} \left\| \phi^{(K_p)} \mathbf{1}_{\{\cdot \leq n_\ell(K_p)\}} \right\|_{\ell^2} > 0.$$

Then $\rho_ \in S_2$ and there exists a diverging subsequence $(K_{p_\ell})_{\ell \geq 1}$ such that the limit*

$$\lim_{\ell \rightarrow +\infty} \phi^{(K_{p_\ell})} = \phi_*$$

exists in ℓ^2 , $\|\phi_\|_{\ell^2} > 0$ and ϕ_* is an eigenvector of \mathcal{M}_0 with eigenvalue $-\rho_*$.*

PROOF. Let $(p_i)_{i \geq 1}$ be a diverging sequence of integers such that

$$\lim_{i \rightarrow +\infty} \left\| \phi^{(K_{p_i})} \mathbf{1}_{\{\cdot \leq n_\ell(K)\}} \right\|_{\ell^2} = \limsup_{p \rightarrow +\infty} \left\| \phi^{(K_p)} \mathbf{1}_{\{\cdot \leq n_\ell(K_p)\}} \right\|_{\ell^2}.$$

For each i , we define a normalized sequences in ℓ^2 by

$$\psi_i(n) = \frac{\phi^{(K_{p_i})}(n) \mathbf{1}_{\{n \leq n_\ell(K_{p_i})\}}}{\left\| \phi^{(K_{p_i})} \mathbf{1}_{\{\cdot \leq n_\ell(K_{p_i})\}} \right\|_{\ell^2}}.$$

It is easy to verify using Proposition 4.1 that $\psi_i \in \text{Dom}(\mathcal{M}_0)$ and

$$\left\| \mathcal{M}_0 \psi_i + \rho^{(K_{p_i})} \psi_i \right\|_{\ell^2} \leq \frac{\mathcal{O}(1)(\log K_{p_i})^2 e^{-a(\log K_{p_i})^2}}{\left\| \phi^{(K_{p_i})} \mathbf{1}_{\{\cdot \leq n_\ell(K_{p_i})\}} \right\|_{\ell^2}}.$$

The first result follows from Proposition B.1 since the right-hand side goes to zero.

The second result follows from Proposition B.2 since the spectrum of \mathcal{M}_0 is discrete and simple by Theorem 7.1. ■

LEMMA 4.4. *Let $\phi^{(K)} \in \text{Dom}(\mathcal{L}_K)$ be a normalized sequence such that*

$$\mathcal{L}_K \phi^{(K)} = -\rho^{(K)} \phi^{(K)}.$$

Let us assume that there exists a diverging subsequence $(K_p)_{p \geq 1}$ such that

$$0 \leq \lim_{p \rightarrow +\infty} \rho^{(K_p)} = \rho_* < +\infty$$

and

$$\lim_{p \rightarrow +\infty} \|\phi^{(K_p)} \mathbb{1}_{\{\cdot \geq n_r(K_p)\}}\|_{\ell^2} > 0.$$

Then $\rho_ \in S_1$ and there exists a diverging subsequence of integers $(p_\ell)_{\ell \geq 1}$ such that*

$$\lim_{\ell \rightarrow +\infty} \frac{Q_{K_{p_\ell}} \phi^{(K_{p_\ell})}}{\|Q_{K_{p_\ell}} \phi^{(K_{p_\ell})}\|_{L^2}} = \psi_*$$

exists in L^2 , $\|\psi_\|_{L^2} = 1$, and ψ_* is an eigenvector of \mathcal{H}_* with eigenvalue $-\rho_*$.*

PROOF. We define for each p a normalized sequence in ℓ^2 by

$$\psi^{(K_p)}(n) = \frac{\phi^{(K_p)}(n) \mathbb{1}_{\{n \geq n_r(K_p)\}}}{\|\phi^{(K_p)} \mathbb{1}_{\{\cdot \geq n_r(K_p)\}}\|_{\ell^2}}.$$

It is easy to verify using Proposition 4.1 that $\psi^{(K_p)} \in \text{Dom}(\mathcal{L}_{K_p})$ and

$$\|\mathcal{L}_{K_p} \psi^{(K_p)} + \rho^{(K_p)} \psi^{(K_p)}\|_{\ell^2} \leq \mathcal{O}(1) \frac{K_p e^{-a(\log K_p)^2}}{\|\phi^{(K_p)} \mathbb{1}_{\{\cdot \geq n_r(K_p)\}}\|_{\ell^2}}. \quad (4.2)$$

From Proposition 4.2 it follows that, for each p , $\psi^{(K_p)}$ decays exponentially fast at infinity. We apply Proposition 6.6, Lemma 6.2, Lemma 6.3, and Theorem 6.1 to conclude that there exists a diverging sequence of integers $(p_i)_{i \geq 1}$ such that the sequence of functions $Q_{K_{p_i}} \psi^{(K_{p_i})}$ converges in L^2 to a normalized function ψ_* .

Let $u \in \mathcal{D}$. We have from (4.2)

$$\lim_{i \rightarrow +\infty} \langle P_{K_{p_i}} u, \mathcal{L}_{K_{p_i}} \psi^{(K_{p_i})} + \rho^{(K_{p_i})} \psi^{(K_{p_i})} \rangle_{\ell^2} = 0$$

hence, since $P_{K_{p_i}} u \in \text{Dom}(\mathcal{L}_{K_{p_i}})$, we obtain

$$\lim_{i \rightarrow +\infty} \langle \mathcal{L}_{K_{p_i}} P_{K_{p_i}} u, \psi^{(K_{p_i})} \rangle_{\ell^2} = -\rho_* \lim_{i \rightarrow +\infty} \langle P_{K_{p_i}} u, \psi^{(K_{p_i})} \rangle_{\ell^2}.$$

From the isometric property of Q (see Lemma 3.1) we get

$$\lim_{i \rightarrow +\infty} \langle Q_{K_{p_i}} \mathcal{L}_{K_{p_i}} P_{K_{p_i}} u, Q_{K_{p_i}} \psi^{(K_{p_i})} \rangle_{L^2}$$

$$= -\rho_* \lim_{i \rightarrow +\infty} \langle Q_{K_{p_i}} P_{K_{p_i}} u, Q_{K_{p_i}} \psi^{(K_{p_i})} \rangle_{L^2}.$$

In other words

$$\lim_{\ell \rightarrow +\infty} \langle \mathcal{L}_{K_{p_\ell}} u, Q_{K_{p_\ell}} \psi^{(K_{p_\ell})} \rangle_{L^2} = -\rho_* \lim_{\ell \rightarrow +\infty} \langle Q_{K_{p_\ell}} P_{K_{p_\ell}} u, Q_{K_{p_\ell}} \psi^{(K_{p_\ell})} \rangle_{L^2}.$$

Using the convergence in L^2 of $(Q_{K_{p_\ell}} \psi^{(K_{p_\ell})})_{\ell \geq 1}$ to ψ_* , Proposition 3.3, and Lemma 3.2, we get

$$\langle \mathcal{H}_* u, \psi_* \rangle_{L^2} = -\rho_* \langle u, \psi_* \rangle_{L^2}, \quad \forall u \in \mathcal{D}.$$

Since \mathcal{D} is dense in the domain of the self-adjoint operator \mathcal{H}_* , we conclude that ψ_* is an eigenvector of \mathcal{H}_* . ■

REMARK 4.1. *Another method to separate the eigenvectors in their dominant components is to use the so-called IMS localization formula. See for example [15]. In [10], convergence of the spectrum of a discrete operator to the spectrum of a continuous one was investigated using discrete pseudo-differential calculus.*

We now state three key propositions.

PROPOSITION 4.5. *Let j be fixed and let $(K_p)_{p \geq 1}$ be a diverging sequence such that*

$$\lim_{p \rightarrow +\infty} \rho_j^{(K_p)} = \rho_*.$$

Let $\phi_j^{(K_p)}$ be a normalized eigenvector of \mathcal{L}_{K_p} with eigenvalue $-\rho_j^{(K_p)}$. Then there exists an infinite sequence of integers $(p_i)_{i \geq 1}$ such that

- $\phi_j^{(K_{p_i})} \mathbb{1}_{\{\cdot < n_\ell(K_{p_i})\}} \xrightarrow{\ell^2} \phi_*$, where ϕ_* is either the null sequence or an eigenvector of \mathcal{M}_0 with eigenvalue $-\rho_*$, namely $\rho_* \in S_2$.
- $Q_{K_{p_i}} \phi_j^{(K_{p_i})} \mathbb{1}_{\{\cdot \geq n_r(K_{p_i})\}} \xrightarrow{L^2} \varphi_*$, where φ_* is either the null function or an eigenvector of \mathcal{H}_* with eigenvalue $-\rho_*$, namely $\rho_* \in S_1$.

Moreover, $\|\phi_\|_{\ell^2}^2 + \|\varphi_*\|_{L^2}^2 = 1$.*

PROOF. We have either

$$\lim_{p \rightarrow +\infty} \left\| \phi_j^{(K_p)} \mathbb{1}_{\{\cdot < n_\ell(K_p)\}} \right\|_{\ell^2} = 0$$

or there exists an infinite sequence of integers $(p_i)_{i \geq 1}$ such that

$$\lim_{i \rightarrow +\infty} \left\| \phi_j^{(K_{p_i})} \mathbb{1}_{\{\cdot < n_\ell(K_{p_i})\}} \right\|_{\ell^2} > 0.$$

The first statement of the proposition follows from Lemma 4.3 applied to the diverging sequence $(K_{p_i})_{i \geq 1}$. Similarly, either

$$\lim_{p \rightarrow +\infty} \left\| \phi_j^{(K_p)} \mathbf{1}_{\{\cdot \geq n_r(K)\}} \right\|_{\ell^2} = 0$$

or there exists an infinite sequence of integers $(p_i)_{i \geq 1}$ such that

$$\lim_{i \rightarrow +\infty} \left\| \phi_j^{(K_{p_i})} \mathbf{1}_{\{\cdot \geq n_r(K_{p_i})\}} \right\|_{\ell^2} > 0.$$

The second statement of the proposition follows from Lemma 4.4 applied to the diverging sequence $(K_{p_i})_{i \geq 1}$. The last statement follows from the normalization $\phi_j^{(K_p)}$ and Proposition 4.1. ■

PROPOSITION 4.6. *Let $\rho \in S_2$. Then there exists a normalized vector $v_\rho \in \text{Dom}(\mathcal{L}_K)$ with eigenvalue $-\rho$ such that*

$$\lim_{K \rightarrow +\infty} \|\mathcal{L}_K v_\rho + \rho v_\rho\|_{\ell^2} = 0.$$

Moreover

$$\lim_{K \rightarrow +\infty} \|v_\rho \mathbf{1}_{\{\cdot > n_\ell(K)\}}\|_{\ell^2} = 0.$$

PROOF. Let v_ρ be a normalized eigenvector corresponding to the eigenvalue $-\rho$ for the operator \mathcal{M}_0 in the space ℓ^2 . From the assumptions we have for $n \leq \lfloor \log K \rfloor$

$$\lambda_n^{(K)} = b'(0) + \mathcal{O}\left(\frac{\log K}{K}\right) \quad \text{and} \quad \mu_n^{(K)} = d'(0) + \mathcal{O}\left(\frac{\log K}{K}\right).$$

Since $\mathcal{M}_0 v_\rho + \rho v_\rho = 0$, the reader can easily check that

$$\|\mathcal{L}_K v_\rho + \rho v_\rho\|_{\ell^2(\mathbb{Z}_{>0} \cap \llbracket 1, \lfloor \log K \rfloor \rrbracket)} \leq \mathcal{O}\left(\frac{\log K}{K}\right).$$

Now using Theorem 7.1 we have

$$v_\rho(n) = \sqrt{n} \left(\frac{d'(0)}{b'(0)} \right)^{\frac{n}{2}} P(n)$$

where P is a certain polynomial. Hence there exists $c_v > 0$ such that for any $n \in \mathbb{Z}_{>0}$, $|v_\rho(n)| \leq c_v \left(\frac{d'(0)}{b'(0)} \right)^{\frac{n}{4}}$. By (2.1) there exists $c > 0$ such $b(x) + d(x) \leq c e^x$ for all $x \geq 0$. This implies that $\lambda_n^{(K_p)} \leq c K_p e^{n/K_p}$ and $\mu_n^{(K_p)} \leq c K_p e^{n/K_p}$. The reader can easily check that

$$\|\mathcal{L}_K v_\rho + \rho v_\rho\|_{\ell^2(\mathbb{Z}_{>0} \cap \llbracket \lfloor \log K \rfloor + 1, \infty \rrbracket)} \leq \mathcal{O}(1) \left(\frac{d'(0)}{b'(0)} \right)^{\frac{\log K}{4}}.$$

It follows that $v_\rho \in \text{Dom}(\mathcal{L}_K)$, and (remember that $b'(0) > d'(0)$)

$$\lim_{K \rightarrow +\infty} \|\mathcal{L}_K v_\rho + \rho v_\rho\|_{\ell^2} = 0.$$

The other statement follows at once from the exponential decay of v . ■

PROPOSITION 4.7. *Let $\rho \in S_1$. Then there exists a sequence of normalized vectors $(\psi_\rho^{(K)})_{K \geq 1} \subset \ell^2$ such that $\psi_\rho^{(K)} \in \text{Dom}(\mathcal{L}_K)$, and*

$$\lim_{K \rightarrow +\infty} \|\mathcal{L}_K \psi_\rho^{(K)} + \rho \psi_\rho^{(K)}\|_{\ell^2} = 0.$$

Moreover

$$\lim_{K \rightarrow +\infty} \|\psi_\rho^{(K)} \mathbf{1}_{\{\cdot < n_r(K)\}}\|_{\ell^2} = 0.$$

PROOF. Let φ_ρ be a real normalized eigenvector of \mathcal{H}_* corresponding to the eigenvalue $-\rho$ (in L^2). Since φ_ρ is a (rescaled) Hermite function (see [2] or Remark 3.1), it satisfies the hypothesis of Proposition 3.3, hence

$$\lim_{K \rightarrow +\infty} \|\mathcal{L}_K \varphi_\rho + \rho \varphi_\rho\|_{L^2} = 0.$$

Using Lemma 3.2 we get

$$\lim_{K \rightarrow +\infty} \|\mathcal{L}_K \varphi_\rho + \rho Q_K P_K \varphi_\rho\|_{L^2} = 0$$

and then

$$\lim_{K \rightarrow +\infty} \|\mathcal{L}_K P_K \varphi_\rho + \rho P_K \varphi_\rho\|_{\ell^2} = 0$$

and

$$\lim_{K \rightarrow +\infty} \|P_K \varphi_\rho\|_{\ell^2} = 1.$$

The first statement follows by letting $\psi_\rho^{(K)}(n) = (P_K \varphi_\rho)(n)$, where P_K is defined in (3.2). The other statement follows from an exponential bound on the decay of φ_ρ . ■

5 Proofs of Theorems 3.5 and 3.7

5.1 Proof of Theorem 3.5

The proof of Theorem 3.5 is an immediate consequence of the following two propositions.

PROPOSITION 5.1. *We have $S_2 \subset G$.*

PROOF. The proof is by contradiction. Let $\rho \in S_2$ and assume $\rho \notin G$. Then there exists $\eta > 0$ be such that for K large enough, $[\rho - \eta, \rho + \eta] \cap G = \emptyset$. It follows from Proposition 4.6 that there exists a normalized vector v in ℓ^2 , such that $v \in \text{Dom}(\mathcal{L}_K)$, and

$$\lim_{K \rightarrow +\infty} \|\mathcal{L}_K v + \rho v\|_{\ell^2} = 0.$$

Therefore, using Proposition B.1, we obtain a contradiction. This ends the proof. ■

PROPOSITION 5.2. *We have $S_1 \subset G$.*

PROOF. The proof is by contradiction. Let $\rho \in S_1$ and assume $\rho \notin G$. Then there exists $\eta > 0$ such that, for K large enough, $[\rho - \eta, \rho + \eta] \cap G = \emptyset$. It follows from Proposition 4.7 that there exists a sequence of normalized vectors $(\psi^{(K)})_{K \geq 1} \subset \ell^2$ such that $\psi^{(K)} \in \text{Dom}(\mathcal{L}_K)$ and

$$\lim_{K \rightarrow +\infty} \|\mathcal{L}_K \psi^{(K)} + \rho \psi^{(K)}\|_{\ell^2} = 0.$$

Therefore, using Proposition B.1, we obtain a contradiction. This ends the proof. ■

5.2 Proof of Theorem 3.7

The first statement follows at once from Proposition 4.5.

The proof of (3.6) is by contradiction. Assume

$$\liminf_{p \rightarrow +\infty} \min \left\{ \left| \rho_{j+1}^{(K_p)} - \rho_* \right|, \left| \rho_{j-1}^{(K_p)} - \rho_* \right| \right\} = 0.$$

Assume $\rho_* \in S_1 \setminus S_2$ and let (p_ℓ) be a diverging sequence of positive integers such that $\rho_i^{(K_{p_\ell})} \rightarrow \rho_*$ and $\rho_{i+1}^{(K_{p_\ell})} \rightarrow \rho_*$ (where $i = j$ or $i = j - 1$). Let $\phi_i^{(K_{p_\ell})}$ be a normalized eigenvector of $\mathcal{L}_{K_{p_\ell}}$ corresponding to the eigenvalue $-\rho_i^{(K_{p_\ell})}$. We define $\phi_{i+1}^{(K_{p_\ell})}$ similarly. We claim that

$$\limsup_{\ell \rightarrow +\infty} \left\| \phi_i^{(K_{p_\ell})} \mathbb{1}_{\{\cdot < n_\ell(K_{p_\ell})\}} \right\|_{\ell^2} = 0.$$

Otherwise Proposition 4.5 would imply that $\rho_* \in S_2$, a contradiction. By Proposition 4.1 we have

$$\lim_{\ell \rightarrow +\infty} \left\| \phi_i^{(K_{p_\ell})} \mathbb{1}_{\{\cdot > n_r(K_{p_\ell})\}} \right\|_{\ell^2} = 1.$$

This implies 1-a).

By a similar argument, we have

$$\lim_{\ell \rightarrow +\infty} \left\| \phi_{i+1}^{(K_{p_\ell})} \mathbb{1}_{\{\cdot > n_r(K_{p_\ell})\}} \right\|_{\ell^2} = 1.$$

By Proposition 4.5, there exists a diverging sequence of integers (ℓ_r) such that

$$\left\| Q_{K_{p_{\ell_r}}} \left(\phi_i^{(K_{p_{\ell_r}})} \mathbb{1}_{\{\cdot > n_r(K_{p_{\ell_r}})\}} \right) - \varphi_* \right\|_{L^2} \rightarrow 0$$

where $\varphi_* \in \text{Dom}(\mathcal{H}_*)$ is a normalized eigenfunction of \mathcal{H}_* corresponding to the eigenvalue $-\rho_*$. By Proposition 4.5 again, there exists a diverging sequence of integers $(r_s)_{s \geq 1}$ such that

$$\left\| Q_{K_{p_{\ell_{r_s}}}} \left(\phi_{i+1}^{(K_{p_{\ell_{r_s}}})} \mathbb{1}_{\{\cdot > n_r(K_{p_{\ell_{r_s}}})\}} \right) - \varphi'_* \right\|_{L^2} \rightarrow 0$$

where $\varphi'_* \in \text{Dom}(\mathcal{H}_*)$ is a normalized eigenfunction of \mathcal{H}_* corresponding to the eigenvalue $-\rho_*$. Since $\phi_i^{(K_{p_{\ell_{r_s}}})}$ and $\phi_{i+1}^{(K_{p_{\ell_{r_s}}})}$ are orthogonal in ℓ^2 , it follows from the previous estimates that

$$\lim_{s \rightarrow +\infty} \left\langle \phi_i^{(K_{p_{\ell_{r_s}}})} \mathbb{1}_{\{\cdot > n_r(K_{p_{\ell_{r_s}}})\}}, \phi_{i+1}^{(K_{p_{\ell_{r_s}}})} \mathbb{1}_{\{\cdot > n_r(K_{p_{\ell_{r_s}}})\}} \right\rangle_{\ell^2} = 0.$$

By Lemma 3.1 we have

$$\left\langle Q_{K_{p_{\ell_{r_s}}}} \left(\phi_i^{(K_{p_{\ell_{r_s}}})} \mathbb{1}_{\{\cdot > n_r(K_{p_{\ell_{r_s}}})\}} \right), Q_{K_{p_{\ell_{r_s}}}} \left(\phi_{i+1}^{(K_{p_{\ell_{r_s}}})} \mathbb{1}_{\{\cdot > n_r(K_{p_{\ell_{r_s}}})\}} \right) \right\rangle_{L^2} \xrightarrow{s \rightarrow +\infty} 0.$$

In other words, $\langle \varphi_*, \varphi'_* \rangle_{L^2} = 0$. This is a contradiction since φ_* and φ'_* are normalized eigenfunctions of \mathcal{H}_* corresponding to the same eigenvalue $-\rho_*$ which is simple.

The case $\rho_* \in S_2 \setminus S_1$ is similar (using again Proposition 4.5), so it is left to the reader.

Let us now assume that $\rho_* \in S_1 \cap S_2$. We now prove (3.7) by contradiction. So we assume that there exists $\delta > 0$ such that

$$\liminf_{p \rightarrow +\infty} \min \left\{ \left| \rho_{j+1}^{(K_p)} - \rho_* \right|, \left| \rho_{j-1}^{(K_p)} - \rho_* \right| \right\} > \delta.$$

Since $\rho_* \in S_1$ and from Proposition 4.7, there exists a sequence of normalized vectors $(\psi_{\rho_*}^{(K_p)})_{p \geq 1} \subset \ell^2$ such that $\psi_{\rho_*}^{(K_p)} \in \text{Dom}(\mathcal{L}_{K_p})$ for all p , and

$$\lim_{p \rightarrow +\infty} \left\| \mathcal{L}_{K_p} \psi_{\rho_*}^{(K_p)} - \rho_* \psi_{\rho_*}^{(K_p)} \right\|_{\ell^2} = 0.$$

We also have (since $\rho_j^{(K_p)} \rightarrow \rho_*$) that

$$\lim_{p \rightarrow +\infty} \left\| \mathcal{L}_{K_p} \psi_{\rho_*}^{(K_p)} - \rho_j^{(K_p)} \psi_{\rho_*}^{(K_p)} \right\|_{\ell^2} = 0.$$

For each p , let $\phi_j^{(K_p)}$ be a normalized eigenvector of \mathcal{L}_{K_p} corresponding to the eigenvalue $-\rho_j^{(K_p)}$. By using Proposition B.2 with $A = \mathcal{L}_{K_p}$, we deduce that there exists a sequence of real numbers $(\theta_p)_p$ such that

$$\lim_{p \rightarrow +\infty} \left\| \psi_{\rho_*}^{(K_p)} - e^{i\theta_p} \phi_j^{(K_p)} \right\|_{\ell^2} = 0. \quad (5.1)$$

Since $\rho_* \in \mathcal{S}_2$ and from Proposition 4.6, there exists a normalized vector $v_{\rho_*} \in \text{Dom}(\mathcal{L}_{K_p})$ for all p such that

$$\lim_{p \rightarrow +\infty} \left\| \mathcal{L}_{K_p} v_{\rho_*} + \rho_* v_{\rho_*} \right\|_{\ell^2} = 0.$$

We also have

$$\lim_{p \rightarrow +\infty} \left\| \mathcal{L}_{K_p} v_{\rho_*} + \rho_j^{(K_p)} v_{\rho_*} \right\|_{\ell^2} = 0.$$

By using Proposition B.2 with $A = \mathcal{L}_{K_p}$, we deduce that there exists a sequence of real numbers $(\theta'_p)_{p \geq 1}$ such that

$$\lim_{p \rightarrow +\infty} \left\| v_{\rho_*} - e^{i\theta'_p} \phi_j^{(K_p)} \right\|_{\ell^2} = 0. \quad (5.2)$$

Moreover, from Propositions 4.7 and 4.6, it follows that

$$\lim_{p \rightarrow +\infty} \left\| \psi_{\rho_*}^{(K_p)} \mathbb{1}_{\{\cdot < n_r(K_p)\}} \right\|_{\ell^2} = 0 \quad \text{and} \quad \lim_{K \rightarrow +\infty} \left\| v_{\rho_*} \mathbb{1}_{\{\cdot > n_\ell(K)\}} \right\|_{\ell^2} = 0$$

which implies (using Proposition 4.1) that

$$\lim_{p \rightarrow +\infty} \langle \psi_{\rho_*}^{(K_p)}, v_{\rho_*} \rangle = 0.$$

This is a contradiction with (5.1) and (5.2). Finally, we prove (3.8) by contradiction. So we assume that there exists a diverging sequence of integers $(p_\ell)_{\ell \geq 1}$ such that

$$\lim_{\ell \rightarrow +\infty} \left| \rho_{j+1}^{(K_{p_\ell})} - \rho_* \right| + \left| \rho_{j-1}^{(K_{p_\ell})} - \rho_* \right| = 0.$$

We now apply Proposition 4.5 with j and K_{p_ℓ} . Hence there exists a diverging sequence of integers $(\ell_s)_{s \geq 1}$ such that

$$\phi_j^{(K_{p_{\ell_s}})} \mathbb{1}_{\{\cdot < n_\ell(K_{p_{\ell_s}})\}} \xrightarrow{\ell^2} \phi_{*,1}$$

and

$$Q_{K_{p_{\ell_s}}} \phi_j^{(K_{p_{\ell_s}})} \mathbb{1}_{\{\cdot \geq n_r(K_{p_{\ell_s}})\}} \xrightarrow{L^2} \varphi_{*,1}$$

and we have $\|\phi_{*,1}\|_{\ell^2}^2 + \|\varphi_{*,1}\|_{L^2}^2 = 1$.

We now apply Proposition 4.5 with $j - 1$ and $K_{p_{\ell_s}}$. Hence there exists a diverging sequence of integers $(s_r)_{r \geq 1}$ such that

$$\phi_{j-1}^{(K_{p_{\ell_{s_r}}})} \mathbb{1}_{\{\cdot < n_{\ell}(K_{p_{\ell_{s_r}}})\}} \xrightarrow{\ell^2} \phi_{*,2}$$

and

$$Q_{K_{p_{\ell_{s_r}}}} \phi_{j-1}^{(K_{p_{\ell_{s_r}}})} \mathbb{1}_{\{\cdot \geq n_r(K_{p_{\ell_{s_r}}})\}} \xrightarrow{L^2} \varphi_{*,2}$$

and we have $\|\phi_{*,2}\|_{\ell^2}^2 + \|\varphi_{*,2}\|_{L^2}^2 = 1$.

We now apply Proposition 4.5 with $j + 1$ and $K_{p_{\ell_{s_r}}}$. Hence there exists a diverging sequence of integers $(r_q)_{q \geq 1}$ such that

$$\phi_{j+1}^{(K_{p_{\ell_{s_r q}}})} \mathbb{1}_{\{\cdot < n_{\ell}(K_{p_{\ell_{s_r q}}})\}} \xrightarrow{\ell^2} \phi_{*,3}$$

and

$$Q_{K_{p_{\ell_{s_r q}}}} \phi_{j+1}^{(K_{p_{\ell_{s_r q}}})} \mathbb{1}_{\{\cdot \geq n_r(K_{p_{\ell_{s_r q}}})\}} \xrightarrow{L^2} \varphi_{*,3}$$

and we have $\|\phi_{*,3}\|_{\ell^2}^2 + \|\varphi_{*,3}\|_{L^2}^2 = 1$.

Moreover, since $\phi_{j-1}^{(K_{p_{\ell_{s_r q}}})}$, $\phi_j^{(K_{p_{\ell_{s_r q}}})}$, $\phi_{j+1}^{(K_{p_{\ell_{s_r q}}})}$ are pairwise orthogonal, we have

$$\langle \phi_{*,m}, \phi_{*,m'} \rangle_{\ell^2} + \langle \varphi_{*,m}, \varphi_{*,m'} \rangle_{L^2} = 0, \quad \forall m \neq m'. \quad (5.3)$$

The linear subspace of ℓ^2 spanned by $\phi_{*,1}, \phi_{*,2}, \phi_{*,3}$ is of dimension at most one because they are eigenvectors of \mathcal{M}_0 for the same simple eigenvalue $-\rho_*$. The linear subspace of L^2 spanned by $\varphi_{*,1}, \varphi_{*,2}, \varphi_{*,3}$ is of dimension at most one because they are eigenfunctions of \mathcal{H}_* for the same simple eigenvalue $-\rho_*$. Therefore the subspace of $\ell^2 \oplus L^2$ spanned by the three vectors $(\phi_{*,1}, \varphi_{*,1})$, $(\phi_{*,2}, \varphi_{*,2})$, $(\phi_{*,3}, \varphi_{*,3})$ is of dimension at most two. However, these three vectors are normalized and pairwise orthogonal by (5.3). We thus arrive at a contradiction.

6 Fréchet-Kolmogorov-Riesz compactness criterion and Dirichlet form

6.1 Fréchet-Kolmogorov-Riesz compactness criterion

We recall the Fréchet-Kolmogorov-Riesz compactness criterion in L^2 .

THEOREM 6.1. *Let $(f_p)_p$ be a normalized sequence in L^2 such that the following two conditions are satisfied.*

- (i) *There exists $\epsilon_0 > 0$ such that there exists a function $R(\epsilon) > 0$ on $(0, \epsilon_0)$ such that*

$$\sup_p \int_{\{|x| > R(\epsilon)\}} |f_p(x)|^2 dx \leq \epsilon.$$

(ii) There exists a positive function α on $]0, 1]$ satisfying

$$\lim_{y \downarrow 0} \alpha(y) = 0$$

and such that for any p and $y \in]-1, 1]$

$$\int |f_p(x+y) - f_p(x)|^2 dx \leq \alpha(|y|).$$

Then one can extract from $(f_p)_p$ a convergent subsequence in L^2 .

We refer to Remark 5 on page 387 in [9]. The following lemmas provide expressions for $R(\epsilon)$ and $\alpha(y)$ in our case. Recall that the potential $V_n(K)$ and $n_r(K)$ have been defined in Section 4 (see (4.1)).

LEMMA 6.2. *Let $C > 0$. Let $\mathcal{F}_{C,K}$ be the set of normalized sequences $(\phi(n))_n$ in ℓ^2 such that $\phi(n) = 0$ for any $n < n_r(K)$ and*

$$\sum_{n=n_r(K)}^{+\infty} (1 \vee V_n(K)) \phi^2(n) \leq C.$$

Then, for any $\phi \in \mathcal{F}_{C,K}$, the function $x \mapsto Q_K \phi(x)$ satisfies condition (i) in Theorem 6.1 with

$$R(\epsilon) = \frac{C}{\epsilon}$$

for any $0 < \epsilon < 1$.

PROOF. We are going to prove that there exist $R_0 > 1$ and $K_0 > 4$ such that for any $K > K_0$, any $R > R_0$ and any $\phi \in \mathcal{F}_{C,K}$

$$\int_{\{|x|>R\}} (Q_K \phi(x))^2 dx \leq \frac{C}{R}.$$

We observe that

$$\int_{\{|x|>R\}} (Q_K \phi(x))^2 dx \leq \sum_{\{n:|n-Kx_*|>R\sqrt{K}-1\}} \phi^2(n).$$

Our aim is to prove that the right-hand side of the above inequality is bounded above by C/R .

It follows from the hypotheses on $\lambda_n^{(K)}$ and $\mu_n^{(K)}$ that there exist constants $K_0 > 4$, $1 > C_2 > 0$, $\Gamma > 0$ and $\zeta > 0$, such that for any $K > K_0$ there exists an integer $\Gamma K > m_{*(K)} > 2Kx_*$ (hence of order K) such that $\mu_n^{(K)} > \zeta n$ for any $n \geq m_{*(K)}$ and

$$\sum_{n=n_r(K)}^{m_{*(K)}} (1 \vee V_n(K)) \phi^2(n) \geq C_2 K^{-1} \sum_{n=n_r(K)}^{m_{*(K)}} (n - Kx_*)^2 \phi^2(n)$$

$$\sum_{n=m_*(K)+1}^{+\infty} (1 \vee V_n(K)) \phi^2(n) \geq C_2 \sum_{n=m_*(K)+1}^{+\infty} n \phi^2(n).$$

These estimates imply the following bounds for any integer $L > 0$

$$\begin{aligned} \sum_{\{n_r(K) \leq n \leq m_*(K)\} \cap \{|n - Kx_*| > L\}} \phi^2(n) &\leq \frac{KC}{C_2 L^2} \mathbb{1}_{\{L < m_*(K)\}} \\ \sum_{\{n > m_*(K)\} \cap \{|n - Kx_*| > L\}} \phi^2(n) &\leq \frac{C}{C_2(L \vee m_*(K))}. \end{aligned}$$

We now replace L by $R\sqrt{K} - 1$ in the above estimates.

Fix $K > 4$. We distinguish two cases according to the value of R .

- 1) $1 \leq R < m_*(K)/\sqrt{K}$. Then $\sqrt{K} - 1 \leq L < m_*(K) \leq \Gamma K$. Since $L = R\sqrt{K} - 1 > R\sqrt{K}/2$ (because $K > 4$), we have

$$\sum_{\{|n - Kx_*| > R\sqrt{K} - 1\}} \phi^2(n) \leq \frac{4C}{R^2} + \frac{C}{C_2 m_*(K)} \leq \frac{4C}{C_2 R^2} + \frac{C\Gamma}{C_2 R^2} \leq \frac{C}{R}$$

if $R > C_2^{-1}(4 + \Gamma)$.

- 2) $R \geq m_*(K)/\sqrt{K}$. Then $L \geq m_*(K)$. We get

$$\sum_{\{|n - Kx_*| > R\sqrt{K} - 1\}} \phi^2(n) \leq \frac{C}{C_2 L} \leq \frac{2C}{R C_2 \sqrt{K}} \leq \frac{C}{R}$$

if $C_2^2 K > 4$.

We define $K_0 = 5 + 4C_2^{-2}$ and $R_0 = 1 + C_2^{-1}(4 + \Gamma)$. The result follows. ■

LEMMA 6.3. *Let $(\phi^{(K)})_{K \in \mathbb{Z}_{>0}}$ be a sequence of normalized elements of ℓ^2 such that $\phi^{(K)}(n) = 0$ for $n \leq n_r(K)$. Assume also that there exists $C > 0$ such that*

$$\sup_{K > 1} \left\{ \sum_{n=n_r(K)}^{+\infty} \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} (\phi^{(K)}(n+1) - \phi^{(K)}(n))^2 + \sum_{n=n_r(K)}^{+\infty} (1 \vee V_n(K)) (\phi^{(K)}(n))^2 \right\} \leq C.$$

Then there exists a positive constant \tilde{C} such that for any $|h| \leq 1$ and any $K > 1$

$$\int [Q_K \phi^{(K)}(x+h) - Q_K \phi^{(K)}(x)]^2 dx \leq \alpha(h) = \tilde{C} |h|.$$

Hence, for any $(\phi^{(K)})_K$ satisfying the above assumptions, the sequence of functions $(Q_K \phi^{(K)})_{K \geq 1}$ satisfies condition ii) in Theorem 6.1 with

$$\alpha(y) = \tilde{C}y.$$

PROOF. It is enough to consider the case $0 < h < 1$. We first consider the case $0 < h \leq 1/\sqrt{K}$. We have

$$\int [Q_K \phi^{(K)}(x+h) - Q_K \phi^{(K)}(x)]^2 dx = \sum_{q \geq 1} \int_{I_q^{(K)}} [Q_K \phi^{(K)}(x+h) - Q_K \phi^{(K)}(x)]^2 dx.$$

Since

$$Q_K \phi^{(K)}(x) = K^{\frac{1}{4}} \sum_{q \geq 1} \phi^{(K)}(q) \mathbf{1}_{I_q^{(K)}}(x)$$

and since the intervals $I_q^{(K)}$ are disjoint, we get

$$\begin{aligned} & \int [Q_K \phi^{(K)}(x+h) - Q_K \phi^{(K)}(x)]^2 dx \\ &= K^{\frac{1}{2}} \sum_{q \geq 1} \int_{I_q^{(K)}} \left(\phi^{(K)}(q) \mathbf{1}_{I_q^{(K)}}(x+h) \right. \\ & \quad \left. + \phi^{(K)}(q+1) \mathbf{1}_{I_{q+1}^{(K)}}(x+h) - \phi^{(K)}(q) \mathbf{1}_{I_q^{(K)}}(x) \right)^2 dx \\ &= K^{\frac{1}{2}} \sum_{q \geq 1} \int_{I_q^{(K)}} \left((\phi^{(K)}(q))^2 \mathbf{1}_{I_q^{(K)}}(x+h) + (\phi^{(K)}(q+1))^2 \mathbf{1}_{I_{q+1}^{(K)}}(x+h) \right. \\ & \quad \left. + (\phi^{(K)}(q))^2 \mathbf{1}_{I_q^{(K)}}(x) - 2(\phi^{(K)}(q))^2 \mathbf{1}_{I_q^{(K)}}(x) \mathbf{1}_{I_q^{(K)}}(x+h) \right. \\ & \quad \left. - 2\phi^{(K)}(q)\phi^{(K)}(q+1) \mathbf{1}_{I_q^{(K)}}(x) \mathbf{1}_{I_{q+1}^{(K)}}(x+h) \right) dx \end{aligned}$$

Let us consider each term separately. Since $\int_{I_q^{(K)}} dx = K^{-\frac{1}{2}}$, we have

$$K^{\frac{1}{2}} \sum_{q \geq 1} \int_{I_q^{(K)}} (\phi^{(K)}(q))^2 dx = \sum_{q \geq 1} (\phi^{(K)}(q))^2.$$

Then we have

$$\begin{aligned} & K^{\frac{1}{2}} \sum_{q \geq 1} \int_{I_q^{(K)}} (\phi^{(K)}(q))^2 \mathbf{1}_{I_q^{(K)}}(x+h) dx \\ &= K^{\frac{1}{2}} \sum_{q \geq 1} (\phi^{(K)}(q))^2 \int_{\frac{q}{\sqrt{K}} - \frac{1}{2\sqrt{K}} - x_*}^{\frac{q}{\sqrt{K}} + \frac{1}{2\sqrt{K}} - x_* - h} dx \\ &= (1 - hK^{\frac{1}{2}}) \sum_{q \geq 1} (\phi^{(K)}(q))^2. \end{aligned}$$

We also have

$$K^{\frac{1}{2}} \sum_{q \geq 1} \int_{I_q^{(K)}} (\phi^{(K)}(q+1))^2 \mathbf{1}_{I_{q+1}^{(K)}}(x+h) dx$$

$$\begin{aligned}
&= K^{\frac{1}{2}} \sum_{q \geq 1} (\phi^{(K)}(q+1))^2 \int_{\frac{q}{\sqrt{K}} + \frac{1}{2\sqrt{K}} - x_* - h}^{\frac{q}{\sqrt{K}} + \frac{1}{2\sqrt{K}} - x_*} dx \\
&= K^{\frac{1}{2}} \sum_{q \geq 1} (\phi^{(K)}(q+1))^2 h \\
&= K^{\frac{1}{2}} h \sum_{q \geq 1} (\phi^{(K)}(q))^2.
\end{aligned}$$

Similarly

$$\begin{aligned}
&- 2K^{\frac{1}{2}} \sum_{q \geq 1} \int_{I_q^{(K)}} (\phi^{(K)}(q))^2 \mathbb{1}_{I_q^{(K)}}(x) \mathbb{1}_{I_q^{(K)}}(x+h) dx \\
&= -2(1 - hK^{\frac{1}{2}}) \sum_{q \geq 1} (\phi^{(K)}(q))^2
\end{aligned}$$

and

$$\begin{aligned}
&- 2K^{\frac{1}{2}} \sum_{q \geq 1} \int_{I_q^{(K)}} \phi^{(K)}(q) \phi^{(K)}(q+1) \mathbb{1}_{I_q^{(K)}}(x) \mathbb{1}_{I_{q+1}^{(K)}}(x+h) dx \\
&= -2hK^{\frac{1}{2}} \sum_{q \geq 1} \phi^{(K)}(q) \phi^{(K)}(q+1).
\end{aligned}$$

We rewrite the last term as follows:

$$\begin{aligned}
&- 2h\sqrt{K} \sum_{q \geq 1} \phi^{(K)}(q) \phi^{(K)}(q+1) \\
&= -2h\sqrt{K} \sum_{q \geq 1} (\phi^{(K)}(q))^2 + 2h\sqrt{K} \sum_{q \geq 1} \phi^{(K)}(q) (\phi^{(K)}(q+1) - \phi^{(K)}(q)).
\end{aligned}$$

Summing up, we get

$$\begin{aligned}
&\int [Q_K \phi^{(K)}(x+h) - Q_K \phi^{(K)}(x)]^2 dx \\
&= \sum_{q \geq 1} (\phi^{(K)}(q))^2 + (1 - h\sqrt{K}) \sum_{q \geq 1} (\phi^{(K)}(q))^2 + K^{\frac{1}{2}} h \sum_{q \geq 1} (\phi^{(K)}(q))^2 \\
&\quad - 2(1 - h\sqrt{K}) \sum_{q \geq 1} (\phi^{(K)}(q))^2 - 2h\sqrt{K} \sum_{q \geq 1} (\phi^{(K)}(q))^2 \\
&\quad + 2h\sqrt{K} \sum_{q \geq 1} \phi^{(K)}(q) (\phi^{(K)}(q+1) - \phi^{(K)}(q)) \\
&= 2h\sqrt{K} \sum_{q \geq 1} \phi^{(K)}(q) (\phi^{(K)}(q+1) - \phi^{(K)}(q)).
\end{aligned}$$

It follows from Cauchy-Schwarz inequality that

$$\int [Q_K \phi^{(K)}(x+h) - Q_K \phi^{(K)}(x)]^2 dx$$

$$\leq 2h\sqrt{K} \left(\sum_{q \geq 1} (\phi^{(K)}(q+1) - \phi^{(K)}(q))^2 \right)^{\frac{1}{2}} \|\phi^{(K)}\|_{\ell^2}.$$

From the assumption of the lemma and using that

$$\inf_{n \geq \lfloor \frac{Kx_*}{3} \rfloor - 1} \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} > \zeta K$$

for some $\zeta > 0$ independent of K , we get

$$K^{\frac{1}{2}} \left(\sum_{q \geq 1} (\phi^{(K)}(q+1) - \phi^{(K)}(q))^2 \right)^{1/2} \leq \sqrt{\frac{C}{\zeta}}.$$

Therefore, since the sequence $\phi^{(K)}$ is normalized, we get

$$\int [Q_K \phi^{(K)}(x+h) - Q_K \phi^{(K)}(x)]^2 dx \leq 2h \sqrt{\frac{C}{\zeta}}.$$

We now consider the case $1 > h > 1/\sqrt{K}$. Let $r = \lfloor h\sqrt{K} \rfloor$ and $h' = h - r/\sqrt{K}$. Note that $0 \leq h' \leq 1/\sqrt{K}$. We have

$$\begin{aligned} & \int [Q_K \phi^{(K)}(x+h) - Q_K \phi^{(K)}(x)]^2 dx \\ &= \int \left[\sum_{j=1}^{r-1} \left(Q_K \phi^{(K)}\left(x + \frac{j+1}{\sqrt{K}}\right) - Q_K \phi^{(K)}\left(x + \frac{j}{\sqrt{K}}\right) \right) \right. \\ & \quad \left. + Q_K \phi^{(K)}\left(x + \frac{r}{\sqrt{K}} + h'\right) - Q_K \phi^{(K)}\left(x + \frac{r}{\sqrt{K}}\right) \right]^2 dx \\ &\leq 2 \int \left[\sum_{j=1}^{r-1} \left(Q_K \phi^{(K)}\left(x + \frac{j+1}{\sqrt{K}}\right) - Q_K \phi^{(K)}\left(x + \frac{j}{\sqrt{K}}\right) \right) \right]^2 dx \\ & \quad + 2 \int \left[Q_K \phi^{(K)}\left(x + \frac{r}{\sqrt{K}} + h'\right) - Q_K \phi^{(K)}\left(x + \frac{r}{\sqrt{K}}\right) \right]^2 dx. \end{aligned}$$

We have already estimated the last term. For the first term we now observe that $e_n^{(K)}\left(x + \frac{j}{\sqrt{K}}\right) = e_{n-j}^{(K)}(x)$. Therefore we can write

$$\begin{aligned} & \sum_{j=1}^{r-1} \left(Q_K \phi^{(K)}\left(x + \frac{j+1}{\sqrt{K}}\right) - Q_K \phi^{(K)}\left(x + \frac{j}{\sqrt{K}}\right) \right) \\ &= \sum_{j=1}^{r-1} \left(\sum_n \phi^{(K)}(n) e_{n-j-1}^{(K)}(x) - \sum_n \phi^{(K)}(n) e_{n-j}^{(K)}(x) \right) \end{aligned}$$

$$\begin{aligned}
&= \sum_{j=1}^{r-1} \left(\sum_p e_p^{(K)}(x) (\phi^{(K)}(p+j+1) - \phi^{(K)}(p+j)) \right) \\
&= \sum_p e_p^{(K)}(x) \sum_{j=1}^{r-1} (\phi^{(K)}(p+j+1) - \phi^{(K)}(p+j)).
\end{aligned}$$

This implies

$$\begin{aligned}
&\int \left[\sum_{j=1}^{r-1} Q_K \phi^{(K)} \left(x + \frac{j+1}{\sqrt{K}} \right) - Q_K \phi^{(K)} \left(x + \frac{j}{\sqrt{K}} \right) \right]^2 dx \\
&= \sum_p \left(\sum_{j=1}^{r-1} (\phi^{(K)}(p+j+1) - \phi^{(K)}(p+j)) \right)^2 \\
&\leq r \sum_p \sum_{j=1}^{r-1} (\phi^{(K)}(p+j+1) - \phi^{(K)}(p+j))^2 \\
&= r^2 \sum_p (\phi^{(K)}(p+1) - \phi^{(K)}(p))^2 \leq \frac{C}{\zeta} \frac{r^2}{K}
\end{aligned}$$

as we have seen before. We observe that $r^2/K \leq h^2 \leq h$ since $h \leq 1$, and the result follows by taking $\tilde{C} = 2\sqrt{\frac{C}{\zeta}} + \frac{C}{\zeta}$. ■

6.2 Dirichlet form for the operator \mathcal{L}_K

We need an estimate on the decay at infinity of the eigenfunctions. Note that since the eigenvalues are real, we can assume that the eigenfunctions are real.

PROPOSITION 6.4. *If ϕ is a normalized sequence in $\text{Dom}(\mathcal{L}_K)$ decaying exponentially fast at infinity, then*

$$\begin{aligned}
&-\langle \phi, \mathcal{L}_K \phi \rangle \\
&= \sum_{n=1}^{+\infty} \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} (\phi(n+1) - \phi(n))^2 + \sum_{n=1}^{+\infty} V_n(K) \phi(n)^2 - \frac{1}{2} \sqrt{\lambda_1^{(K)} \mu_2^{(K)}} \phi(1)^2.
\end{aligned}$$

PROOF. For any fixed positive integer N we have

$$\begin{aligned}
&\sum_{n=1}^N \phi(n) (\mathcal{L}_K \phi)(n) \\
&= \sum_{n=1}^N \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} \phi(n) \phi(n+1) + \sum_{n=2}^N \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}} \phi(n) \phi(n-1)
\end{aligned}$$

$$\begin{aligned}
& - \sum_{n=1}^N (\lambda_n^{(K)} + \mu_n^{(K)}) \phi(n)^2 \\
= & - \frac{1}{2} \sum_{n=1}^N \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} (\phi(n) - \phi(n+1))^2 + \frac{1}{2} \sum_{n=1}^N \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} \phi(n)^2 \\
& + \frac{1}{2} \sum_{n=1}^N \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} \phi(n+1)^2 - \frac{1}{2} \sum_{n=2}^N \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}} (\phi(n) - \phi(n-1))^2 \\
& + \frac{1}{2} \sum_{n=2}^N \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}} \phi(n)^2 + \frac{1}{2} \sum_{n=2}^N \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}} \phi(n-1)^2 \\
& - \sum_{n=1}^N (\lambda_n^{(K)} + \mu_n^{(K)}) \phi(n)^2 \\
= & - \sum_{n=1}^{N-1} \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} (\phi(n) - \phi(n+1))^2 \\
& - \sum_{n=1}^N \left(\lambda_n^{(K)} + \mu_n^{(K)} - \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} - \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}} \mathbb{1}_{\{n>1\}} \right) \phi(n)^2 \\
& - \frac{1}{2} \sqrt{\lambda_N^{(K)} \mu_{N+1}^{(K)}} (\phi(N) - \phi(N+1))^2 + \frac{1}{2} \sqrt{\lambda_N^{(K)} \mu_{N+1}^{(K)}} \phi(N+1)^2 \\
& - \frac{1}{2} \sqrt{\lambda_1^{(K)} \mu_2^{(K)}} \phi_1^2 - \frac{1}{2} \sqrt{\lambda_N^{(K)} \mu_{N+1}^{(K)}} \phi(N)^2.
\end{aligned}$$

Since $\phi \in \text{Dom}(\mathcal{L}_K)$, the functions $\phi \mathbb{1}_{\{n \leq N\}}$ and $\mathcal{L}_K(\phi \mathbb{1}_{\{n \leq N\}})$ converge to ϕ , respectively $\mathcal{L}_K \phi$, in ℓ^2 when N tends to infinity. The result follows by letting N tend to infinity, since $V_n(K)$ is positive for n large enough, and since $\lambda_N^{(K)}$ and $\mu_N^{(K)}$ are exponential in N and ϕ decays exponentially fast by assumption. ■

LEMMA 6.5. *There exists $\xi > 0$ such that for all $K \in \mathbb{Z}_{>0}$, $\inf_{n \geq 1} V_n(K) \geq -\xi$.*

PROOF. Since $(\lambda_n^{(K)})_n$ is an increasing sequence we have

$$\begin{aligned}
V_n(K) &= \lambda_n^{(K)} + \mu_n^{(K)} - \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} - \sqrt{\lambda_{n-1}^{(K)} \mu_n^{(K)}} \mathbb{1}_{\{n>1\}} \\
&\geq \frac{\lambda_n^{(K)} + \mu_n^{(K)}}{2} - \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}}.
\end{aligned}$$

It follows from the general assumptions (see Section 2) that there exists $\tilde{x} \geq 1$ such that all $K \in \mathbb{Z}_{>0}$ and for all $n \geq K\tilde{x}$ we have

$$\frac{\lambda_n^{(K)}}{\mu_n^{(K)}} \leq \frac{1}{5} \quad \text{and} \quad \frac{\lambda_{n+1}^{(K)}}{\mu_n^{(K)}} \leq \frac{5}{4}.$$

For all $n \geq K\tilde{x}$ we have $V_n(K) \geq 0$. When $n \leq K\tilde{x}$, we write

$$\begin{aligned} & \frac{\lambda_n^{(K)} + \mu_n^{(K)}}{2} - \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} \\ &= \frac{(\sqrt{\lambda_n^{(K)}} - \sqrt{\mu_n^{(K)}})^2}{2} + \frac{\lambda_n^{(K)}}{\sqrt{\lambda_n^{(K)}} + \sqrt{\mu_{n+1}^{(K)}}} (\mu_n^{(K)} - \mu_{n+1}^{(K)}). \end{aligned}$$

Now observe that

$$\mu_n^{(K)} - \mu_{n+1}^{(K)} = K \left(d\left(\frac{n}{K}\right) - d\left(\frac{n+1}{K}\right) \right) \geq - \sup_{0 \leq x \leq \tilde{x}+1} d'(x).$$

The rest of the proof is obvious. ■

PROPOSITION 6.6. *Let $\delta > 0$ and ϕ be a real normalized sequence in $\text{Dom}(\mathcal{L}_K)$ decaying exponentially fast, such that $\|\mathcal{L}_K \phi + \rho \phi\|_{\ell^2} \leq \delta$. Then*

$$\begin{aligned} & \sum_{n=1}^{+\infty} \sqrt{\lambda_n^{(K)} \mu_{n+1}^{(K)}} (\phi(n+1) - \phi(n))^2 + \sum_{n=1}^{+\infty} (1 \vee V_n(K)) (\phi(n))^2 \\ & \leq 1 + \rho + \xi + \delta + \frac{1}{2} \sqrt{\lambda_1^{(K)} \mu_2^{(K)}}. \end{aligned}$$

The proof is left to the reader. It is a direct consequence of Proposition 6.4 and Lemma 6.5.

7 Spectral theory of \mathcal{M}_0

Recall that (see (3.5))

$$\begin{aligned} (\mathcal{M}_0 v)(n) &= \\ & \sqrt{b'(0) d'(0) n(n+1)} v(n+1) + \sqrt{b'(0) d'(0) n(n-1)} v(n-1) \mathbb{1}_{\{n>1\}} \\ & - n(b'(0) + d'(0))v(n). \end{aligned}$$

THEOREM 7.1. *The operator \mathcal{M}_0 defined on c_{00} is symmetric for the scalar product of ℓ^2 . We denote by \mathcal{M}_0 its closure which is self-adjoint and bounded above. The spectrum of \mathcal{M}_0 is discrete, all eigenvalues are simple, and we have*

$$\text{Sp}(\mathcal{M}_0) = (d'(0) - b'(0))\mathbb{Z}_{>0} = -S_2.$$

The eigenvector v_m corresponding to the eigenvalue $(d'(0) - b'(0))m$, where $m \in \mathbb{Z}_{>0}$, is given (up to a multiplicative factor) by

$$v_m(n) = \sqrt{n} \left(\frac{d'(0)}{b'(0)} \right)^{\frac{n}{2}} P_m(n) \quad (7.1)$$

where P_m is the monic orthogonal polynomial of degree $m - 1$ associated with the measure q on $\mathbb{Z}_{>0}$ defined by

$$q(n) = n \left(\frac{d'(0)}{b'(0)} \right)^n. \quad (7.2)$$

PROOF. It is easy to verify that \mathcal{M}_0 is a symmetric operator on c_{00} , which is bounded above since from $b'(0) > d'(0)$, we have

$$\inf_n \left(n(b'(0) + d'(0)) - \sqrt{b'(0)d'(0)n(n+1)} - \sqrt{b'(0)d'(0)n(n-1)} \right) > -\infty.$$

It is easy to verify that \mathcal{M}_0 is closable and we denote by \mathcal{M}_0 its closure.

Since for any $m \in \mathbb{Z}_{>0}$, the sequence $v_m(n)$ defined by (7.1) decays exponentially fast with n , it is easy to verify that $v_m \in \text{Dom}(\mathcal{M}_0)$. Note also that if $m \neq m'$, v_m is orthogonal to $v_{m'}$ in ℓ^2 .

By a direct computation one checks that $\mathcal{M}_0 v_1 = (d'(0) - b'(0))v_1$ (recall that $P_1(n) = 1$). It is left to the reader to check that

$$\mathcal{M}_0 v_m(n) = \sqrt{n} \left(\frac{b'(0)}{d'(0)} \right)^{\frac{n}{2}} Q_m(n)$$

where Q_m is a polynomial in n in which the coefficient of n^{m-1} is

$$m(d'(0) - b'(0)).$$

To check that the v_m are eigenvectors, we use a recursive argument. Assume that $m \geq 2$ and for $1 \leq k \leq m - 1$

$$\mathcal{M}_0 v_k = k(d'(0) - b'(0))v_k.$$

We can write

$$\mathcal{M}_0 v_m = m(d'(0) - b'(0))v_m + r_m$$

with

$$r_m = \sqrt{n} \left(\frac{b'(0)}{d'(0)} \right)^{\frac{n}{2}} R_m$$

where R_m is a polynomial in n of degree at most $m - 2$. Therefore

$$r_m \in \text{Span}\{v_1, \dots, v_{m-1}\}.$$

From our recursive assumption, the symmetry of \mathcal{M}_0 , and the orthogonality of the v_k (following from the orthogonality of the P_k), we get that for any $1 \leq k \leq m - 1$

$$0 = \langle v_k, \mathcal{M}_0 v_m \rangle_{\ell^2} = \langle v_k, r_m \rangle_{\ell^2}.$$

Therefore $r_m = 0$. Hence $\mathcal{M}_0 v_m = m(d'(0) - b'(0))v_m$, and we can proceed with the recursion.

We now prove that the v_m form a basis of ℓ^2 . Assume the contrary, namely there exists $u \in \ell^2$ of norm one such that for any m

$$\sum_{n=1}^{+\infty} \bar{u}(n) v_m(n) = 0.$$

We observe that the sequence

$$w(n) = \frac{1}{\sqrt{n}} \left(\frac{b'(0)}{d'(0)} \right)^{\frac{n}{2}} u(n)$$

belongs to $\ell^2(q)$, with q defined in (7.2). Therefore our assumption on u implies that w is orthogonal to all the polynomials in $\ell^2(q)$. Let us show that the set of polynomials is dense in $\ell^2(q)$. It is sufficient to prove that the measure q is the solution of a determinate moment problem, see [8, Corollary 2.50, p. 30]. Following [16, Proposition 1.5, p. 88], it is enough to prove that the moments of order m , denoted by γ_m of q , satisfy the following property: there exists $C > 0$ such that, for any $m \in \mathbb{Z}_{>0}$,

$$\gamma_m = \sum_{n=1}^{+\infty} n^{m+1} \left(\frac{d'(0)}{b'(0)} \right)^n \leq C^m m!.$$

The proof is left to the reader. Therefore the set of all polynomials is dense in $\ell^2(q)$ implying $w = 0$ and we get a contradiction with the existence of a u nonzero orthogonal to all the v_m in ℓ^2 . Therefore, the v_m 's form a basis of ℓ^2 .

We now observe that \mathcal{M}_0 is bounded above. The proof is similar to that of Proposition 6.4 and left to the reader. Since the v_m 's form a basis of ℓ^2 , for any $B > 0$ we have $\ker(\mathcal{M}_0^\dagger - B) = \{0\}$. Hence \mathcal{M}_0 is self adjoint (see for instance [14, Prop. 3.9, p. 43]) and the spectrum is given by

$$\text{Sp}(\mathcal{M}_0) = (d'(0) - b'(0))\mathbb{Z}_{>0}.$$

This ends the proof. ■

A Local maximum principle and consequences thereof

We will state and prove a maximum/minimum principle in a form which is well suited for our purposes. We start with a proposition giving elementary inequalities following from the order on the real line.

PROPOSITION A.1. *Assume $a > 0$, $c > 0$ and $b > a + c$. Let $u, w \in \mathbb{R}$.
If $v > 0$ is such that $au + cw - bv \geq 0$, then $v < \max\{u, w\}$.
If $v < 0$ is such that $au + cw - bv \leq 0$, then $v > \min\{u, w\}$.
Moreover, if $u \geq v \geq w$ are such that $au + cw - bv \geq 0$, then $v \leq \frac{a}{b-c}u$.*

PROOF. If $v > 0$ we have $0 < (b - a - c)v \leq a(u - v) + c(w - v)$ leads to a contradiction if $v \geq \max\{u, w\}$. The case $v < 0$ is similar. The last statement is trivial since $bv \leq au + cw \leq au + cv$. ■

PROPOSITION A.2. *Let $1 < n_1 < n_2$ be integers such $n_2 > n_1 + 1$. Let (α_n) be a finite sequence of strictly positive real numbers defined for $n_1 - 1, \dots, n_2$. Let (β_n) be a finite sequence of strictly positive real numbers defined for n_1, \dots, n_2 . Let (u_n) be a finite sequence of real numbers defined for $n_1 - 1, \dots, n_2 + 1$. Assume that, for all $n_1 \leq n \leq n_2$, we have $\beta_n > \alpha_n + \alpha_{n-1}$.*

If $\alpha_n u_{n+1} + \alpha_{n-1} u_{n-1} - \beta_n u_n \geq 0$, then the sequence (u_n) has no positive local maxima for $n \in \{n_1 + 1, \dots, n_2 - 1\}$. Moreover, if there exists some $u_n > 0$ then the maximum is attained only at the boundary, that is, on the set $\{n_1, n_2\}$.

If $\alpha_n u_{n+1} + \alpha_{n-1} u_{n-1} - \beta_n u_n \leq 0$, then the sequence (u_n) has no positive local minima for $n \in \{n_1 + 1, \dots, n_2 - 1\}$, and if there exists some $u_n < 0$ then the minimum is attained only at the boundary, that is, on the set $\{n_1, n_2\}$.

PROOF. It follows from Proposition A.1. ■

PROPOSITION A.3. *Let $1 < n_1 < n_2$ be integers such $n_2 > n_1 + 1$. Let (α_n) be a finite sequence of strictly positive real numbers defined for $n_1 - 1, \dots, n_2$. Let (β_n) be a finite sequence of strictly positive real numbers defined for n_1, \dots, n_2 . Let (u_n) be a finite sequence of real numbers defined for $n_1 - 1, \dots, n_2 + 1$. Assume that, for all $n_1 \leq n \leq n_2$, we have $\beta_n > \alpha_n + \alpha_{n-1}$.*

If $\alpha_n u_{n+1} + \alpha_{n-1} u_{n-1} - \beta_n u_n \geq 0$, $u_{n_1+1} > 0$ and $u_{n_1+1} \geq u_{n_1}$, then the sequence (u_n) is increasing.

If $\alpha_n u_{n+1} + \alpha_{n-1} u_{n-1} - \beta_n u_n \leq 0$, $u_{n_1+1} < 0$ and $u_{n_1+1} \leq u_{n_1}$, then the sequence (u_n) is decreasing.

Finally, if (u_n) is a positive sequence then there cannot be two local (positive) minima separated by a distance larger than one.

PROOF. It follows recursively from Proposition A.1. ■

B Quasi-eigenvalues and quasi-eigenvectors of self-adjoint operators

The following results are used in the proofs of Lemma 4.3, Propositions 5.1 and 5.2, Theorem 3.7

PROPOSITION B.1. *Let A be a self-adjoint operator in a Hilbert space \mathcal{H} with domain $\text{Dom}(A)$. Assume there exists $u \in \text{Dom}(A)$ of norm 1, $\omega \in \mathbb{R}$ and $\epsilon > 0$ such that*

$$\|Au - \omega u\| \leq \epsilon.$$

Then

$$\mathrm{Sp}(A) \cap [\omega - \epsilon, \omega + \epsilon] \neq \emptyset.$$

PROOF. We will assume that $\omega \notin \mathrm{Sp}(A)$, otherwise the result is trivial. The proof is then by contradiction. If R_ω denotes the resolvent of A at ω , we have

$$u = -R_\omega (A u - \omega u)$$

hence

$$1 \leq \epsilon \|R_\omega\|.$$

The result follows from the estimate (a direct consequence of the spectral decomposition)

$$\|R_\omega\| \leq \frac{1}{d(\omega, \mathrm{Sp}(A))}$$

where d denotes the Euclidean distance on the real line. If $\mathrm{Sp}(A) \cap [\omega - \epsilon, \omega + \epsilon] = \emptyset$, since $\mathrm{Sp}(A)$ is closed, then

$$\delta = d(\omega, \mathrm{Sp}(A)) > \epsilon$$

and we get

$$1 \leq \frac{\epsilon}{\delta} < 1$$

which is a contradiction. ■

PROPOSITION B.2. *Let A be a self-adjoint operator in a Hilbert space \mathcal{H} with domain $\mathrm{Dom}(A)$. Assume there exists $u \in \mathrm{Dom}(A)$ of norm 1, $\omega \in \mathbb{R}$ and $\epsilon > 0$ such that*

$$\|A u - \omega u\| \leq \epsilon.$$

Assume A has discrete spectrum with eigenvalues of multiplicity one, and let $\delta > 0$ denote the minimum distance between two consecutive eigenvalues. Then if $\epsilon < \delta$ there is a $\lambda \in \mathrm{Sp}(A)$ with a normalized eigenvector e such that $|\omega - \lambda| \leq \epsilon$ and

$$\|u - e\| \leq \frac{2\epsilon}{\delta - \epsilon}.$$

PROOF. We will denote by P_z the one-dimensional spectral projector of A corresponding to $z \in \mathrm{Sp}(A)$.

Since $\epsilon < \delta$, using Proposition B.1 we conclude that there is only one eigenvalue of A in $[\omega - \epsilon, \omega + \epsilon]$ and we denote by λ this eigenvalue and by \tilde{e} one of the corresponding eigenvectors (they all differ only by a phase factor). Let

$$v = \omega u - A u.$$

Since $u = P_\lambda u + (\mathrm{Id} - P_\lambda)u$, we get from the spectral decomposition

$$A(\mathrm{Id} - P_\lambda)u - \omega(\mathrm{Id} - P_\lambda)u = (\mathrm{Id} - P_\lambda)v.$$

This implies that

$$\|(\text{Id} - P_\lambda) u\| \leq \frac{\epsilon}{\delta - \epsilon}.$$

Since $P_\lambda u$ is proportional to \tilde{e} , we can write

$$u = \beta \tilde{e} + (\text{Id} - P_\lambda) u$$

with $\beta \in \mathbb{C}$. Since u and \tilde{e} are of norm one, and \tilde{e} and $(\text{Id} - P_\lambda) u$ are orthogonal, we get

$$1 = |\beta|^2 + \|(\text{Id} - P_\lambda) u\|^2$$

which implies

$$1 \geq |\beta| \geq 1 - \frac{\epsilon}{\delta - \epsilon}.$$

Let $\beta = |\beta| \exp(i\theta)$, we define $e = e^{i\theta} \tilde{e}$, and the result follows. ■

Acknowledgments: This work has been supported by the Chair “Modélisation mathématique et biodiversité” of Veolia–Ecole polytechnique–Muséum national d’Histoire naturelle–Fondation de l’Ecole polytechnique. We thank two anonymous referees for their comments which helped to improve the paper.

References

- [1] F.J. Ayala, M.E. Gilpin, J.G. Ehrenfeld. Competition between species: Theoretical models and experimental tests. *Theoretical Pop. Biol.* **4**(1973), 331–356.
- [2] D. Babusci, G. Dattoli, E. Sabia, S. Licciardi. *Mathematical Methods for Physicists*. World Scientific, 2019.
- [3] M. Bertoldi, L. Lorenzi. *Analytical Methods for Markov Semigroups*. Chapman & Hall/CRC, 2007.
- [4] F. Chatelin. *Spectral Approximation of Linear Operators*. SIAM, 2011.
- [5] J.-R. Chazottes, P. Collet, S. Méléard. Sharp asymptotics for the quasi-stationary distribution of birth-and-death processes. *Probab. Theory & Rel. Fields* **164** (2015), 285-332.
- [6] P. Collet, S. Martínez, J. San Martín. *Quasi-Stationary Distributions*. Probability and its Applications. Springer, 2013.
- [7] E. B. Davies. *Spectral theory and differentiable operators*. Cambridge Studies in Advanced Mathematics **42**. Cambridge University Press, 1995.

- [8] P. Deift. *Orthogonal Polynomials and Random Matrices: A Riemann-Hilbert Approach*. American Mathematical Society Providence, Rhode Island 2000.
- [9] H. Hanche-Olsen, H. Holden. The Kolmogorov-Riesz compactness theorem. *Expositiones Mathematicae* **28**(2010), 385-394.
- [10] M. Klein, E. Rosenberger. Harmonic approximation of difference operators J. *Funct. Anal.* **257**, (2009), 3409–3453.
- [11] S. Méléard, D. Villemonais. Quasi-stationary distributions and population processes. *Probab. Surv.* **9** (2012), 340–410.
- [12] R. M. Nisbet, W. S. C. Gurney. *Modeling fluctuating populations*. Wiley-Interscience, Chichester, 1982.
- [13] G. A. Pavliotis *Stochastic Processes and Applications: Diffusion Processes, the Fokker-Planck and Langevin Equations*. Texts in Applied Mathematics **60**. Springer, 2014.
- [14] K. Schmüdgen. *Unbounded Self-adjoint Operators on Hilbert Space*. Graduate Texts in Mathematics book series **265**, Springer, 2012.
- [15] B. Simon. Semi-classical analysis of low lying eigenvalue. I. *Ann. Inst. H. Poincaré*, section A, **38**, no 3, (1983), 295–307.
- [16] B. Simon. The classical moment problem as a self-adjoint finite difference operator. *Advances in Mathematics* **137**, issue 1, (1998), 82-203.
- [17] F. E. Smith. Population dynamics in *Daphnia magna* and a new model for population growth. *Ecology* **44**(1963), 651-663.
- [18] K. Yosida. *Functional analysis*. Reprint of the sixth (1980) edition. Classics in Mathematics. Springer-Verlag, Berlin, 1995.