



HAL
open science

Skew Cyclic and Skew Constacyclic Codes over a mixed alphabet

Karthick Gowdhaman, Cruz Mohan, Chinnapillai Durairajan, Selda Çalkavur,
Patrick Solé

► **To cite this version:**

Karthick Gowdhaman, Cruz Mohan, Chinnapillai Durairajan, Selda Çalkavur, Patrick Solé. Skew Cyclic and Skew Constacyclic Codes over a mixed alphabet. 2023. hal-04308277

HAL Id: hal-04308277

<https://hal.science/hal-04308277v1>

Preprint submitted on 27 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Skew Cyclic and Skew Constacyclic Codes over a mixed alphabet

^aKarthick Gowdhaman · ^bCruz Mohan · ^cChinnapillai
Durairajan · ^dSelda Çalkavur · ^ePatrick Solé

Received: date / Accepted: date

Abstract In this note, we study skew cyclic and skew constacyclic codes over the mixed alphabet $\mathcal{R} = \mathbb{F}_q\mathcal{R}_1\mathcal{R}_2$ where $\mathcal{R}_1 = \mathbb{F}_q + u\mathbb{F}_q$ with $u^2 = u$ and $\mathcal{R}_2 = \mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q$ with $u^2 = u, v^2 = v, uv = vu = 0$. Such codes consist of the juxtaposition of three codes of the same size over respectively $\mathbb{F}_q, \mathcal{R}_1$, and \mathcal{R}_2 .

We investigate the generator polynomial for skew cyclic codes over \mathcal{R} . Furthermore, we discuss the structural properties of the skew cyclic and skew constacyclic codes over \mathcal{R} . We also study their q -ary images under suitable Gray maps.

Keywords Linear codes · Skew-cyclic codes · Gray map · Skew Constacyclic codes

Mathematics Subject Classification (2000) 94B05 · 94B15 · 94B35 · 94B60

1 Introduction

Constacyclic codes constitute a remarkable generalization of cyclic codes and form an important class of linear codes in the coding theory. They were introduced by Berlekamp to construct the analogue of BCH codes in the Lee metric [9].

In 2007, Boucher et al. [14] was the first one to study cyclic codes using a non commutative ring over the skew polynomials ring $\mathbb{F}[x; \theta]$ where θ denotes the automorphism of the finite field \mathbb{F} and they produced many numerical examples which improved the tables of best known codes. The advantage of skew polynomials ring is that the polynomial $x^n - 1$ has more factors in skew polynomials ring than in commutative rings. Later, in [15] Boucher et al. generalised this idea to skew constacyclic codes. In 2011, Siap et al. [34] studied skew cyclic codes of arbitrary length, and established a strong connection

^aSRM Institute of Science and Technology, Ramapuram Campus, Chennai, Tamil Nadu, India.

^{a,b,c}Bharathidasan University, Tiruchirapalli, Tamil Nadu, India.

^bBishop Heber College, Affiliated to Bharathidasan University, Tiruchirapalli, Tamil Nadu, India.

^dKocaeli University, Kocaeli, Turkey

^eI2M, (CNRS, Aix-Marseille University, Centrale Marseille), Marseilles, France

^akarthygowtham@gmail.com, ^bcruzmohan@gmail.com, ^ccdurai66@rediffmail.com, ^dselda.calkavur@kocaeli.edu.tr, ^epatrick.sole@telecom-paris.fr

with classical cyclic and quasi-cyclic codes. In 2012, Abualrub et al. [1] studied θ -cyclic codes over the non-chain ring $\mathbb{F}_2 + u\mathbb{F}_2, v^2 = v$ with respect to Euclidean and Hermitian inner products, and Jitman et al. [27] studied skew constacyclic codes over finite chain rings, and gave the generators of Euclidean and Hermitian dual codes. Later, these codes over non-chain rings are extensively studied. For instance, the rings $\mathbb{F}_3 + v\mathbb{F}_3$ in [3], $\mathbb{F}_q + v\mathbb{F}_q, v^2 = v$ in [21], $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q, u^2 = u, v^2 = v, uv = vu = 0$ in [4] are considered as alphabet for skew cyclic codes. Also, Yao et al. [33] and Dertli & Cengellenmis [17] studied these codes over $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q, u^2 = u, v^2 = v, uv = vu$. In 2017, Gao et al. [20] obtained the structure of skew constacyclic codes over the non-chain ring $\mathbb{F}_q + v\mathbb{F}_q, v^2 = v$ and they obtained skew $(-1 + 2v)$ -constacyclic codes. Islam and Prakash have determined the structural properties of skew constacyclic codes over $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q, u^2 = u, v^2 = v, uv = vu$ in [24], and $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q, u^2 = u, v^2 = v, uv = vu = 0$ in [25]. In 2019, Bhardwaj and Raka [12] studied the skew constacyclic codes over the ring $\mathbb{F}_q[u, v] \langle f(u), g(v), uv - vu \rangle$ by using two non trivial automorphisms.

In another tack, Borges et al. [10] introduced $\mathbb{Z}_2\mathbb{Z}_4$ -linear codes that is to say codes over the **mixed alphabet** $\mathbb{Z}_2\mathbb{Z}_4$, where some coordinates are binary and the rest is quaternary. They have obtained their dual codes as well as their generator matrices. In continuation, Fernandez-Cordoba et al. [19] determined the rank and kernel of $\mathbb{Z}_2\mathbb{Z}_4$ -linear codes. These codes found an engineering application in the area of steganography [29]. Additive codes over the mixed alphabet $\mathbb{Z}_2\mathbb{Z}_{2^s}$ were considered in [5]. Then, the mixed alphabet $\mathbb{Z}_p\mathbb{Z}_{p^s}$, and, more generally $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ was studied in [7, 31, 35, 36]. On the other hand, to the progress of cyclic codes on mixed alphabets, in 2014, Abualrub et al. [2] defined $\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic codes as $\mathbb{Z}_4[x]$ -submodule of $\mathbb{Z}_2[x]/\langle x^r - 1 \rangle \times \mathbb{Z}_4[x]/\langle x^s - 1 \rangle$ and derived the unique set of generators, and minimal spanning set for these codes where s is an odd integer. Also, Borges et al. [11] found generator polynomials and duals for $\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic codes. After introducing the new mixed alphabets $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes, where $u^2 = 0$ in [6], Aydogdu et al. [8] were also investigated constacyclic codes over mixed alphabets by defining them as $\mathbb{Z}_2[u][x]$ -submodules of $\mathbb{Z}_2[x]/\langle x^\alpha - 1 \rangle \times \mathbb{Z}_2[u][x]/\langle x^\beta - (1 + u) \rangle$. They obtained some optimal binary linear codes as the Gray images of $\mathbb{Z}_2\mathbb{Z}_2[u]$ -cyclic codes. Meanwhile, [32] studied the algebraic properties of $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive cyclic and constacyclic codes with the unit $1 + u$, respectively. Therefore, in continuation of these studies, the expected generalization should be $\mathbb{Z}_{2^r}\mathbb{Z}_{2^s}[u]$ -additive cyclic and constacyclic codes, where $u^2 = 0$.

Motivated by the two kinds of above studies, here we consider a mixed alphabet $\mathcal{R} = \mathbb{F}_q\mathcal{R}_1\mathcal{R}_2$ where $\mathcal{R}_1 = \mathbb{F}_q + u\mathbb{F}_q$ with $u^2 = u$ and $\mathcal{R}_2 = \mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q$ with $u^2 = u, v^2 = v, uv = vu = 0$ and study θ_t -cyclic and (θ_t, α) -cyclic codes over \mathcal{R} . We completely determine the algebraic structure of these codes. We study their q -ary images under Gray maps, and give some numerical examples in short lengths.

The material is organized as follows. The next section collects some background material. Section 3 studies Gray maps. Section 4 is dedicated to skew cyclic codes, and Section 5 to skew constacyclic codes. Section 6 concludes the article.

2 Preliminaries

Let \mathbb{F}_q denote the finite field of characteristic p with q elements, where $q = p^m$ for odd prime p and positive integer m . The set \mathbb{F}_q^n of all ordered n -tuples over \mathbb{F}_q forms a vector space with the usual component-wise addition and scalar multiplication of vectors. A non-empty subset \mathcal{C} of \mathbb{F}_q^n is called a code of length n over \mathbb{F}_q and it is called a linear code if \mathcal{C} is a subspace of \mathbb{F}_q^n . From now onward, we denote $\mathcal{R}_1 = \mathbb{F}_q + u\mathbb{F}_q$, with $u^2 = u$ and $\mathcal{R}_2 = \mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q$, with $u^2 = u, v^2 = v, uv = vu = 0$, where $q = p^m$ for odd prime p and positive integer m .

Note that \mathcal{R}_1 and \mathcal{R}_2 are finite non-chain rings. Let $a + ub + vc$ be an element of \mathcal{R}_2 . Then we define two

maps η and δ as follows:

$$\begin{aligned}\eta : \mathcal{R}_2 &\rightarrow \mathbb{F}_q, \quad \delta : \mathcal{R}_2 \rightarrow \mathcal{R}_1, \\ \eta(a + ub + vc) &= a, \quad \delta(a + ub + vc) = a + ub,\end{aligned}$$

It is clear that η and δ are ring homomorphisms. We consider the ring \mathcal{R} ,

$$\mathcal{R} = \mathbb{F}_q \mathcal{R}_1 \mathcal{R}_2 = \{(x, y, z) \mid x \in \mathbb{F}_q, y \in \mathcal{R}_1 \text{ and } z \in \mathcal{R}_2\}$$

We define a \mathcal{R}_2 -multiplication in this ring as follows,

$$\begin{aligned}\star : \mathcal{R}_2 \times \mathcal{R} &\rightarrow \mathcal{R} \\ r \star (x, y, z) &= (\eta(r)x, \delta(r)y, rz)\end{aligned}$$

This is a well defined multiplication and it can be extended componentwise to $\mathcal{R}_\gamma = \mathbb{F}_q^{n_1} \times \mathcal{R}_1^{n_2} \times \mathcal{R}_2^{n_3}$ by

$$\begin{aligned}\star : \mathcal{R}_2 \times \mathcal{R}_\gamma &\rightarrow \mathcal{R}_\gamma \\ r \star (x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}, z_1, \dots, z_{n_3}) &= (\eta(r)x_1, \dots, \eta(r)x_{n_1}, \delta(r)y_1, \dots, \delta(r)y_{n_2}, rz_1, \dots, rz_{n_3})\end{aligned}$$

where $(x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}, z_1, \dots, z_{n_3}) \in \mathcal{R}_\gamma$. By this multiplication, \mathcal{R}_γ is a \mathcal{R}_2 module. A non-empty subset \mathcal{C} of \mathcal{R}_γ is said to be a \mathcal{R} -linear code of length (n_1, n_2, n_3) if \mathcal{C} is an \mathcal{R}_2 -submodule of \mathcal{R}_γ . Now we define the inner product as follows,

$$\langle c, c' \rangle = \sum_1^{n_1} x_i x'_i + \sum_1^{n_2} y_j y'_j + \sum_1^{n_3} z_k z'_k$$

where $c = (x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}, z_1, \dots, z_{n_3})$, $c' = (x'_1, \dots, x'_{n_1}, y'_1, \dots, y'_{n_2}, z'_1, \dots, z'_{n_3})$ are in \mathcal{R}_γ . Let \mathcal{C} be a \mathcal{R} -linear code of length (n_1, n_2, n_3) . Then the dual code of \mathcal{C} is defined as,

$$\mathcal{C}^\perp = \{c' \in \mathcal{R}_\gamma \mid \langle c, c' \rangle = 0 \forall c \in \mathcal{C}\}$$

3 Decomposition and Properties of Gray Maps

Recall that, $\mathcal{R}_1 = \mathbb{F}_q + u\mathbb{F}_q$, with $u^2 = u$. Consider the idempotent orthogonal elements $e_1 = u$ and $e_2 = 1 - u$, then

$$\mathcal{R}_1 = e_1 \mathcal{R}_1 \oplus e_2 \mathcal{R}_1 \cong e_1 \mathbb{F}_q \oplus e_2 \mathbb{F}_q$$

$\because e_1 e_2 = 0, e_1^2 = e_1, e_1 + e_2 = 1$. Hence, $\mathcal{R}_1 = \{ae_1 + be_2 \mid a, b \text{ in } \mathbb{F}_q\}$. We now define the Gray map,

$$\begin{aligned}\varphi_1 : \mathcal{R}_1 &\rightarrow \mathbb{F}_q^2 \\ \varphi_1(ae_1 + be_2) &= (a, b)\end{aligned}$$

It can be extended to the length n by

$$\begin{aligned}\varphi_1 : \mathcal{R}_1^n &\rightarrow \mathbb{F}_q^{2n} \\ \varphi_1((a_1, \dots, a_n)e_1 + (b_1, \dots, b_n)e_2) &= (a_1, \dots, a_n, b_1, \dots, b_n)\end{aligned}$$

Note that it is a linear map. We define the Gray weight of a codeword in \mathcal{R}_1 as

$$wt_G(ae_1 + be_2) = wt_H(a, b)$$

where wt_H denotes the Hamming weight. If x, y lie in \mathcal{R}_1^n , then their mutual distance is given by

$$d_G(x, y) = \sum_1^n wt_G(x_i - y_i) = \sum_1^{2n} wt_H(\varphi_1(x) - \varphi_1(y)) = d_H(\varphi_1(x), \varphi_1(y)).$$

Hence φ_1 is a weight preserving map. A non-empty subset \mathcal{C} of \mathcal{R}_1^n is said to be a linear code of length n if \mathcal{C} is \mathcal{R}_1 submodule of \mathcal{R}_1^n .

For $i \in \{1, 2\}$, $A_i \subseteq \mathcal{R}_1$

$$A_1 \oplus A_2 = \{a_1 + a_2 \mid a_i \in A_i\} \text{ and } A_1 \otimes A_2 = \{(a_1, a_2) \mid a_i \in A_i\}.$$

Let \mathcal{C}_e be a linear code of length n over \mathcal{R}_1 . Then we define,

$$\begin{aligned} \mathcal{C}_{e_1} &= \{y_1 \in \mathbb{F}_q^n \mid e_1 y_1 + e_2 y_2 \in \mathcal{C}_e, \text{ for some } y_2 \in \mathbb{F}_q^n\} \\ \mathcal{C}_{e_2} &= \{y_2 \in \mathbb{F}_q^n \mid e_1 y_1 + e_2 y_2 \in \mathcal{C}_e, \text{ for some } y_1 \in \mathbb{F}_q^n\} \end{aligned}$$

Therefore any linear code \mathcal{C}_e over \mathcal{R}_1 can be represented as $\mathcal{C}_e = e_1 \mathcal{C}_{e_1} \oplus e_2 \mathcal{C}_{e_2}$ and $\varphi_1(\mathcal{C}_e) = \mathcal{C}_{e_1} \otimes \mathcal{C}_{e_2}$. Hence \mathcal{C}_{e_1} and \mathcal{C}_{e_2} are \mathbb{F}_q -linear codes. Also note that $\varphi_1(\mathcal{C}_e^\perp) = \varphi_1(\mathcal{C}_e)^\perp$.

Recall that, $\mathcal{R}_2 = \mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q$, with $u^2 = u, v^2 = v, uv = vu = 0$. Let $o_1 = (1 - u - v), o_2 = u, o_3 = v$ be idempotent orthogonal elements in \mathcal{R}_2 , then

$$\mathcal{R}_2 = o_1 \mathcal{R}_2 \oplus o_2 \mathcal{R}_2 \oplus o_3 \mathcal{R}_2 \cong o_1 \mathbb{F}_q \oplus o_2 \mathbb{F}_q \oplus o_3 \mathbb{F}_q$$

$\because o_i o_j = 0 (i \neq j), o_i^2 = o_i, o_1 + o_2 + o_3 = 1$. Hence any element in \mathcal{R}_2 can be written as $ao_1 + bo_2 + o_3c_3$. We now define a weight preserving linear Gray map φ_2 ,

$$\begin{aligned} \varphi_2 : \mathcal{R}_2 &\rightarrow \mathbb{F}_q^3 \\ \varphi_2(ao_1 + bo_2 + co_3) &= (a, b, c) \end{aligned}$$

It can be extended to length n by the formula

$$\varphi_2((a_1, \dots, a_n)o_1 + (b_1, \dots, b_n)o_2 + (c_1, \dots, c_n)o_3) = (a_1, \dots, a_n, b_1, \dots, b_n, c_1, \dots, c_n).$$

We define the Gray weight of a codeword in \mathcal{R}_2 as

$$wt_G(ao_1 + bo_2 + co_3) = wt_H(a, b, c)$$

where wt_H denotes the Hamming weight. If x, y are in \mathcal{R}_2^n , then their Gray distance is given by

$$d_G(x, y) = \sum_1^n wt_G(x_i - y_i) = \sum_1^{3n} wt_H(\varphi_2(x) - \varphi_2(y)) = d_H(\varphi_2(x), \varphi_2(y)).$$

For $i \in \{1, 2, 3\}$, $A_i \subseteq \mathcal{R}_2$

$$A_1 \oplus A_2 \oplus A_3 = \{a_1 + a_2 + a_3 \mid a_i \in A_i\} \text{ and } A_1 \otimes A_2 \otimes A_3 = \{(a_1, a_2, a_3) \mid a_i \in A_i\}.$$

Let \mathcal{C}_o be a linear code of length n over \mathcal{R}_2 . We define,

$$\begin{aligned} \mathcal{C}_{o_1} &= \{z_1 \in \mathbb{F}_q^n \mid o_1 z_1 + o_2 z_2 + o_3 z_3 \in \mathcal{C}_o, \text{ for some } z_2, z_3 \in \mathbb{F}_q^n\} \\ \mathcal{C}_{o_2} &= \{z_2 \in \mathbb{F}_q^n \mid o_1 z_1 + o_2 z_2 + o_3 z_3 \in \mathcal{C}_o, \text{ for some } z_1, z_3 \in \mathbb{F}_q^n\} \\ \mathcal{C}_{o_3} &= \{z_3 \in \mathbb{F}_q^n \mid o_1 z_1 + o_2 z_2 + o_3 z_3 \in \mathcal{C}_o, \text{ for some } z_1, z_2 \in \mathbb{F}_q^n\} \end{aligned}$$

Then any linear code \mathcal{C}_o over \mathcal{R}_2 can be represented as $\mathcal{C}_o = o_1\mathcal{C}_{o_1} \oplus o_2\mathcal{C}_{o_2} \oplus o_3\mathcal{C}_{o_3}$ and $\varphi_2(\mathcal{C}_o) = \mathcal{C}_{o_1} \otimes \mathcal{C}_{o_2} \otimes \mathcal{C}_{o_3}$, where $\mathcal{C}_{o_1}, \mathcal{C}_{o_2}$ and \mathcal{C}_{o_3} are \mathbb{F}_q -linear codes. Also note that $\varphi_2(\mathcal{C}_o^\perp) = \varphi_2(\mathcal{C}_o)^\perp$. Henceforth, we define the Gray map φ on \mathcal{R} using the maps defined previously,

$$\begin{aligned} \varphi : \mathcal{R} &\rightarrow \mathbb{F}_q^6 \\ \varphi(x, y, z) &= (x, \varphi_1(y), \varphi_2(z)) \end{aligned}$$

now we can extend this map to \mathcal{R}_γ ,

$$\varphi(x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}, z_1, \dots, z_{n_3}) = (x_1, \dots, x_{n_1}, \varphi_1(y_1), \dots, \varphi_1(y_{n_2}), \varphi_2(z_1), \dots, \varphi_2(z_{n_3}))$$

then the Gray weight of an element in \mathcal{R}_γ can be denoted by $wt_G(\alpha) = wt_H(\varphi(\alpha))$. Any linear code \mathcal{C} of \mathcal{R}_γ can be represented by $\mathcal{C} = \mathcal{C}_1 \otimes \mathcal{C}_e \otimes \mathcal{C}_o$, where $\mathcal{C}_1, \mathcal{C}_e$ and \mathcal{C}_o are linear code over $\mathbb{F}_q, \mathcal{R}_1$ and \mathcal{R}_2 . Let $G_{\mathbb{F}_q}$ be the generator matrix for linear code over \mathbb{F}_q . The generator matrix $G_{\mathcal{R}_1}$ for a linear code over \mathcal{R}_1 is denoted by

$$G_{\mathcal{R}_1} = \begin{bmatrix} e_1 G_{e_1} \\ e_2 G_{e_2} \end{bmatrix}$$

where G_{e_i} is the generator matrix for the linear code \mathcal{C}_{e_i} , for $i = \{1, 2\}$. The generator matrix $G_{\mathcal{R}_2}$ for the linear code over \mathcal{R}_2 is

$$G_{\mathcal{R}_2} = \begin{bmatrix} o_1 G_{o_1} \\ o_2 G_{o_2} \\ o_3 G_{o_3} \end{bmatrix}$$

where G_{o_i} is the generator matrix for the linear code \mathcal{C}_{o_i} , for $i = \{1, 2, 3\}$. Using the generator matrices above, we can say that the generator matrix G for the linear code over \mathcal{R} is,

$$G = \begin{bmatrix} G_{\mathbb{F}_q} & 0 & 0 \\ 0 & G_{\mathcal{R}_1} & 0 \\ 0 & 0 & G_{\mathcal{R}_2} \end{bmatrix}$$

Note that the minimum distance of \mathcal{C} is $\min\{d_H(\mathcal{C}_1), d_H(\varphi_1(\mathcal{C}_e)), d_H(\varphi_2(\mathcal{C}_o))\}$. The following Theorem provides the weight preserving nature of the Gray map,

Theorem 1 *The Gray map φ defined above is linear and weight preserving.*

Proof Let $x = (x_1, x_2, x_3), x' = (x'_1, x'_2, x'_3)$ be in \mathcal{R}_γ where $x_1, x'_1 \in \mathbb{F}_q^{n_1}, x_2, x'_2 \in \mathcal{R}_1^{n_2}, x_3, x'_3 \in \mathcal{R}_2^{n_3}$

$$\begin{aligned} \varphi(x + x') &= \varphi(x_1 + x'_1, x_2 + x'_2, x_3 + x'_3) \\ &= (x_1 + x'_1, \varphi_1(x_2 + x'_2), \varphi_2(x_3 + x'_3)) \\ &= (x_1, \varphi_1(x_2), \varphi_2(x_3)) + (x'_1, \varphi_1(x'_2), \varphi_2(x'_3)) (\because \varphi_1 \text{ and } \varphi_2 \text{ are linear}) \\ &= \varphi(x) + \varphi(x') \end{aligned}$$

Using the linear map φ ,

$$d_G(x, x') = wt_G(x - x') = wt_H(\varphi(x) - \varphi(x')) = d_H(\varphi(x), \varphi(x'))$$

Hence φ is a weight preserving linear map.

The following theorem gives the parameters of the Gray image of a linear code.

Theorem 2 Let $\mathcal{C} \subseteq \mathcal{R}_\gamma$ be $(n_1 + n_2 + n_3, d_G)$ linear code then $\varphi(\mathcal{C})$ is $(n_1 + 2(n_2) + 3(n_3), d_H)$ linear code over \mathbb{F}_q , where $d_G = d_H$.

Proof The proof can be extended from the proof of Theorem 1.

The following Theorem characterizes $\varphi(\mathcal{C})$,

Theorem 3 Let $\mathcal{C} \subseteq \mathcal{R}_\gamma$ be linear, then $\varphi(\mathcal{C}) = \mathcal{C}_1 \otimes_{i=1}^{i=2} \mathcal{C}_{e_i} \otimes_{j=1}^{j=3} \mathcal{C}_{o_j}$, $|\mathcal{C}| = |\mathcal{C}_1| \prod_{i=1}^{i=2} |\mathcal{C}_{e_i}| \prod_{j=1}^{j=3} |\mathcal{C}_{o_j}|$.

Proof Let

$$\varphi(x, y, z) = (x, \varphi_1(y), \varphi_3(z)) = (a_1, \dots, a_6) \in \varphi(\mathcal{C}) \subseteq \mathbb{F}_q^6.$$

Note that φ is bijective and $\mathcal{C} = \mathcal{C}_1 \otimes \mathcal{C}_e \otimes \mathcal{C}_o$ is linear. So $a_1 = x \in \mathcal{C}_1$. Also note that $\varphi_1(y) = (a_2, a_3)$, $\varphi_2(z) = (a_4, a_5, a_6)$. Since φ_i 's are bijective, $a_2e_1 + a_3e_2 \in e_1\mathcal{C}_{e_1} \oplus e_2\mathcal{C}_{e_2} = \mathcal{C}_e$. Hence $(a_2, a_3) \in \mathcal{C}_{e_1} \otimes \mathcal{C}_{e_2}$ and similarly $(a_4, a_5, a_6) \in \mathcal{C}_{o_1} \otimes \mathcal{C}_{o_2} \otimes \mathcal{C}_{o_3}$. The converse holds in a similar way. The second part of the statement follows from the fact that φ is bijective.

The following Theorem furnishes the decomposition of the dual of the linear code \mathcal{C} .

Theorem 4 Let $\mathcal{C} = \mathcal{C}_1 \otimes \mathcal{C}_e \otimes \mathcal{C}_o$ be a linear code over \mathcal{R} then $\mathcal{C}^\perp = \mathcal{C}_1^\perp \otimes \mathcal{C}_e^\perp \otimes \mathcal{C}_o^\perp$ where $\mathcal{C}_1^\perp, \mathcal{C}_e^\perp$ and \mathcal{C}_o^\perp are duals for the respective linear codes.

Proof Let $\mathcal{C}^\perp = \{c' \in \mathcal{R}_\gamma \mid \langle c, c' \rangle = 0 \text{ for all } c \in \mathcal{C}\} = \{(x', y', z') \in \mathcal{R}_\gamma \mid x' \in \mathbb{F}_q^{n_1}, y' \in \mathcal{R}_1^{n_2}, z' \in \mathcal{R}_2^{n_3}\}$. Let $c = (x, y, z) \in \mathcal{C} = \mathcal{C}_1 \otimes \mathcal{C}_e \otimes \mathcal{C}_o$. Then

$$\langle c, c' \rangle = xx' + yy' + zz' = 0.$$

So $x' \in \mathcal{C}_1^\perp, y' \in \mathcal{C}_e^\perp, z' \in \mathcal{C}_o^\perp$ and so $\mathcal{C}^\perp \subseteq \mathcal{C}_1^\perp \otimes \mathcal{C}_e^\perp \otimes \mathcal{C}_o^\perp$. Since $|\mathcal{C}^\perp| = |\mathcal{C}_1^\perp| |\mathcal{C}_e^\perp| |\mathcal{C}_o^\perp|$, the statement holds.

The next result shows that the Gray maps is compatible with duality.

Theorem 5 If $\mathcal{C} \subseteq \mathcal{R}_\gamma$ is linear, then $\varphi(\mathcal{C}^\perp) = \varphi(\mathcal{C})^\perp$.

Proof Let $(x, y, z) \in \mathcal{C}$ and $(x', y', z') \in \mathcal{C}^\perp$ where $x \in \mathcal{C}_1, y \in \mathcal{C}_e, z \in \mathcal{C}_o$ and $x' \in \mathcal{C}_1^\perp, y' \in \mathcal{C}_e^\perp, z' \in \mathcal{C}_o^\perp$, then $\langle (x|y|z, x'|y'|z') \rangle = 0$. Using Theorem 4, $\mathcal{C}_1^\perp, \mathcal{C}_e^\perp$ and \mathcal{C}_o^\perp are duals for $\mathcal{C}_1, \mathcal{C}_e$ and \mathcal{C}_o . Now, we have $\varphi(x, y, z) = (x, \varphi_1(y), \varphi_2(z)), \varphi(x', y', z') = (x', \varphi_1(y'), \varphi_2(z'))$, then the inner product is given by

$$\begin{aligned} \langle \varphi(x, y, z), \varphi(x', y', z') \rangle &= \langle (x, \varphi_1(y), \varphi_2(z)), (x', \varphi_1(y'), \varphi_2(z')) \rangle \\ &= \langle (x, 0, 0), (x', 0, 0) \rangle + \langle (0, \varphi_1(y), 0), (0, \varphi_1(y'), 0) \rangle \\ &\quad + \langle (0, 0, \varphi_2(z)), (0, 0, \varphi_2(z')) \rangle \\ &= 0 \quad (\because \varphi_1(\mathcal{C}^\perp) = \varphi_1(\mathcal{C})^\perp, \varphi_2(\mathcal{C}^\perp) = \varphi_2(\mathcal{C})^\perp) \end{aligned}$$

So, $\varphi(\mathcal{C}^\perp) \subseteq \varphi(\mathcal{C})^\perp$. Since the cardinality is same, the statement holds.

The following result provides the self duality nature of the linear code and its Gray image,

Corollary 1 Let $\mathcal{C} \subseteq \mathcal{R}_\gamma$ be linear, then \mathcal{C} is self-dual iff $\varphi(\mathcal{C})$ is self-dual. Further $\varphi(\mathcal{C})$ is a self-orthogonal over \mathbb{F}_q iff \mathcal{C} is self-orthogonal.

Proof Let \mathcal{C} be a self-dual linear code of length n over \mathcal{R} . That is $\mathcal{C} = \mathcal{C}^\perp$. Then $\varphi(\mathcal{C}) = \varphi(\mathcal{C}^\perp)$, and hence by Theorem 5, we have $\varphi(\mathcal{C}) = (\varphi(\mathcal{C}))^\perp$. Thus $\varphi(\mathcal{C})$ is a self-dual linear code of length $n_1 + 2n_2 + 3n_3$ over \mathbb{F}_q . Conversely, let $\varphi(\mathcal{C})$ be a self-dual linear code of length $n_1 + 2n_2 + 3n_3$ over \mathbb{F}_q . Then $\varphi(\mathcal{C}) = (\varphi(\mathcal{C}))^\perp$, and hence by Theorem 5, we have $\varphi(\mathcal{C}) = \varphi(\mathcal{C}^\perp)$. Since φ is bijection, $\mathcal{C} = \mathcal{C}^\perp$. Therefore, \mathcal{C} is a self-dual linear code over \mathcal{R}_γ . Similarly the self-orthogonal case holds.

4 Skew Cyclic \mathcal{R} -codes

Let θ_t be a non-trivial Frobenius automorphism defined by

$$\theta_t : \mathbb{F}_q \rightarrow \mathbb{F}_q, \theta_t(a) = a^{p^t},$$

where t divides m . It can be extended to \mathcal{R}_1 and \mathcal{R}_2 by

$$\theta_t(a + ub) = \theta_t(a) + u\theta_t(b), \theta_t(a + ub + vc) = \theta_t(a) + u\theta_t(b) + v\theta_t(c).$$

Since $t|m$, the order of automorphism θ_t is $\frac{m}{t}$. We define a polynomial ring $\mathcal{R}_i[x, \theta_t]$ ($1 \leq i \leq 2$) as follows,

$$\mathcal{R}_i[x, \theta_t] = \{a_1 + \cdots + a_n x^n \mid a_j \in \mathcal{R}_i, 1 \leq j \leq n\}$$

Clearly $\mathcal{R}_i[x, \theta_t]$ is a ring with respect to usual addition and the multiplication defined by

$$ax^m bx^n = a\theta_t^m(a)x^{m+n}$$

Note that it is a non commutative ring unless θ_t is an identity map. A non empty set \mathcal{C} is said to be linear code of length n_i over \mathcal{R}_i if it is a \mathcal{R}_i submodule of $\mathcal{R}_i^{n_i}$. Using the above polynomial rings above, we extend the polynomial ring to \mathcal{R} by

$$\mathcal{R}[x, \theta_t] = \{(a(x), b(x), c(x)) : a(x) \in \mathbb{F}_q[x], b(x) \in \mathcal{R}_1[x], c(x) \in \mathcal{R}_2[x]\}.$$

It can be seen that $\mathcal{R}[x, \theta_t]$ is a $\mathcal{R}_2[x; \theta_t]$ submodule with respect to usual addition and multiplication defined by

$$\begin{aligned} \star : \mathcal{R}_2[x] \times \mathcal{R}[x, \theta_t] &\rightarrow \mathcal{R}[x, \theta_t] \\ (ax^s) \star (b_1 x^i, b_2 x^j, b_3 x^k) &= (\eta(a)x^s b_1 x^i, \delta(a)x^s b_2 x^j, ax^s b_3 x^k) \\ &= (\eta(a)\theta_t^s(b_1)x^{s+i}, \delta(a)\theta_t^s(b_2)x^{s+j}, a\theta_t^s(b_3)x^{s+k}) \end{aligned}$$

However under associative and distributive laws, the multiplication can be extended to $\mathcal{R}_\gamma[x; \theta_t] = \frac{\mathcal{F}_q[x; \theta_t]}{\langle x^{n_1} - 1 \rangle} \times \frac{\mathcal{R}_1[x; \theta_t]}{\langle x^{n_2} - 1 \rangle} \times \frac{\mathcal{R}_2[x; \theta_t]}{\langle x^{n_3} - 1 \rangle}$ as follows,

$$\star : \mathcal{R}_2[x; \theta_t] \times \mathcal{R}_\gamma[x; \theta_t] \rightarrow \mathcal{R}_\gamma[x; \theta_t]$$

$$r(x) \star (f_1(x) + \langle x^{n_1} - 1 \rangle, f_2(x) + \langle x^{n_2} - 1 \rangle, f_3(x) + \langle x^{n_3} - 1 \rangle) = (\eta(r(x))f_1(x) + \langle x^{n_1} - 1 \rangle, \delta(r(x))f_2(x) + \langle x^{n_2} - 1 \rangle, r(x)f_3(x) + \langle x^{n_3} - 1 \rangle)$$

Definition 1 [14] A non-trivial \mathcal{R} -submodule \mathcal{C} of \mathcal{R}^n is called a θ_t -cyclic code if for any $c = (c_0, c_1, \dots, c_{n-1}) \in \mathcal{C}$, $\sigma_1(c) = (\theta_t(c_{n-1}), \theta_t(c_0), \dots, \theta_t(c_{n-2})) \in \mathcal{C}$. The operator σ_1 is called a θ_t -cyclic shift operator on \mathcal{R}^n .

Definition 2 A non-trivial \mathcal{R}_2 -submodule \mathcal{C} of \mathcal{R}_γ is called a θ_t -cyclic code if for any $c = (c_{0,1}, c_{1,1}, \dots, c_{n_1-1,1}, c_{0,e}, c_{1,e}, \dots, c_{n_2-1,e}, c_{0,o}, c_{1,o}, \dots, c_{n_3-1,o}) \in \mathcal{C}$, $\sigma(c) = (\theta(c_{n_1-1,1}), \theta(c_{0,1}), \dots, \theta(c_{n_1-2,1}), \theta(c_{n_2-1,e}), \theta(c_{0,e}), \dots, \theta(c_{n_2-2,e}), \theta(c_{n_3-1,o}), \theta(c_{0,o}), \dots, \theta(c_{n_3-2,o})) \in \mathcal{C}$. The operator σ is called as θ_t -cyclic shift operator on \mathcal{R}^n .

The following result yields the relationship between the θ_t -cyclic codes over \mathcal{R} and \mathbb{F}_q ,

Theorem 6 Let $\mathcal{C} = \mathcal{C}_1 \otimes \mathcal{C}_e \otimes \mathcal{C}_o \subseteq \mathcal{R}_\gamma$ be linear. Then \mathcal{C} is a θ_t -cyclic code if and only if $\mathcal{C}_1, \mathcal{C}_e$ and \mathcal{C}_o are θ_t -cyclic codes of length n_1, n_2 and n_3 over $\mathbb{F}_q, \mathcal{R}_1$ and \mathcal{R}_2 respectively.

Proof Let $C = C_1 \otimes C_e \otimes C_o$ be a θ_t -cyclic code over \mathcal{R} . Let $z = (z_1, z_e, z_o) \in C$ that is

$$z = (z_{0,1}, z_{1,1}, \dots, z_{n_1-1,1}, z_{0,e}, z_{1,e}, \dots, z_{n_2-1,e}, z_{0,o}, z_{1,o}, \dots, z_{n_3-1,o}) \in C.$$

Then $\sigma(z) = (\theta(z_{n_1-1,1}), \theta(z_{0,1}), \dots, \theta(z_{n_1-2,1}), \theta(z_{n_2-1,e}), \theta(z_{0,e}), \dots, \theta(z_{n_2-2,e}), \theta(z_{n_3-1,o}), \theta(z_{0,o}), \dots, \theta(z_{n_3-2,o})) = (\sigma(z_1), \sigma(z_e), \sigma(z_o)) \in C$. From this we can conclude that,

$$\sigma(z_1) \in C_1, \sigma(z_e) \in C_e \text{ and } \sigma(z_o) \in C_o$$

Hence C_1, C_e and C_o are θ_t -cyclic code of length n_i . The converse holds in a similar way.

We recall the following Theorem from [4].

Theorem 7 [4] *Let $C_o = o_1C_{o_1} \oplus o_2C_{o_2} \oplus o_3C_{o_3}$ be a linear code over \mathcal{R}_2 of length n_3 then C_o is θ_t -cyclic code iff $C_{o_i} (1 \leq i \leq 3)$ is a θ_t -cyclic code of length n_3 over \mathbb{F}_q .*

The analogue of this result in our setting is as follows.

Theorem 8 [21] *Let $C_e = e_1C_{e_1} \oplus e_2C_{e_2}$ be a linear code over \mathcal{R}_1 of length n_2 then C_e is θ_t -cyclic code iff $C_{e_i} (1 \leq i \leq 2)$ is a θ_t -cyclic code of length n_2 over \mathbb{F}_q .*

Theorem 9 *$C = C_1 \otimes C_e \otimes C_o$ be a linear code of length $\gamma = n_1 + n_2 + n_3$, then C is θ_t -cyclic iff $C_1, C_{e_i}, C_{o_j} (1 \leq i \leq 2, 1 \leq j \leq 3)$ are θ_t -cyclic code of length n_1, n_2, n_3 over \mathbb{F}_q respectively.*

Proof We obtain the proof on combining proofs of Theorem 6,7 and 8.

These notions are well-behaved with respect to duality as the next result shows.

Theorem 10 *If C is a θ_t -cyclic code of length n , then its dual C^\perp is also a θ_t -cyclic code.*

Proof From Theorem 9, $C_1, C_{e_i}, C_{o_j} (1 \leq i \leq 2, 1 \leq j \leq 3)$ are θ_t -cyclic codes over \mathbb{F}_q . Then $C_1^\perp, C_{e_i}^\perp, C_{o_j}^\perp (1 \leq i \leq 2, 1 \leq j \leq 3)$ are θ_t -cyclic codes over \mathbb{F}_q from [16] and once again by using Theorem 9, C^\perp becomes a θ_t -cyclic code.

Recall the following result from [34].

Lemma 1 [34] *Let C be a θ_t -cyclic code of length n over \mathbb{F}_q . Then there exists a polynomial $f(x) \in \mathbb{F}_q[x; \theta_t]$ such that $C = \langle f(x) \rangle$ and $x^n - 1 = g(x)f(x)$ in $\mathbb{F}_q[x; \theta_t]$.*

By assuming $o(\theta_t)|n$, the counterpart follows;

Theorem 11 *Let $C = C_1 \otimes C_e \otimes C_o$ be a θ_t -cyclic code of length n over \mathcal{R} and order of θ_t divides n . Then $C = \langle B_1, B_e, B_o \rangle$ where $B_1 = \langle (f_1(x), 0, 0) \rangle, B_e = \langle (0, f_e(x), 0) \rangle$, and $B_o = \langle (b_1(x), b_e(x), f_o(x)) \rangle$, such that $C_1 = \langle f_1(x) \rangle, C_e = \langle f_e(x) \rangle, C_o = \langle f_o(x) \rangle, b_1(x) \in C_1$ and $b_2(x) \in C_e$.*

Proof Let $C = C_1 \otimes C_e \otimes C_o$ be a θ_t -cyclic code of length $\gamma = n_1 + n_2 + n_3$ over \mathcal{R} . Then by Theorem 6, C_1, C_e, C_o are θ_t -cyclic codes of length n_i over $\mathbb{F}_q, \mathcal{R}_1$ and \mathcal{R}_2 . Define a homomorphism from C to \mathcal{R} as follows,

$$\begin{aligned} \psi : C &\mapsto \mathcal{R} \\ \psi(c_1(x), c_e(x), c_o(x)) &= (0, 0, c_o(x)) \end{aligned}$$

Define

$$\ker(\psi) = \{(c_1(x), c_e(x), 0) : c_1(x) \in \mathcal{C}_1, c_e(x) \in \mathcal{C}_e\}$$

$$\mathcal{I} = \{(c_1(x), c_e(x)) \in \mathbb{F}_q[x; \theta_t] \times \mathcal{R}_1[x; \theta_t] : (c_1(x), c_e(x), 0) \in \ker(\psi)\}.$$

Clearly $\mathcal{I} = \mathcal{I}_1 \times \mathcal{I}_e$ forms a submodule of $\mathbb{F}_q[x; \theta_t] \times \mathcal{R}_1[x; \theta_t]$. Therefore there exist a polynomial $f_1(x)$ and $f_e(x)$ in $\mathbb{F}_q[x; \theta_t]$ and $\mathcal{R}_1[x; \theta_t]$ respectively, generating \mathcal{I}_1 and \mathcal{I}_e with $f_1(x)|x^{n_1} - 1$ and $f_e(x)|x^{n_e} - 1$. Thus $\mathcal{I} = \langle (f_1(x), 0), (0, f_e(x)) \rangle$, then for any $(c_1(x), c_e(x), 0) \in \ker(\psi)$, $(c_1(x), c_e(x)) = v(x) \star (f_1(x), 0), (0, f_e(x))$ for some $v(x) \in \mathcal{R}_1[x; \theta_t]$. Finally it leads to $\ker(\psi) = \langle (f_1(x), 0, 0), (0, f_e(x), 0) \rangle$. The fact that \mathcal{C} is a submodule implies that $\psi(\mathcal{C})$ is a submodule. By using the first isomorphism theorem,

$$\mathcal{C}/\ker(\psi) \cong \psi(\mathcal{C}).$$

Let $(b_1(x), b_e(x), f_o(x)) \in \mathcal{C}$ then $\psi(b_1(x), b_e(x), f_o(x)) = (0, 0, f_o(x))$. From this any θ_t -cyclic code of length n can be represented by $\mathcal{C} = \langle (f_1(x), 0, 0), (0, f_e(x), 0), (b_1(x), b_e(x), f_o(x)) \rangle$ where $f_1(x)|(x^{n_1} - 1)$, $f_e(x)|(x^{n_2} - 1)$ and $f_o(x)|(x^{n_3} - 1)$.

Further, we have \mathcal{C} is θ_t -cyclic then \mathcal{C}_k where $k \in \{1, e_1, e_2, o_1, o_2, o_3\}$ is skew θ_t -cyclic code over \mathbb{F}_q with respective lengths. From Theorem 3, $|\mathcal{C}| = |\mathcal{C}_1| \prod_{i=1}^{i=2} |C_{e_i}| \prod_{j=1}^{j=3} |C_{o_j}|$ since each \mathcal{C}_k is θ_t -cyclic it is generated by a polynomial $f_k(x)$ thus $|\mathcal{C}| = q^{\gamma - \sum_{i=1}^6 \epsilon_i}$ where $\gamma = n_1 + 2(n_2) + 3(n_3)$. The following Theorem provides the generator polynomials for θ_t -cyclic codes over \mathbb{F}_q .

Theorem 12 *Let $\mathcal{C} = \mathcal{C}_1 \otimes \mathcal{C}_e \otimes \mathcal{C}_o$ be a skew cyclic code over \mathcal{R} of length $\gamma = n_1 + n_2 + n_3$. Then there exists a polynomial*

- (i) $f_1(x) \in \mathbb{F}_q[x; \theta_t]$ such that $\mathcal{C}_1 = \langle f_1(x) \rangle$ and $x^{n_1} - 1 = g_1(x)f_1(x)$.
- (ii) $f_e(x) \in \mathcal{R}_1[x; \theta_t]$ such that $\mathcal{C}_e = \langle f_e(x) \rangle$ and $x^{n_2} - 1 = g_e(x)f_e(x)$ where $f_e(x) = \sum_{i=1}^2 e_i f_{e_i}(x)$.
- (iii) $f_o(x) \in \mathcal{R}_2[x; \theta_t]$ such that $\mathcal{C}_o = \langle f_o(x) \rangle$ and $x^{n_3} - 1 = g_o(x)f_o(x)$ where $f_o(x) = \sum_{i=1}^3 o_i f_{o_i}(x)$.

Proof Let \mathcal{C} be a θ_t -cyclic code of length $\gamma = n_1 + n_2 + n_3$. From Theorem 6, we have that $\mathcal{C}_1, \mathcal{C}_e$ and \mathcal{C}_o are θ_t -cyclic codes. Using Lemma 1, (i) follows.

The the proof of (ii) is as follows, let $\mathcal{C}_e = e_1 \mathcal{C}_{e_1} \oplus e_2 \mathcal{C}_{e_2}$ be a θ_t -cyclic code of length n_2 over \mathcal{R}_1 . Theorem 7 says that, \mathcal{C}_{e_1} and \mathcal{C}_{e_2} are θ_t -cyclic codes of length n_2 over \mathbb{F}_q . Lemma 1 says that we have $\mathcal{C}_i = \langle f_{e_i}(x) \rangle$ and $x^{n_2} - 1 = g_{e_i}(x)f_{e_i}(x)$ in $\mathbb{F}_q[x; \theta_t]$ for $i \in \{1, 2\}$. Then $e_i f_{e_i}(x) \in \mathcal{C}$ for $i \in \{1, 2\}$. Also for any $f_e(x) \in \mathcal{C}$, we have $f_e(x) = \sum_{i=1}^2 e_i h_{e_i}(x) f_{e_i}(x)$ where $h_{e_i}(x) \in \mathbb{F}_q[x; \theta_t]$ for $i \in \{1, 2\}$. Thus $f_e(x) \in \langle e_1 f_{e_1}(x), e_2 f_{e_2}(x) \rangle$. Therefore, $\mathcal{C} = \langle e_1 f_{e_1}(x), e_2 f_{e_2}(x) \rangle$. As $x^{n_2} - 1 = g_{e_i}(x)f_{e_i}(x)$ in $\mathbb{F}_q[x; \theta_t]$ for $i \in \{1, 2\}$. Let $f_e(x) = e_1 f_{e_1}(x) + e_2 f_{e_2}(x) \in \mathcal{R}_1[x; \theta_t]$. Then $f_e(x) \in \mathcal{C}$. On the other hand $e_i f_{e_i}(x) = e_i f_e(x) \in \langle f_e(x) \rangle$ for $i = 1, 2$. Consequently, $\mathcal{C} = \langle f_e(x) \rangle$. Further, $[\sum_{i=1}^2 e_i g_{e_i}(x)]f_e(x) = \sum_{i=1}^2 e_i g_{e_i}(x) f_{e_i}(x) = \sum_{i=1}^2 e_i (x^{n_2} - 1) = x^{n_2} - 1$. Then $x^{n_2} - 1 = g_e(x)f_e(x)$ in $\mathcal{R}_1[x; \theta_t]$, where $g_e(x) = \sum_{i=1}^2 e_i g_{e_i}(x)$. Thus (ii) is follows. (iii) is similar to the proof of (ii).

5 Skew Constacyclic Code over \mathcal{R}

In this section, we study skew θ_t -constacyclic codes over \mathcal{R} . We choose a unit element $\alpha \in \mathcal{R}_2^*$ such that α satisfying the condition $\alpha^2 = 1, (\alpha = 1, -1, \dots)$.

Definition 3 Let $\alpha_i \in \mathbb{F}_{p^t}/\{0\}$, a linear code $\mathcal{C} \subseteq \mathcal{R}_\gamma[x, \theta]$ is called skew $\alpha = \alpha_1 + u\alpha_2 + v\alpha_3$ -constacyclic code if it is invariant under the cyclic shift operator λ_α that is whenever

$$\begin{aligned} c &= (x_0, x_1, \dots, x_{n_1-1}, y_0, y_1, \dots, y_{n_2-1}, z_0, z_1, \dots, z_{n_3-1}) \in \mathcal{C} \\ \lambda_\alpha(c) &= (\alpha_1\theta_t(x_{n_1-1}), \theta_t(x_0), \dots, \theta_t(x_{n_1}), (\alpha_1 + u\alpha_2)\theta_t(y_{n_1-1}), \theta_t(y_1), \dots, \theta_t(y_{n_2-2}), \\ &\quad (\alpha_1 + u\alpha_2 + v\alpha_3)\theta_t(z_{n_3-1}), \theta_t(z_0), \dots, \theta_t(z_{n_3-2})) \in \mathcal{C} \end{aligned}$$

The following two results translate symmetry conditions into algebraic constraints. The proof of the first result is immediate and omitted.

Theorem 13 Let $R_{n,\lambda} = R[x; \theta_t]/\langle x^n - \lambda \rangle$. A linear code \mathcal{C} of length n over R is (θ_t, λ) -cyclic code if and only if \mathcal{C} is a left $R[x; \theta_t]$ -submodule of $R_{n,\lambda}$.

Theorem 14 A code \mathcal{C} is skew α -cyclic code over $\mathcal{R}_\gamma = \frac{\mathbb{F}_q[x, \theta_t]}{x^{n_1} - \alpha} \times \frac{\mathcal{R}_1[x, \theta_t]}{x^{n_2} - \alpha} \times \frac{\mathcal{R}_2[x, \theta_t]}{x^{n_3} - \alpha}$ iff \mathcal{C} is a left $\mathcal{R}_2[x, \theta_t]$ module over \mathcal{R}_γ .

Proof Let \mathcal{C} be a skew α -cyclic code. Then by definition $x \star (f(x)|g(x)|h(x)) \in \mathcal{C}$

$$\begin{aligned} x \star (f(x)|g(x)|h(x)) &= (\theta_t(f_0)x + \theta_t(f_1)x^2 + \dots + \alpha_1\theta_t(f_{n_1-1}), \theta_t(g_0)x + \theta_t(g_1)x^2 + \dots + (\alpha_1 + u\alpha_2)\theta_t(g_{n_2-1}), \\ &\quad \theta_t(h_0)x + \theta_t(h_1)x^2 + \dots + (\alpha_1 + u\alpha_2 + v\alpha_3)\theta_t(h_{n_3-1})) \in \mathcal{C} \end{aligned}$$

And by using linearity of \mathcal{C} ,

$$r(x) \star (g_1(x)|g_2(x)|g_3(x)) \in \mathcal{C}$$

for some $r(x) \in \mathcal{R}_2[x, \theta_t]$. Hence \mathcal{C} is an left $\mathcal{R}_2[x, \theta_t]$ submodule over \mathcal{R}_γ . Conversely, assume that \mathcal{C} is an left $\mathcal{R}_2[x, \theta_t]$ submodule over \mathcal{R}_γ , then we have $x \star (f(x)|g(x)|h(x)) \in \mathcal{C}$ implies \mathcal{C} is skew α -cyclic code.

Theorem 15 Let $\mathcal{C}_o \subseteq \mathcal{R}_2^n$ be a skew $\alpha = \alpha_1 + u\alpha_2 + v\alpha_3$ -cyclic code of length n iff $\mathcal{C}_{o_1}, \mathcal{C}_{o_2}$ and \mathcal{C}_{o_3} are skew $\alpha_1, \alpha_1 + \alpha_2, \alpha_1 + \alpha_3$ -cyclic codes over \mathbb{F}_q of length n .

Proof Let \mathcal{C}_o be a skew α -cyclic code, Let $a = x o_1 + y o_2 + z o_3 \in \mathcal{C}_o$ where $x = (x_0, x_1, \dots, x_{n_1-1}) \in \mathcal{C}_{o_1}, y = (y_0, y_1, \dots, y_{n_2-1}) \in \mathcal{C}_{o_2}$ and $z = (z_0, z_1, \dots, z_{n_3-1}) \in \mathcal{C}_{o_3}$. Then we have by definition, $\lambda_\alpha(x) \in \mathcal{C}_o$,

$$\begin{aligned} \lambda_\alpha(o_1(x_0, x_1, \dots, x_{n_1-1}) + o_2(y_0, y_1, \dots, y_{n_2-1}) \\ + o_3(z_0, z_1, \dots, z_{n_3-1})) &= ((\alpha_1 + u\alpha_2 + v\alpha_3) \star o_1(\theta_t(x_{n_1-1}), \theta_t(x_0), \dots, \theta_t(x_{n_1-2})) + \\ &\quad (\alpha_1 + u\alpha_2 + v\alpha_3) \star o_2(\theta_t(y_{n_2-1}), \theta_t(y_0), \dots, \theta_t(y_{n_2-2})) + \\ &\quad (\alpha_1 + u\alpha_2 + v\alpha_3) \star o_3(\theta_t(z_{n_3-1}), \theta_t(z_0), \dots, \theta_t(z_{n_3-2}))) \\ &\implies \lambda_{\alpha_1}(x) + \lambda_{\alpha_1 + \alpha_2}(y) + \lambda_{\alpha_1 + \alpha_3}(z) \in \mathcal{C}_o \\ &\implies \lambda_{\alpha_1}(x) \in \mathcal{C}_{o_1}, \lambda_{\alpha_1 + \alpha_2}(y) \in \mathcal{C}_{o_2}, \lambda_{\alpha_1 + \alpha_3}(z) \in \mathcal{C}_{o_3} \end{aligned}$$

Hence, $\mathcal{C}_{o_1}, \mathcal{C}_{o_2}$ and \mathcal{C}_{o_3} are skew $\alpha_1, \alpha_1 + \alpha_2, \alpha_1 + \alpha_3$ -cyclic codes over \mathbb{F}_q of length n .

Conversely, assume that $\mathcal{C}_{o_1}, \mathcal{C}_{o_2}$ and \mathcal{C}_{o_3} are skew $\alpha_1, \alpha_1 + \alpha_2, \alpha_1 + \alpha_3$ -cyclic codes over \mathbb{F}_q of length n . Let m_0, m_1, \dots, m_{n-1} be an element in \mathcal{C}_o , where $m_i = o_1 x_i + o_2 y_i + o_3 z_i$ such that $x = (x_0, x_1, \dots, x_{n_1-1}) \in \mathcal{C}_{o_1}, y = (y_0, y_1, \dots, y_{n_2-1}) \in \mathcal{C}_{o_2}$ and $z = (z_0, z_1, \dots, z_{n_3-1}) \in \mathcal{C}_{o_3}$. Then we have $\lambda_{\alpha_1}(x) \in \mathcal{C}_{o_1}, \lambda_{\alpha_1 + \alpha_2}(y) \in \mathcal{C}_{o_2}$ and $\lambda_{\alpha_1 + \alpha_3}(z) \in \mathcal{C}_{o_3}$. So we get,

$$\begin{aligned} o_1 \lambda_{\alpha_1}(x) + o_2 \lambda_{\alpha_1 + \alpha_2}(y) + o_3 \lambda_{\alpha_1 + \alpha_3}(z) &= o_1 \lambda_{\alpha_1}(x_0, x_1, \dots, x_{n_1-1}) + o_2 \lambda_{\alpha_1 + \alpha_2}(y_0, y_1, \dots, y_{n_2-1}) \\ &\quad + o_3 \lambda_{\alpha_1 + \alpha_3}(z_0, z_1, \dots, z_{n_3-1}) \in \mathcal{C} \\ &= \lambda_\alpha(m_0, m_1, \dots, m_{n-1}) \in \mathcal{C} \end{aligned}$$

Hence, \mathcal{C} is skew α -cyclic code over \mathcal{R}_2^n

Theorem 16 Let \mathcal{C}_e be a skew $\alpha = \alpha_1 + u\alpha_2$ -cyclic code over \mathcal{R}_1 iff \mathcal{C}_{e_1} and \mathcal{C}_{e_2} are skew $\alpha_1 + \alpha_2$ and α_1 -cyclic codes over \mathbb{F}_q .

Proof The proof is similar to Theorem 15 taking mod v to the above condition.

Theorem 17 Let \mathcal{C} be a skew $\alpha = \alpha_1 + u\alpha_2 + v\alpha_3$ -cyclic code over \mathcal{R} of length $\gamma = n_1 + n_2 + n_3$ iff $\mathcal{C}_1, \mathcal{C}_e$ and \mathcal{C}_o are $\alpha_1, \alpha_1 + u\alpha_2$ and $\alpha_1 + u\alpha_2 + v\alpha_3$ -cyclic codes over $\mathbb{F}_q, \mathcal{R}_1$ and \mathcal{R}_2 respectively.

Proof Let $\mathcal{C}_1, \mathcal{C}_e$ and \mathcal{C}_o be $\alpha_1, \alpha_1 + u\alpha_2$ and $\alpha_1 + u\alpha_2 + v\alpha_3$ -cyclic. Consider $x = (x_0, x_1, \dots, x_{n_1-1}), y = (y_0, y_1, \dots, y_{n_2-1})$ and $z = (z_0, z_1, \dots, z_{n_3-1})$. Consider $\alpha_1 + u\alpha_2 = \beta$. Then we have,

$$(x, y, z) \in \mathcal{C} \implies (\lambda_{\alpha_1}(x), \lambda_{\beta}(y), \lambda_{\alpha}(z)) \in \mathcal{C}$$

Hence \mathcal{C} is skew α -cyclic. The converse part holds similarly.

Theorem 18 Let \mathcal{C} be a skew α -cyclic code of length $\gamma = n_1 + n_2 + n_3$ iff \mathcal{C}_1 is skew α_1 -cyclic code of length n_1 , \mathcal{C}_{e_1} & \mathcal{C}_{e_2} are $\alpha_1 + \alpha_2, \alpha_1$ -cyclic codes of length n_2 and $\mathcal{C}_{o_1}, \mathcal{C}_{o_2}$ & \mathcal{C}_{o_3} are skew $\alpha_1, \alpha_1 + \alpha_2, \alpha_1 + \alpha_3$ -cyclic codes over \mathbb{F}_q of length n_3 .

Proof Using Theorem 15,16,17 the result follows.

Theorem 19 Let \mathcal{C} be a skew $\alpha = \alpha_1 + u\alpha_2 + v\alpha_3$ -cyclic code over \mathcal{R} of length $\gamma = n_1 + n_2 + n_3$ iff $\mathcal{C}_1^\perp, \mathcal{C}_e^\perp$ and \mathcal{C}_o^\perp are $(\alpha_1)^{-1}, (\alpha_1 + u\alpha_2)^{-1}$ and $(\alpha_1 + u\alpha_2 + v\alpha_3)^{-1}$ -cyclic.

Proof Let \mathcal{C} be a skew α -cyclic code, [Lemma 3.1,[27]] says that \mathcal{C}^\perp is skew $(\alpha_1 + u\alpha_2 + v\alpha_3)^{-1}$ -cyclic code. From Theorem 17, we have $\mathcal{C}_1^\perp, \mathcal{C}_e^\perp$ and \mathcal{C}_o^\perp are skew $(\alpha_1)^{-1}, (\alpha_1 + u\alpha_2)^{-1}$ and $(\alpha_1 + u\alpha_2 + v\alpha_3)^{-1}$ -cyclic.

Corollary 2 Let $\mathcal{C} = \mathcal{C}_1 \otimes \mathcal{C}_e \otimes \mathcal{C}_o$. be a skew $\alpha = \alpha_1 + u\alpha_2 + v\alpha_3$ -cyclic code over \mathcal{R} of length $\gamma = n_1 + n_2 + n_3$. Then there exists a polynomial.

- (i) $f_1(x) \in \mathbb{F}_q[x; \theta_t]$ such that $\mathcal{C}_1 = \langle f_1(x) \rangle$ and $x^{n_1} - \alpha_1 = g_1(x)f_1(x)$.
- (ii) $f_e(x) \in \mathcal{R}_1[x; \theta_t]$ such that $\mathcal{C}_e = \langle f_e(x) \rangle$ and $x^{n_2} - (\alpha_1 + u\alpha_2) = g_e(x)f_e(x)$.
- (iii) $f_o(x) \in \mathcal{R}_2[x; \theta_t]$ such that $\mathcal{C}_o = \langle f_o(x) \rangle$ and $x^{n_3} - (\alpha_1 + u\alpha_2 + v\alpha_3) = g_o(x)f_o(x)$.

Proof The proof is similar to the proof of Theorem 12.

Theorem 20 Let $\mathcal{C} = \mathcal{C}_1 \otimes \mathcal{C}_e \otimes \mathcal{C}_o$ be a θ_t -constacyclic code of length γ over \mathcal{R} . Then $\mathcal{C} = \langle \mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_o \rangle$ where $\mathcal{B}_1 = \langle (f_1(x), 0, 0) \rangle, \mathcal{B}_2 = \langle (0, f_e(x), 0) \rangle$, and $\mathcal{B}_o = \langle (b_1(x), b_e(x), f_o(x)) \rangle$.

Proof The proof is similar to proof of Theorem 11.

Example 1 Let $q = 9$ and $\mathbb{F}_9 = \mathbb{F}_3[z]$ with $z^2 + 1 = 0$ then consider the ring $R_\gamma = \frac{\mathbb{F}_9[x, \theta_t]}{x^4 - 1} \times \frac{\mathcal{R}_1[x, \theta_t]}{x^5 - 1} \times \frac{\mathcal{R}_2[x, \theta_t]}{x^5 - 1}$ and θ_3 be the Frobenius automorphism that is $\theta_3(a) = a^3$ where $a \in \mathbb{F}_9$.

$$x^4 - 1 = (x + 1)(x + 2)(x + z)(x + 2z) \in \mathbb{F}_9[x, \theta_3]$$

$$x^5 - 1 = (x + 2)(x^4 + x^3 + x^2 + x + 1) \in \mathbb{F}_9[x, \theta_3]$$

$f_1(x) = \langle (x + 1) \rangle, f_e(x) = \langle e_1(x + 2) + e_2(x + 2) \rangle, f_o(x) = \langle o_1(x + 2) + o_2(x + 2) + o_3(x + 2) \rangle$ from Theorem 12 we have f_i divides $x^{n_i} - 1$ for $(i = 1, e, o)$ thus we have a code with parameter $[29, 18, 2]$ over \mathbb{F}_9 .

Example 2 Let $q = 25$ and $\mathbb{F}_9 = \mathbb{F}_3[z]$ with $z^2 + z + 1 = 0$ then consider the ring $R_\gamma = \frac{\mathbb{F}_{25}[x, \theta_t]}{x^4 - 1} \times \frac{\mathcal{R}_1[x, \theta_t]}{x^6 - 1} \times \frac{\mathcal{R}_2[x, \theta_t]}{x^4 - 1}$ and θ_5 be the Frobenius automorphism that is $\theta_5(a) = a^5$ where $a \in \mathbb{F}_{25}$.

$$x^4 - 1 = (x + 2)(x + 3)(x + z)(x + z + 1) \in \mathbb{F}_{25}[x, \theta_5]$$

$$x^6 - 1 = (x^2 - 1)(x^2 + x + 1)(x^2 - x + 1) \in \mathbb{F}_{25}[x, \theta_5]$$

$f_1(x) = \langle (x + 2) \rangle, f_e(x) = \langle e_1(x^2 - 1) + e_2(x^2 - 1) \rangle, f_o(x) = \langle o_1(x + z + 1) + o_2(x + z + 1) + o_3(x + z + 1) \rangle$ from Theorem 12 we have f_i divides $x^{n_i} - 1$ for $(i = 1, e, o)$ thus we have a code with parameter $[28, 20, 2]$ over \mathbb{F}_{25} .

6 Conclusion and open problems

In this note, we have studied the algebraic and metric structure of skew cyclic and skew constacyclic codes over a special mixed alphabet. Thus, our codes have a structure of module over the largest of the three alphabets \mathcal{R}_2 . Codes over the product ring $\mathbb{F}_q \times \mathcal{R}_1 \times \mathcal{R}_2$ would be modules over that larger ring. The two algebraic structures are different and should not be confused.

The present work leads itself to two paths of generalization: consider different mixed alphabets or replace the concepts of cyclicity by that of quasi-cyclicity. The former path seems easier than the latter, in view of the many examples of rings that have been used as alphabets of cyclic codes in recent years. On the other hand, the structure of quasi-cyclic codes is always more subtle than that of cyclic codes.

References

1. Abualrub, T., Aydin, N. and Seneviratne, P. (2012) ‘On θ -cyclic codes over $F_2 + vF_2$ ’, *The Australasian Journal of Combinatorics*, Vol. 54, pp.115-126.
2. T. Abualrub, I. Siap and N. Aydin, $\mathbb{Z}_2\mathbb{Z}_4$ -Additive cyclic codes, *IEEE Trans. Inform. Theory*, **60** (2014), 1508-1514.
3. M. Ashraf and G. Mohammad, *Skew cyclic codes* $\mathbb{F}_3 + v\mathbb{F}_3$, *Int. J. Inf. Coding Theory* **24** (2014), 218-225.
4. M. Ashraf and G. Mohammad, *Skew cyclic codes* $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q$, *Asian-European Journal of Mathematics* **115** (2018), 1850072.
5. I. Aydogdu and I. Siap, The structure of $\mathbb{Z}_2\mathbb{Z}_2^s$ -additive codes: Bounds on the minimum distance, *Appl. Math. Inf. Sci.*, **7**(6) (2013), 2271-2278.
6. I. Aydogdu, T. Abualrub and I. Siap, On $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes, *Int. J. Comput. Math.*, **92** (2015), 1806-1814.
7. I. Aydogdu and I. Siap, On $\mathbb{Z}_{p^r}\mathbb{Z}_p^s$ -additive codes, *Linear Multilinear Algebra*, **63** (2015), 2089-2102.
8. I. Aydogdu, T. Abualrub and I. Siap, The $\mathbb{Z}_2\mathbb{Z}_2[u]$ -cyclic and constacyclic codes, *IEEE Trans. Inform. Theory*, **63**(8) (2016), 4883-4893.
9. E.R Berlekamp, Negacyclic codes for the Lee metric in R.C. Bose, TA Dowling, *Combinatorial Mathematics*, University of North Carolian Press (1969), 298–316.
10. J. Borges, C. Fernandez-Cordoba, J. Pujol and J. Rifa, $\mathbb{Z}_2\mathbb{Z}_4$ -linear codes: generator matrices and duality, *Des. Codes Cryptogr.*, **54** (2010), 167-179.
11. J. Borges, C. Fernandez-Cordoba and R. Ten-Valls, $\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic codes, generator polynomials and dual codes, *IEEE Trans. Inform. Theory*, **62** (2016), 6348-6354.
12. S. Bhardwaj and M. Raka, *Skew constacyclic codes over a non-chain ring* $\mathbb{F}_q[u,v]/\langle f(u),g(v),uv - vu \rangle$, arXiv:1905.12933 [cs.IT] (2019), (preprint).
13. W. Bosma and J. Cannon, *Handbook of Magma Functions*, Version 2.12, Volume 1, Sydney, (2006).
14. D. Boucher, W. Geiselmann and F. Ulmer, *Skew cyclic codes*, *Appl. Algebra Eng. Comm.* **18** (2007), 379-389.
15. D. Boucher, P. Solé and F. Ulmer, *Skew constacyclic codes over Galois rings*, *Adv. Math. Commun.* **2**(3) (2011), 273-292.
16. D. Boucher and F. Ulmer, *Coding with skew polynomial rings*, *J. Symb. Comput.* **44**(12) (2009), 1644-1656.
17. A. Dertli and Y. Cengellenmis, *Skew cyclic codes over* $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q$, *Journal of Science and Arts.* (2017), **2**(39), pp. 215-222.
18. A. Dertli and Y. Cengellenmis, *On the linear codes over the ring R_p* , *Discrete Mathematics, Algorithms and Applications* (2016), **8**(2), 1650036.
19. C. Fernandez-Cordoba, J. Pujol, and M. Villanueva, $\mathbb{Z}_2\mathbb{Z}_4$ -linear codes: Rank and kernel, *Des. Codes Cryptogr.*, **56**(1) (2010), 43–59.
20. J. Gao, F. Ma and F. Fu, *Skew constacyclic codes over the ring* $\mathbb{F}_q + v\mathbb{F}_q$, *Appl. Comput. Math.* **6**(3) (2017), 286-295.
21. F. Gursoy, I. Siap and B. Yildiz, *Construction of skew cyclic codes over* $\mathbb{F}_q + v\mathbb{F}_q$, *Adv. Math. Commun.* **8**(3) (2014), 313-322.
22. Hammons, A. R. Jr., Kumar, P. V., Calderbank, A. R., Sloane, N. J. A. and Solé, P. (1994) The \mathbb{Z}_4 - linearity of Kerdock, Preparata, Goethals, and related codes, *IEEE Transactions on Information Theory*, Vol. 40, No. 2, pp.301-319.
23. R. Hill, *A First Course in Coding Theory*, Oxford Applied Linguistics, Clarendon Press, 1986.
24. H. Islam and O. Prakash, *Skew cyclic and skew* $(\alpha_1 + u\alpha_2 + v\alpha_3 + uv\alpha_4)$ -constacyclic codes over $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q$, *Int. J. Inf. Coding Theory* **5**(2) (2018), 101-116.
25. H. Islam and O. Prakash, *A note on skew constacyclic codes over* $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q$, *Discrete Mathematics, Algorithms and Applications* **11**(3) (2019), 1950030.

26. H. Islam and O. Prakash, On $\mathbb{Z}_p\mathbb{Z}_p[u, v]$ -additive cyclic and constacyclic codes, (2019), <http://arxiv.org/abs/1905.06686v1>.
27. S. Jitman, S. Ling and P. Udomkavanich, *Skew constacyclic codes over finite chain ring*, Adv. Math. Commun. **6** (2012), 39-63.
28. J-F. Qian, L-N. Zhang and S-X. Zhu, $(1 + u)$ constacyclic and cyclic codes over $\mathbb{F}_2 + u\mathbb{F}_2$, Appl. Math. Lett. **19**(8) (2006), 820-823.
29. H. Rifa-Pous, J. Rifa, and L. Ronquillo, $\mathbb{Z}_2\mathbb{Z}_4$ -additive perfect codes in steganography, Adv. Math. Commun., **5**(3) (2011), 425-433.
30. M. Shi, L. Qian, L. Sok, N. Aydin and P. Sole, *On constacyclic codes over $\mathbb{Z}_4[u]/\langle u^2 - 1 \rangle$ and their Gray images*, Finite Fields Appl. **45** (2017), 86-95.
31. M. Shi, R. Wu and D. S. Krotov, On $\mathbb{Z}_p\mathbb{Z}_{p^k}$ -additive codes and their duality. IEEE Trans. Inf. Theory, **65**(6) (2018), 3842-3847.
32. B. Srinivasulu and B. Maheshanand, The $\mathbb{Z}_2(\mathbb{Z}_2 + u\mathbb{Z}_2)$ -additive cyclic codes and their duals, Discrete Math. Algorithm. Appl., **8** (2016), 1793-8317.
33. T. Yao, M. Shi and P. Sole, *On Skew cyclic codes over $\mathbb{F}_q + u\mathbb{F}_q + v\mathbb{F}_q + uv\mathbb{F}_q$* , J. Algebra Comb. Discrete Struct. Appl. **2**(3) (2015), 163-168.
34. I. Siap, T. Abualrub, N. Aydin and P. Seneviratne, *Skew cyclic codes of arbitrary length*, Int. J. Inf. Coding Theory **2** (2011), 10-20.
35. T. Yao and S. Zhu, $\mathbb{Z}_p\mathbb{Z}_{p^s}$ -additive cyclic codes are asymptotically good, Cryptogr. Commun. **12** (2020), 253-264.
36. T. Yao, S. Zhu and X. Kai, Asymptotically good $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive cyclic codes, Finite Fields Appl., **63** (2020), <https://doi.org/10.1016/j.ffa.2020.101633>.
37. H. Yu, S. Zhu and X. Kai, $(1 - uv)$ -constacyclic codes over $\mathbb{F}_p + u\mathbb{F}_p + v\mathbb{F}_p + uv\mathbb{F}_p$, J. Syst. Sci. Complex. **27**(5) (2014), 811-816.
38. S. Zhu, Y. Wang and M. Shi *Some results on cyclic codes over $\mathbb{F}_2 + u\mathbb{F}_2$* , IEEE Trans. Inform. Theory **56**(4) (2010), 1680-1684.
39. X. Zheng and B. Kong, *Cyclic codes and $\lambda_1 + \lambda_2u + \lambda_3v + \lambda_4uv$ -constacyclic codes over $\mathbb{F}_p + u\mathbb{F}_p + v\mathbb{F}_p + uv\mathbb{F}_p$* , Adv. Math. Commun. **306** (2017), 86-91.