



HAL
open science

PGPR-Soil Microbial Communities' Interactions and Their Influence on Wheat Growth Promotion and Resistance Induction against *Mycosphaerella graminicola*

Erika Samain, Jérôme Duclercq, Essaïd Ait Barka, Michael Eickermann, Cédric Ernenwein, Candice Mazoyon, Vivien Sarazin, Frédéric Dubois, Thierry Aussenac, Sameh Selim

► To cite this version:

Erika Samain, Jérôme Duclercq, Essaïd Ait Barka, Michael Eickermann, Cédric Ernenwein, et al.. PGPR-Soil Microbial Communities' Interactions and Their Influence on Wheat Growth Promotion and Resistance Induction against *Mycosphaerella graminicola*. *Biology*, 2023, 12 (1416), pp.2-18. 10.3390/biology12111416 . hal-04309786

HAL Id: hal-04309786

<https://hal.science/hal-04309786>

Submitted on 27 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Article

PGPR-Soil Microbial Communities' Interactions and Their Influence on Wheat Growth Promotion and Resistance Induction against *Mycosphaerella graminicola*

Erika Samain ^{1,2}, Jérôme Duclercq ³ , Essaïd Ait Barka ⁴ , Michael Eickermann ⁵ , Cédric Ernenwein ², Candice Mazoyon ³, Vivien Sarazin ⁶ , Frédéric Dubois ³ , Thierry Aussenac ⁷  and Sameh Selim ^{1,*} 

- ¹ AGHYLE UP 2018.C101, SFR Condorcet FR CNRS 3417, UniLaSalle, 19 Rue Pierre Waguët, BP 30313, CEDEX, 60026 Beauvais, France; erika.samain@hotmail.fr
- ² SDP, 1 Rue Quesnay, CEDEX, 02000 Laon, France; cedric.ernenwein@rovensanext.com
- ³ Unité Ecologie et Dynamique des Systèmes Anthropisés (EDYSAN, UMR7058 CNRS), Université de Picardie Jules Verne (UPJV), 80039 Amiens, France; jerome.duclercq@u-picardie.fr (J.D.); candice.mazoyon@outlook.fr (C.M.); frederic.dubois@u-picardie.fr (F.D.)
- ⁴ Unité de Recherche RIBP EA4707 USC INRAE 1488, SFR Condorcet FR CNRS 3417, Université de Reims Champagne Ardenne, 51100 Reims, France; ea.barka@univ-reims.fr
- ⁵ Luxembourg Institute of Science and Technology, 4422 Belvaux, Luxembourg; michael.eickermann@list.lu
- ⁶ AgroStation, 68700 Aspach-le-Bas, France; vivien.sarazin@agrostation.fr
- ⁷ UP Institut Polytechnique UniLaSalle, Université d'Artois, ULR 7519, 19 Rue Pierre Waguët, BP 30313, CEDEX, 60026 Beauvais, France; thierry.aussenac@unilasalle.fr
- * Correspondence: sameh.selim@unilasalle.fr; Tel.: +33-3-4406-3825

Simple Summary: Plant resistance inducers, such as plant-growth-promoting rhizobacteria (PGPR), are among the most important alternatives to fungicides because they employ different modes of action, conferring protection against biotic and abiotic stresses while promoting plant growth. However, the loss of their efficacy under field conditions is a subject of debate and may be attributed to the influence of environmental factors, genetic diversity and native soil microbial communities. Additionally, the inoculation of plants with PGPR can impact the complex and dynamic balance of soil microbial communities. Recently, we published the efficiency of three PGPR in single and co-inoculation to induce wheat resistance mechanisms against Septoria leaf blotch and drought stress. Here, we investigated the interactions between PGPR and soil microbial communities.

Abstract: The efficiency of plant-growth-promoting rhizobacteria (PGPR) may not be consistently maintained under field conditions due to the influence of soil microbial communities. The present study aims to investigate their impact on three PGPR-based biofertilizers in wheat. We used the PGPR *Paenibacillus* sp. strain B2 (PB2), PB2 in co-inoculation with *Arthrobacter agilis* 4042 (Mix 2), or with *Arthrobacter* sp. SSM-004 and *Microbacterium* sp. SSM-001 (Mix 3). Inoculation of PB2, Mix 2, and Mix 3 into non-sterile field soil had a positive effect on root and aboveground dry biomass, depending on the wheat cultivar. The efficiency of the PGPR was further confirmed by the protection they provided against *Mycosphaerella graminicola*, the causal agent of Septoria leaf blotch disease. PB2 exhibited protection of $\geq 37.8\%$, while Mix 2 showed $\geq 47.9\%$ protection in the four cultivars tested. These results suggest that the interactions between PGPR and native soil microbial communities are crucial for promoting wheat growth and protection. Additionally, high-throughput sequencing of microbial communities conducted 7 days after PGPR inoculations revealed no negative effects of PB2, Mix 2, and Mix 3 on the soil microbial community structure. Interestingly, the presence of *Arthrobacter* spp. appeared to mitigate the potential negative effect of PB2 on bacterial community and foster root colonization by other beneficial bacterial strains.

Keywords: *Mycosphaerella graminicola*; *Paenibacillus* sp. strain B2; *Arthrobacter* spp.; *Microbacterium* spp.; PGPR co-inoculation; induced systemic resistance; growth promotion; soil microbial communities



Citation: Samain, E.; Duclercq, J.; Ait Barka, E.; Eickermann, M.; Ernenwein, C.; Mazoyon, C.; Sarazin, V.; Dubois, F.; Aussenac, T.; Selim, S. PGPR-Soil Microbial Communities' Interactions and Their Influence on Wheat Growth Promotion and Resistance Induction against *Mycosphaerella graminicola*. *Biology* **2023**, *12*, 1416. <https://doi.org/10.3390/biology12111416>

Academic Editor: Bernard R. Glick

Received: 21 October 2023

Revised: 3 November 2023

Accepted: 8 November 2023

Published: 10 November 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Thousands to millions of microbial species, including fungi and bacteria, live in the rhizosphere, interact with plant roots, and are influenced by plant species and their growth stage, which is determined by root exudates [1]. These interactions can be pathogenic, neutral, or beneficial, as PGPR (Plant Growth-Promoting Rhizobacteria). PGPRs serve as biofertilizers by enhancing nutrient accessibility to plants, as biostimulators by secreting plant hormones important for plant growth, as antagonists to pathogens, or by inducing systemic resistance (ISR) in the plant. PGPR-mediated ISR is akin to pathogen-induced systemic acquired resistance (SAR), bolstering the plant's immune system against abiotic and biotic stresses. For instance, *Septoria tritici* leaf blotch (STB) disease in wheat, caused by *Mycosphaerella graminicola*, can lead to crop losses exceeding 40% [2,3]. ISR promotes the formation of physical barriers such as callose and lignin, the synthesis of protective compounds in plants, including reactive oxygen species, phytoalexins, and phenolic compounds [4,5]. Consequently, PGPR-mediated ISR is regarded as one of the most promising alternatives to fungicides, which are a subject of environmental and health-related controversies [6].

However, PGPR-induced ISR represents a quantitative, non-specific resistance with a broad-spectrum effect, which has often been demonstrated as highly effective under controlled conditions, but is not consistently maintained under field conditions [6,7]. Notably, almost all studies conducted under controlled conditions are limited in their scope, even when conducted under sterile conditions that restrict the range of strains and plant genotypes studied. Factors such as environmental conditions, genetic diversity, and the influence of native microbial communities on PGPR may provide possible explanations for the reduced efficiency of PGPR in field conditions [8]. Indeed, the soil can influence PGPR colonization, primarily through its physiochemical and existing microbial communities. These communities can have direct effects, such as trophic competitions and antagonistic or synergistic interactions, and indirect effects on plant growth and root exudation [5,9–11]. Inoculated PGPR must be competent, survive, proliferate, and act efficiently on plant growth and protection [12]. Native microbial communities often form competitive environments with a wide range of species, which can impact the viability and characteristics of PGPR inoculants as biofertilizer and biocontrol agents [8]. The successful colonization of this environment relies on the rhizosphere competence of the PGPR inoculum [13]. Therefore, understanding the population dynamics of the microbial community is crucial when applying microbial inoculants to establish links between plant growth promotion and protection against pathogens. In contrast to the effects of microbial communities on PGPR when used as biocontrol products, the potential environmental consequences of PGPR inoculants on native microbial communities have been relatively overlooked. Nevertheless, the introduction of high densities of viable, efficient, and competitive microbes may, at least temporarily, impact the complex and dynamic balance of soil microbial communities and the composition of taxonomic groups [14]. Ensuring the safety of microbial inoculation to the environment is a critical aspect in the development of potential biocontrol agents.

In this study, we focused on *Paenibacillus* sp. strain B2 (PB2), which is known to produce the cyclo-lipopeptide, paenimyxin, which serves as an antagonist against Gram-positive, Gram-negative bacteria, and fungi. PB2 has been shown to induce transient changes in soil bacterial community structure within 7 days of application [11,15] and to trigger ISR against *M. graminicola* in wheat [5,6]. Moreover, when used in co-inoculation with *Arthrobacter agilis* 4042 (Mix 2) or with *Arthrobacter* SSM-004 and *Microbacterium* SSM-001 (Mix 3), PB2 exhibited the ability to promote wheat growth and enhance its tolerance to drought stress, in addition to its protective effect against pathogens [10,16]. However, these previous results were obtained under controlled and sterile conditions. The objectives of our current work are to investigate the impact of soil microbial communities on the activities of PB2 when used alone or in co-inoculation regarding wheat growth and protection against *M. graminicola*. Additionally, we aim to examine the effects of these inoculations on bacterial and fungal communities in the rhizosphere and within wheat roots.

2. Materials and Methods

2.1. Microorganisms and Inoculum Preparation

Paenibacillus sp. strain B2 [17] was kindly provided by Dr. van Tuinen from INRA Dijon, France. *Arthrobacter agilis* was provided by C. Ernenwein of SDP, Laon, France, and *Microbacterium* sp. strain SSM1 and *Arthrobacter* sp. SSM4 were isolated in the author's laboratory.

PGPR cultures were prepared as described by Selim et al. [11]. Briefly, to prepare the final bacterial inocula, cells were harvested at $OD_{0.5}$, centrifuged at $2655 \times g$ for 10 min at 4 °C, washed twice, and then suspended in a sterile solution of 10 mM $MgSO_4$ (Sigma, St. Louis, MO, USA). Bacterial cell vitality was confirmed by plating 100 μ L of the inoculum on Luria-Bertani (LB, Sigma, St. Louis, MO, USA). Spores of *M. graminicola* strain IPO 323 (provided by Dr. F. Suffert, INRA Grignon) were collected from liquid cultures via centrifugation at $2655 \times g$ for 5 min at 15 °C, washed twice with sterile distilled water, and suspended in 10 mM $MgSO_4$ (Sigma, St. Louis, MO, USA) containing 0.1% Tween 20 surfactant. Fungal spore vitality was assessed by plating 100 μ L of the inoculum on potato dextrose agar (PDA, Sigma, St. Louis, MO, USA).

2.2. Soil Physicochemical Analysis

The soil used in this study was collected from the top 30 cm layer of an agricultural field at Institut Polytechnique UniLaSalle, Beauvais, France. This soil is a silt-loam soil (granulometric composition: silt 68.9%, clay 20.2%, sand 8.9%). Its composition includes organic matter at 1.8%, limestone at 0.2% and it has a pH of 7.1. Additionally, its nutrient content is as follows: 0.99 g N·kg⁻¹, 119 mg P₂O₅·kg⁻¹, and 194 mg K₂O·kg⁻¹.

2.3. Plant Material and Growth Conditions

Four wheat cultivars, Alixan, Altigo, Cellule, and Hyfi, with different degrees of resistance to STB, 4, 5.5, 6.5, and 7, respectively, on a scale of 1 (totally susceptible) to 9 (totally resistant; [18]) were used. The wheat seeds were disinfected according to Samain et al. [5], with some modifications. This involved an overnight incubation in a solution of the antibiotics oxytetracycline, streptomycin, penicillin, and ampicillin (100 mg·L⁻¹ each). The seeds were then submerged in a 10% calcium hypochlorite solution for 10 min and washed three times in sterile Milli-Q water after each disinfection step. The sterilized seeds were pre-germinated on a 0.5% (*w/v*) water agar medium and incubated for 24 h at 4 °C, followed by 48 h at 20 °C, and finally, 24 h at 4 °C in the dark. The germinated seeds were then transferred to an inoculum consisting of a single PGPR strain or a mixture containing an equal amount of each PGPR strain, resulting in a final concentration of 10⁶ CFU·mL⁻¹ in 10 mM $MgSO_4$. One mL of the inoculum was added per grain for one hour with gentle shaking. For the non-inoculated condition, seeds were submerged in 10 mM $MgSO_4$. After inoculation, the seeds were transferred into 250 mL pots filled with either sterile or non-sterile soil, which was a mixture of silt-loam soil and sand in a 1:1 ratio (*v/v*). The pots were placed in a phytotron with conditions set at 18 °C (+/−2 °C), 40% humidity, and a 16 h photoperiod with a photon flux density of 185 μ mol m⁻² s⁻¹ provided by white fluorescent tubes (Philips Master Cool White 80 W/865, Lamotte Beuvron, France). The plants were irrigated three times per week with 50 mL of distilled water per pot. At the 3-leaf growth stage, the plants were inoculated with 1 mL/plant, which contained 10⁶ spores of *M. graminicola* strain IPO323 and 0.1% (*v/v*) tween 20.

2.4. Wheat Resistance Induction against *M. graminicola* and Wheat Biomasses

Seventeen days after infection, the leaves were collected and freeze-dried to evaluate the protective effect against *M. graminicola* in response to the inoculation with PB2 or PGPR mixtures. To quantify the infection level of *M. graminicola*, we conducted DNA extraction and qPCR analysis following the method described by Selim et al. [2]. Briefly, DNA was extracted using the DNeasy 96 Plant Kit (Qiagen, Germantown, MA, USA) according to the manufacturer's protocol. To quantify the infection level of *M. graminicola*, we used

specific primers and a TaqMan Minor Groove Binder probe targeting a 63-bp fragment of the *M. graminicola* β -tubulin-specific gene (For; GCCTTCCTACCCCACCATGT, Rev; CCTGAATCGCGCATCGTTA, Probe; FAM-TTACGCCAAGACATTC-MGB, GeneBank accession number AY547264; [19]). TaqMan[®] assays were carried out using 12.5 μ L Universal TaqMan[®] PCR Master Mix (Life Technologies SAS, Villebon sur Yvette, France), 0.3 μ M of each primer, 0.2 μ M probe, 200 ng DNA, and water to a volume of 25 μ L. The qPCR conditions were set as follows: 10 min at 95 °C, followed by 40 cycles of 15 s at 95 °C, and 1 min at 60 °C. All qPCR experiments were performed using the StepOnePlus Real-Time PCR System (Applied Biosystems, Waltham, MA, USA). To calibrate the qPCR analysis, we used serial dilutions of the cloned target sequence of the β -tubulin gene of *M. graminicola*, ranging from 10² to 10⁷ copies, as previously described [3]. The results were expressed as the β -tubulin copy number per 100 ng of leaf DNA (BCN_{100ng}). In addition, we calculated the dry biomass of roots and leaves of each wheat cultivar in each modality at the 6-week-old plant stage.

2.5. Taxonomic Analysis of Microbial Communities

DNA was extracted from endophytic microorganisms in both the root (100 g) and rhizosphere (300 g) of wheat plants. The rhizospheric soil was obtained by harvesting the soil that surrounds plant roots after they have been removed from their growing substrate. Samples were collected from non-inoculated and PGPR-inoculated plants (PB2 alone or in co-inoculation), and from plants that were non-infested or infested with *M. graminicola*. We used the DNeasy 96 Plant kit (Qiagen, Germantown, MA, USA) for root samples and the QIAamp DNA Mini kit (Qiagen, Hilden, Germany) for soil samples, following the respective manufacturer's protocols. As in [20], the concentration of DNA was quantified fluorometrically using the AccuBlue High Sensitivity dsDNA Quantification Kit (Biotium, Fremont, CA, USA) and a monochromator based multimode microplate reader (Infinite M1000 PRO, Tecan System, Morgan Hill, CA, USA). For the amplification of the 16S_{V3-V4} rRNA regions, we used primers Bakt_341F (5'-CCTACGGGNGGCWGCAG-3') and Bakt_805R (5'-GACTACHVGGGTATCTAATCC-3'; [21]). To target the fungal internal transcribed spacer (ITS) region, we used primers ITS2_KYO1 (5'-CTRYGTTCTTCATCGDT-3') and ITS2_KYO2 (5'-TTYRCTRCTCGTTCTTCATC-3'; [22]). Both forward and reverse primers were designed to include overhang sequences compatible with Illumina Nextera XT index sequencing adapters. We used 5 ng of DNA per sample and verified the success of the PCR via agarose gel electrophoresis with GelGreen Nucleic Acid Gel Stain as the fluorescent intercalator, and the presence of bands of the right size on the gel with a MUPID One LED Illuminator (Advance, Suisun City, CA, USA). All amplicons were quantified using the AMPure XP Beads Kit (Beckman Coulter, Carlsbad, CA, USA) and the AccuBlue High Sensitivity dsDNA Quantitation Kit. Illumina Nextera XT Index sequencing adapters were incorporated into the amplicons through PCR. The final libraries were repurified using AMPure XP beads and quantified using the AccuBlue High Sensitivity dsDNA Quantitation Kit and the Infinite M1000 PRO microplate reader. To ensure library size and the absence of primer-dimer contamination, a 1 μ L sample of a 1:50 dilution of the final library was run on a Bioanalyzer DNA 1000 chip using a Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA, USA). Purified libraries were pooled at equal molarity, denatured with freshly prepared 0.2 N NaOH, diluted to 4 pM using pre-chilled hybridization buffer, spiked with 5% of a pre-made PhiX control library (PhiX control v2, Illumina, San Diego, CA, USA), and loaded into a MiSeq v2 Reagent Kit (500 Cycles PE, Illumina, USA) for sequencing in a MiSeq sequencer (Illumina).

2.6. Bioinformatic Analysis

Illumina sequence reads into FastQ format were analyzed with the FROGS (Find Rapidly OTU with Galaxy Solution) pipeline [23] on the Galaxy instance (v.2.3.0) of the Genotoul bioinformatics platform (<http://sigenae-workbench.toulouse.inra.fr>) (accessed on 13 September 2022). Sequences containing ambiguous bases (N) or those lacking

specific primers were removed. We identified and trimmed primer sequences that had less than 10% of differences using Cutadapt (v.1.18; [24]). Sequence clustering was performed using the SWARM algorithm (v2.1.5, [25]). This involved an initial denoising step to create fine sequence clusters with minimal differences ($d = 1$), followed by a second step with an aggregation distance of three. The representative sequences for each cluster or OTU (Operational Taxonomic Units) were subjected to chimera detection using the VSEARCH algorithm for subsequent removal [26]. Taxonomic identification of each OTU was performed up to the species level using the RDPClassifier and BLAST tools. These tools were applied to the non-redundant small sub-unit database from SILVA (v123) for bacterial communities and against the ITS Warcup Training Set (v2; [27]) for the fungal communities. The high-throughput sequencing of the 16S_{V3-V4} rRNA and ITS regions resulted in 10,293,301 and 3,487,769 reads, respectively.

2.7. Soil Microbial Metabolic Profiles

The metabolic potential of soil communities was assessed using community-level physiological profiles (CLPP) with Biolog EcoPlate (Biolog Inc., Hayward, CA, USA; [28]). Soil samples were analyzed in triplicate using an EcoPlate consisting of 31 different carbon sources and a blank control. Each well of the EcoPlate was inoculated with 100 μ L of a soil suspension (prepared by mixing 1 g of fresh soil in 10 mL of physiological water), containing approximately 10^5 colony-forming units. The plates were incubated for 196 h at 25 °C using an OmniLog[®] system (Biolog Inc.). To measure the rate of carbon source utilization, the reduction of tetrazolium was monitored. Tetrazolium is a redox color indicator that changes from colorless to purple, its change was detected at a wavelength of 590 nm. Data were recorded at 15 min intervals over a 192-h period and were stored as OmniLog units in Biolog data analysis software (v1.7). To calculate values for each well, blank values were subtracted from each well in the plate. To make comparisons between samples, only one absorption time point at 50 h was used, as recommended in [29]. At this time point, the metabolic potential of the soil microbial communities in each sample was calculated by determining the average color development in the wells (AWCD). This calculation involved summing the optical density data for all wells and dividing the sum by 31 (the total number of substrates). To assess the functional richness of the microbial community in the soil, we counted the total number of wells in one replicate with an absorbance greater than 25 OmniLog units.

2.8. Data Analysis and Statistics

Each experiment was biologically replicated twice, and each experimental condition within each experiment included a minimum of five replicates, except for high-throughput sequencing, where three technical replicates were used. To test the significance of PGPR inoculation and wheat genotype on plant growth and protection against *M. graminicola*, we performed variance analysis (ANOVA) and used Tukey's multiple range test for separating treatment means, with a significance level set at $p \leq 0.05$.

From the OTU \times sample abundance tables, we calculated taxonomic richness and Shannon indices to assess α -diversity. To ensure comparability of taxonomic richness across samples, we rarefied the number of bacterial and fungal OTUs using the *rrarefy* function from the *vegan* R package [30]. Since the dataset did not meet the assumptions of normality, all diversity indices were compared between treatments using the R package *agricolae* [31], applying a non-parametric Kruskal–Wallis one-way analysis of variance, followed by a Conover–Iman post hoc test whenever significant. p -values were adjusted using the Benjamini–Hochberg FDR procedure in R with the *p.adjust* function. To examine differences in species composition between communities, we employed permutational multivariate analysis of variance (PERMANOVA) with 999 permutations. We examined variations in bacterial and fungal community structures using Nonmetric Multidimensional Scaling (NMDS) based on Jaccard distances, utilizing the *metaMDS* function from the *vegan*

R package. For this analysis, we selected the 50% most abundant bacterial or fungal genera with a 30% best axis fit, using the *ordiselect* function from the *vegan* R package.

For the CLPP analysis, since the dataset did not meet ANOVA assumptions, we compared means between treatment groups using the R package *agricolae* and a Kruskal–Wallis test ($p < 0.05$), followed by pairwise Wilcoxon rank sum tests with Holm’s p-adjust method for multiple comparisons.

All statistical tests were performed with the R software (v4.2.2, The R Foundation, <https://www.r-project.org/>).

3. Results

3.1. Effects of PGPR Mixture on Wheat Biomass under Field Soil

The wheat cultivars Alixan and Cellule inoculated with PB2 showed no effects on dry biomass of roots and leaves. In contrast, Hyfi and Altigo cultivars responded differently to this inoculation. Specifically, Hyfi exhibited a notable 28.3% increase in leaf biomass and a remarkable 67% increase in root biomass when compared to non-inoculated wheat. As for Altigo, while its leaf biomass remained unaffected, there was a significant 77% increase in root biomass compared to non-inoculated plants (Figure 1). Furthermore, inoculation with Mix 2 led to increases in leaf and root biomass in all wheat cultivars, except Hyfi, where only the root biomass showed a notable increase. In contrast, the use of Mix 3 resulted in a significant increase in leaf biomass, but this effect was observed only in the Cellule and Altigo cultivars (Figure 1).




Figure 1. Effect of root inoculation with *Paenibacillus* sp. strain B2 (PB2), Mix 2, composed of PB2 and *Arthrobacter agilis*, or Mix 3, composed of PB2, *Microbacterium* sp. strain SSM1 and *Arthrobacter* sp. SSM4 PB2, on wheat growth in non-sterile field soils. Plants were inoculated with PGPR by dipping pre-germinated seeds in a suspension totaling 10^6 CFU·mL⁻¹. Dry biomasses of leaves (A) and roots (B) were determined 6 weeks after sowing in four wheat cultivars (Alixan, Altigo, Cellule, Hyfi). Values shown are means with SD ($n = 5$). Different lower-case letters indicate significant differences between groups according to Tukey’s test at $p \leq 0.05$.

3.2. Resistance Induction in Wheat against *M. graminicola*

The level of protection induced in response to root inoculation with PGPR was determined by comparing BCN_{100ng} values in the leaves of the wheat plants, which served as controls. These control plants were non-inoculated with PGPR but infested with *M. graminicola*. We considered the protection efficiency at 40% as a minimum limit to accept the PGPR’ product as a resistance inducer. In sterile soil, BCN_{100ng} values were 10903, 379,

396, and 4444 for Alixan, Altigo, Cellule, and Hyfi, respectively. In non-sterile soil, these values were 5571, 244, 242, and 2358, respectively.

With PB2 inoculation, the induced protection against *M. graminicola* showed a significant decrease in non-sterile soil compared to sterile soil for the Alixan and Altigo cultivars. For Alixan, the protection decreased from 75.2% to 46.9%, and for Altigo, it decreased from 78.4% to 37.8%. However, no significant difference was observed for the Cellule and Hyfi cultivars, with the protection efficiency exceeding 57% (Figure 2).




Figure 2. Protection efficiency induced by *Paenibacillus* sp. strain B2 (PB2), Mix 2, composed of PB2 and *Arthrobacter agilis*, and Mix 3, composed of PB2, *Microbacterium* sp. strain SSM1, and *Arthrobacter* sp. SSM4 PB2, against *M. graminicola*, strain IPO323 in non-sterile field soil, in four wheat cultivars (Alixan, Altigo, Cellule, Hyfi) at the 3-leaf growth stage, represented as percent reduction in *M. graminicola* β -tubulin copy number in 100 ng leaf DNA (BCN_{100ng} DNA) extracted from the third leaf (L3) 17 days after infection with *M. graminicola*. Plants were inoculated with Mix 2, Mix 3, and PB2 by dipping pre-germinated seeds in a suspension of 10^6 CFU·mL⁻¹. The dotted line represents the minimal protection efficiency accepted and determined by 40%. Values shown are means with SD ($n = 5$). Different lower-case letters indicate significant differences between groups according to Tukey's test at $p \leq 0.05$.

Regarding Mix 2, only the cultivar Hyfi displayed a significant decrease in protection, decreasing from 77.5% in sterile soil to 47.9% in non-sterile soil. However, in sterile soil, the protection efficiency ranged from 62.4% to 83%, and in non-sterile soil, it ranged from 47.9% to 63.6%, with no significant difference among cultivars (Figure 2).

In sterile soil, mix 3 exhibited high and stable protection efficiency across all cultivars, with values exceeding 73.9%. However, under non-sterile soil conditions, Altigo and Hyfi maintained protection efficiencies of over 55% without significant decreases compared to sterile soil. In contrast, Alixan and Cellule experienced significant reduction, with protection efficiencies of 33% and 20%, respectively (Figure 2).

3.3. Influence of PGPR Inoculation on Microbial Communities

3.3.1. α -Diversity

The impact of PGPR inoculation, whether alone or in co-inoculation, on rhizospheric and endophytic bacterial communities of two wheat cultivars (Alixan and Cellule) was assessed using three PGPR modalities, PB2, Mix 2, and Mix 3, and compared with non-inoculated plants.

Permutative multivariate analysis of variance (PermANOVA) revealed that the α -diversity of the bacterial and fungal communities under investigation were primar-

ily influenced by the type of compartment (rhizospheric vs. endophytic, $F = 115.92$, $p < 0.001$ and $F = 544.47$, $p < 0.001$, respectively, Table S1). Among the bacterial and fungal communities, the α -diversity of these communities was found to be significantly greater in the rhizosphere, characterized by a higher diversity and species richness compared to the roots. No significant differences in species richness were observed in either rhizospheric or endophytic bacterial communities between the two cultivars or following PGPR inoculation, except for a reduction in species richness in the rhizospheric community of Alixan, which occurred after PB2 inoculation (Table 1).

Table 1. Estimation of the richness and diversity (Shannon index) of 16S rRNA gene libraries from sequencing analysis. Pre-germinated seeds were immersed in a PGPR suspension of 10^6 CFU·mL⁻¹. The genetic structure of bacterial communities was observed 7 days after sowing in an agricultural field soil. Values shown are means with SD ($n = 3$). Different lower-case letters indicate significant differences between groups according to Kruskal–Wallis test at $p \leq 0.05$.

Analysis Target	Cultivar	Modality	Richness			Shannon		
			Mean	SD	Sign	Mean	SD	Sign
Rhizosphere	Alixan	Non-inoculated	2454.7	222.23	b	6.435	0.01	f
		PB2	1776.7	442.10	c	6.134	0.27	bcf
		Mix2	2664.3	293.14	b	6.298	0.16	f
	Cellule	Mix3	2445.0	846.29	b	6.273	0.14	cf
		Non-inoculated	2904.3	399.74	b	6.316	0.01	f
		PB2	2735.7	225.90	b	6.330	0.11	f
		Mix2	2793.0	146.43	b	6.271	0.12	cf
		Mix3	2583.3	605.81	b	6.258	0.14	cf
Endosphere	Alixan	Non-inoculated	848.0	173.35	a	5.626	0.08	ad
		PB2	606.7	524.89	a	5.242	0.52	de
		Mix2	1069.3	178.21	a	5.922	0.08	abc
	Cellule	Mix3	820.7	350.44	a	5.820	0.27	ab
		Non-inoculated	818.5	95.46	a	5.681	0.07	a
		PB2	533.0	137.96	a	5.172	0.31	e
		Mix2	977.0	327.11	a	5.828	0.21	ab
		Mix3	670.3	422.44	a	5.649	0.35	a

The richness of fungal species in the rhizosphere was similar in the two wheat cultivars, Alixan and Cellule. While the richness of the Cellule cultivar remained unaffected by the inoculations, the richness of the Alixan cultivar was reduced by the inoculation with Mix 2 and Mix 3. Conversely, no effect of inoculation or cultivar type on endophytic species richness was detected (Table 2).

The Shannon index of the bacterial community in the rhizosphere also showed no difference between the two cultivars, whether under non-inoculated or PGPR-inoculated conditions. This similarity extended to the endophytic communities as well, except for Cellule plants inoculated with PB2, where a significantly lower index was compared to non-inoculated wheat plants (Table 1). The Shannon index of rhizospheric fungal communities was not influenced by the cultivar type or inoculations, except for those with PB2, which exhibited a lower value than in the non-inoculated situation (for both cultivar types). In the roots, this index remained unaffected by any modality (Table 2).

The bacterial communities in the rhizosphere of non-inoculated wheat were predominantly composed of *Proteobacteria*, which collectively accounted for 43% of the bacterial species in the rhizosphere. Other dominant phyla in this bacterial community included *Patescibacteria* (16%), *Acidobacteria* (15%), *Actinobacteria* (7%), *Planctomycetes* (6%), and *Chloroflexi* (5%).

The endophytic communities were primarily characterized by an enrichment of *Proteobacteria* (46%), along with notable representations of *Actinobacteria* (29%) and *Chloroflexi* (13%, Figure 3).

Table 2. Estimation of richness and diversity (Shannon index) of ITS gene libraries from sequencing analysis. Pre-germinated seeds were immersed in a PGPR suspension of 10^6 CFU·mL⁻¹. The genetic structure of fungal communities was observed 7 days after sowing in an agricultural field soil. Values shown are means with SD ($n = 3$). Different lower-case letters indicate significant differences between groups according to the Kruskal–Wallis test at $p \leq 0.05$.

Analysis Target	Cultivar	Modality	Richness			Shannon		
			Mean	SD	Sign	Mean	SD	Sign
Rhizosphere	Alixan	Non-inoculated	270.67	13.32	c	4.34	0.18	d
		PB2	241.00	18.52	bc	3.22	1.29	bf
		Mix2	217.67	13.05	b	4.16	0.34	de
		Mix3	223.67	23.12	b	3.80	0.63	def
	Cellule	Non-inoculated	246.67	33.08	bc	4.37	0.06	d
		PB2	239.00	7.81	b	3.30	1.15	bef
		Mix2	233.00	8.89	b	4.16	0.15	de
		Mix3	237.67	46.69	b	4.12	0.29	def
Endosphere	Alixan	Non-inoculated	26.67	9.45	a	2.15	0.03	ac
		PB2	25.33	15.28	a	2.15	0.35	ac
		Mix2	28.33	6.35	a	2.74	0.14	ab
		Mix3	17.33	6.35	a	2.11	0.70	ac
	Cellule	Non-inoculated	23.00	8.49	a	2.55	0.42	abc
		PB2	23.67	2.31	a	1.77	0.41	c
		Mix2	28.00	8.89	a	2.61	0.29	abc
		Mix3	18.00	11.27	a	2.01	0.46	ac




Figure 3. Relative abundance of bacterial 16 rRNA sequences of the different phyla for each modality. The abundance is expressed as a percentage and was calculated using the values of 16S rRNA copy numbers and nearest genome sizes for each bacterial OTU (Operational Taxonomic Unit) in the rhizosphere and endophyte compartments of Alixan and Cellule cultivars inoculated with *Paenibacillus* sp. strain B2 (PB2), Mix 2, composed of PB2 and *Arthrobacter agilis*, or Mix 3, composed of PB2, *Microbacterium* sp. strain SSM1 and *Arthrobacter* sp. SSM4 PB2, and non-inoculated control. Pre-germinated seeds were immersed in a PGPR suspension with a final concentration of 10^6 CFU·mL⁻¹. The genetic structure of the bacterial communities was observed 7 days after sowing in an agricultural field soil.

PermANOVA analysis clearly showed that the structure of the bacterial community was strongly associated with the community’s origin (rhizospheric or endophytic, $F = 544.48, p < 0.001$). Additionally, it was influenced by the cultivar ($F = 9.68, p = 0.005$) and,

to a lesser extent, by the type of inoculation ($F = 3.86$, $p = 0.017$, Table S1). Combinations that included both the origin of the bacterial community and the inoculation ($F = 3.14$, $p = 0.036$), as well as combinations involving all three factors (origin, inoculation, and variety, $F = 4.06$, $p = 0.008$), also showed significant effects on community structuring.

The fungal community was primarily composed of *Ascomycota*, *Basidiomycota*, and *Zygomycota*, with the cultivar type not having a significant influence. However, perMANOVA analysis (Table S1) revealed that the proportions of these different phyla were strongly influenced by the origin of this community (rhizospheric or endophytic). Inoculation with PGPR had an impact on the fungal community by increasing the presence of *Ascomycota* at the expense of *Basidiomycota*. This effect was more pronounced in co-inoculations in the rhizosphere. In the roots, a similar response was observed, but it was the *Zygomycota* that were most affected by the inoculations (Figure 4).




Figure 4. Relative abundance of fungal ITS sequences of different phyla for each modality. The abundance is expressed as a percentage and was calculated using the values of ITS copy numbers and genome sizes of the closest hits to each fungal OTU (Operational Taxonomic Unit), in the rhizosphere and endophyte compartments of Alixan and Cellule cultivars inoculated with *Paenibacillus* sp. strain B2 (PB2), Mix 2, composed of PB2 and *Arthrobacter agilis*, or Mix 3, composed of PB2, *Microbacterium* sp. strain SSM1, and *Arthrobacter* sp. SSM4 PB2, and non-inoculated control. Pre-germinated seeds were immersed in a PGPR suspension with a final concentration of 10^6 CFU·mL⁻¹. The genetic structure of the fungal communities was observed 7 days after sowing in an agricultural field soil.

3.3.2. β -Diversity

The non-metric multidimensional scaling (NMDS) analysis (stress = 0.126) revealed that rhizospheric bacterial communities were distributed along two axes. The first axis was characterized by the relative abundances of *Proteobacteria* genera such as *Pseudomonas*, *Pseudorhodobacter*, *Rhizobium*, *Rhodobacter*, and *Skermanella*, while the second axis was associated with the relative abundances of certain genera of *Actinobacteria* and *Patescibacteria*, evolving in opposite directions. Inoculation with PB2 primarily influenced bacterial communities in the rhizosphere along axis 1, with more pronounced effects observed in Alixan plants (Figure 5A). Inoculation with Mix 2 also impacted bacterial communities in the rhizosphere of both cultivars, primarily along axis 2. In contrast, inoculation with Mix 3 showed no significant effect on rhizospheric communities.




Figure 5. Non-metric multidimensional scaling (NMDS) of microbial communities. Rhizospheric (A) and endophytic (B) bacterial communities. The 50% most abundant bacterial genera with 30% best axis fit are shown and colored according to their phylum (*Acidobacteria* = blue, *Actinobacteria* = black, *Bacteroidetes* = yellow, *Chloroflexi* = green, *Patescibacteria* = violet, and *Proteobacteria* = red). Rhizospheric (C) and endophytic (D) fungal communities. The 50% most abundant bacterial genera with 30% best axial fit are shown and colored according to their phylum (*Ascomycota* = red, *Basidiomycota* = blue, *Chytridiomycota* = gray, and *Zygomycota* = orange). PB2, *Paenibacillus* sp. strain B2 (PB2); Mix 2, PGPR mixture composed of PB2 and *Arthrobacter agilis*; Mix 3, PGPR mixture composed of PB2, *Microbacterium* sp. strain SSM1, and *Arthrobacter* sp. SSM4.

The distribution of endophytic bacterial communities was characterized by two axes in the NMDS ordination (stress = 0.116, Figure 5B). Axis 1 was defined by the relative abundance of various genera of *Proteobacteria*, while axis 2 was associated with *Actinobacteria* and *Patescibacteria*. Inoculation with PB2 affected the bacterial community along axis 1 when applied to the cultivar Cellule, while the community shifted along axis 2 in the case of the cultivar Alixan. Inoculations with Mix 2 and Mix 3 also led to changes in endophytic communities, with the direction of evolution along axis 1 being opposite to that observed with PB2.

In the NMDS analysis (stress = 0.265) of the endophytic fungal community structures in the rhizosphere of both cultivars, their similarity was evident (Figure 5C). Inoculation with both cultivars with PB2 slightly influenced these communities along axis 2, with the cultivar Alixan exhibiting more notable changes. Inoculation with Mix 2 also resulted in a shift along axis 2 of the ordination, with a very similar community composition between the two cultivars, characterized by the relative abundances of *Peziza*, *Spizellomyces*, and *Trametes*. The change in relative abundances observed in Alixan, following inoculation with Mix 3, resulted in a fungal genus assemblage that resembled a mixture of the responses observed in plants inoculated with PB2 and those inoculated with Mix 2.

Conversely, the inoculation of the cultivar Cellule with Mix 3 had a pronounced impact on the rhizospheric community along axis 1. This shift was characterized by the relative abundance of the genera such as *Alternaria*, *Rhizoctonia*, and *Reddellomyces*. Examination of the endophytic communities (stress = 0.218) revealed that, despite significant differences between the two cultivars, PGPR inoculations yielded very similar communities (Figure 5D).

3.3.3. Microbial Community Function

While the number of substrates metabolized by the microbial communities in the rhizosphere remained unaffected by cultivar type or bacterial inoculations (Figure 6A), the metabolic activity, as indicated by the average well color development (AWCD), was notably higher in the rhizosphere communities of the Cellule cultivar compared to Alixan cultivar (Figure 6B). In both genotypes, inoculation with PB2 led to enhanced metabolic capabilities in the rhizospheric communities for utilizing various EcoPlate substrates. In the Alixan cultivar, this metabolic activity was unaffected by the Mix 2 inoculation, but PB2 and Mix 3 inoculations led to increased metabolic activity. These changes were characterized by the increased utilization of carbohydrates, amino acids, and amines/amides, alongside the decreased utilization of carboxylic and acetic acids (Figure 6C). In contrast, the metabolic activity of the rhizosphere communities in the Cellule cultivar was solely increased by PB2 inoculation, which was linked to an elevated degradation of carbohydrates.




Figure 6. Metabolic potential of rhizospheric soil communities. Estimation of functional richness (A), average well color development (AWCD) expressed in OmniLog units (B), and principal component analysis (PCA) of carbon substrates metabolization (C) in the rhizospheric soil communities of Alixan and Cellule cultivars inoculated with *Paenibacillus* sp. strain B2 (PB2), Mix 2, composed of PB2 and *Arthrobacter agilis*, or Mix 3, composed of PB2, *Microbacterium* sp. strain SSM1, and *Arthrobacter* sp. SSM4 PB2, and non-inoculated control. Pre-germinated seeds were dipped in a PGPR suspension with a final concentration of 10^6 CFU·mL⁻¹. Values presented are means with SD ($n = 3$). Different lowercase letters indicate significant differences between groups according to the Kruskal–Wallis test at $p \leq 0.05$.

4. Discussion

To successfully employ PGPR strains as biofertilizer and biocontrol agents, it is essential to establish a beneficial relationship between the inoculants and the plants. It is widely

acknowledged that PGPR must interact with plants to influence plant physiology [32]. This interaction depends on abiotic factors, such as physicochemical characteristics of soil, and biotic factors such as native microbial communities [9].

In our previous publications [5,10,16], we observed that PB2, Mix 2, and Mix 3, in sterile soil conditions, significantly promoted growth and induced resistance, with some variability based on the cultivar. In the current study, conducted in non-sterile field soils, PGPR strains, in general, maintained their activities in promoting wheat growth and protecting against *M. graminicola*. These effects were cultivar-dependent, and specifically, the Mix 2 treatments resulted in increased dry biomass of roots and leaves across all four wheat cultivars tested. The timing of PGPR inoculation is an important factor to consider in enabling the assimilation of the inoculating microbes into the plant rhizosphere. In our previous studies, we demonstrated that an additional inoculation with these PGPR one or two weeks after the initial one did not provide any additional benefits for wheat protection or growth promotion. However, the introduced PGPR, whether through seed incubation in a bacterial suspension for one hour or via seed coating, were detected in both ecto- and endo-roots across all wheat growth stages using specific primers [5,6,10,16]. As described in several studies, introducing PGPR as early as possible into the indigenous microbial population supports their establishment and may, in part, account for the sustained activities of PB2, Mix 2, and Mix 3 under non-sterile conditions [33,34]. Furthermore, coating the seed with the endophytic PB2, AA, SSM-001, and SSM-004 bacteria fosters the development of selected communities of beneficial bacteria within the germination structures, representing one of the earliest colonization events [35].

However, our previous results in sterile soil indicated that an increase in leaf biomass was only observed following Mix 2 inoculation in Altigo and Cellule cultivars, with no increase noted with PB2 alone [5,10]. These results suggest that interactions between PGPR and microbial communities positively impact wheat growth, as confirmed by the protection conferred by PB2 ($\geq 37.8\%$) and Mix 2 ($\geq 47.9\%$) against *M. graminicola* across all four cultivars tested in non-sterile field soils. Positive interactions (commensalism, mutualism, and synergism) can enable the functioning of a population [36]. Moreover, our study revealed that the genotype effect observed under sterile soil for Mix 2 on wheat growth promotion and protection against *M. graminicola* did not persist in non-sterile field soils. Nonetheless, our understanding of the factors influencing the rhizosphere competence of PGPR inoculants remains limited, and only a few studies have reported positive interactions between native microbial communities and bacterial inoculants [8,37,38].

In contrast, Mix 3 exhibited a decline in its plant growth-promoting activities and offered limited protection against *M. graminicola* compared to its performance in sterile soil [16]. However, the activity of most of PGPR strains tends to decrease in non-sterile soils when compared to sterile soils, as demonstrated by Gholami et al. (2009) [39] in maize with *P. putida* strain R-168, *P. fluorescens* strain R-93, *P. fluorescens* DSM 50090, *P. putida* DSM291, *A. lipoferum* DSM 1691, and *A. brasilense* DSM 1690. Negative interactions between introduced microbial inoculants and native microbial communities may result in the exclusion of the inoculant from the indigenous community [40,41] or trigger various allelopathic events [42,43]. Additionally, native microbial communities often represent highly competitive communities that have adapted to their environment, which can influence the biofertilization and biocontrol activities of the bacterial inoculants [8]. Moreover, the efficacy of PGPR inoculation is contingent upon the indigenous microbial communities and the specific soil used [9,44]. These factors, in turn, are influenced by root morphology, the growth stage of the plant, soil physical and chemical properties, and root exudates [36].

Ideally, microbial inputs used as biostimulants or biocontrol agents should have minimal or controlled effects on the environment. This includes factors like their dispersal, persistence, microbial function, and cycling [45]. A major concern is the impact of introduced microorganisms on the existing microbiome, which may result from direct ecological interactions, such as competition or inhibition [46]. Introduced microorganisms can indirectly affect native rhizospheric microbial communities by altering plant traits and root

exudation [47], and they can even influence the innate endophytic community by modulating plant responses [48]. In this study, both the rhizospheric and endophytic bacterial and fungal communities exhibited high diversity, which probably reflects the development of the plant under non-limiting conditions [20,49]. Although PGPR inoculation did not significantly impact the diversity of rhizospheric or endophytic bacterial communities, the taxonomic composition of these communities notably changed in response to PB2 inoculation. This shift in bacterial rhizospheric communities may be attributed to the production of paenimycin by PB2 [11], which has antagonistic effects against some Gram-positive and Gram-negative bacteria and certain fungi [11,15]. In fact, PB2 influenced the structuring of rhizospheric communities, promoting a more beneficial community with specific *Proteobacteria*, including *Pseudomonas*, *Pseudorhodobacter*, *Rhizobium*, *Rhodobacter*, and *Skermanella*; these bacteria are often recognized as PGPR themselves [50–53]. Furthermore, the maintenance of high endophytic bacterial diversity in inoculated plants suggests the preservation or enhancement of specific endophytes within the plant in response to bacterial inoculation [54,55]. In our study, the taxonomic changes observed in endophytic communities were primarily due to PB2 inoculation, which redirected the community towards a higher abundance of *Proteobacteria*. Fungal diversity was slightly reduced in the rhizosphere of both cultivars upon inoculation with PB2, resulting in the promotion of saprophytic fungi, such as *Peziza* [56], some of which may also contribute to improved plant performance [57].

The presence of *Arthrobacter agilis* 4042 in combination with PB2 (Mix 2) restricted the impact of PB2 to bacterial communities only and fostered an alternative bacterial rhizospheric structure primarily characterized by methylotrophic bacteria. Methylotrophic bacteria are well-known for their positive effects on plant growth [58] and their association with abiotic stress tolerance [59]. This effect was mirrored in the endophytic bacterial community, where although diversity remained unaffected, it maintained a distribution similar to that in the non-inoculated plants. When *Arthrobacter* sp. SSM-004 and *Microbacterium* sp. SSM-001 were combined with PB2 (Mix 3), it further constrained the effect of PB2 in shaping a more favorable bacterial community. Mix 3 also impacted rhizospheric fungi, resulting in a higher abundance of some plant pathogenic fungi, such as the genera *Alternaria* and *Rhizoctonia* [60,61]. This alteration was also observed in the endophytic community, with the induction of fungal strains of the genus *Coniothyrium*. Some strains from this genus can be used to control fungal diseases [62–64].

Moreover, the taxonomic changes observed after the bacterial inoculations did not affect the functionality of the soil. The high diversity observed at both rhizospheric and endophytic levels suggests that functional redundancy is able to compensate for these taxonomic changes [20]. Finally, our study primarily focused on the influence of PGPR inoculation 7 days after sowing, but microbial communities also depend on the growth stage of wheat. The influence of microbial communities on PGPR activities and vice versa varies with the sampling time, soil properties, plant development stage, and microbial strains used [65]. Further analyses conducted at various stages of wheat growth would offer a better understanding of the long-term effectiveness of PGPR. Additionally, it is essential to conduct a comprehensive investigation into the underlying mechanisms of positive and negative interactions between introduced microbial inoculants and native microbial communities, which is indeed the focus of our future research projects.

5. Conclusions

The present study highlighted the impact of indigenous soil microbial communities on the efficacy of PGPR inoculation using pre-germinated seeds in non-sterile agricultural cropland. Inoculation with Mix 2 demonstrated positive interactions, enhancing both PGPR biofertilization and biocontrol activities. Importantly, these positive effects were observed across different wheat genotypes, unlike the outcomes with Mix 3. The inoculation of PGPR had a significant influence on shaping bacterial and fungal communities, aiming to maximize their potential benefits for both the soil and the plant. Notably, the presence of

Arthrobacter spp. Within Mix 2 seemed to mitigate any potential negative effects of PB2, favoring root colonization by other beneficial bacteria known for their positive contributions to plant growth and defense mechanisms against abiotic stress.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/biology12111416/s1>. Table S1: Permutative multivariate analysis of variance (PerMANOVA) of the bacterial and fungal communities.

Author Contributions: E.S., S.S., C.M. and J.D. carried out the experimental work. E.S. wrote the manuscript. S.S., J.D. and M.E. revised the manuscript. J.D., V.S. and F.D. analyzed the microbial genetic diversity. S.S., T.A., C.E. and E.A.B. managed the experimental design and the PhD research program and S.S., T.A. and C.E. supervised the project. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the SDP society and the ANRT “Association Nationale de la Recherche et de la Technologie” under the convention N° 56/2016 through doctoral fellowships (E.S.) and by the European Regional Development Fund (ERDF) under the ERDF/FSE 2014–2020 PhD grant (C.M.).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The raw sequence data for 16S_{V3-V4} rRNA and ITS region are available at figshare: <https://doi.org/10.6084/m9.figshare.24496414> (accessed on 13 September 2022).

Acknowledgments: We are grateful to SDP for financing this work, with the support of the French ‘Association National Recherche Technologie (ANRT)’, under convention N° 56/2016.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Nihorimbere, V.; Ongena, M.; Smargiassi, M.; Thonart, P. Beneficial Effect of the Rhizosphere Microbial Community for Plant Growth and Health. *Biotechnol. Agron. Soc. Environ.* **2011**, *15*, 327–337.
2. Selim, S.; Roisin-Fichter, C.; Andry, J.-B.; Bogdanow, B.; Sambou, R. Real-Time PCR to Study the Effect of Timing and Persistence of Fungicide Application and Wheat Varietal Resistance on *Mycosphaerella graminicola* and Its Sterol 14 α -Demethylation-Inhibitor-Resistant Genotypes. *Pest Manag. Sci.* **2014**, *70*, 60–69. [[CrossRef](#)] [[PubMed](#)]
3. Selim, S.; Jean-Baptiste, A.; Boris, B.; Céline, R.-F. Accuracy of Real-Time PCR to Study *Mycosphaerella Graminicola* Epidemic in Wheat: From Spore Arrival to Fungicide Efficiency. In *Fungicides—Beneficial and Harmful Aspects*; INTECH Open Access Publisher: London, UK, 2011; Volume 12, pp. 219–238.
4. Benhamou, N. Elicitor-Induced Plant Defence Pathways. *Trends Plant Sci.* **1996**, *1*, 233–239. [[CrossRef](#)]
5. Samain, E.; van Tuinen, D.; Jeandet, P.; Aussenac, T.; Selim, S. Biological Control of Septoria Leaf Blotch and Growth Promotion in Wheat by *Paenibacillus* Sp. Strain B2 and *Curtobacterium Plantarum* Strain EDS. *Biol. Control* **2017**, *114*, 87–96. [[CrossRef](#)]
6. Samain, E.; Aussenac, T.; Selim, S. The Effect of Plant Genotype, Growth Stage, and *Mycosphaerella graminicola* Strains on the Efficiency and Durability of Wheat-Induced Resistance by *Paenibacillus* Sp. Strain B2. *Front. Plant Sci.* **2019**, *10*, 587. [[CrossRef](#)]
7. Walters, D.; Walsh, D.; Newton, A.; Lyon, G. Induced Resistance for Plant Disease Control: Maximizing the Efficacy of Resistance Elicitors. *Phytopathology* **2005**, *95*, 1368–1373. [[CrossRef](#)]
8. Van Veen, J.A.; Overbeek, L.S.V.; Elsas, J.D.V. Fate and Activity of Microorganisms Introduced into Soil. *Microbiol. Mol. Biol. Rev.* **1997**, *61*, 15.
9. Ramos, B.; García, J.A.L.; Probanza, A.; Barrientos, M.L.; Gutierrez Mañero, F.J. Alterations in the Rhizobacterial Community Associated with European Alder Growth When Inoculated with PGPR Strain *Bacillus licheniformis*. *Environ. Exp. Bot.* **2003**, *49*, 61–68. [[CrossRef](#)]
10. Samain, E.; Ernenwein, C.; Aussenac, T.; Selim, S. Effective and Durable Systemic Wheat-Induced Resistance by a Plant-Growth-Promoting Rhizobacteria Consortium of *Paenibacillus* Sp. Strain B2 and *Arthrobacter* Spp. Strain AA against *Zymoseptoria Tritici* and Drought Stress. *Physiol. Mol. Plant Pathol.* **2022**, *119*, 101830. [[CrossRef](#)]
11. Selim, S.; Negrel, J.; Govaerts, C.; Gianinazzi, S.; van Tuinen, D. Isolation and Partial Characterization of Antagonistic Peptides Produced by *Paenibacillus* sp. Strain B2 Isolated from the Sorghum Mycorrhizosphere. *Appl. Environ. Microbiol.* **2005**, *71*, 6501–6507. [[CrossRef](#)]

12. Cattelan, A.J.; Hartel, P.G.; Fuhrmann, J.J. Screening for Plant Growth–Promoting Rhizobacteria to Promote Early Soybean Growth. *Soil Sci. Soc. Am. J.* **1999**, *69*, 1670–1680. [CrossRef]
13. Weller, D.M. Biological Control of Soilborne Plant Pathogens in the Rhizosphere with Bacteria. *Annu. Rev. Phytopathol.* **1988**, *26*, 379–407. [CrossRef]
14. Jones, M.L.; Rivett, D.W.; Pascual-García, A.; Bell, T. Relationships between Community Composition, Productivity and Invasion Resistance in Semi-Natural Bacterial Microcosms. *eLife* **2021**, *10*, e71811. [CrossRef] [PubMed]
15. Selim, S.; Martin-Laurent, F.; Rouard, N.; Gianinazzi, S.; van Tuinen, D. Impact of a New Biopesticide Produced by *Paenibacillus* Sp. Strain B2 on the Genetic Structure and Density of Soil Bacterial Communities. *Pest Manag. Sci.* **2007**, *63*, 269–275. [CrossRef]
16. Samain, E.; Ernenwein, C.; Aussenac, T.; Selim, S. Efficacy and Durability of *Paenibacillus* Sp. Strain B2 in Co-Inoculation with *Arthrobacter* Sp. SSM-004 and *Microbacterium* Sp. SSM-001 for Growth Promoting and Resistance Induction in Wheat against *Mycosphaerella Graminicola* and Drought Stress. *Plant Pathol. Microbiol.* **2022**, *13*, 1000603.
17. Budi, S.W.; Tuinen, D.V.; Martinotti, G.; Gianinazzi, S. Isolation from the Sorghum Bicolor Mycorrhizosphere of a Bacterium Compatible with Arbuscular Mycorrhiza Development and Antagonistic towards Soilborne Fungal Pathogens. *Appl. Environ. Microbiol.* **1999**, *65*, 5148–5150. [CrossRef]
18. Arvalis Institut du Végétal Blé Tendre—Maladies; Comportement Vis-à-Vis Des Maladies. *Choisir* **2017**. Available online: https://www.arvalis.fr/sites/default/files/imported_files/premiers_resultats_bth_alsace_1263678181362291022.pdf (accessed on 3 November 2023).
19. Bearchell, S.J.; Fraaije, B.A.; Shaw, M.W.; Fitt, B.D. Wheat Archive Links Long-Term Fungal Pathogen Population Dynamics to Air Pollution. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 5438–5442. [CrossRef]
20. Alahmad, A.; Decocq, G.; Spicher, F.; Kheirbeik, L.; Kobaissi, A.; Tetu, T.; Dubois, F.; Duclercq, J. Cover Crops in Arable Lands Increase Functional Complementarity and Redundancy of Bacterial Communities. *J. Appl. Ecol.* **2019**, *56*, 651–664. [CrossRef]
21. Herlemann, D.P.; Labrenz, M.; Jürgens, K.; Bertilsson, S.; Waniek, J.J.; Andersson, A.F. Transitions in Bacterial Communities along the 2000 Km Salinity Gradient of the Baltic Sea. *ISME J.* **2011**, *5*, 1571–1579. [CrossRef]
22. Toju, H.; Tanabe, A.S.; Yamamoto, S.; Sato, H. High-Coverage ITS Primers for the DNA-Based Identification of Ascomycetes and Basidiomycetes in Environmental Samples. *PLoS ONE* **2012**, *7*, e40863. [CrossRef] [PubMed]
23. Escudíe, F.; Auer, L.; Bernard, M.; Mariadassou, M.; Cauquil, L.; Vidal, K.; Maman, S.; Hernandez-Raquet, G.; Combes, S.; Pascal, G. FROGS: Find, Rapidly, OTUs with Galaxy Solution. *Bioinformatics* **2018**, *34*, 1287–1294. [CrossRef] [PubMed]
24. Martin, M. Cutadapt Removes Adapter Sequences from High-Throughput Sequencing Reads. *EMBnet J.* **2011**, *17*, 10–12. [CrossRef]
25. Mahé, F.; Rognes, T.; Quince, C.; de Vargas, C.; Dunthorn, M. Swarm: Robust and Fast Clustering Method for Amplicon-Based Studies. *PeerJ* **2014**, *2*, e593. [CrossRef]
26. Rognes, T.; Flouri, T.; Nichols, B.; Quince, C.; Mahé, F. VSEARCH: A Versatile Open Source Tool for Metagenomics. *PeerJ* **2016**, *18*, e2584. [CrossRef]
27. Deshpande, V.; Wang, Q.; Greenfield, P.; Charleston, M.; Porrás-Alfaro, A.; Kuste, C.R.; Cole, J.R.; Midgley, D.; Tran-Dinh, N. Fungal Identification Using a Bayesian Classifier and the Warcup Training Set of Internal Transcribed Spacer Sequences. *Mycologia* **2015**, *108*, 1–5. [CrossRef]
28. Langrand, J.; Lounès-Hadj Sahraou, A.; Duclercq, J.; Raveau, R.; Laruelle, F.; Bert, V.; Fontaine, J. Coriander (*Coriandrum sativum*) Cultivation Combined with Arbuscular Mycorrhizal Fungi Inoculation and Steel Slag Application Influences Trace Elements-Polluted Soil Bacterial Functioning. *Plants* **2023**, *12*, 618. [CrossRef]
29. Weber, K.P.; Grove, J.A.; Gehder, M.; Anderson, W.A.; Legge, R.L. Data Transformations in the Analysis of Community-Level Substrate Utilization Data from Microplates. *J. Microbiol. Methods* **2007**, *69*, 461–469. [CrossRef]
30. Oksanen, J.; Blanchet, F.G.; Kindt, R.; Legendre, P.; Minchin, P.R.; O’hara, R.B. Community Ecology Package. *R Package Version* **2013**, *2*, 1–295.
31. Mendiburu, F. De Agricolae: Statistical Procedures for Agricultural Research. 2021. Available online: <https://rdr.io/cran/agricolae/> (accessed on 13 September 2022).
32. Chanway, C.P.; Turkington, R.; Holl, F.B. Ecological Implications of Specificity between Plants and Rhizosphere Micro-Organisms. *Adv. Ecol. Res.* **1991**, *21*, 121–169. [CrossRef]
33. Hastings, A.; Harrison, S. Metapopulation Dynamics and Genetics. *Ann. Rev. Ecol. Syst.* **1994**, *25*, 167–188. [CrossRef]
34. Levins, R. Extinction. *Lect. Math. Life Sci.* **1976**, *2*, 75–107.
35. Sturz, A.V.; Novak, J. Endophytic communities of Rhizobacteria and the Strategies Required to Create Yield Enhancing Associations with Crops. *Appl. Soil Ecol.* **2000**, *15*, 183–190. [CrossRef]
36. Trabelsi, D.; Mhamdi, R. Microbial Inoculants and Their Impact on Soil Microbial Communities: A Review. *BioMed Res. Int.* **2013**, *2013*, 863240. [CrossRef] [PubMed]
37. Ciccillo, F.; Fiore, A.; Bevivino, A.; Dalmastri, C.; Tabacchioni, S.; Chiarini, L. Effects of Two Different Application Methods of Burkholderia Ambifaria MCI 7 on Plant Growth and Rhizospheric Bacterial Diversity. *Environ. Microbiol.* **2002**, *4*, 238–245. [CrossRef] [PubMed]
38. Schreiter, S.; Ding, G.-C.; Heuer, H.; Neumann, G.; Sandmann, M.; Grosch, R.; Kropf, S.; Smalla, K. Effect of the Soil Type on the Microbiome in the Rhizosphere of Field-Grown Lettuce. *Front. Microbiol.* **2014**, *5*, 144. [CrossRef]

39. Gholami, A.; Shahsavani, S.; Nezarat, S. The Effect of Plant Growth Promoting Rhizobacteria (PGPR) on Germination, Seedling Growth and Yield of Maize. *Int. J. Biol. Life Sci.* **2009**, *1*, 35–40.
40. Backer, R.; Rokem, J.S.; Ilangumaran, G.; Lamont, J.; Praslickova, D.; Ricci, E.; Subramanian, S.; Smith, D.L. Plant Growth-Promoting Rhizobacteria: Context, Mechanisms of Action, and Roadmap to Commercialization of Biostimulants for Sustainable Agriculture. *Front. Plant Sci.* **2018**, *9*, 1473. [[CrossRef](#)]
41. Nelson, L.M. Plant Growth Promoting Rhizobacteria (PGPR): Prospects for New Inoculants. *Crop Manag.* **2004**, *3*, 301–305. [[CrossRef](#)]
42. Sturz, A.V.; Christie, B.R. Endophytic Bacteria of Red Clover as Agents of Allelopathic Clover-Maize Syndromes. *Soil Biol. Biochem.* **1996**, *28*, 583–588. [[CrossRef](#)]
43. Sturz, A.V.; Christie, B.R. Endophytic Bacterial Systems Governing Red Clover Growth and Development. *Ann. Appl. Biol.* **1995**, *126*, 285–290. [[CrossRef](#)]
44. Gagliardi, J.V.; Buyer, J.S.; Angle, J.S.; Russek-Cohen, E. Structural and Functional Analysis of Whole-Soil Microbial Communities for Risk and Efficacy Testing Following Microbial Inoculation of Wheat Roots in Diverse Soils. *Soil Biol.* **2001**, *33*, 25–40. [[CrossRef](#)]
45. Bashan, Y.; Kloepper, J.W.; De-Bashan, L.E.; Nannipieri, P. A Need for Disclosure of the Identity of Microorganisms, Constituents, and Application Methods When Reporting Tests with Microbe-Based or Pesticide-Based Products. *Biol. Fertil. Soils* **2016**, *52*, 283–284. [[CrossRef](#)]
46. Mawarda, P.C.; Le Roux, X.; Van Elsland, J.D.; Salles, J.F. Deliberate Introduction of Invisible Invaders: A Critical Appraisal of the Impact of Microbial Inoculants on Soil Microbial Communities. *Soil Biol. Biochem.* **2020**, *148*, 107874. [[CrossRef](#)]
47. Mitter, E.K.; Tosi, M.; Obregón, D.; Dunfield, K.E.; Germida, J.J. Rethinking Crop Nutrition in Times of Modern Microbiology: Innovative Biofertilizer Technologies. *Front. Sustain. Food Syst.* **2021**, *5*, 606815. [[CrossRef](#)]
48. Ardanov, P.; Sessitsch, A.; Häggman, H.; Kozyrovska, N.; Pirttilä, A.M. Methylobacterium-Induced Endophyte Community Changes Correspond with Protection of Plants against Pathogen Attack. *PLoS ONE* **2012**, *7*, e46802. [[CrossRef](#)]
49. Karimi, B.; Maron, P.A.; Chemidlin-Prevost Boure, N.; Bernard, N.; Gilbert, D.; Ranjard, L. Microbial Diversity and Ecological Networks as Indicators of Environmental Quality. *Environ. Chem. Lett.* **2017**, *15*, 265–281. [[CrossRef](#)]
50. Ajilogba, C.F.; Olanrewaju, O.S.; Babalola, O.O. Plant Growth Stage Drives the Temporal and Spatial Dynamics of the Bacterial Microbiome in the Rhizosphere of Vigna Subterranea. *Front. Microbiol.* **2022**, *13*, 825377. [[CrossRef](#)]
51. Çakmakçı, R.; Dönmez, F.; Aydın, A.; Şahin, F. Growth Promotion of Plants by Plant Growth-Promoting Rhizobacteria under Greenhouse and Two Different Field Soil Conditions. *Soil Biol. Biochem.* **2006**, *38*, 1482–1487. [[CrossRef](#)]
52. Liu, X.; Jiang, X.; He, X.; Zhao, W.; Cao, Y.; Guo, T.; Li, T.; Ni, H.; Tang, X. Phosphate-Solubilizing *Pseudomonas* Sp. Strain P34-L Promotes Wheat Growth by Colonizing the Wheat Rhizosphere and Improving the Wheat Root System and Soil Phosphorus Nutritional Status. *J. Plant Growth Regul.* **2019**, *38*, 1314–1324. [[CrossRef](#)]
53. Yanni, Y.G.; Dazzo, F.B.; Squartini, A.; Zanardo, M.; Zidan, M.I.; Elsadany, A.E.Y. Assessment of the Natural Endophytic Association between Rhizobium and Wheat and Its Ability to Increase Wheat Production in the Nile Delta. *Plant Soil* **2016**, *407*, 367–383. [[CrossRef](#)]
54. Podolich, O.V.; Ardanov, P.; Zaets, I.; Pirttilä, A.M.; Kozyrovska, N. Reviving of the Endophytic Bacterial Community as a Putative Mechanism of Plant Resistance. *Plant Soil* **2015**, *388*, 367–377. [[CrossRef](#)]
55. Podolich, O.V.; Ardanov, P.; Voznyuk, T.M.; Kovalchuk, M.V.; Danylchenko, O.V.; Lashevskiy, V.V.; Lyastchenko, S.A.; Kozyrovska, N.O. Endophytic Bacteria from Potato in Vitro Activated by Exogenic Non-Pathogenic Bacteria. *Biopolym. Cell* **2007**, *23*, 21–27. [[CrossRef](#)]
56. Yin, C.; Schlatter, D.C.; Kroese, D.R.; Paulitz, T.C.; Hagerty, C.H. Responses of Soil Fungal Communities to Lime Application in Wheat Fields in the Pacific Northwest. *Front. Microbiol.* **2021**, *12*, 576763. [[CrossRef](#)]
57. Ou, T.; Zhang, M.; Gao, H.; Wang, F.; Xu, W.; Liu, X.; Wang, L.; Wang, R.; Xie, J. Study on the Potential for Stimulating Mulberry Growth and Drought Tolerance of Plant Growth-Promoting Fungi. *Int. J. Mol. Sci.* **2023**, *24*, 4090. [[CrossRef](#)] [[PubMed](#)]
58. Man, Y.; Wang, J.; Tam, N.F.Y.; Wan, X.; Huang, W.; Zheng, Y.; Yang, Y. Responses of Rhizosphere and Bulk Substrate Microbiome to Wastewater-Borne Sulfonamides in Constructed Wetlands with Different Plant Species. *Sci. Total Environ.* **2020**, *706*, 135955. [[CrossRef](#)] [[PubMed](#)]
59. Kumar, M.; Kour, D.; Yadav, A.N.; Saxena, R.; Rai, P.K.; Jyoti, A.; Tomar, R.S. Biodiversity of Methylophilic Microbial Communities and Their Potential Role in Mitigation of Abiotic Stresses in Plants. *Biologia* **2019**, *74*, 287–308. [[CrossRef](#)]
60. Turzhanova, A.; Khapilina, O.N.; Tumenbayeva, A.; Shevtsov, V.; Raiser, O.; Kalendar, R. Genetic Diversity of *Alternaria* Species Associated with Black Point in Wheat Grains. *PeerJ* **2020**, *8*, e9097. [[CrossRef](#)]
61. Yin, C.; Hulbert, S.H.; Schroeder, K.L.; Mavrodi, O.; Mavrodi, D.; Dhingra, A.; Schillinger, W.F.; Paulitz, T.C. Role of Bacterial Communities in the Natural Suppression of Rhizoctonia Solani Bare Patch Disease of Wheat (*Triticum aestivum* L.). *Appl. Environ. Microbiol.* **2013**, *79*, 7428–7438. [[CrossRef](#)]
62. Jones, E.E.; Rabeendran, N.; Stewart, A. Biocontrol of *Sclerotinia sclerotiorum* Infection of Cabbage by *Coniothyrium minitans* and *Trichoderma* spp. *Biocontrol Sci. Technol.* **2014**, *24*, 1363–1382. [[CrossRef](#)]
63. Ojaghian, M.R. Biocontrol of Potato White Mold Using *Coniothyrium minitans* and Resistance of Potato Cultivars to *Sclerotinia sclerotiorum*. *Plant Pathol. J.* **2010**, *26*, 346–352. [[CrossRef](#)]

64. Sun, Q.; Zhang, P.; Zhao, Z.; Li, X.; Sun, X.; Jiang, W. Continuous Wheat/Soybean Cropping Influences Soybean Yield and Rhizosphere Microbial Community Structure and Function. *Agronomy* **2022**, *13*, 28. [[CrossRef](#)]
65. Kozdrój, J.; Trevors, J.T.; van Elsas, J.D. Influence of Introduced Potential Biocontrol Agents on Maize Seedling Growth and Bacterial Community Structure in the Rhizosphere. *Soil Biol. Biochem.* **2004**, *36*, 1775–1784. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.