



HAL
open science

Deformation and hydrothermal alteration insights of a high strain sheared zone in a gneissic basement (Black Forest, Germany). An analogue for fractured geothermal crystalline reservoirs

Benjamin Avakian, Béatrice Ledéser, Ronan Hebert, Philippe Claeys, Steven Goderis, Ghislain Trullenque, Sébastien Potel

► To cite this version:

Benjamin Avakian, Béatrice Ledéser, Ronan Hebert, Philippe Claeys, Steven Goderis, et al.. Deformation and hydrothermal alteration insights of a high strain sheared zone in a gneissic basement (Black Forest, Germany). An analogue for fractured geothermal crystalline reservoirs. 28e Réunion des Sciences de la Terre, SGF; Géosciences Rennes, Oct 2023, Rennes (FR), France. hal-04311377

HAL Id: hal-04311377

<https://hal.science/hal-04311377v1>

Submitted on 5 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Deformation and hydrothermal alteration insights of a high strain sheared zone in a gneissic basement (Black Forest, Germany). An analogue for fractured geothermal crystalline reservoirs.

Benjamin Avakian^{*1,2,3}, Béatrice Ledéser¹, Ronan Hebert¹, Philippe Claeys², Steven Goderis², Ghislain Trullenque³, and Sébastien Potel³

¹CY Cergy Paris Université – Geosciences Environnement Cergy – France

²Vrije Universiteit [Brussel] – Belgique

³UniLaSalle – B2R – France

Enhanced geothermal systems in basement rocks consist in the exploitation of the hot brines circulating within the fracture network as the main fluid pathway within the reservoir. This is the case at Soultz-sous-Forêts and Rittershoffen (France). The fracture network must be clearly documented before developing a geothermal installation to reduce the risks of targeting low permeability fractures or induced seismicity related to reservoir stimulation. A potential sub-surface analogue to Soultz-sous-Forêts' reservoir is found in a highly strained zone rich in clay minerals in Shauinsland's mine (Black Forest, Germany). The analogue is crosscut by an ore vein that disappears close to the core of the fault zone and reappears on its other side. Petrographical and petrophysical characterizations were done to identify deformation and hydrothermal alteration markers such as grain size reduction, fracturation, illitization process and precipitation of carbonate veins. This was coupled with a geochemical study focusing on the effect of hydrothermal alteration on the chemical composition of the deformed host rock. Micro X-ray fluorescence element maps were gained at thin section scale (3x4 centimeters) to identify fluid circulation features within the host rock. Major element concentrations were quantified using powdered samples taken along the fault zone to identify hydrothermal alteration stages. Porosity, carbonate content and magnetic susceptibility increase towards the fault core and tend to be correlated with clay content. Fe precipitated massively in the core zone and Ca in veins of the damage zone. High fluid flow was channelled in the core zone which is no longer permeable as indicated by the high clay content. Combined with mechanical data and numerical modelling, this may help reduce the risks of "dry well" or generate seismic events during geothermal stimulation.

Mots-Clés: Enhanced geothermal systems, analogue, fault zone, fluid/rock interaction, permeability

*Intervenant