



HAL
open science

FROM INGHAM TO NAZAROV'S INEQUALITY: A SURVEY ON SOME TRIGONOMETRIC INEQUALITIES

Philippe Jaming, Chadi Saba

► **To cite this version:**

Philippe Jaming, Chadi Saba. FROM INGHAM TO NAZAROV'S INEQUALITY: A SURVEY ON SOME TRIGONOMETRIC INEQUALITIES. 2023. hal-04314308

HAL Id: hal-04314308

<https://hal.science/hal-04314308>

Preprint submitted on 29 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FROM INGHAM TO NAZAROV'S INEQUALITY: A SURVEY ON SOME TRIGONOMETRIC INEQUALITIES

PHILIPPE JAMING & CHADI SABA

ABSTRACT. The aim of this paper is to give an overview of some inequalities about L^p -norms ($p = 1$ or $p = 2$) of harmonic (periodic) and non-harmonic trigonometric polynomials. Among the material covered, we mention Ingham's Inequality about L^2 norms of non-harmonic trigonometric polynomials, the proof of the Littlewood conjecture by Mc Gehee, Pigno and Smith on the lower bound of the L^1 norm of harmonic trigonometric polynomials as well as its counterpart in the non-harmonic case due to Nazarov. For the latter one, we give a quantitative estimate that completes our recent result with an estimate of L^1 -norms over small intervals.

We also give some stronger lower bounds when the frequencies satisfy some more restrictive conditions (lacunary Fourier series, "multi-step arithmetic sequences").

Most proofs are close to existing ones and some open questions are mentioned at the end.

CONTENTS

1. Introduction	1
2. L^2 estimates	4
2.1. Ingham's inequalities	4
2.2. The condition $T > 1$	7
2.3. Sequences with large gaps	8
3. L^1 estimates with integer frequencies	11
3.1. The L^1 norm of the Dirichlet kernel	11
3.2. Lacunary trigonometric polynomials	12
3.3. The proof of Littlewood's conjecture by Mc Gehee, Pigno and Smith	19
3.4. Strongly multidimensional sets	24
3.5. Newman's extremal trigonometric polynomial	34
4. L^1 estimates with real frequencies	36
4.1. Some results for the Besicovitch norm	36
4.2. Nazarov's theorem	38
5. Some open problems	48
References	49

1. INTRODUCTION

The aim of this paper is to give an overview of the main estimates of L^p -norms of harmonic and non-harmonic trigonometric polynomials, when $p = 2$ and $p = 1$. In many fields of mathematics, ranging from number theory (*see e.g.* the previous survey [1]) to signal processing and PDEs, one is lead to investigate such norms. Our main motivation comes from the use of Ingham's inequality (lower and upper estimates of L^2 -norms of non-harmonic trigonometric polynomials) in control theory of PDEs. We refer the interested reader to the book by Komornik and Loreti [14]

2020 *Mathematics Subject Classification.* 42A75.

Key words and phrases. Ingham's Inequality, Littlewood problem, non-harmonic Fourier series, lacunary series.

Let us not be more precise. We will here restrict attention to $p = 2$ or $p = 1$ and investigate L^p -norm estimates of (harmonic) trigonometric polynomials

$$\int_{-1/2}^{1/2} \left| \sum_{k=-N}^N c_k e^{2i\pi n_k t} \right|^p dt \quad c_k \in \mathbb{C}, n_k \in \mathbb{N}$$

or non-harmonic trigonometric polynomials

$$\frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{k=-N}^N c_k e^{2i\pi \lambda_k t} \right|^p dt \quad T > 0, c_k \in \mathbb{C}, \lambda_k \in \mathbb{R}$$

as well as their limits when $T \rightarrow +\infty$

$$\limsup_{T \rightarrow +\infty} \frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{k=-N}^N c_k e^{2i\pi \lambda_k t} \right|^p dt \quad T > 0, c_k \in \mathbb{C}, \lambda_k \in \mathbb{R}$$

(that is Besicovitch norms). Note for future use that, when the λ_k 's are all integers, we recover the $L^p([-1/2, 1/2])$ -norms.

The simplest case $p = 2$ for harmonic trigonometric polynomials known by Parseval's relation states that

$$\int_{-1/2}^{1/2} \left| \sum_{k=-N}^N c_k e^{2i\pi k t} \right|^2 dt = \sum_{k=-N}^N |c_k|^2.$$

The situation becomes much complicated for non-harmonic trigonometric polynomials but is well understood thanks to a deep result by Ingham:

Theorem A (Ingham). *Let $T > 1$, then there are two constants A and B such that*

- for every sequence $(\lambda_k)_{k \in \mathbb{Z}}$ with $\lambda_{k+1} - \lambda_k \geq 1$,
 - for every complex sequence $(c_k)_{k=-N, \dots, N}$
- for every integer $N \geq 1$,

$$(1.1) \quad A \sum_{k=-N}^N |c_k|^2 \leq \frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{k=-N}^N c_k e^{2i\pi \lambda_k t} \right|^2 dt \leq B \sum_{k=-N}^N |c_k|^2.$$

The condition $T > 1$ is optimal in the following sense; When $T = 1$, there is a sequence $(\lambda_k)_{k \in \mathbb{Z}}$ with $\lambda_{k+1} - \lambda_k \geq 1$ and a sequence $(c_k) \in \ell^2$ such that the lower bound in (1.1) does not hold for every N for any $A > 0$.

Note that the upper bound in (1.1) holds for every $T > 0$, the restriction $T > 1$ is only needed for the lower bound, which is the most interesting one for control theory. In the next section, we will present Ingham's proof and also compute the constants A and B . We will also present an argument due to Haraux that allows to obtain the lower bound with T arbitrarily small when $|\lambda_{k+1} - \lambda_k| \rightarrow +\infty$ when $k \rightarrow \pm\infty$.

Once the L^2 -theory settled, we will move to the L^1 -theory which is much richer. To start with, the behavior of the L^1 -norm of a trigonometric polynomial may depend on the frequencies. There are two extreme cases

– the frequencies form an arithmetic sequence, e.g. the Dirichlet kernel, one might expect the L^1 -norm to be small in view of the estimate: when $N \rightarrow +\infty$

$$\int_{-1/2}^{1/2} \left| \sum_{k=-N}^N e^{2i\pi k t} \right| dt = \frac{4}{\pi^2} \ln N + O(1),$$

– the frequencies form a geometric sequence, or more generally a lacunary sequence ($n_{k+1} \geq q n_k$ with $q > 1$) then the L^1 -norms are much large: for N large enough

$$\int_{-1/2}^{1/2} \left| \sum_{k=0}^N e^{2i\pi n_k t} \right| dt \geq C \sqrt{N}.$$

We will prove these two estimates in Section 3. In view of those estimates, Littlewood conjectured that the Dirichlet kernel has worse possible behavior, namely that

$$L_N := \inf_{n_0 < n_1 < \dots < n_N} \int_{-1/2}^{1/2} \left| \sum_{k=0}^N e^{2i\pi n_k t} \right| dt \geq C \log N$$

for some constant $C \leq \frac{4}{\pi^2}$. The first non-trivial estimate was obtained by Cohen [5] who proved that

$$L_N \geq C(\ln N / \ln \ln N)^{1/8}$$

for $N \geq 4$. Subsequent improvements are due to Davenport [6], Fournier [8] and crucial contributions by Pichorides [19, 20, 21, 22] leading to $L_N \geq C \ln N / (\ln \ln N)^2$. Finally, Littlewood's conjecture was proved independently by Konyagin [15] and Mc Gehee, Pigno, Smith [16] in 1981. In both papers, Littlewood's conjecture is actually obtained as a corollary of a stronger result (and they are not consequences of one another). The second one is the one we will focus on here and is the following result:

Theorem B (Mc Gehee, Pigno & Smith [16]). *For $n_1 < n_2 < \dots < n_N$ integers and a_1, \dots, a_N complex numbers,*

$$\int_{-1/2}^{1/2} \left| \sum_{k=1}^N a_k e^{2i\pi n_k t} \right| dt \geq C_{MPS} \sum_{k=1}^N \frac{|a_k|}{k}$$

where C_{MPS} is a universal constant ($C_{MPS} = 1/30$ would do).

We will present its proof below.

We would like to insist that this is the worse possible behavior. To start, as we already mentioned, in the lacunary case, the lower bound is \sqrt{N} . We will show a recent result of Hanson [10] who considered a family of sets (so-called "strongly multi-dimensional sets") of which the simplest example is a multi-step arithmetic sequence. By that, we mean a sequence $n_{jM+k} = jD + kd$, $j \in \mathbb{Z}$, $k = 0, \dots, M-1$ and $Md \ll D$. In other words, we take an arithmetic sequence with large step D and, after each element of the sequence, we add a small piece of an arithmetic sequence with small step. It is then shown that

$$\int_{-1/2}^{1/2} \left| \sum_{j=0}^N \sum_{k=0}^M e^{2i\pi(jD+kd)t} \right| dt \geq C \ln N \ln M.$$

We also present an example of Newman that constructed a_0, \dots, a_N all of modulus 1 such that

$$\int_{-1/2}^{1/2} \left| \sum_{k=0}^N a_k e^{2i\pi k t} \right| dt \geq \sqrt{N} - C.$$

This shows that it is possible to be far away from the lower bound (and actually near to the best possible upper bound).

The last part of this survey is devoted to L^1 -norms of non-harmonic trigonometric polynomials. The best result to day is the following:

Theorem C. *Let $\lambda_0 < \lambda_2 < \dots < \lambda_N$ be N distinct real numbers and a_0, \dots, a_N be complex numbers. Then*

(i) *we have*

$$\lim_{T \rightarrow +\infty} \frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{k=0}^N a_k e^{2i\pi \lambda_k t} \right| dt \geq \frac{1}{26} \sum_{k=1}^N \frac{|a_k|}{k+1}.$$

(ii) *If further a_0, \dots, a_N all have modulus larger than 1, $|a_k| \geq 1$ then*

$$\lim_{T \rightarrow +\infty} \frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{k=0}^N a_k e^{2i\pi \lambda_k t} \right| dt \geq \frac{4}{\pi^3} \ln N.$$

(iii) Assume further that for $k = 0, \dots, N-1$, $\lambda_{k+1} - \lambda_k \geq 1$, then, for every $T > 1$, there exists a constant $C(T)$ such that, for every $a_0, \dots, a_N \in \mathbb{C}$,

$$\frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{k=0}^N a_k e^{2i\pi\lambda_k t} \right| dt \geq C(T) \sum_{k=1}^N \frac{|a_k|}{k+1}.$$

Moreover,

- (a) for $T \geq 72$ we can take $C(T) = \frac{1}{122}$;
 (b) for $1 < T \leq 2$, $C(T) = O((T-1)^{15/2})$.

Let us comment on this theorem. To start with, the theorem is essentially a quantitative version of a result of F. Nazarov [17] and is due to the authors together with K. Kellay [13]. Only the case $1 < T \leq 2$ has not been presented before. To be more precise, when the λ_k 's are integers, this is of course the result of Mc Gehee, Pigno and Smith while Point 2 comes from a slight modification of their argument by Stegeman and Yabuta. Note that the constants are the same as those in Theorem B (and are actually a bit better). While it is obvious that this inequality implies Theorem B (with the same constant), there is an elegant argument by Hudson and Leckband [11] that allows to show that the converse is also true. We will present this argument below. Point 3 is due to Nazarov with non-explicit constant. A direct proof of this theorem is given in [13] with the explicit constants mentioned above. Only the case of small T has not been presented so far and is thus the main novelty of the present paper. It is obtained by a very mild modification of Nazarov's proof. Note also that once we have established Point 3 for some T_0 , it is valid for all $T \geq T_0$ so that we actually recover Nazarov's original result (with much worse constants than the ones stated above).

The remaining of the paper is organised as follows: Section 2 is devoted to the L^2 case and we prove the inequalities in Ingham's Theorem in Section 2.1 and the necessity of the condition $T > 1$ in Section 2.2. Further, Haraux's argument is presented in Section 2.3.

In Section 3, we investigate the L^1 norms when the frequencies are integers. We start with the classical asymptotic estimate of the Dirichlet kernel and devote Section 2.2 to lacunary trigonometric polynomials. Section 3.3 is devoted to Mc Gehee, Pigno, Smith's proof of Littlewood's conjecture. In Section 3.4 we present Hanson's result on strongly multidimensional sequences and we conclude in Section 3.5 with the result of Newman.

Section 4 is devoted to the case of non-harmonic trigonometric polynomials. We start in Section 4.1 with the argument of Hudson and Leckband and then extend the case of lacunary trigonometric polynomials to the non-harmonic setting and the Besicovitch L^1 -norms. We conclude this section with the proof of the quantitative version of Nazarov's theorem for small T .

In the last section, we present a few open questions.

2. L^2 ESTIMATES

2.1. Ingham's inequalities. The aim of this section is to show the following: let $\Lambda = \{\lambda_k\}_{k \in \mathbb{Z}} \subset \mathbb{R}$ be a 1-separated sequence, $|\lambda_k - \lambda_\ell| \geq 1$ if $k \neq \ell$. Let $\mathcal{P}(\Lambda)$ be the set of (non-harmonic) trigonometric polynomials

$$\mathcal{P}(\Lambda) = \left\{ P(t) := \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t} : (a_k)_{k \in \mathbb{Z}} \subset \mathbb{C} \text{ with finite support} \right\}.$$

Note that, if $P \in \mathcal{P}(\Lambda)$ is given, then the a_k 's are determined by

$$a_k = \lim_{T \rightarrow +\infty} \frac{1}{T} \int_{-T/2}^{T/2} P(t) e^{-2i\pi\lambda_k t} dt.$$

We can then define two natural norms on $\mathcal{P}(\Lambda)$, namely

$$\|P\|_{L^2([-T/2, T/2])} := \left(\frac{1}{T} \int_{-T/2}^{T/2} |P(t)|^2 dt \right)^{\frac{1}{2}}$$

and

$$\|P\|_{\ell^2} := \left(\sum_{k \in \mathbb{Z}} |a_k|^2 \right)^{\frac{1}{2}}.$$

Our aim in this section is to show that, when $T > 1$, these two norms are equivalent. This is done by proving two inequalities. The first one is the direct inequality:

Proposition 2.1 (Ingham's direct inequality). *Let $(a_k)_{k \in \mathbb{Z}}$ be a finitely supported sequence of complex numbers and $(\lambda_k)_{k \in \mathbb{Z}}$ be a sequence of real numbers with $\lambda_{k+1} - \lambda_k \geq 1$. For every $T > 0$,*

$$(2.1) \quad \frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t} \right|^2 dt \leq 2 \frac{T+1}{T} \sum_{k \in \mathbb{Z}} |a_k|^2.$$

Proof. We consider the function H on \mathbb{R} defined by

$$h(x) = \begin{cases} \cos \pi x & \text{when } |x| \leq \frac{1}{2} \\ 0 & \text{otherwise} \end{cases}.$$

As h is real and even, its Fourier transform is given by

$$\hat{h}(t) = 2 \int_0^{1/2} \cos \pi x \cos 2\pi x t dx = \frac{2 \cos \pi t}{\pi (1 - 4t^2)}$$

with the understanding that $\hat{h}(1/2) = \frac{1}{2}$. From this, one shows that $\hat{h}(x) \geq \frac{1}{2}$ for $|x| \leq \frac{1}{2}$. Finally, let

$$g(x) = h * h(x) = \begin{cases} \frac{\sin \pi |x| - \pi(|x| - 1) \cos \pi x}{2\pi} & \text{when } |x| \leq 1 \\ 0 & \text{otherwise} \end{cases}.$$

One easily shows that g is even, non-negative, supported in $[-1, 1]$ and that $g(0) = \frac{1}{2}$. Further its Fourier transform is $\hat{g}(t) = \hat{h}(t)^2$. In particular, $\hat{g}(t) \geq 0$ and $\hat{g}(t) \leq \frac{1}{4}$ for $|t| \leq \frac{1}{2}$.

But then, if $(a_k)_{k \in \mathbb{Z}}$ is finitely supported and $P(t) = \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t}$,

$$\begin{aligned} \int_{-1/2}^{1/2} |P(t)|^2 dt &\leq 4 \int_{-1/2}^{1/2} \hat{g}(t) |P(t)|^2 dt \leq 4 \int_{\mathbb{R}} \hat{g}(x) \left| \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t} \right|^2 dt \\ &= 4 \sum_{j, k \in \mathbb{Z}} a_j \bar{a}_k \int_{\mathbb{R}} \hat{g}(t) e^{2i\pi(\lambda_j - \lambda_k)t} dt \\ &= 4 \sum_{j, k \in \mathbb{Z}} a_j \bar{a}_k g(\lambda_j - \lambda_k). \end{aligned}$$

Note that the sums are actually finite. Further, if $j \neq k$ then $|\lambda_j - \lambda_k| \geq 1$ and, as g is supported in $[-1, 1]$, we then have $g(\lambda_j - \lambda_k) = 0$. This implies that

$$\int_{-1/2}^{1/2} |P(t)|^2 dt \leq 4g(0) \sum_{j \in \mathbb{Z}} |a_j|^2$$

so that the inequality is proven for $T = 1$ since $4g(0) = 2$.

For $T < 1$ we simply write

$$\frac{1}{T} \int_{-T/2}^{T/2} |P(t)|^2 dt \leq \frac{1}{T} \int_{-1/2}^{1/2} |P(t)|^2 dt \leq \frac{2}{T} \sum_{j \in \mathbb{Z}} |a_j|^2.$$

To conclude, notice first that, if $I = [a - 1/2, a + 1/2]$ and $P(t) = \sum_{j \in \mathbb{Z}} a_j e^{2i\pi\lambda_j t}$ then

$$\begin{aligned} \int_I |P(t)|^2 dt &= \int_{-1/2}^{1/2} |P(a+t)|^2 dt = \int_{-1/2}^{1/2} \left| \sum_{j \in \mathbb{Z}} a_j e^{2i\pi\lambda_j a} e^{2i\pi\lambda_j t} \right|^2 dt \\ &\leq 2 \sum_{j \in \mathbb{Z}} |a_j e^{2i\pi\lambda_j a}|^2 = 2 \sum_{j \in \mathbb{Z}} |a_j|^2 \end{aligned}$$

from the case $T = 1$.

Now let $T > 1$ and cover the interval $[-T/2, T/2]$ by $K = \lceil T \rceil \leq T + 1$ intervals I_1, \dots, I_K of length 1. Then

$$\frac{1}{T} \int_{-T/2}^{T/2} |P(t)|^2 dt \leq \frac{1}{T} \sum_{j=1}^K \int_{I_j} |P(t)|^2 dt \leq 2 \frac{T+1}{T} \sum_{j \in \mathbb{Z}} |a_j|^2.$$

This completes the proof. \square

We now show that a converse inequality also holds, but this time with the extra condition that $T > 1$:

Proposition 2.2 (Ingham's converse inequality). *Let $(a_k)_{k \in \mathbb{Z}}$ be a finitely supported sequence of complex numbers and $(\lambda_k)_{k \in \mathbb{Z}}$ be a sequence of real numbers with $\lambda_{k+1} - \lambda_k \geq 1$. For every $T > 1$,*

$$(2.2) \quad \frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t} \right|^2 dt \geq C(T) \sum_{k \in \mathbb{Z}} |a_k|^2$$

with

$$C(T) = \begin{cases} \frac{\pi^2 T^2 - 1}{8 T^3} & \text{for } 1 < T \leq 2 \\ \frac{3\pi^2}{64} & \text{for } T \geq 2 \end{cases}.$$

Proof. We will do so in three steps. We first establish this inequality for $1 < T \leq 2$.

Let H be defined by $H(x) = \mathbb{1}_{[-1/2, 1/2]} \cos \pi x$ and notice that

$$\widehat{H}(t) = \frac{2 \cos \pi t}{\pi (1 - 4t^2)}.$$

Note that H being non-negative, \widehat{H} is maximal at 0 (which may be checked directly).

Next consider $H * H$ and notice that, as H is non-negative, even, continuous with support $[-1/2, 1/2]$, then $H * H$ is non-negative, even, continuous with support $[-1, 1]$ and its Fourier transform is

$$\widehat{H * H} = \widehat{H}^2.$$

Next $H \in H^1(\mathbb{R})$ with $H' = -\pi \mathbb{1}_{[-1/2, 1/2]} \sin \pi x$ and

$$\widehat{H}'(t) = 4it \frac{\cos \pi t}{1 - 4t^2}$$

thus

$$\widehat{H' * H'}(t) = -(2\pi t)^2 \widehat{H}^2(t)$$

We now consider $G_T = \pi^2 T^2 H * H + H' * H'$ so that G_T is continuous, real valued, even and supported in $[-1, 1]$.

$$\widehat{G}_T(t) = \pi^2 (T^2 - 4t^2) \widehat{H}^2(t)$$

is even (so G_T is the Fourier transform of \widehat{G}_T) and in L^1 . Further \widehat{G} is non-negative on $[-T/2, T/2]$ and negative on $\mathbb{R} \setminus [-T/2, T/2]$.

This implies that

$$\begin{aligned} \int_{-T/2}^{T/2} \widehat{G}_T(t) \left| \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t} \right|^2 dt &\geq \int_{\mathbb{R}} \widehat{G}_T(t) \left| \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t} \right|^2 dt \\ &= \sum_{k, \ell \in \mathbb{Z}} a_k \bar{a}_\ell \int_{\mathbb{R}} \widehat{G}_T(t) e^{2i\pi(\lambda_k - \lambda_\ell)t} dt \\ &= \sum_{k, \ell \in \mathbb{Z}} a_k \bar{a}_\ell G_T(\lambda_k - \lambda_\ell) = \sum_{k \in \mathbb{Z}} |a_k|^2 G_T(0). \end{aligned}$$

In the last line, we use that $|\lambda_k - \lambda_\ell| \geq 1$ when $k \neq \ell$ thus $G_T(\lambda_k - \lambda_\ell) = 0$.

Now, for $\xi \in [-T/2, T/2]$,

$$\widehat{G}_T(\xi) = \pi^2 (T^2 - 4\xi^2) \widehat{H}^2(\xi) \leq \pi^2 (T^2 - 4\xi^2) \widehat{H}^2(0) \leq 4T^2$$

while

$$G_T(0) = \pi^2 \int_{-1/2}^{1/2} T^2 \cos^2 \pi t - \sin^2 \pi t dt = \frac{\pi^2}{2} (T^2 - 1)$$

which leads to

$$(2.3) \quad \int_{-T/2}^{T/2} \left| \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t} \right|^2 dt \geq \frac{\pi^2 T^2 - 1}{8 T^2} \sum_{k \in \mathbb{Z}} |a_k|^2.$$

For $2 \leq T \leq 6$, we simply write

$$\int_{-T/2}^{T/2} \left| \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t} \right|^2 dt \geq \int_{-1}^1 \left| \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t} \right|^2 dt \geq \frac{3\pi^2}{32} \sum_{k \in \mathbb{Z}} |a_k|^2$$

where the second inequality is (2.3) with $T = 2$, establishing (2.2) with $C = \frac{3\pi^2}{32T} \geq \frac{\pi^2}{64}$.

Now let $T \geq 6$ and $M_T = [T/2]$ so that $M_T \geq \frac{T}{2} - 1 \geq \frac{T}{3}$. For $j = 0, \dots, M_T - 1$, let $t_j = -T/2 + j + 1$

so that the intervals $[t_j - 1, t_j + 1[$ are disjoint and $\bigcup_{j=0}^{M_T-1} [t_j - 1, t_j + 1[\subset [-T/2, T/2]$ thus

$$\begin{aligned} \int_{-T/2}^{T/2} \left| \sum_{k \in \mathbb{Z}} b_k e^{2i\pi\lambda_k t} \right|^2 dt &\geq \sum_{j=0}^{M_T-1} \int_{t_j-1}^{t_j+1} \left| \sum_{k \in \mathbb{Z}} b_k e^{2i\pi\lambda_k t} \right|^2 dt \\ &= \sum_{j=0}^{M_T-1} \int_{-1}^1 \left| \sum_{k \in \mathbb{Z}} b_k e^{2i\pi\lambda_k t_j} e^{2i\pi\lambda_k t} \right|^2 dt. \end{aligned}$$

Now, apply (2.3) with $a_k = b_k e^{2i\pi\lambda_k t_j}$ and $T = 2$ to get

$$\frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{j=1}^N b_k e^{2i\pi\lambda_k t} \right|^2 dt \geq \frac{3\pi^2}{32} \frac{M_T}{T} \sum_{k=1}^N |a_k|^2 \geq \frac{\pi^2}{32} \sum_{k=1}^N |a_k|^2,$$

establishing (2.2) with $C = \frac{\pi^2}{32}$. □

Finally, we notice that, with a change of variable, and a simple limiting argument to remove the condition on the support of (a_k) , we have just proved the following;

Theorem 2.3 (Ingham). *Let $\gamma > 0$ and $T > \frac{1}{\gamma}$. There exist two constants $C = C(\gamma T)$, $B = B(\gamma T)$ such that*

- for every sequence of real numbers $\Lambda = \{\lambda_k\}_{k \in \mathbb{Z}}$ such that $\lambda_{k+1} - \lambda_k \geq \gamma$;
- for every sequence $(a_k)_{k \in \mathbb{Z}} \in \ell^2(\mathbb{Z}, \mathbb{C})$,

$$(2.4) \quad C \sum_{k \in \mathbb{Z}} |a_k|^2 \leq \frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{k \in \mathbb{Z}} a_k e^{2i\pi\lambda_k t} \right|^2 dt \leq B \sum_{k \in \mathbb{Z}} |a_k|^2.$$

2.2. **The condition $T > 1$.** We now show that the condition $T > 1$ is mandatory to establish (2.2) for every Λ and every $P \in \mathcal{P}(\Lambda)$.

Proposition 2.4 (Ingham). *There exists a real sequence $\{\lambda_k\}_{k \in \mathbb{Z}}$ with $\lambda_{k+1} - \lambda_k \geq 1$ and a sequence $(a_k)_{k \in \mathbb{Z}}$ with $\sum_{k \in \mathbb{Z}} |a_k|^2 = 1$ such that,*

$$(2.5) \quad \int_{-1/2}^{1/2} \left| \sum_{k=-N}^N a_k e^{2i\pi\lambda_k t} \right|^2 dt \rightarrow 0$$

when $N \rightarrow +\infty$.

In other words, for (2.2) to hold for $T = 1$ for this sequence (λ_k) and every sequence (a_k) then $C(1) = 0$ so that the condition $T > 1$ is necessary for this inequality to be interesting.

Proof. Let $\alpha > 0$ and define, for $|z| < 1$,

$$g_\alpha(z) = (1+z)^{-\alpha} = \frac{\exp\left(-i\alpha \arctan \frac{\Im(z)}{1+\Re(z)}\right)}{|1+z|^\alpha}, \quad |z| < 1.$$

Of course, we may also write g_α as a power series

$$g_\alpha(z) = \sum_{n=0}^{+\infty} \frac{(-\alpha)_n}{n!} z^n$$

where $(\alpha)_0 = 1$, $(-\alpha)_n = -\alpha(-\alpha-1)\cdots(-\alpha-n+1)$.

Next define

$$\begin{aligned} f_\alpha(r, t) &= 2\Re\left(e^{i\pi(\alpha+1)t} g_\alpha(re^{2i\pi t})\right) \\ &= e^{i\pi(\alpha+1)t} g_\alpha(re^{2i\pi t}) + e^{-i\pi(\alpha+1)t} g_\alpha(re^{-2i\pi t}) \\ &= \sum_{n=0}^{+\infty} \frac{(-\alpha)_n}{n!} r^n e^{2i\pi\left(n+\frac{\alpha+1}{2}\right)t} + \sum_{n=0}^{+\infty} \frac{(-\alpha)_n}{n!} r^n e^{-2i\pi\left(n+\frac{\alpha+1}{2}\right)t}. \end{aligned}$$

Now set $\Lambda = \{\lambda_j\}_{j \in \mathbb{Z}}$ with $\lambda_j = j + \frac{\alpha+1}{2}$ when $j \geq 0$ and $\lambda_j = j + 1 - \frac{\alpha+1}{2}$ for $j \leq -1$, then $\lambda_{j+1} - \lambda_j \geq 1$ (and even = 1 excepted for $\lambda_0 - \lambda_{-1} = 1 + \alpha$). In particular, if we set

$$P_{m,r}(t) = \sum_{n=0}^m \frac{(-\alpha)_n}{n!} r^n e^{2i\pi\left(n+\frac{\alpha+1}{2}\right)t} + \sum_{n=0}^m \frac{(-\alpha)_n}{n!} r^n e^{-2i\pi\left(n+\frac{\alpha+1}{2}\right)t} := \sum_{k \in \mathbb{Z}} a_{m,r}(k) e^{2i\pi\lambda_k t} \in \mathcal{P}(\Lambda)$$

and $P_{m,r} \rightarrow f_\alpha$ when $m \rightarrow +\infty$, uniformly over $t \in [-1/2, 1/2]$.

Further, Parseval's relation reads

$$\sum_{n=0}^{+\infty} \left| \frac{(-\alpha)_n}{n!} r^n \right|^2 = \int_{-1/2}^{1/2} \left| \sum_{n=0}^{+\infty} \frac{(-\alpha)_n}{n!} r^n e^{2i\pi n t} \right|^2 dt = \int_{-1/2}^{1/2} |g_\alpha(re^{2i\pi t})|^2 dt$$

thus

$$\lim_{m \rightarrow +\infty} \sum_{k \in \mathbb{Z}} |a_{m,r}(k)|^2 = \lim_{m \rightarrow +\infty} 2 \sum_{n=0}^{+\infty} \left| \frac{(-\alpha)_n}{n!} r^n \right|^2 = 2 \int_{-1/2}^{1/2} |g_\alpha(re^{2i\pi t})|^2 dt.$$

It follows that, if we had

$$(2.6) \quad \int_{-1/2}^{1/2} |P_{m,r}(t)|^2 dt \geq C \sum_{k \in \mathbb{Z}} |a_{m,r}(k)|^2$$

then, letting $m \rightarrow +\infty$, we would also have

$$(2.7) \quad \int_{-1/2}^{1/2} |f_\alpha(r, t)|^2 dt \geq 2C \int_{-1/2}^{1/2} |g_\alpha(re^{2i\pi t})|^2 dt$$

for every $0 < r < 1$ and every $0 < \alpha < 1$.

But, if we fix $t \in]-1/2, 1/2[$ then, when $r \rightarrow 1$,

$$g_\alpha(re^{\pm 2i\pi t}) = \frac{1}{(1 + re^{\pm 2i\pi t})^\alpha} \rightarrow \frac{1}{(1 + e^{\pm 2i\pi t})^\alpha} = \frac{e^{\mp i\alpha\pi t}}{2^\alpha \cos^\alpha \pi t}$$

(this is where we use that $T \leq 1$) while

$$|g_\alpha(re^{\pm 2i\pi t})|^2 = \frac{1}{((1-r)^2 + 4r \cos^2 \pi t)^\alpha} \leq \frac{1}{4 \cos^{2\alpha} \pi t}$$

for $\frac{1}{2} < r < 1$. Similar bounds follow for $f_\alpha(r, t)$:

$$f_\alpha(r, t) = e^{i\pi(\alpha+1)t} g_\alpha(re^{2i\pi t}) + e^{-i\pi(\alpha+1)t} g_\alpha(re^{-2i\pi t}) \rightarrow \frac{e^{i\pi t} + e^{-i\pi t}}{2^\alpha \cos^\alpha \pi t} = \frac{1}{2^{\alpha-1} \cos^{\alpha-1} \pi t}$$

while

$$|f_\alpha(r, t)|^2 \leq \frac{1}{\cos^{2\alpha} \pi t}.$$

When $2\alpha < 1$ the majorants are integrable so that we can let $r \rightarrow 1$ in (2.7). This leads to

$$(2.8) \quad 2^{2-2\alpha} \int_{-1/2}^{1/2} \frac{dt}{\cos^{2\alpha-2} \pi t} \geq 2^{1-2\alpha} C \int_{-1/2}^{1/2} \frac{dt}{\cos^{2\alpha} \pi t}.$$

Letting $\alpha \rightarrow \frac{1}{2}$ the right hand side goes to $+\infty$ while the left hand side stays bounded. We obtain the desired contradiction. \square

2.3. Sequences with large gaps.

Lemma 2.5 (Haraux). *Let $\Lambda \subset \mathbb{R}$ be a sequence such that $\gamma(\Lambda) = \inf_{k \neq \ell \in \mathbb{Z}} |\lambda_k - \lambda_\ell| > 1$ and let $T > \gamma(\Lambda)^{-1}$.*

Assume that there exists $0 < C \leq 1 \leq B$ such that, for every $(a_\lambda)_{\lambda \in \Lambda}$ finitely supported sequence of complex numbers,

$$C \sum_{\lambda \in \Lambda} |a_\lambda|^2 \leq \frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{\lambda \in \Lambda} a_\lambda e^{2i\pi \lambda t} \right| dt \leq B \sum_{\lambda \in \Lambda} |a_\lambda|^2.$$

Let $\mu \in \mathbb{R} \setminus \Lambda$ and, for sake of simplicity, assume that $\gamma(\Lambda \cup \{\mu\}) \geq 1$. Let $0 < \delta < \min\left(T, \frac{1}{4}\right)$, then there is a D with $0 < D < C$ such that, for every $(a_\lambda)_{\lambda \in \Lambda \cup \{\mu\}}$

$$C \sum_{\lambda \in \Lambda} |a_\lambda|^2 \leq \frac{1}{T + \delta} \int_{-\frac{T+\delta}{2}}^{\frac{T+\delta}{2}} \left| \sum_{\lambda \in \Lambda \cup \{\mu\}} a_\lambda e^{2i\pi \lambda t} \right| dt$$

Moreover, D can be taken of the form

$$D = \kappa \frac{C}{B} \delta^4$$

for some absolute constant $\kappa > 0$ ($\kappa = 8^{-4}$ would do).

Note that, as δ is arbitrarily small, as $T > \gamma(\Lambda)^{-1}$ is arbitrarily near to $\gamma(\Lambda \setminus \{\lambda_0\})^{-1}$ so is $T + \delta$.

Proof. Set

$$f(t) = \sum_{\lambda \in \Lambda \cup \{\mu\}} a_\lambda e^{2i\pi \lambda t}.$$

For $0 < \delta < \frac{1}{4}$, we define

$$\begin{aligned} g_\delta(t) &= f(t) - \frac{1}{\delta} \int_{-\delta/2}^{\delta/2} e^{-2i\pi\mu s} f(t+s) ds \\ &= \sum_{\lambda \in \Lambda} a_\lambda \left(1 - \frac{\sin \pi(\lambda - \mu)\delta}{\pi(\lambda - \mu)\delta} \right) e^{2i\pi\lambda t} := \sum_{\lambda \in \Lambda} b_\lambda(\delta) e^{2i\pi\lambda t}. \end{aligned}$$

The first observation is that, as $|\lambda - \mu| \geq 1$,

$$\left| 1 - \frac{\sin \pi(\lambda - \mu)\delta}{\pi(\lambda - \mu)\delta} \right|^2 \geq \eta_\delta(\mu) := \sup_{s \geq \pi\delta} \left| 1 - \frac{\sin s}{s} \right|^2 > 0.$$

It is not difficult to see that, if $\delta \leq \frac{1}{4}$, $\eta_\delta \geq \frac{\delta^4}{100}$. This is what needs to be adjusted if we do not assume that $\text{dist}(\mu, \Lambda) = \min_{\lambda \in \Lambda} |\lambda - \mu| \geq 1$.

Further, as for $\lambda \neq \lambda' \in \Lambda$, then $|\lambda - \lambda'| \geq \gamma(\Lambda)$, the hypothesis of the lemma states that, as $T > \frac{1}{\gamma(\Lambda)}$,

$$(2.9) \quad \frac{1}{T} \int_{-T/2}^{T/2} |g_\delta(t)|^2 dt \geq C \sum_{\lambda \in \Lambda} |b_\lambda(\delta)|^2 \geq C\eta_\delta \sum_{\lambda \in \Lambda} |a_\lambda|^2 \geq \frac{C\delta^4}{100} \sum_{\lambda \in \Lambda} |a_\lambda|^2$$

On the other hand, from Cauchy-Schwarz,

$$\begin{aligned} |g_\delta(t)|^2 &\leq 2|f(t)|^2 + 2 \left| \frac{1}{\delta} \int_{-\delta/2}^{\delta/2} e^{-2i\pi\lambda_0 s} f(t+s) ds \right|^2 \\ &\leq 2|f(t)|^2 + 2 \frac{1}{\delta} \int_{-\delta/2}^{\delta/2} |f(t+s)|^2 ds \\ &= 2|f(t)|^2 + \frac{1}{\delta} \int_{t-\delta}^{t+\delta} |f(s)|^2 ds. \end{aligned}$$

It follows that

$$\begin{aligned} \frac{1}{T} \int_{-T/2}^{T/2} |g_\delta(t)|^2 dt &\leq \frac{2}{T} \int_{-T/2}^{T/2} |f(t)|^2 dt + \frac{2}{\delta} \frac{1}{T} \int_{-T/2}^{T/2} \int_{t-\delta}^{t+\delta} |f(s)|^2 ds dt \\ &= \frac{2}{T} \int_{-T/2}^{T/2} |f(t)|^2 dt + \frac{2}{T} \int_{-T/2-\delta}^{T/2+\delta} |f(s)|^2 \frac{1}{\delta} \int_{\max(-T/2, s-\delta/2)}^{\min(T/2, s+\delta/2)} dt ds \\ &\leq \frac{2}{T} \int_{-T/2}^{T/2} |f(t)|^2 dt + \frac{2}{T} \int_{-T/2-\delta/2}^{T/2+\delta/2} |f(s)|^2 ds \\ &\leq 4 \frac{1}{T} \int_{-\frac{T+\delta}{2}}^{\frac{T+\delta}{2}} |f(s)|^2 ds \leq 8 \frac{1}{T+\delta} \int_{-\frac{T+\delta}{2}}^{\frac{T+\delta}{2}} |f(s)|^2 ds \end{aligned}$$

since we assumed that $T + \delta \leq 2T$.

With (2.9), we have thus shown that

$$(2.10) \quad \sum_{\lambda \in \Lambda} |a_\lambda|^2 \leq \frac{800}{C\delta^4} \frac{1}{T+\delta} \int_{-\frac{T+\delta}{2}}^{\frac{T+\delta}{2}} |f(s)|^2 ds.$$

It remains to add $|a_\mu|^2$ on the left hand side. As $a_\mu = f - \sum_{\lambda \in \Lambda} a_\lambda e^{2i\pi\lambda t}$,

$$\begin{aligned} |a_\mu|^2 &= \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} |a_\mu|^2 dt \leq 2 \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} |f(t)|^2 dt + 2 \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} \left| \sum_{\lambda \in \Lambda} a_\lambda e^{2i\pi\lambda t} \right|^2 dt \\ &\leq 4 \frac{1}{T+\delta} \int_{-\frac{T+\delta}{2}}^{\frac{T+\delta}{2}} |f(t)|^2 dt + 2B \sum_{\lambda \in \Lambda} |a_\lambda|^2 \end{aligned}$$

where we used again that $T + \delta \leq 2T$ and the hypothesis of the lemma. But then, from (2.10), we get

$$|a_\mu|^2 \leq \left(4 + \frac{1600B}{C\delta^4}\right) \frac{1}{T + \delta} \int_{-\frac{T+\delta}{2}}^{\frac{T+\delta}{2}} |f(s)|^2 ds.$$

Adding this to (2.10), we finally get that

$$\begin{aligned} \sum_{\lambda \in \Lambda \cup \{\mu\}} |a_\lambda|^2 &\leq \left(4 + \frac{800(2B+1)}{C\delta^4}\right) \frac{1}{T + \delta} \int_{-\frac{T+\delta}{2}}^{\frac{T+\delta}{2}} |f(s)|^2 ds \\ &\leq \frac{1}{\kappa} \frac{B}{C\delta^4} \frac{1}{T + \delta} \int_{-\frac{T+\delta}{2}}^{\frac{T+\delta}{2}} |f(s)|^2 ds \end{aligned}$$

with $\kappa^{-1} = 800 \times 3 + 4$ since $C(\gamma_0 T) \leq B(\gamma_0 T)$, $\delta \leq 1$ and $B(\gamma_0 T) \geq 1$. We thus obtain the inequality with

$$D = \kappa \frac{C}{B} \delta^4$$

as claimed. \square

We can now use this lemma to improve Ingham's Inequality.

Corollary 2.6. *Let Λ be a sequence such that $\gamma(\Lambda) = \min_{\lambda \neq \lambda' \in \Lambda} |\lambda - \lambda'| \geq 1$ and let $F = (\lambda_1, \dots, \lambda_n) \subset \Lambda$.*

Let $S > \frac{1}{\gamma(\Lambda \setminus F)}$. Then there exists two constants $0 < D < 1 < B$ such that, for every finitely supported sequence $(a_\lambda)_{\lambda \in \Lambda}$,

$$(2.11) \quad D \sum_{\lambda \in \Lambda} |a_\lambda|^2 \leq \frac{1}{S} \int_{-S/2}^{S/2} \left| \sum_{\lambda \in \Lambda} a_\lambda e^{2i\pi\lambda t} \right|^2 dt \leq B \sum_{\lambda \in \Lambda} |a_\lambda|^2.$$

Proof. Set $\Lambda_0 = \Lambda$ and, for $j = 1, \dots, n$, $\Lambda_j = \Lambda \setminus \{\lambda_1, \dots, \lambda_n\}$ and $\gamma_j = \min_{\lambda \neq \lambda' \in \Lambda_j} |\lambda - \lambda'| \geq 1$.

From Ingham's Theorem, we know that

– for every $2 > T > \frac{1}{\gamma_n}$

$$\frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{\lambda \in \Lambda_n} a_\lambda e^{2i\pi\lambda t} \right|^2 dt \geq C_n \sum_{\lambda \in \Lambda_n} |a_\lambda|^2$$

with $C_n = \frac{\pi^2}{2^6} (T\gamma_n - 1)$;

– for every $2 > T > 0$, and $j = 0, \dots, n$,

$$\frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{\lambda \in \Lambda_j} a_\lambda e^{2i\pi\lambda t} \right|^2 dt \leq B_j \sum_{\lambda \in \Lambda_j} |a_\lambda|^2$$

with $B_j = \frac{6(T\gamma_j + 1)}{T\gamma_j} = 6 \left(1 + \frac{1}{\gamma_j T}\right)$.

In particular, the upper bound in (2.11) is already established.

Now let $\frac{1}{\gamma(\Lambda \setminus F)} < T < S$ and $\delta = \frac{S-T}{n}$. One might take $T = \frac{1}{2} \left(S + \frac{1}{\gamma(\Lambda \setminus F)}\right)$ so that

$$\delta = \frac{1}{2n} \left(S - \frac{1}{\gamma(\Lambda \setminus F)}\right)$$

From Hauraux's lemma, we get

$$\frac{1}{T + \delta} \int_{-\frac{T+\delta}{2}}^{\frac{T+\delta}{2}} \left| \sum_{\lambda \in \Lambda_{n-1}} a_\lambda e^{2i\pi\lambda t} \right|^2 dt \geq C_{n-1} \sum_{\lambda \in \Lambda_{n-1}} |a_\lambda|^2$$

with $C_{n-1} = \kappa \delta^4 \frac{C_n}{B_n}$. We can therefore apply Haraux's lemma again, till we reach Λ_0 . At each step, the constant C_{n-j} is replaced by $C_{n-j-1} = \kappa \delta^4 \frac{C_{n-j}}{B_{n-j}}$ while the integral ranges over $\left[-\frac{T+(j+1)\delta}{2}, \frac{T+(j+1)\delta}{2}\right]$.

This finally leads to

$$\frac{1}{T+n\delta} \int_{-\frac{T+n\delta}{2}}^{\frac{T+n\delta}{2}} \left| \sum_{\lambda \in \Lambda_{n-1}} a_\lambda e^{2i\pi\lambda t} \right|^2 dt \geq C_0 \sum_{\lambda \in \Lambda} |a_\lambda|^2$$

with

$$C_0 = \kappa^n \delta^{4n} \frac{C_n}{B_1 B_2 \cdots B_n}.$$

As $T+n\delta = S$, this establishes (2.11). \square

Note that, taking $\kappa = 8^{-4}$ we obtain

$$D = \frac{\pi^2}{2^6} \left(\frac{3\delta^4}{2^{13}} \right)^n \frac{\gamma_n T - 1}{\prod_{j=0}^n \left(1 + \frac{1}{\gamma_j T} \right)}.$$

3. L^1 ESTIMATES WITH INTEGER FREQUENCIES

3.1. The L^1 norm of the Dirichlet kernel. Recall that the Dirichlet kernel is given by

$$D_N(t) = \sum_{k=-N}^N e^{2i\pi kt} = \frac{\sin(2N+1)\pi t}{\sin \pi t}.$$

The following is a classical estimate of the L^1 -norm of this kernel, which is called the *Lebesgue constant*.

Lemma 3.1. *When $N \rightarrow +\infty$,*

$$\|D_N\|_1 := \int_{-1/2}^{1/2} |D_N(t)| dt = \frac{4}{\pi^2} \ln N + O(1).$$

Proof. The proof is based on the following inequality which is easy to establish and whose proof is left to the reader: for $-\frac{\pi}{2} \leq s \leq \frac{\pi}{2}$, $|s| \left(1 - \frac{s^2}{3}\right) \leq |\sin s| \leq |s|$. In particular, for $|t| \leq \frac{1}{2}$,

$$\frac{1}{\pi|t|} \leq \frac{1}{|\sin \pi t|} \leq \frac{1}{\pi|t|} \frac{1}{1 - \pi^2 t^2/6} \leq \frac{1}{\pi|t|} + \frac{\pi|t|}{3}$$

since $\frac{1}{1-u} \leq 1 + 2u$ for $|u| \leq \frac{1}{2}$. It follows that

$$\begin{aligned} \int_{-1/2}^{1/2} \frac{|\sin(2N+1)\pi t|}{|\pi t|} dt &\leq \int_{-1/2}^{1/2} \frac{|\sin(2N+1)\pi t|}{\sin \pi t} dt \\ &\leq \int_{-1/2}^{1/2} \frac{|\sin(2N+1)\pi t|}{|\pi t|} dt + \int_{-1/2}^{1/2} |\sin(2N+1)\pi t| \frac{\pi|t|}{3} dt. \end{aligned}$$

As

$$\int_{-1/2}^{1/2} |\sin(2N+1)\pi t| \frac{\pi|t|}{3} dt \leq \int_{-1/2}^{1/2} \frac{\pi|t|}{3} dt = \frac{\pi}{12}$$

we get

$$\|D_N\|_1 = \int_{-1/2}^{1/2} \frac{|\sin(2N+1)\pi t|}{|\pi t|} dt + O(1).$$

Using parity and the change of variable $s = (2N+1)\pi t$ we obtain

$$\int_{-1/2}^{1/2} \frac{|\sin(2N+1)\pi t|}{|\pi t|} dt = 2 \int_0^{1/2} \frac{|\sin(2N+1)\pi t|}{|\pi t|} dt = \frac{2}{\pi} \int_0^{N\pi+\pi/2} \frac{|\sin s|}{s} ds.$$

Note that

$$\int_{N\pi}^{N\pi+\pi/2} \frac{|\sin s|}{s} ds \leq \int_{N\pi}^{N\pi+\pi/2} \frac{ds}{s} = \ln \frac{N\pi + \pi/2}{N\pi} = O(1/N)$$

while

$$\int_0^\pi \frac{|\sin s|}{s} ds \leq \int_0^\pi 1 ds = \pi.$$

It remains to estimate

$$\begin{aligned} \frac{2}{\pi} \int_\pi^{N\pi} \frac{|\sin s|}{s} ds &= \frac{2}{\pi} \sum_{j=1}^{N-1} \int_{j\pi}^{(j+1)\pi} \frac{|\sin s|}{s} ds \\ &= \frac{2}{\pi} \sum_{j=1}^{N-1} \int_0^\pi \frac{|\sin s|}{s + j\pi} ds = \frac{2}{\pi} \sum_{j=1}^{N-1} \int_0^\pi \frac{\sin s}{s + j\pi} ds \end{aligned}$$

as $\sin s \geq 0$ on $[0, \pi]$. But then

$$\frac{2}{(j+1)\pi} = \int_0^\pi \frac{\sin s}{\pi + j\pi} ds \leq \int_0^\pi \frac{\sin s}{s + j\pi} ds \leq \int_0^\pi \frac{\sin s}{j\pi} ds = \frac{2}{j\pi}.$$

Writing

$$\sum_{j=1}^{N-1} \frac{1}{j+1} = \sum_{j=1}^N \frac{1}{j} - 1 = \sum_{j=1}^{N-1} \frac{1}{j} - 1 + \frac{1}{N},$$

it follows that

$$\frac{2}{\pi} \int_\pi^{N\pi} \frac{|\sin s|}{s} ds = \frac{4}{\pi^2} \sum_{j=1}^N \frac{1}{j} + O(1).$$

Recalling that $\sum_{j=1}^N \frac{1}{j} = \ln N + O(1)$ the result follows. \square

3.2. Lacunary trigonometric polynomials. In this section, we consider a sequence of integers (n_k) such that $\frac{n_{k+1}}{n_k} \geq q > 1$. Such sequences are called q -lacunary in the sense of Hadamard (or simply lacunary).

First note that a q -lacunary sequence is a finite union of q' -lacunary sequences with $q' \geq 3$. Indeed, if $q \geq 3$ there is nothing to prove and for $1 < q < 3$, take N an integer such that $q^N \geq 3$ and write $n_k^{(\ell)} = n_{\ell+kN}$ then $(n_k^{(\ell)})_k$ is q^N -Lacunary and $\{n_k\} = \bigcup_{\ell=0}^{N-1} \{n_k^{(\ell)}\}$. Next, for $q \geq 3$, q -lacunary sequences have a particular arithmetic property:

Lemma 3.2. *Let $q \geq 3$ and $(n_k)_{k \geq 0}$ a sequence such that $n_0 \geq 1$ and $n_{k+1} \geq qn_k$. Consider two finite sequences $\varepsilon_\ell, \eta_\ell \in \{-1, 0, 1\}$ for $\ell = 0, \dots, m$ and assume that*

$$(3.1) \quad \sum_{\ell=0}^m \varepsilon_\ell n_\ell = \sum_{\ell=0}^m \eta_\ell n_\ell$$

then $\varepsilon_\ell = \eta_\ell$ for every ℓ .

In other words, an integer can be represented in at most one way as $\sum \pm n_\ell$. Such a sequence is called *quasi-independent*.

Proof. First observe that $n_j \leq \frac{1}{q^{m-j}} n_m = \frac{q^j}{q^m} n_m$ for $j = 0, \dots, m$.

Assume that (3.1) holds and define $v_\ell = \varepsilon_\ell - \eta_\ell$ so that

$$\sum_{\ell=0}^m v_\ell n_\ell = 0.$$

Assume towards a contradiction that there is an ℓ such that $v_\ell \neq 0$. Without loss of generality, we may assume that the largest such ℓ is m and, up to exchanging ε_ℓ and η_ℓ , that $v_m \geq 1$.

Observe that $v_\ell \in \{-2, -1, 0, 1, 2\}$ so that we obtain the desired contradiction writing

$$\begin{aligned} 0 = \sum_{\ell=0}^m v_\ell n_\ell &= v_m n_m + \sum_{\ell=0}^{m-1} v_\ell n_\ell \geq n_m - 2 \sum_{\ell=0}^{m-1} n_\ell \geq n_m - 2 \sum_{\ell=0}^{m-1} \frac{q^\ell}{q^m} n_m \\ &= \left(1 - \frac{2}{q^m} \frac{q^m - 1}{q - 1}\right) n_m = \frac{q^{m+1} - 3q^m + 2}{(q-1)q^m} n_m > 0 \end{aligned}$$

since $q \geq 3$ and $n_m > 0$. \square

Note that this result is valid when the n_k 's are real, not only for integers.

The aim of this section is to prove that trigonometric polynomials with lacunary frequencies have large L^1 -norms of which the following estimate is a particular case: there exists a constant $C > 0$ such that, for every N ,

$$\int_{-1/2}^{1/2} \left| \sum_{k=1}^N e^{2i\pi n_k t} \right| dt \geq C\sqrt{N}.$$

This follows from a more general theorem which estimates L^p -norms of lacunary Fourier series. The aim of this section is to present this result. To do so, we follow closely [25, Chapter V.8] which goes through Rademacher series. First let us introduce those series.

To start, let us recall that $\mathcal{D}_k = \{[j2^{-k-1}, (j+1)2^{-k-1}[, j = 0, \dots, 2^{k+1} - 1\}$ the dyadic intervals of generation k and $\mathcal{D} = \bigcup_{k \geq 0} \mathcal{D}_k$ the set of all dyadic intervals. Also, if $I, J \in \mathcal{D}$ then either $I \cap J = \emptyset$ or $I \subset J$ or $J \subset I$. The Rademacher functions of generation k are then functions that take alternative values $+1$ and -1 on successive intervals in \mathcal{D}_k , that is

$$r_k(t) = \sum_{j=0}^{2^{k+1}-1} (-1)^j \mathbb{1}_{[j2^{-k-1}, (j+1)2^{-k-1}[} = \text{sign}(\sin(2\pi 2^j t)).$$

The first observation is that, if $I \in \mathcal{D}_\ell$ and $k > \ell$ then r_k takes the value $+1$ on half of I and -1 on the other half so that $\int_I r_k = 0$. A first consequence is that r_k is orthogonal to r_ℓ in $L^2([0, 1])$ since r_ℓ is constant on each $I \in \mathcal{D}_\ell$ so that $\int_I r_k r_\ell = 0$ and \mathcal{D}_ℓ is a covering of $[0, 1]$. Moreover, as $|r_k| = 1$, the family $(r_k)_{k \geq 0}$ is an orthonormal sequence in $L^2([0, 1])$.

In particular, we now fix a sequence $(c_j)_{j \geq 0}$ such that $\sum_{k=0}^{+\infty} |c_k|^2$ converges, we can define

$$f = \sum_{k=0}^{+\infty} c_k r_k$$

and this series converges in $L^2([0, 1])$ thus $f \in L^2([0, 1])$. We actually have a bit better:

Theorem 3.3. *If $\sum_{k=0}^{+\infty} |c_k|^2 < +\infty$ then there exists $f \in L^2([0, 1])$ defined by*

$$f = \sum_{k=0}^{+\infty} c_k r_k$$

and this series converges both in $L^2([0, 1])$ and almost everywhere.

Proof. The L^2 convergence has already been established. Further, let $F = \int f$ be the indefinite integral of f and let $E \subset [0, 1]$ be the set of Lebesgue points of f so that $|E| = 1$ and on E , F' exists and is finite.

Now let, $S_n[f]$ be the n -th partial sum of this series

$$S_n[f](x) = \sum_{k=0}^n c_k r_k(x).$$

As $S_n[f] \rightarrow f$ in $L^2([0, 1])$, for every $0 \leq a < b \leq 1$,

$$\left| \int_a^b (f(x) - S_n[f](x)) dx \right| \leq \int_0^1 |f(x) - S_n[f](x)| dx \leq \left(\int_0^1 |f(x) - S[f](x)|^2 dx \right)^{1/2} \rightarrow 0.$$

We have just shown that, if I is an interval, then $\int_I S_n[f] \rightarrow \int_I f$ thus also, if we fix $j \geq 1$, $\int_I (S_n[f] - S_{j-1}[f]) \rightarrow \int_I (f - S_{\ell-1}[f])$. On the other hand, if $I \in \mathcal{D}_{\ell-1}$ and $k \geq \ell$, then $\int_I r_k = 0$ so that $\int_I S_n[f] = \int_I S_{\ell-1}[f]$. Letting $n \rightarrow +\infty$ we obtain that

$$\int_I f(x) dx = \int_I S_{\ell-1}[f](x) dx \quad \text{for every } I \in \mathcal{D}_{\ell-1}.$$

Next, let $x_0 \in E$ not a dyadic rational ($x_0 \neq \frac{p}{2^q}$, $p, q \in \mathbb{N}$) and let $I_k =]j2^{-k}, (j+1)2^{-k}[$ be such that $x_0 \in E \cap I_k$. Then, as $S_{k-1}[f]$ is constant over I_k

$$S_{k-1}[f](x_0) = \frac{1}{|I_k|} \int_{I_k} S_k[f](x) dx = \frac{1}{|I_k|} \int_{I_k} f(x) dx \rightarrow F'(x_0)$$

when $k \rightarrow +\infty$. □

The second result is that f is actually in every L^p space:

Theorem 3.4. *Let $(c_k) \in \ell^2$ and $f = \sum_{k=0}^{+\infty} c_k r_k$. Then, for $1 \leq p < +\infty$, $f \in L^p([0, 1])$. Moreover, there exists A_p, B_p , depending on p only, such that*

$$A_p \left(\sum_{k=0}^{+\infty} |c_k|^2 \right)^{\frac{1}{2}} \leq \left(\int_0^1 |f(x)|^p dx \right)^{\frac{1}{p}} \leq B_p \left(\sum_{k=0}^{+\infty} |c_k|^2 \right)^{\frac{1}{2}}.$$

Proof. Let us first notice that the theorem holds for $p = 2$ since

$$\gamma := \left(\int_0^1 |f(x)|^2 dx \right)^{\frac{1}{2}} = \left(\sum_{k=0}^{+\infty} |c_k|^2 \right)^{\frac{1}{2}}$$

i.e. the inequalities are equalities with $A_2 = B_2 = 1$.

Next, let us notice that this implies the lower bound when $p > 2$ with $A_p = 1$ since then, with Hölder

$$\left(\int_0^1 |f(x)|^p dx \right)^{\frac{1}{p}} \geq \left(\int_0^1 |f(x)|^2 dx \right)^{\frac{1}{2}} = \gamma$$

and also implies the upper bound with $B_p = 1$ for $p < 2$ since now Hölder implies that

$$\left(\int_0^1 |f(x)|^p dx \right)^{\frac{1}{p}} \leq \left(\int_0^1 |f(x)|^2 dx \right)^{\frac{1}{2}} = \gamma$$

Further, as Hölder also implies that if $2(m-1) < p \leq 2m$ for some integer $m \geq 2$, then

$$\left(\int_0^1 |f(x)|^p dx \right)^{\frac{1}{p}} \leq \left(\int_0^1 |f(x)|^{2m} dx \right)^{\frac{1}{2m}}$$

so that the upper bound

$$(3.2) \quad \left(\int_0^1 |f(x)|^{2k} dx \right)^{\frac{1}{2k}} \leq B_{2k} \gamma$$

would also implies the upper bound for $2(m-1) < p \leq 2m$ with $B_p \leq B_{2k}$.

Next, let us show that the upper bound for $p = 4$ implies the lower bound for $p < 2$. Assume for the moment that we are able to prove that

$$\left(\int_0^1 |f(x)|^4 dx \right)^{\frac{1}{4}} \leq B_4 \gamma.$$

Let $1 \leq q < 2$ and write $2 = qt + 4(1-t)$, that is, take $t = \frac{2}{4-q}$. Then, from Hölder

$$\begin{aligned} \gamma^2 &= \int_0^1 |f(x)|^2 dx = \int_0^1 |f(x)|^{qt} |f(x)|^{4(1-t)} dx \leq \left(\int_0^1 |f(x)|^q dx \right)^t \left(\int_0^1 |f(x)|^4 dx \right)^{1-t} \\ &\leq (B_4 \gamma)^{4(1-t)} \left(\int_0^1 |f(x)|^q dx \right)^t = (B_4 \gamma)^{2-qt} \left(\int_0^1 |f(x)|^q dx \right)^t \end{aligned}$$

thus

$$\left(\int_0^1 |f(x)|^q dx \right)^{\frac{1}{q}} \geq B_4^{1-\frac{4-q}{q}} \gamma.$$

So it remains to prove (3.2). Notice also that it is enough to prove this inequality with real c_k 's. The constant in the complex case is then multiplied by 2: write $f = f_r + i f_i$ where $f_r = \sum \Re(c_k) r_k$ and $f_i = \sum \Im(c_k) r_k$. Then

$$\|f\|_{2m} \leq \|f_r\|_{2m} + \|f_i\|_{2m} \leq B_{2m}^{\mathbb{R}} \left[\left(\sum_{k=0}^{+\infty} |\Re(c_k)|^2 \right)^{\frac{1}{2}} + \left(\sum_{k=0}^{+\infty} |\Im(c_k)|^2 \right)^{\frac{1}{2}} \right] \leq 2 B_{2m}^{\mathbb{R}} \gamma$$

since $|\Re(c_k)|, |\Im(c_k)| \leq |c_k|$.

To conclude, we write

$$\int_0^1 S_n[f](x)^{2m} dx = \sum_{\ell_0 + \dots + \ell_n = 2m} A_{\ell_0, \dots, \ell_n} c_0^{\ell_0} \dots c_n^{\ell_n} \int_0^1 r_0^{\ell_0}(x) \dots r_n^{\ell_n}(x) dx$$

where $\ell_j \geq 0$ for every j and

$$A_{\ell_0, \dots, \ell_n} = \frac{(\ell_0 + \dots + \ell_n)!}{\ell_0! \dots \ell_n!}.$$

Now observe that

$$\int_0^1 r_0^{\ell_0}(x) \dots r_n^{\ell_n}(x) dx \begin{cases} 1 & \text{all the } \ell_j \text{'s are even} \\ 0 & \text{otherwise} \end{cases}$$

and that

$$\left(\sum_{k=0}^n c_k^2 \right)^m = \sum_{\ell_0 + \dots + \ell_n = m} A_{\ell_0, \dots, \ell_n} (c_0^2)^{\ell_0} \dots (c_n^2)^{\ell_n}.$$

Further, when $\ell_0 + \dots + \ell_n = m$,

$$\frac{A_{2\ell_0, \dots, 2\ell_n}}{A_{\ell_0, \dots, \ell_n}} = \frac{(m+1)(m+2) \dots 2m}{\prod_{j=0}^n (\ell_j + 1)(\ell_j + 2) \dots 2\ell_j} \leq \frac{(m+1)(m+2) \dots 2m}{2^m} \leq m^m$$

(with the convention that on the denominator is $(\ell_j + 1)(\ell_j + 2) \dots 2\ell_j = 1$ when $\ell_j = 0$). It follows that

$$\int_0^1 S_n[f](x)^{2m} dx \leq m^m \left(\sum_{k=0}^n |c_k|^2 \right)^m.$$

As $S_n[f] \rightarrow f$ a.e., we conclude that

$$\left(\int_0^1 |f(x)|^{2m} dx \right)^{\frac{1}{2m}} \leq m^{1/2} \left(\sum_{k=0}^{+\infty} |c_k|^2 \right)^{\frac{1}{2}}$$

that is $B_{2m} = 2m^{1/2}$. □

The estimate $B_{2m} = 2m^{1/2}$ allows to improve a bit the result:

Corollary 3.5. *Let $(c_k) \in \ell^2$ and $f = \sum_{k=0}^{+\infty} c_k r_j$. Then, for every $\mu > 0$, $\exp(\mu|f|^2) \in L^1([0, 1])$.*

Proof. Let us fix $\mu > 0$. We first show that if $\gamma := \|c_j\|_2$ is small enough, then $\exp(\mu|f|^2) \in L^1([0, 1])$. Indeed

$$(3.3) \quad \int_0^1 \exp(\mu|f(x)|^2) dx = \sum_{m=0}^{+\infty} \frac{\mu^m}{m!} \int_0^1 |f(x)|^{2m} dx \leq \sum_{m=0}^{+\infty} \frac{m^m}{m!} (4\mu\gamma^2)^m.$$

But $\frac{m^m}{m!} \leq \sum_{n=0}^{+\infty} \frac{m^n}{n!} = e^m$ so that

$$\int_0^1 \exp(\mu|f(x)|^2) dx \leq \sum_{m=0}^{+\infty} (4e\mu\gamma^2)^m = \frac{1}{1 - 4e\mu\gamma^2} < +\infty$$

provided $\gamma^2 < \frac{1}{4e\mu}$.

Next, take any $f \in L^1(0, 1)$, and apply the first part to $f - S_n[f] = \sum_{j=n+1}^{+\infty} c_j r_j$. As $\gamma_n^2 := \sum_{j=n+1}^{+\infty} |c_j|^2 \rightarrow 0$,

for n large enough $\gamma_n^2 < \frac{1}{8e\mu}$ thus $\exp(2\mu|f - S_n[f]|^2) \in L^1([0, 1])$.

Finally, as $|f|^2 \leq 2|f - S_n[f]|^2 + 2|S_n[f]|^2$, we have

$$\exp(\mu|f|^2) \leq \exp(2\mu|f - S_n[f]|^2) \exp(2\mu|S_n[f]|^2) \in L^1$$

since $|S_n[f]| \in L^\infty$ thus also $\exp(2\mu|S_n[f]|^2) \in L^\infty$. □

Next, we consider series of the form

$$\sum_{j=0}^{+\infty} c_j e^{2i\pi j t} r_j(x).$$

The idea is that such series are of the form $\sum \pm c_j e^{2i\pi j t}$, that is, choosing $x \in (0, 1)$ at random, we randomly change the sign of c_j . Our first result is the following:

Theorem 3.6. *Let $(c_k) \in \ell^2$ and $f_x(t) = \sum_{k=0}^{+\infty} c_k r_k(x) e^{2i\pi k t}$. Then, for almost every $x \in (0, 1)$, the series converges almost everywhere in $t \in (0, 1)$ and $f_x \in L^p([0, 1])$ for every $1 \leq p < +\infty$.*

Proof. Let E be the set of $(x, t) \in [0, 1]^2$ where the series defining f converges.

According to Theorem 3.3, for every $t \in [0, 1]$, the set $E_t^2 = \{(x, t) \in E\}$ has measure $|E_x| = 1$. It follows that $|E| = 1$ but then, for almost every $x \in [0, 1]$, $E_x^1 = \{(x, t) \in E\}$ has also measure $|E_x^1| = 1$.

Next, set $\gamma = \|c_k\|_2$ and fix $n \geq 1$. As in (3.3),

$$(3.4) \quad \frac{\mu^n}{n!} \int_0^1 |f_x(t)|^{2n} dx = \sum_{m=0}^{+\infty} \frac{\mu^m}{m!} \int_0^1 |f_x(t)|^{2m} dx = \int_0^1 \exp(\mu|f_x(t)|^2) dx \leq \frac{1}{1 - 4e\mu\gamma^2}$$

provided $\mu < \frac{1}{4e\gamma^2}$. It follows that

$$\int_0^1 \int_0^1 |f_x(t)|^{2n} dt dx = \int_0^1 \int_0^1 |f_x(t)|^{2n} dx dt \leq \frac{n!}{(1 - 4e\mu\gamma^2)\mu^n} < +\infty.$$

But then, for every n , there is a set $F_n \subset [0, 1]$ with $|F_n| = 0$ such that, if $x \in [0, 1] \setminus F_n$, $\int_0^1 |f_x(t)|^{2n} dt < +\infty$. Setting $F = \bigcup F_n$, $|F| = 0$ and, for every $x \in [0, 1] \setminus F$, for every n , $f_x \in L^{2n}$. Using the inclusion

of $L^{2n}([0, 1]) \subset L^p([0, 1])$ when $p \leq 2n$, we obtain that, for almost every x , $f_x \in L^p([0, 1])$ for every $p \geq 1$, as claimed. \square

We can now prove the main result of this section:

Theorem 3.7. *Let $q > 1$ and $(n_j)_{j \geq 0}$ be a q -lacunary sequence of integers, $n_0 \geq 1$ and $n_{j+1} \geq qn_j$. Let $1 \leq p < +\infty$. There are two constants $A_{p,q}, B_{p,q}$ such that, if $(c_j)_{j \geq 0} \in \ell^2$, then $g(t) = \sum_{j \geq 0} c_j e^{2i\pi n_j t}$ is in $L^p([0, 1])$ with*

$$(3.5) \quad A_{p,q} \left(\sum_{j=0}^{+\infty} |c_j|^2 \right)^{\frac{1}{2}} \leq \left(\int_0^1 \left| \sum_{j \geq 0} c_j e^{2i\pi n_j t} \right|^p dt \right)^{\frac{1}{p}} \leq B_{p,q} \left(\sum_{j=0}^{+\infty} |c_j|^2 \right)^{\frac{1}{2}}.$$

Remark 3.8. Note that a simple change of variable also shows that, for every integer M ,

$$(3.6) \quad A_{p,q} \left(\sum_{j=0}^{+\infty} |c_j|^2 \right)^{\frac{1}{2}} \leq \left(\frac{1}{M} \int_{-M/2}^{M/2} \left| \sum_{j \geq 0} c_j e^{2i\pi \frac{n_j}{M} t} \right|^p dt \right)^{\frac{1}{p}} \leq B_{p,q} \left(\sum_{j=0}^{+\infty} |c_j|^2 \right)^{\frac{1}{2}}.$$

Also, we may assume that $q \rightarrow A_{p,q}, B_{p,q}$ are continuous.

Proof. The beginning of the proof is the same as for Theorem 3.4. Parseval's identity shows that (3.5) is satisfied when $p = 2$ with $A_{2,q} = B_{2,q} = 1$. The lower bound is then automatically satisfied for $p \geq 2$ with $A_{p,q} = 1$ while the upper bound is satisfied for $p \leq 2$ with $B_{2,q} = 1$. Finally, if we establish the upper bound for $p > 2$, using Hölder's inequality in the same way as in the proof of Theorem 3.4, the lower bound follows for $p < 2$ with $A_{2,q} = B_{4,q}^{1-\frac{4-p}{p}}$. Also, it is enough to prove the upper bound when $p = 2m$, $m \geq 2$ and then, if $2(m-1) < p \leq 2m$, $B_{p,q} \leq B_{2k,q}$. Another reduction is that, by homogeneity, it is enough to prove the theorem when $\sum_{j=0}^{+\infty} |c_j|^2 = 1$.

A further restriction is that it is enough to prove the theorem for $q \geq 3$. Indeed, for $1 < q < 3$, we introduce an integer N_q such that $q^{N_q} \geq 3$ and write $n_k^{(\ell)} = n_{kN_q + \ell}$ for $\ell = 0, \dots, N_q - 1$. Then $n_{k+1}^{(\ell)} \geq q^{N_q} n_k^{(\ell)}$. If the theorem is established when $q \geq 3$ then, for each ℓ , the upper bound in (3.5) reads

$$\left(\int_0^1 \left| \sum_{k \geq 0} c_{kN_q + \ell} e^{2i\pi n_k^{(\ell)} t} \right|^p dt \right)^{\frac{1}{p}} \leq B_{p,q^{N_q}} \left(\sum_{k=0}^{+\infty} |c_{kN_q + \ell}|^2 \right)^{\frac{1}{2}}.$$

But then, with the triangular inequality in L^p ,

$$\begin{aligned} \left(\int_0^1 \left| \sum_{j \geq 0} c_j e^{2i\pi n_j t} \right|^p dt \right)^{\frac{1}{p}} &= \left(\int_0^1 \left| \sum_{\ell=0}^{N_q-1} \sum_{k \geq 0} c_{kN_q + \ell} e^{2i\pi n_k^{(\ell)} t} \right|^p dt \right)^{\frac{1}{p}} \\ &\leq \sum_{\ell=0}^{N_q-1} \left(\int_0^1 \left| \sum_{k \geq 0} c_{kN_q + \ell} e^{2i\pi n_k^{(\ell)} t} \right|^p dt \right)^{\frac{1}{p}} \\ &\leq B_{p,q^{N_q}} \sum_{\ell=0}^{N_q-1} \left(\sum_{k=0}^{+\infty} |c_{kN_q + \ell}|^2 \right)^{\frac{1}{2}} \\ &\leq N_q^{1/2} B_{p,q^{N_q}} \left(\sum_{\ell=0}^{N_q-1} \sum_{k=0}^{+\infty} |c_{kN_q + \ell}|^2 \right)^{\frac{1}{2}} \\ &= N_q^{1/2} B_{p,q^{N_q}} \left(\sum_{j \geq 0} |c_j|^2 \right)^{\frac{1}{2}}, \end{aligned}$$

where we have used Cauchy-Schwarz in \mathbb{R}^{N_q} in the next to last line.

A last reduction comes from the observation that, for every k

$$\int_0^1 \exp(\mu|g(t)|^2) dt = \sum_{n=0}^{+\infty} \frac{\mu^n}{n!} \int_0^1 |g(t)|^{2n} dt \geq \frac{\mu^m}{m!} \int_0^1 |g(t)|^{2m} dt.$$

It is therefore enough to prove that there is a $\mu(q)$ and a $C > 0$ such that, if $\mu < \mu(q)$

$$(3.7) \quad \int_0^1 \exp(\mu|g(t)|^2) dt \leq C$$

which would then imply that

$$\int_0^1 |g(t)|^{2k} dt \leq C \frac{m!}{\mu^m}$$

as desired.

In order to prove (3.7), let us introduce

$$f_x(t) = \sum_{j \geq 0} c_j r_{n_j}(x) e^{2i\pi n_j t}.$$

Integrating (3.4) with respect to t and using Fubini, we deduce that

$$\int_0^1 \int_0^1 \exp(\mu|f_x(t)|^2) dt dx \leq K := \frac{1}{1 - 4e\mu\gamma^2}.$$

But then, there is an x_0 (that we can assume not to be a dyadic rational $x_0 \neq 2^j/k$) such that

$$\int_0^1 \exp(\mu|f_{x_0}(t)|^2) dt \leq K.$$

Next, we consider the Riesz product

$$P_k(x) = \prod_{j=0}^k (1 + r_{n_j}(x_0) \cos 2\pi n_j t) = \prod_{k=0}^m \left(1 + r_{n_k}(x_0) \frac{e^{2\pi n_k t} + e^{-2\pi n_k t}}{2} \right) = \sum_{j \in \mathbb{Z}} \gamma_j e^{2i\pi j t}$$

where the Fourier coefficients have the following property:

$$- \gamma_0 = 1$$

$$- \gamma_j = 0 \text{ if } j \text{ is an integer that is not of the form } \sum \pm n_\ell, \text{ in particular when } |j| > \sum_{k=0}^n n_k$$

$$- \text{if } j = \sum \varepsilon_\ell n_\ell \text{ with } \varepsilon_\ell \in \{-1, 0, 1\}. \text{ As } q > 3, \text{ this } \varepsilon_\ell \text{'s are unique. Then } \gamma_j = \prod_{\varepsilon_j \neq 0} \frac{r_{n_j}(x_0)}{2}. \text{ In}$$

particular, $\gamma_{n_j} = \frac{r_{n_j}(x_0)}{2}$ for $j = 0, \dots, k$ and $\gamma_{n_j} = 0$ for $j > k$.

As a consequence, the partial sums of the Fourier series of g are given by

$$S_{n_k}[g](t) := \sum_{j=0}^k c_j e^{2i\pi n_j t} = \sum_{j=0}^k c_j r_j(x_0)^2 e^{2i\pi n_j t} = 2 \int_0^1 f_{x_0}(s) P_k(t-s) ds.$$

Note that $P_k \geq 0$ and $\int_0^1 P_k(t) dt = \gamma_0 = 1$ so that $\nu_k = P_k(t) dt$ is a probability measure. As $\varphi(s) = \exp(\mu s^2)$ is increasing and convex, we apply Jensen's inequality (with the measure ν_k) to obtain

$$\varphi\left(\frac{1}{2} S_{n_k}[g](t)\right) \leq \varphi\left(\int_0^1 |f_{x_0}(s)| P_k(t-s) ds\right) \leq \int_0^1 \varphi(|f_{x_0}(s)|) |P_k(t-s)| ds.$$

Integrating over $[0, 1]$ and using Fubini, we get

$$\int_0^1 \varphi\left(\frac{1}{2} S_{n_k}[g](t)\right) dt \leq \int_0^1 \varphi(|f_{x_0}(s)|) \int_0^1 P_k(t-s) dt ds = \int_0^1 \varphi(|f_{x_0}(s)|) ds \leq K.$$

Letting $k \rightarrow +\infty$, we obtain

$$\int_0^1 \exp\left(\frac{\mu}{2}|g(t)|^2\right) dt \leq K$$

as claimed (up to $\mu/2$ replacing μ). \square

3.3. The proof of Littlewood's conjecture by Mc Gehee, Pigno and Smith. We will now give the proof of the Littlewood conjecture, following closely [4].

Theorem 3.9 (Mc Gehee, Pigno, Smith). *There exists a constant $C_{MPS} \leq \frac{4}{\pi^2}$ such that, if $(n_k)_{k \in \mathbb{N}}$ is an increasing sequence of integers and (a_k) is a complex sequence with finite support, then*

$$C_{MPS} \sum_{k=0}^{+\infty} \frac{|a_k|}{k+1} \leq \int_0^1 \left| \sum_{k=0}^{+\infty} a_k e^{2i\pi n_k t} \right| dt.$$

The proof given below will give $C_{MPS} = \frac{1}{96}$ which is not the best possible. We avoid minor technicalities which would lead to $C_{MPS} = \frac{1}{30}$ (see e.g. [7, 24]). The proof in [13] requires the introduction of various parameters and a cumbersome optimization of those parameters in the final step and leads to $C_{MPS} = \frac{1}{26}$.

Note also that the sum starts at 0, we have so far been unable to obtain a lower bound when the sum runs over \mathbb{Z} .

3.3.1. Strategy of the proof.

Notation 3.10. For a function $F \in L^2([0, 1])$ and $s \in \mathbb{Z}$, we write

$$\widehat{F}(s) = c_s(F) = \int_0^1 F(t) e^{-2i\pi s t} dt$$

for the Fourier coefficients of F . Its Fourier series is then

$$F(t) = \sum_{s \in \mathbb{Z}} c_s(F) e^{2i\pi s t}$$

and Parseval's relation reads

$$\int_0^1 |F(t)|^2 dt = \sum_{s \in \mathbb{Z}} |c_s(F)|^2$$

and can also be written in the form

$$\int_0^1 F(t)G(t) dt = \sum_{s \in \mathbb{Z}} c_s(F)c_{-s}(G).$$

We fix a trigonometric polynomial

$$(3.8) \quad \phi(t) = \sum_{k=0}^N a_k e^{2i\pi n_k t} \quad \text{and} \quad S = \sum_{k=0}^N \frac{|a_k|}{k+1}.$$

We then write $|a_k| = a_k u_k$ with u_k complex numbers of modulus 1 and we introduce

$$(3.9) \quad T_0(t) = \sum_{k=0}^N \frac{u_k}{k+1} e^{-2i\pi n_k t}.$$

Then by Parseval identity, written in the second form we have

$$(3.10) \quad S := \sum_{k=0}^N \frac{|a_k|}{k+1} = \sum_{k=0}^N \widehat{\phi}(n_k) \widehat{T_0}(-n_k) = \int_0^1 \phi(t) T_0(t) dt.$$

so that

$$S \leq \|\phi\|_1 \|T_0\|_{L^\infty([0,1])}.$$

The issue is that, typically we have no control over the L^∞ norm of T_0 other than the trivial and explosive control by $\sum \frac{1}{k+1}$, so we will correct T_0 into another test function T_1 as follows;

(1) The L^∞ norm of the corrected function T_1 is controlled by a constant C , $\|T_1\|_\infty \leq C$.

(2) $\widehat{T}_1(-n_k)$ only differs a little from $\widehat{T}_0(-n_k)$ for $0 \leq k \leq N$, say $|\widehat{T}_1(-n_k) - \widehat{T}_0(-n_k)| \leq \frac{\widehat{T}_0(-n_k)}{2} = \frac{1}{2(k+1)}$, while we impose no condition on the behavior of $\widehat{T}_1(n)$ for $n \neq -n_k$.

We would then conclude as follows:

$$\left| S - \int_0^1 \phi(t)T_1(t) dt \right| = \left| \int_0^1 \phi(t)(T_0 - T_1)(t) dt \right| = \left| \sum_{k=0}^N \widehat{\phi}(\lambda_k)(\widehat{T}_0(-n_k) - \widehat{T}_1(-n_k)) \right|$$

with Parseval. With the triangular inequality and our expected estimate $|\widehat{T}_1(-n_k) - \widehat{T}_0(-n_k)| \leq \frac{1}{2(k+1)}$, we then conclude that

$$\left| S - \int_0^1 \phi(t)T_1(t) dt \right| \leq \sum_{k=1}^N |\widehat{\phi}(\lambda_k)| |\widehat{T}_0(-n_k) - \widehat{T}_1(-n_k)| \leq \sum_{k=1}^N \frac{|a_k|}{2(k+1)} = \frac{S}{2}.$$

But then

$$\frac{1}{2}S \leq \left| \int_0^1 \phi(t)T_1(t) dt \right| \leq \|T_1\|_\infty \int_0^1 |\phi(t)| dt \leq C \int_0^1 |\phi(t)| dt$$

which is the expected result with $A = \frac{1}{2C}$.

The proof of the Theorem B thus amounts to proving the following lemma:

Lemma 3.11. *There exists a universal constant C and a $T_1 \in L^\infty$ such that*

(1) $\|T_1\|_\infty \leq C$

(2) $|\widehat{T}_1(-n_k) - \widehat{T}_0(-n_k)| \leq \frac{1}{2}|\widehat{T}_0(-n_k)| = \frac{1}{2(k+1)}$ for $0 \leq k \leq N$

where T_0 is the function given by (3.9).

The remaining of this section is devoted to the proof of this lemma.

3.3.2. *Proof of Lemma 3.11.* First note that, up to eventually adding extra zeros to the sequence (a_k) and adding $\lambda_{N+j} = \lambda_N + j$ to the sequence (λ_k) , we may assume that $N = 2^{m+1} - 2$.

We start by decomposing T_0 into a sum of dyadic blocs on which the amplitude $|\widehat{T}_0(-\lambda_k)| = \frac{1}{k+1}$ is more or less constant. More precisely, for $j = 0, \dots, m$ we set $I_j = [2^j, 2^{j+1}[$ and

$$f_j = \sum_{k+1 \in I_j} \frac{u_k}{k+1} e^{-2i\pi n_k t}.$$

The function T_0 now appears as the partial sum of order m of the series $\sum f_j$, in other words

$$T_0 = \sum_{j=0}^m f_j.$$

Let us start with a simple lemma:

Lemma 3.12. *With the above notations, we have*

(1) $\|f_j\|_{L^\infty([0,1])} \leq 1$.

(2) $\|f_j\|_{L^2([0,1])} \leq 2^{-j/2}$.

Proof. By construction, $|I_j| = 2^j$ hence

$$\|f_j\|_\infty \leq \sum_{k+1 \in I_j} \frac{1}{k+1} \leq \frac{|I_j|}{2^j} = 1.$$

On the other hand, Parseval implies that

$$\|f_j\|_2^2 = \sum_{k+1=2^j}^{2^{j+1}-1} \frac{1}{(k+1)^2} \leq \frac{2^j}{4^j} = 2^{-j}$$

as claimed. \square

Now, write the Fourier series of each $|f_j| \in L^2([0, 1])$

$$|f_j| = \sum_{s \in \mathbb{Z}} c_s(|f_j|) e^{2i\pi s t}.$$

To each $|f_j|$, we associate $h_j \in L^2([-\pi, \pi])$ defined via its Fourier series as

$$h_j(t) = c_0(|f_j|) + 2 \sum_{s=1}^{\infty} c_s(|f_j|) e^{2i\pi s t}.$$

Lemma 3.13. *For $0 \leq j \leq n$, the following properties hold*

- (1) $\operatorname{Re}(h_j) = |f_j| \leq 1$;
- (2) $\|h_j\|_{L^2([0,1])} \leq \sqrt{2} \|f_j\|_{L^2([0,1])}$.

Proof. First, as $|f_j|$ is real valued, $c_0(|f_j|)$ is also real while $\overline{c_s(|f_j|)} = c_{-s}(|f_j|)$ for every $s \geq 1$. Hence

$$\overline{h_j}(t) = c_0(|f_j|) + 2 \sum_{s=1}^{\infty} c_{-s}(|f_j|) e^{-ist}$$

and thus

$$\operatorname{Re}(h_j) = \frac{h_j + \overline{h_j}}{2} = c_0(|f_j|) + \sum_{s \neq 0} c_s(|f_j|) e^{ist} = |f_j| \leq 1$$

by lemma 3.12.

By Parseval's identity and again $\overline{c_s(|f_j|)} = c_{-s}(|f_j|)$,

$$\begin{aligned} \|h_j\|_2^2 &= \left| c_0(|f_j|) \right|^2 + 4 \sum_{s=1}^{\infty} \left| c_s(|f_j|) \right|^2 = \left| c_0(|f_j|) \right|^2 + 2 \sum_{s=1}^{\infty} \left| c_s(|f_j|) \right|^2 + 2 \sum_{s=1}^{\infty} \left| c_{-s}(|f_j|) \right|^2 \\ &\leq 2 \sum_{s \in \mathbb{Z}} |c_s(|f_j|)|^2 = 2 \|f_j\|_2^2 \end{aligned}$$

as claimed. \square

We now define a sequence $(F_j)_{j=0, \dots, n}$ inductively through

$$F_0 = f_0 \quad \text{and} \quad F_{j+1} = F_j e^{-\eta h_{j+1}} + f_{j+1}$$

where $0 < \eta \leq 1$ is a real number to be adjusted later on. Further set

$$E_\eta := \sup_{0 < x \leq 1} \frac{x}{1 - e^{-\eta x}} = \frac{1}{\eta} \sup_{0 < x \leq \eta} \frac{x}{1 - e^{-x}} = \frac{1}{1 - e^{-\eta}} \leq \frac{2}{\eta}.$$

Lemma 3.14. *For $0 \leq j \leq n$, $\|F_j\|_\infty \leq \frac{2}{\eta}$.*

Proof. By definition of E_η , if $C \leq E_\eta$ and $0 \leq x \leq 1$, then $C e^{-\eta x} + x \leq E_\eta e^{-\eta x} + x \leq E_\eta$.

We can now prove by induction over j that $|F_j| \leq E_\eta$. First, when $j = 0$, from Lemma 3.12 we get

$$\|F_0\|_\infty = \|f_0\|_\infty \leq 1 \leq E_\eta.$$

Assume now that $\|F_j\|_\infty \leq E_\eta$, then

$$\begin{aligned} |F_{j+1}(t)| &= |F_j(t) e^{-\eta h_{j+1}(t)} + f_{j+1}(t)| \leq |F_j(t)| e^{-\eta \Re(h_{j+1}(t))} + |f_{j+1}(t)| \\ &= |F_j(t)| e^{-\eta |f_{j+1}(t)|} + |f_{j+1}(t)|. \end{aligned}$$

As $|f_{j+1}(t)| \leq 1$ and $|F_j(t)| \leq E_\eta$, we get $|F_{j+1}(t)| \leq E_\eta$. It remains to prove that $E_\eta \leq \frac{2}{\eta}$. To do so, it suffices to see that

$$e^{-y} \leq 1 - \left(\frac{e-1}{e}\right)y \quad \text{for } 0 \leq y \leq 1,$$

yielding the result immediately. \square

Lemma 3.15. For $0 \leq \ell \leq n$ and $j = 0, \dots, k$, let $g_{j,k} = e^{-\eta H_{j,k}}$ with

$$H_{j,k} = \begin{cases} h_{j+1} + \dots + h_k & \text{when } j < k \\ 0 & \text{when } j = k \end{cases}.$$

Then

$$F_k = \sum_{j=0}^k f_j g_{j,k}.$$

Proof. By induction on k , when $k = 0$, $H_{0,0} = 0$ thus $g_{0,0} = 1$ and, indeed, we have

$$F_0 = f_0 = f_0 g_{0,0}.$$

Assume now that the formula has been established at rank $k-1$ and let us show that $F_k = \sum_{j=0}^k f_j g_{j,k}$. By construction, we have

$$F_k = F_{k-1} e^{-\eta h_k} + f_k = \left(\sum_{j=0}^{k-1} f_j g_{j,k-1} \right) e^{-\eta h_k} + f_k.$$

with the induction hypothesis. It remains to notice that $g_{k,k} = e^{-\eta H_{k,k}} = 1$ and that, for $j = 0, \dots, k-1$, $H_{j,k} = H_{j,k-1} + h_k$ thus $g_{j,k} = g_{j,k-1} e^{-\eta h_k}$ so that, indeed, we have $F_k = \sum_{j=0}^k f_j g_{j,k}$ as claimed. \square

Recall that

$$T_0 = \sum_{j=0}^m f_j$$

and we set

$$T_1^\eta = F_m = \sum_{j=0}^m f_j g_{j,m}$$

where the dependence on η comes from the definition of the $g_{j,n}$'s, in particular

$$\|T_1^\eta\|_\infty \leq E_\eta.$$

The first part of Lemma 3.11 is thus established and it remains to prove the second part. To do so, we start by some intermediary results;

Lemma 3.16. If $H \in H^\infty$ (Hardy space) and $\operatorname{Re}(H) \geq 0$, then $e^{-H} \in H^\infty$ and

$$\|e^{-H} - 1\|_2 \leq \|H\|_2.$$

Proof. Since H^∞ is a Banach algebra, the partial sums $\sum_{k=0}^n (-1)^k \frac{H^k}{k!}$ of e^{-H} are elements of H^∞ . Moreover, since H is bounded, these sommes converge uniformly toward e^{-H} , with $e^{-H} \in H^\infty$. Finally, if $z \in \mathbb{C}$ and $\Re(z) \geq 0$,

$$|e^{-z} - 1| = \left| \int_0^1 z e^{-tz} dt \right| \leq \int_0^1 |z| e^{-t\Re(z)} dt \leq |z|.$$

In our case $z = H(t)$, and we have

$$|e^{-H(t)} - 1| \leq |H(t)|$$

and by integration we have the desired inequality. \square

Next let us introduce the following notations:

- (1) Let $f \in L^1([0, 1])$, the spectrum of f , denoted by $\text{spec}(f)$ is the set of non zero Fourier coefficients, more precisely

$$\text{spec}(f) = \{n \in \mathbb{Z} : \widehat{f}(n) \neq 0\}.$$

It is easy to show that, if $f, g \in L^2([0, 1])$, then

$$\text{spec}(fg) \subset \text{spec}(f) + \text{spec}(g) = \{\lambda + \mu : \lambda \in \text{spec}(f), \mu \in \text{spec}(g)\}.$$

- (2) Let A be a subset of $[1, N[$, we denote by Λ_A the set of $n_j, j \in A$ i.e

$$\Lambda_A = \{n_j, j \in A\}.$$


Lemma 3.17. Let $k+1 \in I_\ell = [2^\ell, 2^{\ell+1}[$ then $\widehat{f_j g_{j,m}}(-n_k) = 0$ if $j < \ell$.

Proof. We must show that $-n_k \notin \text{spec}(f_j g_{j,m})$. By contradiction, we suppose that $-n_k \in \text{sp}(f_j g_j)$.

But since $\text{spec}(f_j) \subset -\Lambda_{I_j}$ and, from Lemma 3.16, $\text{spec}(g_{j,m}) \subset \mathbb{N}$, thus

$$\text{spec}(f_j g_{j,m}) \subset -\Lambda_{I_j} + \mathbb{N}.$$

However, since $j < \ell$, I_j is to the left of I_ℓ , then $-\Lambda_{I_\ell}$ is completely to the left of $-\Lambda_{I_j} + \mathbb{N}$ (since $k+1 \in I_\ell$, then $-n_k \in -\Lambda_{I_\ell}$ by definition of Λ_{I_ℓ}).



□

Proof of (2) in Lemma 3.11. Let $k \in [0, N[$ and $\ell \leq m$ such that $k+1 \in I_\ell$. Hence

$$\widehat{T_1}(-n_k) = \sum_{j=0}^m \widehat{f_j g_{j,m}}(-n_k) = \sum_{j=\ell}^m \widehat{f_j g_{j,m}}(-n_k)$$

with Lemma 3.17.

On the other hand, $\widehat{T_0}(-n_k) = \frac{u_k}{k+1}$ while $\widehat{f_j}(-n_k) = \frac{u_k}{k+1}$ if $k+1 \in I_j$ i.e. if $j = \ell$ and $\widehat{f_j}(-n_k) = 0$ otherwise. So we can write

$$\widehat{T_0}(-n_k) = \sum_{j=\ell}^m \widehat{f_j}(-n_k),$$

hence

$$\left| \widehat{T_1}(-n_k) - \widehat{T_0}(-n_k) \right| = \sum_{j=\ell}^m [f_j(\widehat{g_{j,m}} - 1)](-n_k) = \sum_{j=\ell}^m c_{-n_k} [f_j(g_{j,m} - 1)].$$

But

$$\begin{aligned} |c_{-n_k} [f_j(g_{j,m} - 1)]| &\leq \|f_j(g_{j,m} - 1)\|_{L^1([0,1])} \\ &\leq \|f_j\|_{L^2([0,1])} \|g_{j,m} - 1\|_{L^2([0,1])} = \|f_j\|_{L^2([0,1])} \|e^{-\eta H_{j,m}} - 1\|_{L^2([0,1])} \end{aligned}$$

by definition of $g_{j,m}$. Using Lemma 3.16 and then Lemma 3.15, we obtain

$$\begin{aligned} |c_{-n_k} [f_j(g_{j,m} - 1)]| &\leq \eta \|f_j\|_2 \|H_{j,m}\|_2 \leq \eta \|f_j\|_2 \sum_{r=j+1}^m \|h_r\|_2 \\ &\leq \eta \|f_j\|_2 \sqrt{2} \sum_{r=j+1}^m \|f_r\|_2 \end{aligned}$$

with Lemma 3.13. Then, from Lemma 3.12, we obtain

$$|c_{-n_k} [f_j(g_{j,m} - 1)]| \leq \eta 2^{-j/2} \sqrt{2} \sum_{r=j+1}^{+\infty} 2^{-r/2} = \eta 2^{-j/2} \sqrt{2} \frac{2^{-(j+1)/2}}{1 - 2^{-1/2}}.$$

We conclude that

$$\left| \widehat{T}_1(-\lambda_k) - \widehat{T}_0(-\lambda_k) \right| \leq 3\eta \sum_{j=\ell}^m 2^{-j} \leq 6\eta 2^{-\ell}.$$

On the other hand, as $k+1 \in I_\ell = [2^\ell, 2^{\ell+1}[$,

$$|\widehat{T}_0(-\lambda_k)| = \frac{1}{k+1} > 2^{-\ell-1}$$

so that

$$\left| \widehat{T}_1(-\lambda_k) - \widehat{T}_0(-\lambda_k) \right| \leq 12\eta |\widehat{T}_0(-\lambda_k)|.$$

Choosing $\eta = \frac{1}{24}$ gives the result. \square

Note that, when $\eta = \frac{1}{24}$, $E_\eta \leq 48$ so that we can take $C = 48$ in Lemma 3.11, leading to $C_{MPS} = \frac{1}{96}$ in Theorem 3.9.

3.4. Strongly multidimensional sets. Let $\delta > 0$ and $(m, n) \in \mathbb{N}^2$. A subset A of \mathbb{Z} is $(\delta; m, n)$ -strongly 2-dimensional if there exists numbers d and D with $D > (2 + \delta)d$ such that

$$A = \bigcup_{k \in I} (A_k + kD)$$

for some set I containing at least m integers and subsets $A_k \subseteq \{-d, \dots, d\}$ verifying $|A_k| \geq n$.

Theorem 3.18 (Hanson [10]). *Let $\delta > 0$ and m, n be two positive integers satisfying*

$$(3.11) \quad n \geq \pi^3 2^{21} C_{MPS}^3 \ln(n)^3 \quad \text{and} \quad m \geq \pi^3 2^{21} C_{MPS}^3 \ln(n)^3 \ln(m)^3,$$

where C_{MPS} is the constant in theorem B. Suppose A is $(\delta; m, n)$ strongly 2-dimensional subset of \mathbb{Z} . Then

$$\int_0^1 \left| \sum_{a \in A} e^{2i\pi a t} \right| dt \geq \frac{C_{MPS}^2}{(2^9 \pi)^2 \left(2 + \ln\left(1 + \frac{2}{\delta}\right) \right)} \ln(m) \ln(n)$$

Combining this result with Theorem 3.3 in [23], we see that this theorem is also best possible up to the constant.

Given a set $I \subseteq \mathbb{Z}$, a positive integer q , and an arbitrary integer s , we define

$$I(q; s) = \{k \in I : k \equiv s \pmod{q}\}.$$

The proof of Theorem 3.18 is a direct consequence of two lemmas. The first one is the following:

Lemma 3.19. *Let I be a set of integers with $|I| \geq 8$. Then there are positive integers q and s such that*

$$\frac{|I|^{\frac{1}{3}}}{8} \leq |I(q; s)| \leq q^{1/2}.$$

Proof. For each $j \geq 1$, we choose any s_j such that $|I(4^j; s_j)|$ is maximal. But, on one hand,

$$I = \bigcup_{s=0}^{4^j-1} I(4^j, s)$$

and on the other hand, for j fixed, the sets $I(4^j; s)$ are disjoint, so at least one of them has cardinality larger than $4^{-j}|I|$. In particular,

$$(3.12) \quad |I(4^j, s_j)| \geq 4^{-j}|I|.$$

For $j = 1$, we thus have $|I(4; s_1)| \geq \frac{|I|}{4} \geq 2$. On the other hand, if $j \equiv s \pmod{kp}$ then $j \equiv s \pmod{p}$ so that, for any s

$$I(4^m; s) \subset I(4^\ell; s) \quad \text{for } \ell < m,$$

and, for sufficiently large j we have $|I(4^j; s_j)| = 1 \leq 2^j$. Therefore, there exists a minimal j_0 such that $|I(4^{j_0}; s_{j_0})| \leq 2^{j_0}$. Let $q = 4^{j_0}$, and $s = s_{j_0}$ then using (3.12) and the definition of j_0

$$\frac{|I|}{q} \leq |I(q; s)| = |I(4^{j_0}; s_{j_0})| \leq 2^{j_0} = q^{\frac{1}{2}}.$$

In particular

$$(3.13) \quad |I|^{\frac{1}{3}} \leq q^{\frac{1}{2}}.$$

By minimality of $j_0 - 1$

$$\frac{q^{\frac{1}{2}}}{2} = 2^{j_0-1} \leq \left| I \left(4^{j_0-1}; s_{j_0-1} \right) \right| \leq \sum_{r=0}^3 \left| I \left(4^{j_0}; s_{j_0-1} + r4^{j_0-1} \right) \right| \leq 4|I(4^{j_0}; s_{j_0})| = 4|I(q; s)|$$

by definition of s_{j_0} . We thus get

$$|I(q; s)| \geq \frac{q^{\frac{1}{2}}}{8} \geq \frac{|I|^{\frac{1}{3}}}{8},$$

with (3.13). □

Lemma 3.20. *Let $\delta > 0$ and let d and D be positive integers with $(2 + \delta)d < D$. Suppose I is a finite set of integers, and let*

$$F(t) = \sum_{k \in I} f_k(t) e^{2i\pi Dkt}$$

where

$$f_k(t) = \sum_{|n| \leq d} a_{n,k} e^{2i\pi nt}$$

Let q and s with $q > 4\pi$ and suppose $I(q; s) = \{k_1, \dots, k_J\}$ then we have

$$\|F\|_{L^1([0,1])} \geq \frac{1}{32\pi(2 + \ln(1 + \frac{2}{\delta}))} \sum_{j=1}^J \|f_{k_j}\|_{L^1([0,1])} \left(\frac{C_{MPS}}{2j} - \frac{2\pi d}{qD} \right)$$

We split the remaining of this section into two parts. In the first one, we show that Lemmas 3.19 and 3.20 imply Theorem 3.11. In the second one, we prove the Lemma 3.20.

3.4.1. *Proof of Theorem 3.18.* Let $\delta > 0$, m, n be two integers satisfying the conditions of the theorem and let A be strongly (δ, m, n) -regular. Thus, there are two integers d, D with $D > (2 + \delta)d$, such that we can write

$$A = \bigcup_{k \in I} (A_k + kD),$$

with $|I| \geq m$ and $A_k \subset \{-d, \dots, d\}$ with $|A_k| \geq n$. We can then write

$$F(t) := \sum_{a \in A} e^{2i\pi at} = \sum_{k \in I} f_k(t) e^{2i\pi Dkt}$$

with

$$f_k(t) = \sum_{a \in A_k} e^{2i\pi at} = \sum_{n=-d}^d a_{n,k} e^{2i\pi nt}$$

with $a_{n,k} = 1$ if $n \in A_k$ and $a_{n,k} = 0$ otherwise.

Assume first that there exists $k_1 \in I$ such that

$$(3.14) \quad \|f_{k_1}\|_{L^1([0,1])} \geq \frac{C_{MPS}}{2^9 \pi} \ln(m) \ln(n).$$

We then choose $q \geq \frac{16\pi}{7C_{MPS}}$ in such a way that there is an s such that $I(q, s) = \{k_1\}$. Hence by Lemma 3.20,

$$\begin{aligned} \|F\|_{L^1((0,1))} &\geq \frac{\|f_{k_1}\|_{L^1((0,1))}}{32\pi \left(2 + \ln\left(1 + \frac{2}{\delta}\right)\right)} \left(\frac{C_{MPS}}{2} - \frac{2\pi d}{qD}\right) \\ &\geq \frac{C_{MPS}\|f_{k_1}\|_{L^1((0,1))}}{2^6\pi \left(2 + \ln\left(1 + \frac{2}{\delta}\right)\right)} \left(1 - \frac{7d}{8D}\right). \end{aligned}$$

As $D > 2d$, using (3.14), we conclude that, in this case

$$\|F\|_{L^1((0,1))} \geq \left(\frac{C_{MPS}}{2^9\pi}\right)^2 \frac{\ln n \ln m}{2 + \ln\left(1 + \frac{2}{\delta}\right)}$$

which establishes the theorem.

We will thus assume that, for each $k \in I$,

$$(3.15) \quad \|f_k\|_{L^1((0,1))} \leq \frac{C_{MPS}}{2^9\pi} \ln(m) \ln(n).$$

Note that, from Theorem B,

$$\|f_k\|_{L^1((0,1))} \geq C_{MPS} \ln(n)$$

so that $2^9\pi \leq \ln(m)$, in particular, $m \geq 8$. We then take q and s given by Lemma 3.19 applied to the set I so that $J = |I(q; s)|$ satisfies

$$(3.16) \quad \frac{m^{\frac{1}{3}}}{8} \leq J \leq q^{\frac{1}{2}}.$$

We write

$$I(q; s) = \{k_1 < \dots < k_J\}$$

From Lemma 3.20, we get that

$$\begin{aligned} \|F\|_{L^1((0,1))} &\geq \frac{1}{2^5\pi \left(2 + \ln\left(1 + \frac{2}{\delta}\right)\right)} \sum_{j=1}^J \|f_{k_j}\|_{L^1} \left(\frac{C_{MPS}}{2j} - \frac{2\pi d}{qD}\right) \\ &= \frac{1}{2^5\pi \left(2 + \ln\left(1 + \frac{2}{\delta}\right)\right)} (T_1 - T_2) \end{aligned}$$

with

$$T_1 = \frac{C_{MPS}}{2} \sum_{j=1}^J \frac{\|f_{k_j}\|_{L^1}}{j} \quad \text{and} \quad T_2 = \frac{2\pi d}{qD} \sum_{j=1}^J \|f_{k_j}\|_{L^1}.$$

Next, as $\|f_{k_j}\|_{L^1} \geq C_{MPS} \ln(n)$,

$$T_1 \geq \frac{C_{MPS}^2 \ln(n)}{2} \sum_{j=1}^J \frac{1}{j} \geq \frac{C_{MPS}^2 \ln(n)}{2} \ln J \geq \frac{C_{MPS}^2 \ln(n) \ln(m)}{8}$$

with (3.16).

On the other hand, from (3.15), we get

$$T_2 \leq \frac{2\pi J d C_{MPS}}{qD} \ln(m) \ln(n) \leq \frac{2\pi C_{MPS}}{q^{\frac{1}{2}}} \ln(m) \ln(n),$$

since $d \leq \frac{D}{2}$ and $J \leq q^{\frac{1}{2}}$ with (3.15). Further, (3.15) also implies that

$$q^{\frac{1}{2}} \geq \frac{m^{\frac{1}{3}}}{8} \geq 2^4 \pi C_{MPS} \ln(m) \ln(n)$$

with (3.11), leading to $T_2 \leq \frac{1}{2^{13}\pi}$.

We have established that

$$\begin{aligned} \|F\|_{L^1((0,1))} &\geq \frac{1}{2^5\pi \left(2 + \ln\left(1 + \frac{2}{\delta}\right)\right)} \left(\frac{C_{MPS}^2 \ln(m) \ln(n)}{2^3} - \frac{1}{2^{13}\pi} \right) \\ &= \frac{C_{MPS}^2}{(2^9)^2\pi \left(2 + \ln\left(1 + \frac{2}{\delta}\right)\right)} \left(2^{10} \ln(m) \ln(n) - \frac{1}{C_{MPS}^2} \right) \\ &\geq \frac{C_{MPS}^2}{(2^9\pi)^2 \left(2 + \ln\left(1 + \frac{2}{\delta}\right)\right)} \ln(m) \ln(n) \end{aligned}$$

since $\ln(m) \ln(n) C_{MPS}^2 \geq \frac{1}{2^{10}\pi - 1}$.

3.4.2. *Proof of Lemma 3.20.* The rest of this chapter consist in proving Lemma 3.20. the proof is divided into several lemmas.

The first one is a simple lemma about numerical integration of trigonometric polynomials:

Lemma 3.21. *Let N be a positive integer. Then for any trigonometric polynomial f of degree d*

$$\left| \|f\|_{L^1((0,1))} - \frac{1}{N} \sum_{j=0}^{N-1} \left| f\left(\frac{j}{N}\right) \right| \right| \leq \frac{2\pi d}{N} \|f\|_{L^1((0,1))}.$$

Note that, as f is 1-periodic, writing $N = R + S$, $\frac{1}{N} \sum_{j=0}^{N-1} \left| f\left(\frac{j}{N}\right) \right| = \frac{1}{R+S} \sum_{j=-R}^{S-1} \left| f\left(\frac{j}{R+S}\right) \right|$.

Proof. We write, using the triangular and reverse triangular inequalities

$$\begin{aligned} \left| \int_0^1 |f(t)| dt - \frac{1}{N} \sum_{j=0}^{N-1} \left| f\left(\frac{j}{N}\right) \right| \right| &= \left| \sum_{j=0}^{N-1} \int_{\frac{j}{N}}^{\frac{j+1}{N}} |f(t)| dt - \left| f\left(\frac{j}{N}\right) \right| \right| \\ &\leq \sum_{j=0}^{N-1} \int_{\frac{j}{N}}^{\frac{j+1}{N}} \left| f(t) - f\left(\frac{j}{N}\right) \right| dt = \sum_{j=0}^{N-1} \int_{\frac{j}{N}}^{\frac{j+1}{N}} \left| \int_{\frac{j}{N}}^t f'(s) ds \right| dt \\ &\leq \sum_{j=0}^{N-1} \int_{\frac{j}{N}}^{\frac{j+1}{N}} \int_{\frac{j}{N}}^t |f'(s)| ds dt = \sum_{j=0}^{N-1} \int_{\frac{j}{N}}^{\frac{j+1}{N}} |f'(s)| \int_s^{\frac{j+1}{N}} dt ds \\ &\leq \frac{1}{N} \sum_{j=0}^{N-1} \int_{\frac{j}{N}}^{\frac{j+1}{N}} |f'(s)| ds = \int_0^1 |f'(s)| ds \end{aligned}$$

where we have used Fubini and the bound $\int_s^{\frac{j+1}{N}} dt \leq \frac{1}{N}$ when $\frac{j}{N} \leq s \leq \frac{j+1}{N}$. We conclude with Bernstein's inequality, $\|f'\|_{L^1((0,1))} \leq 2\pi d \|f\|_{L^1((0,1))}$. \square

For a finitely supported sequence $(A(k))_{k \in \mathbb{Z}}$ we define its discrete Fourier transform (or \mathbf{Z} -Fourier transform) as

$$\mathcal{F}_d[A](t) = \sum_{k \in \mathbb{Z}} A(k) e^{-2i\pi kt}.$$

If A, B are two finitely supported sequences, their convolution is the sequence $A * B$ defined by

$$A * B(k) = \sum_{n \in \mathbb{Z}} A(k-n) B(n).$$

The Convolution Theorem is also valid here: $\mathcal{F}_d[A * B](t) = \mathcal{F}_d[A](t) \mathcal{F}_d[B](t)$. Two classical examples are

– the Dirichlet kernel: set $d_L = \mathbb{1}_{-L, \dots, L}$ so that

$$D_L(t) = \mathcal{F}_d[d_L](t) = \sum_{|k| \leq L} e^{2i\pi kt} = \frac{\sin(\pi(2L+1)t)}{\sin(\pi t)};$$

– the Fejer kernel: set $f_L(k) = \left(1 - \frac{|k|}{L+1}\right) \mathbb{1}_{-L, \dots, L}(k)$ so that

$$F_L(t) = \mathcal{F}_d[f_L](t) = \sum_{|k| \leq L} \left(1 - \frac{|k|}{L+1}\right) e^{2i\pi kt} = \frac{1}{L+1} \frac{(\sin(\pi(L+1)t))^2}{(\sin(\pi t))^2}.$$

Lemma 3.22. *Let M, N, R, S be integers with $2 \leq M < N$. Then there exists a function $K_{M,N}$ with the following properties:*

- (1) $K_{M,N}(k) = 1$ for $|k| \leq N$,
- (2) $K_{M,N}(k) = 0$ for $|k| \geq N + 2M$
- (3) when $R + S \geq 2N + 4M$, $\frac{1}{R+S} \sum_{j=-R}^{S-1} \left| \mathcal{F}_d[K_{M,N}] \left(\frac{j}{R+S} \right) \right| \leq 16\pi(2 + \ln(1 + N/M))$

Proof. Define

$$\begin{aligned} K_{M,N}(k) &= \frac{1}{M} d_{N+M} * f_{M-1}(k) = \frac{1}{M} \sum_{n \in \mathbb{Z}} \mathbb{1}_{\{-M-N, \dots, M+N\}}(k-n) \mathbb{1}_{\{-M+1, \dots, M+1\}}(n) \\ &= \frac{1}{M} \sum_{\substack{|n| \leq M-1 \\ |n-k| \leq N+M}} \left(1 - \frac{|n|}{M}\right). \end{aligned}$$

First, for $|k| \leq N$, if $|n| \leq M-1$, then $|n-k| \leq |n| + |k| \leq N + M - 1$, so that

$$K_{M,N}(k) = \frac{1}{M} \sum_{|n| \leq M-1} \left(1 - \frac{|n|}{M}\right) = 1,$$

since

$$\begin{aligned} \sum_{|n| \leq M-1} \left(1 - \frac{|n|}{M}\right) &= \sum_{n=1-M}^0 \left(1 + \frac{n}{M}\right) + \sum_{n=1}^{M-1} \left(1 - \frac{n}{M}\right) \\ &= \frac{M+1}{2} + \frac{M-1}{2} = M. \end{aligned}$$

On the other hand, if $|k| \geq N + 2M$ and $|n| \leq M-1$ then $|k-n| \geq |k| - |n| \geq N + M + 1$ so that the sum defining $K_{M,N}$ is empty and $K_{M,N} = 0$.

To prove the last item, the Convolution Theorem shows that

$$\mathcal{F}_d[K_{M,N}](t) = \frac{1}{M} D_{N+M}(t) F_{M-1}(t).$$

As D_{N+M} and F_{M-1} are both even, so is $K_{M,N}$ thus

$$\int_0^1 |\mathcal{F}_d[K_{M,N}](t)| dt = 2 \int_0^{\frac{1}{2}} |\mathcal{F}_d[K_{M,N}](t)| dt = 2(I_1 + I_2 + I_3)$$

where

$$I_1 = \frac{1}{M} \int_0^{\frac{1}{N+M}} |D_{N+M}(t) F_{M-1}(t)| dt$$

$$I_2 = \frac{1}{M} \int_{\frac{1}{N+M}}^{\frac{1}{M}} |D_{N+M}(t) F_{M-1}(t)| dt,$$

$$I_3 = \frac{1}{M} \int_{\frac{1}{M}}^{\frac{1}{2}} |D_{N+M}(t) F_{M-1}(t)| dt.$$

We have

$$|D_{N+M}(t)| \leq 2(N+M) + 1, \quad 0 \leq F_{M-1}(t) \leq M, \quad \int_0^1 F_{M-1}(t) dt = 1.$$

It follows that

$$I_1 \leq \frac{1}{M} \int_0^{\frac{1}{N+M}} (2(M+N) + 1)M dt = 2 + \frac{1}{M+N} \leq 3.$$

since M, N are positive integers.

Using the explicit expressions of D_{N+M} and F_{M-1} , we have

$$\begin{aligned} I_2 &= \frac{1}{M^2} \int_{\frac{1}{N+M}}^{\frac{1}{M}} \frac{|\sin(\pi(2(N+M)+1)t)| \sin^2(\pi Mt)}{\sin^3(\pi t)} dt \leq \frac{1}{M^2} \int_{\frac{1}{N+M}}^{\frac{1}{M}} \frac{\sin^2(\pi Mt)}{\sin^3(\pi t)} dt \\ &\leq \frac{\pi^2}{8} \int_{\frac{1}{N+M}}^{\frac{1}{M}} \frac{dt}{t} = \frac{\pi^2}{8} \ln\left(1 + \frac{N}{M}\right). \end{aligned}$$

using that $\sin \pi t \leq \pi t$ for $t \geq 0$ and that $\sin \pi t \geq 2t$ for $0 \leq t \leq \frac{1}{2}$.

Finally, for I_3 , we do the same computation to bound

$$I_3 \leq \frac{1}{M^2} \int_{\frac{1}{M}}^{\frac{1}{2}} \frac{\sin^2 \pi Mt}{t^3} dt = \int_1^{\frac{M}{2}} \frac{\sin^2 \pi s}{s^3} ds \leq \int_1^{\frac{M}{2}} \frac{ds}{s^3} \leq \frac{1}{2}.$$

Grouping all terms and slightly upper bounding the numerical constants, we obtain

$$\int_0^1 |\mathcal{F}_d[K_{M,N}](t)| dt \leq 8 \left(2 + \ln\left(1 + \frac{N}{M}\right)\right).$$

By Lemma 3.21 (and the 1-periodicity of $\mathcal{F}_d[K_{M,N}]$), we obtain

$$\frac{1}{R+S} \sum_{j=-R}^{S-1} \left| \mathcal{F}_d[K_{M,N}]\left(\frac{j}{2R+1}\right) \right| \leq \left(1 + \frac{2\pi d}{R+S}\right) \|\mathcal{F}_d[K_{M,N}]\|_{L^1([0,1])}.$$

But $d = N + 2M - 1$ and $R + S \geq 2N + 4M$ so

$$\frac{2\pi d}{R+S} \leq \frac{\pi(2N + 4M - 2)}{2N + 4M} \leq \pi.$$

Hence we get

$$\begin{aligned} \frac{1}{R+S} \sum_{j=-R}^{S-1} \left| \mathcal{F}_d[K_{M,N}]\left(\frac{j}{R+S}\right) \right| &\leq (1 + \pi) \|\mathcal{F}_d[K_{M,N}]\|_{L^1([0,1])} \\ &\leq 16\pi \left(2 + \ln\left(1 + \frac{N}{M}\right)\right), \end{aligned}$$

concluding the proof. \square

Lemma 3.23. *Let R, S be positive integers and $K = (K(-R), \dots, K(S-1)) \in \mathbb{C}^{R+S}$. We extend K into*

- *an $R+S$ -periodic sequence $K^{(p)}(j(R+S) + \ell) = K(\ell)$, for $\ell = -R, \dots, S-1$ and $j \in \mathbb{Z}$;*
- *a finitely supported sequence $K^{(0)}$ by setting $K^{(0)}(\ell) = K(\ell)$ for $\ell = -R, \dots, S-1$ and $K^{(0)}(\ell) = 0$ for $\ell \geq S$ and for $\ell \leq -R-1$.*

Then

$$\int_0^1 \left| \sum_{m \in \mathbb{Z}} a_m K^{(p)}(m) e^{2i\pi mt} \right| dt \leq \frac{1}{R+S} \sum_{\ell=-R}^{S-1} \left| \mathcal{F}_d[K^{(0)}]\left(\frac{\ell}{R+S}\right) \right| \int_0^1 \left| \sum_{m \in \mathbb{Z}} a_m e^{2i\pi mt} \right| dt$$

Proof. Write elements of \mathbb{C}^{R+S} as (a_{-R}, \dots, a_{S-1}) and the scalar product $\langle a, b \rangle = \sum_{\ell=-R}^{S-1} a_\ell \overline{b_\ell}$. For $j =$

$-R, \dots, S-1$, denote by $e_k := \left[\frac{1}{\sqrt{R+S}} e^{2i\pi \frac{k\ell}{R+S}} \right]_{\ell=-R}^{S-1}$ so that $(e_k)_{k=-R, \dots, S-1}$ is an orthonormal basis of \mathbb{C}^{R+S} .

Write $A_\ell = \sum_{n \in \mathbb{Z}} \overline{a_{n(R+S)+\ell}} e^{-2i\pi(n(R+S)+\ell)t}$ for $\ell = -R, \dots, S-1$ and $A = (A_\ell) \in \mathbb{C}^{R+S}$. Then, by periodicity of $K^{(p)}$

$$\begin{aligned} \sum_{m \in \mathbb{Z}} a_m K^{(p)}(m) e^{2i\pi m t} &= \sum_{\ell=-R}^{S-1} K(\ell) \sum_{j \in \mathbb{Z}} a_{j(R+S)+\ell} e^{2i\pi(j(R+S)+\ell)t} = \langle K, A \rangle = \sum_{k=-R}^{S-1} \langle K, e_k \rangle \overline{\langle A, e_k \rangle} \\ &= \frac{1}{R+S} \sum_{k=-R}^{S-1} \left(\sum_{\ell=-R}^{S-1} K(\ell) e^{-2i\pi k \frac{\ell}{R+S}} \right) \left(\sum_{\ell=-R}^{S-1} \sum_{n \in \mathbb{Z}} a_{n(R+S)+\ell} e^{2i\pi(n(R+S)+\ell)t} e^{2i\pi k \frac{\ell}{R+S}} \right) \\ &= \frac{1}{R+S} \sum_{k=-R}^{S-1} \mathcal{F}_d[K_0] \left(\frac{k}{R+S} \right) \sum_{\ell=-R}^{S-1} \sum_{n \in \mathbb{Z}} a_{n(R+S)+\ell} e^{2i\pi[(n(R+S)+\ell)t + \ell \frac{k}{R+S}]}. \end{aligned}$$

Noticing that $e^{2i\pi \ell \frac{k}{R+S}} = e^{2i\pi[(n(R+S)+\ell) \frac{k}{R+S}]}$, we may write

$$\begin{aligned} \sum_{m \in \mathbb{Z}} a_m K^{(p)}(m) e^{2i\pi m t} &= \frac{1}{R+S} \sum_{k=-R}^{S-1} \mathcal{F}_d[K^{(0)}] \left(\frac{k}{R+S} \right) \sum_{\ell=-R}^{S-1} \sum_{n \in \mathbb{Z}} a_{n(R+S)+\ell} e^{2i\pi(n(R+S)+\ell)(t + \frac{k}{R+S})} \\ &= \frac{1}{R+S} \sum_{k=-R}^{S-1} \mathcal{F}_d[K^{(0)}] \left(\frac{k}{R+S} \right) \sum_{m \in \mathbb{Z}} a_m e^{2i\pi m(t + \frac{k}{R+S})}. \end{aligned}$$

From this, we deduce that

$$\begin{aligned} \int_0^1 \left| \sum_{m \in \mathbb{Z}} a_m K^{(p)}(m) e^{2i\pi m t} \right| dt &\leq \frac{1}{R+S} \sum_{k=-R}^{S-1} \left| \mathcal{F}_d[K^{(0)}] \left(\frac{k}{R+S} \right) \right| \int_0^1 \left| \sum_{m \in \mathbb{Z}} a_m e^{2i\pi m(t + \frac{k}{R+S})} \right| dt \\ &= \frac{1}{R+S} \sum_{k=-R}^{S-1} \left| \mathcal{F}_d[K^{(0)}] \left(\frac{k}{R+S} \right) \right| \int_0^1 \left| \sum_{m \in \mathbb{Z}} a_m e^{2i\pi m u} \right| du, \end{aligned}$$

with the change of variable $u = t + \frac{j}{R+S}$ and periodicity of $u \rightarrow \sum_m a_m e^{2i\pi m u}$. \square

Lemma 3.24. *Let d, D and q be positive integers with $(2+2\delta)d+4 \leq D$ for some $\delta > 0$ and $q \geq 4\pi$.*

Suppose I is a finite set of integers and, for each $k \in I$ let f_k be a trigonometric polynomial of degree at most d . Then for any integer s , we have:

$$\int_0^1 \left| \sum_{k \in I(q;s)} f_k(t) e^{2i\pi D k t} \right| dt \leq 32\pi(2 + \ln(1 + 2/\delta)) \int_0^1 \left| \sum_{k \in I} f_k(t) e^{2i\pi D k t} \right| dt.$$

Proof. Assume we can prove the lemma for $s = 0$, that is, for any sequence $(f_k)_{k \in \mathbb{Z}}$ of trigonometric polynomials of degree at most d and any finite set I ,

$$\int_0^1 \left| \sum_{q\ell \in I} f_{\ell q}(t) e^{2i\pi D \ell q t} \right| dt \leq 32\pi(2 + \ln(1 + 2/\delta)) \int_0^1 \left| \sum_{k \in I} f_k(t) e^{2i\pi D k t} \right| dt.$$

Then, replacing I with $I - s$ and replacing (f_k) with (f_{k+s}) we get

$$\begin{aligned} \int_0^1 \left| \sum_{k \in I(q;s)} f_k(t) e^{2i\pi D \ell q t} \right| dt &= \int_0^1 \left| \sum_{s+\ell q \in I} f_{s+\ell q}(t) e^{2i\pi D \ell q t} \right| dt \\ &\leq 32\pi(2 + \ln(1 + 2/\delta)) \int_0^1 \left| \sum_{k \in I-s} f_{s+k}(t) e^{2i\pi D k t} \right| dt \\ &= 32\pi(2 + \ln(1 + 2/\delta)) \int_0^1 \left| e^{-2i\pi D s t} \sum_{\ell \in I} f_\ell(t) e^{2i\pi D k t} \right| dt \end{aligned}$$

which is the desired estimate since $|e^{-2i\pi D s t}| = 1$. So there is no loss of generality is assuming $s = 0$.

Next, we write $f_k = \sum_{-d \leq \ell \leq d} a_\ell^k e^{2i\pi(Dk+\ell)t}$ and

$$F(t) = \sum_{k \in I} \sum_{-d \leq \ell \leq d} a_\ell^k e^{2i\pi(Dk+\ell)t} = \sum_m a_m e^{2i\pi m t}$$

where

$$(3.17) \quad \begin{cases} a_m = a_\ell^k & \text{when } m = Dk + \ell, \quad k \in I \quad \text{and} \quad |\ell| \leq d \\ 0 & \text{otherwise} \end{cases}.$$

Let $N = d$, $M = \lceil \frac{\delta d}{2} \rceil$ the smallest integer larger than $\frac{\delta d}{2}$ and let $K_{M,N}$ the sequence from Lemma 3.22. Since

$$N + 2M \leq d + 2 \left(\frac{\delta d}{2} + 1 \right) = d + \delta d + 2 \leq \frac{D}{2}$$

then we have

$$\text{supp}(K_{M,N}) = [-N - 2M, N + 2M] \subseteq \left[-\frac{D}{2}, \frac{D}{2} \right].$$

Further $K_{M,N}^{(p)}(m) = 1$ for $m \in [-d, d]$. Next, $K_{M,N}^{(p)}$ to be the qD -periodic sequence defined by $K_{M,N}^{(p)}(jqD + \ell) = K_{M,N}(\ell)$ for $\ell = -N - 2M, \dots, N + 2M$ and $K_{M,N}^{(p)}(k) = 0$ for all other k 's. We take $R, S > N + 2M$ such that $R + S = qD$ and then $K_{M,N}^{(p)}(k) = 0$ for $k = -R, \dots, -N - 2M - 1$ and for $k = N + 2M + 1, \dots, S - 1$.

From Lemma 3.23, we get

$$\begin{aligned} \int_0^1 \left| \sum_m a_m K_{M,N}^{(p)}(m) e^{2i\pi m t} \right| dt &\leq \frac{1}{R+S} \sum_{j=-R}^{S-1} \left| \mathcal{F}_d[K_{M,N}] \left(\frac{j}{R+S} \right) \right| \int_0^1 \left| \sum_{m \in \mathbb{Z}} a_m e^{2i\pi m u} \right| du \\ &\leq 16\pi \left(2 + \ln \left(1 + \frac{N}{M} \right) \right) \|F\|_{L^1([0,1])} \end{aligned}$$

with Lemma 3.22.

As $\text{supp}(K_{M,N}) \subseteq \left[-\frac{D}{2}, \frac{D}{2} \right]$ and $K_{M,N}^{(p)}$ is periodic of period $R + S = qD$ then

$$(3.18) \quad K_{M,N}^{(p)}(m) \neq 0 \quad \text{if} \quad m = jqD + \ell' \quad \text{for} \quad j \in \mathbb{Z} \quad \text{and} \quad |\ell'| \leq \frac{D}{2}.$$

Combining (3.17) and (3.18) we have that $a_m K_{M,N}^{(p)}(m) \neq 0$ only when

$$m = Dk + \ell = jqD + \ell'.$$

Hence $|jqD - Dk| = |\ell - \ell'| \leq d + \frac{D}{2} < D$. But this can only happen when $jq = k$, which then also implies $\ell = \ell'$. In particular, $m = jqD + \ell$ with $|\ell| \leq d$ and then

$$a_m K_{M,N}^{(p)}(m) = a_{jqD+\ell} K_{M,N}^{(p)}(jqD + \ell) = a_{jqD+\ell} K_{M,N}^{(p)}(\ell) = a_\ell^{jqD}.$$

It follows that

$$\begin{aligned}
\sum_{m \in \mathbb{Z}} a_m K_{M,N}(m) e^{2i\pi m t} &= \sum_{j \in \mathbb{Z}} \sum_{-d \leq \ell \leq d} a_{jqD+\ell} e^{2i\pi[(jqD+\ell)t]} \\
&= \sum_{\substack{k=0 \\ k \in I}}^{\text{mod } q} \sum_{-d \leq \ell \leq d} a_{kD+\ell} e^{2i\pi k D t} e^{2i\pi \ell t} \\
&= \sum_{k \in I(q;0)} e^{2i\pi k D t} \sum_{-d \leq \ell \leq d} a_{kD+\ell} e^{2i\pi \ell t} \\
&= \sum_{k \in I(q;0)} f_k(t) e^{2i\pi k D t}.
\end{aligned}$$

Finally we get

$$\begin{aligned}
\int_0^1 \left| \sum_{k \in I(q;0)} f_k(t) e^{2i\pi k D t} \right| dt &= \int_0^1 \left| \sum_m a_m K_{M,N}(m) e^{2i\pi m t} \right| dt \\
&\leq 32\pi (2 + \ln(1 + N/M)) \|F\|_{L^1([0,1])} \\
&\leq 32\pi (2 + \ln(1 + 2/\delta)) \|F\|_{L^1([0,1])}
\end{aligned}$$

since $\frac{N}{M} \leq \frac{2d}{\delta D} \leq \frac{2}{\delta}$. □

We can now prove the lemma.

Proof of lemma 3.20. Write $I(q; s) = \{k_1, \dots, k_J\}$ and write each k_j in the form $k_j = r_j q + s$. Applying lemma 3.24 yields

$$\begin{aligned}
\|F\|_{L^1([0,1])} &\geq \frac{1}{32\pi(2 + \ln(1 + 2/\delta))} \int_0^1 \left| \sum_{j=1}^J f_{k_j}(t) e^{2i\pi k_j D t} \right| dt = \int_0^1 \left| e^{2i\pi D s t} \sum_{j=1}^J f_{k_j}(t) e^{2i\pi r_j q D t} \right| dt \\
&= \frac{1}{qD} \int_0^{qD} \left| \sum_{j=1}^J f_{k_j} \left(\frac{s}{qD} \right) e^{2i\pi r_j s} \right| ds \\
&= \frac{1}{qD} \sum_{m=0}^{qD-1} \int_m^{m+1} \left| \sum_{j=1}^J f_{k_j} \left(\frac{s}{qD} \right) e^{2i\pi r_j s} \right| ds.
\end{aligned}$$

But

$$\sum_{j=1}^J f_{k_j} \left(\frac{s}{qD} \right) e^{2i\pi r_j s} = \sum_{j=1}^J f_{k_j} \left(\frac{m}{qD} \right) e^{2i\pi r_j s} + \sum_{j=1}^J \left[f_{k_j} \left(\frac{s}{qD} \right) - f_{k_j} \left(\frac{m}{qD} \right) \right] e^{2i\pi r_j s}$$

so that

$$\|F\|_{L^1([0,1])} \geq \frac{1}{32(2 + \ln(1 + 2/\delta))} (T_1 - T_2)$$

with

$$T_1 = \frac{1}{qD} \sum_{m=0}^{qD-1} \int_m^{m+1} \left| \sum_{j=1}^J f_{k_j} \left(\frac{m}{qD} \right) e^{2i\pi r_j s} \right| ds,$$

and

$$T_2 = \frac{1}{qD} \sum_{m=0}^{qD-1} \int_m^{m+1} \left| \sum_{j=1}^J \left(f_{k_j} \left(\frac{s}{qD} \right) - f_{k_j} \left(\frac{m}{qD} \right) \right) e^{2i\pi r_j s} \right| ds.$$

It remains to show that

$$T_1 \geq \frac{C_{MPS}}{2} \sum_{j=1}^J \frac{\|f_{k_j}\|_{L^1}}{j} \quad \text{and} \quad T_2 \leq \frac{2\pi d}{qD} \sum_{j=1}^J \|f_{k_j}\|_{L^1}.$$

Let us start with T_1 : using the 1-periodicity in s ,

$$\begin{aligned} T_1 &= \frac{1}{qD} \sum_{m=0}^{qD-1} \int_0^1 \left| \sum_{j=1}^J f_{k_j} \left(\frac{m}{qD} \right) e^{2i\pi r_j s} \right| ds, \\ &\geq \frac{C_{MPS}}{qD} \sum_{m=0}^{qD-1} \sum_{j=1}^J \frac{\left| f_{k_j} \left(\frac{m}{qD} \right) \right|}{j} = \frac{C_{MPS}}{qD} \sum_{j=1}^J \frac{1}{j} \sum_{m=0}^{qD-1} \left| f_{k_j} \left(\frac{m}{qD} \right) \right| \end{aligned}$$

with Theorem B. Applying Lemma 3.21 to f_{k_j} and using that $\frac{2\pi d}{qD} < \frac{1}{2}$ with our hypothesis on q and D , we get

$$\frac{1}{qD} \sum_{m=1}^{qD} \left| f_{k_j} \left(\frac{m-1}{qD} \right) \right| \geq \frac{\|f_{k_j}\|_{L^1}}{2}$$

and the desired estimate of T_1 follows immediately.

Let us now estimate T_2 . For $s \in [m, m+1]$, we have

$$\begin{aligned} \left| \sum_{j=1}^J \left(f_{k_j} \left(\frac{s}{qD} \right) - f_{k_j} \left(\frac{m}{qD} \right) \right) e^{2i\pi r_j s} \right| &= \left| \sum_{j=1}^J \int_{\frac{m}{qD}}^{\frac{s}{qD}} f'_{k_j}(t) e^{2i\pi r_j s} dt \right| \\ &\leq \int_{\frac{m}{qD}}^{\frac{m+1}{qD}} \left| \sum_{j=1}^J f'_{k_j}(t) e^{2i\pi r_j s} \right| dt. \end{aligned}$$

From the 1-periodicity in s , the integral of this quantity over $[m, m+1]$ is the same as the integral over $[0, 1]$. Thus

$$\begin{aligned} T_2 &\leq \frac{1}{qD} \sum_{m=0}^{qD-1} \int_m^{m+1} \int_{\frac{m}{qD}}^{\frac{m+1}{qD}} \left| \sum_{j=1}^J f'_{k_j}(t) e^{2i\pi r_j s} \right| dt ds \\ &= \frac{1}{qD} \sum_{m=0}^{qD-1} \int_{\frac{m}{qD}}^{\frac{m+1}{qD}} \int_0^1 \left| \sum_{j=1}^J f'_{k_j}(t) e^{2i\pi r_j s} \right| ds dt \\ &= \frac{1}{qD} \int_0^1 \int_0^1 \left| \sum_{j=1}^J f'_{k_j}(t) e^{2i\pi r_j s} \right| ds dt \\ &\leq \frac{1}{qD} \sum_{j=1}^J \|f'_{k_j}\|_{L^1([0,1])} \leq \frac{2\pi d}{qD} \sum_{j=1}^J \|f_{k_j}\|_{L^1([0,1])} \end{aligned}$$

with Bernstein's inequality. \square

3.5. Newman's extremal trigonometric polynomial. Finally, let us insist on the fact that all results in this section are worst case scenarios *i.e.* the lowest possible L^1 norm of a trigonometric polynomial with eventually a constraint on the spectrum or on the coefficients.

If one looks at the maximal possible value of the L^1 -norm under the constraint that all coefficients have modulus 1, then, from Cauchy-Schwarz

$$\int_0^1 \left| \sum_{j=0}^N a_j e^{2i\pi jt} \right| dt \leq \left(\int_0^1 \left| \sum_{j=0}^N a_j e^{2i\pi jt} \right|^2 dt \right)^{\frac{1}{2}} = \left(\sum_{j=0}^N |a_j|^2 \right)^{\frac{1}{2}} = \sqrt{N+1}$$

if $|a_j| = 1$ for all j 's. However, if one wants to reach the bound $\sqrt{N+1}$, one would need that equality holds in the Cauchy-Schwarz Inequality so $\left| \sum_{j=0}^N a_j e^{2i\pi jt} \right|$ should be constant. But, it is easy to show that the only trigonometric polynomials with constant modulus are monomials (thus in our case, equality can

only occur when $N = 0$). Indeed, consider the polynomial $P(z) = \sum_{j=0}^N a_j z^j$ with $a_N \neq 0$. Introduce

$P^*(z) = \overline{P(\bar{z})} = \sum_{j=0}^N \overline{a_j} z^j$. Then $|P(z)|^2 = c$ when $|z| = 1$ can be written in the form $P(z)P^*(1/z) = c$ for $|z| = 1$, which is now an equality between meromorphic functions over $\mathbb{C} \setminus \{0\}$. As this is true on $\{|z| = 1\}$ it is true over all of $\mathbb{C} \setminus \{0\}$, in particular, P has only 0 as zero thus $P(z) = a_N z^N$.

So the natural question is, how near to $\sqrt{N+1}$ one can get. A result by E. Beller and D. J. Newman [2] shows that one can obtain $\sqrt{N} - O(1)$. For sake of simplicity, we will only present a previous result by D. J. Newman [18] which contains similar ingredients:

Theorem 3.25 (Newman, [18]). *There is an absolute constant $c > 0$ and an integer N_0 such that, for every $N \geq N_0$ there is a sequence a_0, \dots, a_N with $|a_j| = 1$ for all j and such that*

$$\int_0^1 \left| \sum_{j=0}^N a_j e^{2i\pi jt} \right| dt \geq \sqrt{N} - c.$$

Proof. The example is related to Gauss sums. We fix N and set $\omega = \exp\left(i\frac{\pi}{N+1}\right)$, $a_j = \omega^{j^2}$ and define

$$P(t) = \sum_{j=0}^N a_j e^{2i\pi jt} = \sum_{j=0}^N \exp\left(2i\pi \left(\frac{j^2}{2(N+1)} + jt\right)\right).$$

The theorem is based on the following lemma:

Lemma 3.26. *With the previous notation, when $N \rightarrow +\infty$*

$$\int_0^1 |P(t)|^4 dt = N^2 + O(N^{3/2}).$$

We postpone the proof of the lemma and show how we can conclude with it.

Parseval's identity gives $\|P\|_2 = \sqrt{N+1}$ and then Hölder's inequality implies

$$N \leq N+1 = \int_0^1 |P(t)|^2 dt \leq \left(\int_0^1 |P(t)| dt \right)^{\frac{2}{3}} \left(\int_0^1 |P(t)|^4 dt \right)^{\frac{1}{3}}.$$

From the lemma, we deduce that there is a constant $A > 0$ such that $\int_0^1 |P(t)|^4 dt \leq N^2 + AN^{3/2}$ when N is large enough, so that

$$\begin{aligned} \int_0^1 |P(t)| dt &\geq \frac{N^{3/2}}{(N^2 + AN^{3/2})^{1/2}} = N^{1/2} \frac{1}{(1 + AN^{-1/2})^{1/2}} = N^{1/2} \left(1 - \frac{A}{2} N^{-1/2} + o(N^{-1/2})\right) \\ &\geq N^{1/2} - A \end{aligned}$$

for N large enough. □

It remains to prove the lemma:

Proof of Lemma 3.26. We write

$$\left| \sum_{j=0}^N a_j e^{2i\pi jt} \right|^2 = \left(\sum_{j=0}^N a_j e^{2i\pi jt} \right) \left(\sum_{k=0}^N \overline{a_k} e^{-2i\pi kt} \right) := \sum_{\ell=-N}^N c_\ell e^{2i\pi \ell t}.$$

Note that $c_0 = \sum_{j=0}^N a_j \bar{a}_j = N + 1$, $c_{-\ell} = \bar{c}_\ell$ and, for $\ell \geq 1$,

$$\begin{aligned} c_\ell &= \sum_{k=0}^{N-\ell} a_{\ell+k} \bar{a}_k = \sum_{k=0}^{N-\ell} \omega^{(\ell+k)^2} \omega^{-k^2} = \omega^{\ell^2} \sum_{k=0}^{N-\ell} \omega^{2k\ell} \\ &= \omega^{\ell^2} \frac{1 - \omega^{2(N-\ell+1)\ell}}{1 - \omega^{2\ell}} = -\omega^{-\ell} \frac{\omega^{\ell^2} - \omega^{-\ell^2}}{\omega^\ell - \omega^{-\ell}} \end{aligned}$$

since $\omega^{2(N+1)} = 1$. We thus obtain that

$$|c_\ell|^2 = \frac{\sin^2 \frac{\pi}{N+1} \ell^2}{\sin^2 \frac{\pi}{N+1} \ell}.$$

On the other hand, from Parseval,

$$\int_0^1 |P(t)|^4 dt = \sum_{\ell=-N}^N |c_\ell|^2 = (N+1)^2 + 2 \sum_{\ell=1}^N |c_\ell|^2$$

since $c_0 = N+1$ and $|c_{-\ell}| = |c_\ell|$.

To compute the last sum, we split it into 4 parts

$$\begin{aligned} S_1 &= \sum_{1 \leq \ell \leq \sqrt{N}} |c_\ell|^2 & S_2 &= \sum_{\sqrt{N} < \ell \leq \frac{N+1}{2}} |c_\ell|^2 \\ S_3 &= \sum_{\frac{N+1}{2} < \ell < N+1-\sqrt{N}} |c_\ell|^2 & S_4 &= \sum_{N+1-\sqrt{N} \leq \ell \leq N} |c_\ell|^2. \end{aligned}$$

As $|c_{N+1-\ell}| = |c_\ell|$ we have $S_4 \leq S_1$ and $S_3 \leq S_2$ so that we only need to show that S_1 and S_3 are both $O(N^{3/2})$.

For S_1 , we use the estimate $\left| \frac{\sin \ell \theta}{\sin \theta} \right| \leq \ell$ to bound $|c_\ell|^2 \leq \ell^2$, leading to

$$S_1 \leq \sum_{1 \leq \ell \leq \sqrt{N}} \ell^2 \leq N \sum_{1 \leq \ell \leq \sqrt{N}} 1 = N^{3/2}.$$

Further, as for $0 \leq \theta \leq \frac{\pi}{2}$, $\sin \theta \geq \frac{2}{\pi} \theta$, we bound $|c_\ell|^2 \leq \frac{(N+1)^2}{4\ell^2}$ so that

$$S_2 \leq \frac{(N+1)^2}{4} \sum_{\sqrt{N} < \ell \leq \frac{N+1}{2}} \frac{1}{\ell^2} \leq \frac{(N+1)^2}{4} \int_{\sqrt{N}-1}^{+\infty} \frac{dt}{t^2} = \frac{(N+1)^2}{4(\sqrt{N}-1)} = O(N^{3/2})$$

as claimed. \square

4. L^1 ESTIMATES WITH REAL FREQUENCIES

4.1. Some results for the Besicovitch norm. We will now present three results for the Besicovitch norms of non-harmonic trigonometric polynomials.

Let us start with the analogue of Mc Gehee, Pigno and Smith's result. Let us present a nice argument by Hudson and Leckband that shows that the estimate for the Besicovitch norm follows from Theorem B. The converse is of course true as well and the result was obtained directly in [13] and even allowed to slightly improve the constant.

Theorem 4.1 (Hudson & Leckband [11]). *For $(\lambda_j)_{j \geq 0}$ be real numbers with $\lambda_{j+1} - \lambda_j > 0$ and let $(a_j)_{j \geq 0}$ be a sequence of complex numbers with finite support. Then*

$$\lim_{T \rightarrow +\infty} \frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{j=0}^{+\infty} a_j e^{2i\pi\lambda_j t} \right| dt \geq C_{MPS} \sum_{j=0}^{+\infty} \frac{|a_j|}{j+1}$$

where C_{MPS} is the same constant as in Theorem B.

Moreover, assume that (λ_k) is q -lacunary i.e. $\lambda_0 \geq 1$ and $\lambda_{k+1} \geq q\lambda_k$. Then, for $1 \leq p < +\infty$,

$$A_{p,q} \left(\sum_{j=0}^{+\infty} |a_j|^2 \right)^{1/2} \leq \lim_{T \rightarrow +\infty} \left(\frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{j=0}^{+\infty} a_j e^{2i\pi\lambda_j t} \right|^p dt \right)^{\frac{1}{p}} \leq B_{p,q} \left(\sum_{j=0}^{+\infty} |a_j|^2 \right)^{1/2}$$

with $A_{p,q}, B_{p,q}$ the constants in Theorem 3.7.

Actually, only the first part is in [11]. The second part is here obtained with almost the same proof and seems to be new.

Proof. Let a_0, \dots, a_N be complex numbers, $\lambda_0 < \lambda_1 < \dots < \lambda_N$ be real numbers and

$$\Phi(t) = \sum_{j=0}^N a_j e^{2i\pi\lambda_j t}.$$

Let $\varepsilon > 0$. By a lemma of Dirichlet ([25, p 235], [9]), there is an increasing sequence of integers $(M_n)_{n \geq 1}$ and, for each $n \geq 1$ a finite family of integers $(N_{j,n})_{j=0, \dots, N}$ such that

$$\left| \lambda_j - \frac{N_{j,n}}{M_n} \right| < \frac{\varepsilon}{M_n} \quad \text{for } j = 0, \dots, N$$

which implies that

$$\left| e^{2i\pi\lambda_j t} - e^{2i\pi\frac{N_{j,n}}{M_n}t} \right| \leq 2\pi \left| \lambda_j - \frac{N_{j,n}}{M_n} \right| |t| \leq 2\pi \frac{\varepsilon}{M_n} |t| \quad \text{for } j = 0, \dots, N.$$

Define the M_n -periodic function

$$\Psi_n(t) = \sum_{j=0}^N a_j e^{2i\pi N_{j,n}t/M_n}$$

and note that, for $t \in [-M_n/2, M_n/2]$,

$$|\Phi(t) - \Psi_n(t)| \leq \sum_{j=0}^N |a_j| \left| e^{2i\pi\lambda_j t} - e^{2i\pi\frac{N_{j,n}}{M_n}t} \right| \leq 2\pi\varepsilon \sum_{j=0}^N |a_j|.$$

But then

$$\begin{aligned} \left| \left(\frac{1}{M_n} \int_{-M_n/2}^{M_n/2} |\Phi(t)|^p dt \right)^{1/p} - \left(\frac{1}{M_n} \int_{-M_n/2}^{M_n/2} |\Psi_n(t)|^p dt \right)^{1/p} \right| & \\ & \leq \left(\frac{1}{M_n} \int_{-M_n/2}^{M_n/2} |\Phi(t) - \Psi_n(t)|^p dt \right)^{1/p} \\ & \leq \left(2\pi \sum_{j=0}^N |a_j| \right)^{1/p} \varepsilon^{1/p}. \end{aligned}$$

We can now conclude as follows. First, in the general case, As Ψ_n is M_n -periodic, we may apply Theorem B to obtain

$$\begin{aligned} C_{MPS} \sum_{j=0}^N \frac{|a_j|}{j+1} & \leq \frac{1}{M_n} \int_{-M_n/2}^{M_n/2} |\Psi_n(t)| dt \\ & \leq \frac{1}{M_n} \int_{-M_n/2}^{M_n/2} |\Phi(t)| dt + \frac{1}{M_n} \int_{-M_n/2}^{M_n/2} |\Phi(t) - \Psi_n(t)| dt \\ & \leq \frac{1}{M_n} \int_{-M_n/2}^{M_n/2} |\Phi(t)| dt + 2\pi\varepsilon \sum_{j=0}^N |a_j| \end{aligned}$$

Letting $n \rightarrow +\infty$ and then $\varepsilon \rightarrow 0$ we obtain Theorem 4.1:

$$\lim_{T \rightarrow +\infty} \frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{k=0}^N a_k e^{2i\pi\lambda_k t} \right| dt \geq C_{MPS} \sum_{k=0}^N \frac{|a_k|}{k+1}.$$

Let us now assume further that $\lambda_{k+1} \geq q\lambda_k$. Let \tilde{q} be such that $1 < \tilde{q} < q$ and, assume that ε has been chosen such that $q - (1+q)\varepsilon > \tilde{q}$. Observe that

$$(4.1) \quad \frac{N_{j+1,n}}{M_n} \geq \lambda_{j+1} - \left| \lambda_{j+1} - \frac{N_{j+1,n}}{M_n} \right| \geq q\lambda_j - \frac{\varepsilon}{M_n}$$

$$(4.2) \quad \geq q \frac{N_{j,n}}{M_n} - q \left| \lambda_j - \frac{N_{j,n}}{M_n} \right| - \frac{\varepsilon}{M_n}$$

$$(4.3) \quad \geq q \frac{N_{j,n}}{M_n} - (1+q) \frac{\varepsilon}{M_n},$$

that is $N_{j+1,n} \geq qN_{j,n} - (1+q)\varepsilon \geq \tilde{q}N_{j,n}$.

Applying (3.6) to $n_k = N_{k,n}$ and $M = M_n$ we obtain

$$A_{p,\tilde{q}} \left(\sum_{j=0}^{+\infty} |a_j|^2 \right)^{\frac{1}{2}} \leq \left(\frac{1}{M_n} \int_{-M_n/2}^{M_n/2} \left| \sum_{j \geq 0} a_j e^{2i\pi \frac{N_{j,n}}{M_n} t} \right|^p dt \right)^{\frac{1}{p}} \leq B_{p,\tilde{q}} \left(\sum_{j=0}^{+\infty} |a_j|^2 \right)^{\frac{1}{2}}.$$

From (4.1) we conclude that

$$\begin{aligned} A_{p,\tilde{q}} \left(\sum_{j=0}^{+\infty} |a_j|^2 \right)^{\frac{1}{2}} - \left(2\pi \sum_{j=0}^N |a_j| \right)^{1/p} \varepsilon^{1/p} &\leq \left(\frac{1}{M_n} \int_{-M_n/2}^{M_n/2} \left| \sum_{j \geq 0} a_j e^{2i\pi \frac{N_{j,n}}{M_n} t} \right|^p dt \right)^{\frac{1}{p}} \\ &\leq B_{p,\tilde{q}} \left(\sum_{j=0}^{+\infty} |a_j|^2 \right)^{\frac{1}{2}} + \left(2\pi \sum_{j=0}^N |a_j| \right)^{1/p} \varepsilon^{1/p}. \end{aligned}$$

The result follows by letting $\varepsilon \rightarrow 0$ and then $\tilde{q} \rightarrow q$. \square

4.2. Nazarov's theorem.

Theorem 4.2 (Quantitative version of Nazarov theorem for small δ). *There exists positive constants c_*, δ_* such that, for $0 < \delta < \delta_*$ the following holds:*

for every $\lambda_0 < \dots < \lambda_N$ of real numbers satisfying $\lambda_{k+1} - \lambda_k \geq 1$ and a_0, \dots, a_N complex numbers,

$$(4.4) \quad \sum_{k=0}^N \frac{|a_k|}{k+1} \leq \frac{c_*}{\delta^{15/2}} \int_{-\frac{1+\delta}{2}}^{\frac{1+\delta}{2}} \left| \sum_{k=0}^N a_k e^{2i\pi\lambda_k t} \right| dt.$$

First, we start by some preliminary notations and results. Let $I = I_\delta := [-\frac{1+\delta}{2}, \frac{1+\delta}{2}]$.

Let us fix $(\lambda_k)_{k=0, \dots, N} \subset \mathbb{R}$ with $\lambda_{k+1} - \lambda_k \geq 1$ for every k , $(a_k)_{k=0, \dots, N}$ a sequence of complex numbers and write $|a_k| = a_k u_k$ with $|u_k| = 1$. Let

$$S = \sum_{k=0}^N \frac{|a_k|}{k+1} \quad \text{and} \quad \phi(t) = \sum_{k=0}^N a_k e^{2i\pi\lambda_k t},$$

so that we must find C_δ such that:

$$(4.5) \quad \|\phi\|_{L^1(I)} \geq C_\delta S.$$

We define

$$S_\delta = \sum_{k=0}^N \frac{|a_k|}{k+N_\delta} \quad \text{and} \quad T_\delta(t) = \sum_{k=0}^N \frac{u_k}{k+N_\delta} e^{-2i\pi\lambda_k t}$$

where N_δ is a large integer that we will adjust through the proof. This integer will be of the form $N_\delta = 2^{m_\delta}$. We will prove that

$$(4.6) \quad \|\phi\|_{L^1(I)} \geq B_\delta S_\delta$$

and, as $k + N_\delta \leq (k + 1)N_\delta$, $S_\delta \geq \frac{S}{N_\delta}$ so that we obtain the desired inequality (4.5) with a constant $C_\delta = \frac{B_\delta}{N_\delta}$.

The meaning of N_δ is the following. Consider:

– a new sequence of frequencies $(\tilde{\lambda}_j)_{j \in \mathbb{Z}}$ such that $\tilde{\lambda}_j = \lambda_{j-N_\delta}$ for $j = N_\delta, \dots, N_\delta + N$. and then $\tilde{\lambda}_j = \lambda_0 + j - N_\delta$ for $j < N_\delta$ and $\tilde{\lambda}_j = \lambda_N + j - N$ for $j > N$. In particular, we still have $|\tilde{\lambda}_j - \tilde{\lambda}_k| \geq 1$. In other words, the sequence $(\lambda_j)_{j=0, \dots, N}$ is completed into a sequence $(\lambda_j)_{j \in \mathbb{Z}}$ that is still 1-separated and then shifting it by N_δ .

– A new sequence of complex numbers $(\tilde{a}_j)_{j \in \mathbb{Z}}$ with $\tilde{a}_j = a_{j-N_\delta}$ for $j = N_\delta, \dots, N_\delta + N$ and $\tilde{a}_j = 0$ for other j 's. In other words, the sequence $(a_j)_{j=0, \dots, N}$ is completed into a sequence $(a_j)_{j \in \mathbb{Z}}$ by 0-padding it and then shifting it by N_δ .

Then (4.6) reads

$$\int_{I_\delta} \left| \sum_{j \in \mathbb{Z}} \tilde{a}_j e^{2i\pi \tilde{\lambda}_j t} \right| dt \geq B_\delta \sum_{j \in \mathbb{Z}} \frac{|\tilde{a}_j|}{j + N_\delta}$$

with the convention that $0/0 = 0$.

Note that, up to adding 0 terms at the end of the sequence a_j , we may assume that $N + N_\delta$ is of the form $2^{n_\delta} - 1$ for some integer n_δ . This will allow us to write

$$\sum_{k=0}^N = \sum_{j=m_\delta}^{n_\delta} \sum_{2^j \leq r + N_\delta < 2^{j+1}}.$$

Next, as in Ingham's proof, we will introduce an auxiliary function. Again, we consider

$$h(t) = \begin{cases} \cos(\pi t) & \text{if } |t| \leq \frac{1}{2} \\ 0 & \text{otherwise} \end{cases}$$

whose Fourier transform is given by

$$\hat{h}(\lambda) = \frac{2 \cos(\pi \lambda)}{\pi (1 - 4\lambda^2)}.$$

We will need to smooth a bit this function to obtain a better decay of the Fourier transform and thereby slightly enlarge its support. More precisely, let $p = 10$, $q = 8$ and let

$$f_\delta(t) = \frac{p+q}{\delta} \mathbb{1}_{[-\frac{\delta}{2(p+q)}, \frac{\delta}{2(p+q)}]}(t)$$

and define its $p + q$ -fold self convolution¹

$$g_\delta = *_p *_{q+q} f_\delta.$$

Clearly g_δ is non-negative, even and with support $[-\frac{\delta}{2}, \frac{\delta}{2}]$. Finally, we define φ_δ as

$$(4.7) \quad \varphi_\delta = \frac{\pi}{2} h * g_\delta.$$

In the following lemma, we list the properties needed on φ_δ . They are all established via easy calculus and straight forward Fourier analysis.

Lemma 4.3. *There is a $c_0 > 0$ and a $\delta_0 > 0$ such that, if $0 < \delta < \delta_0$ then,*

$$(1) \quad \widehat{\varphi}_\delta(\lambda) = \frac{\cos(\pi \lambda)}{1 - 4\lambda^2} \operatorname{sinc}^{p+q} \left(\frac{\pi \delta \lambda}{p+q} \right),$$

¹If χ is a function, we write $*_1 \psi = \psi * \psi$ for the convolution of ψ with itself and then define inductively $*_{k+1} \psi = (*_k \psi) * \psi$.

$$(2) \|\varphi_\delta\|_\infty \leq \frac{\pi}{2}.$$

$$(3) \text{ Let } D_\delta = c_0 \delta^{-\frac{p+q}{p+2}} \geq 1 \text{ and } \nu > 0 \text{ then, for } |\lambda| \geq \max(1, D_\delta \nu^{\frac{1}{p+2}}),$$

$$|\widehat{\varphi}(\lambda)| \leq \frac{\nu}{|\lambda|^q}$$

$$(4) \text{ Let } \gamma_\delta = \left(\operatorname{sinc} \frac{\pi\delta}{p+q} \right)^{p+q} \text{ then, for } |\lambda| \geq 1,$$

$$|\widehat{\varphi}_\delta(\lambda)| \leq \frac{\gamma_\delta}{4\lambda^2 - 1}.$$

From now on, we will assume that $0 < \delta < \frac{7}{3\pi} < 1$ so that $D_\delta \geq 1$. In particular, when $|\lambda| \geq D_\delta$, we have $|\widehat{\varphi}(\lambda)| \leq |\lambda|^{-3}$.

Proof of Lemma 4.3. The first one is simple Fourier analysis.

For the second one, we write

$$\|\varphi_\delta\|_\infty \leq \frac{\pi}{2} \|g_\delta\|_1 \|h\|_\infty \leq \frac{\pi}{2} \|f_\delta\|_1^{p+q} \|h\|_\infty$$

and use simple calculus to conclude.

For the third one, we notice that $4\lambda^2 - 1 \geq 3\lambda^2$ when $\lambda \geq 1$ thus

$$|\widehat{\varphi}_\delta(\lambda)| = \left| \frac{\cos(\pi\lambda)}{4\lambda^2 - 1} \left(\frac{\sin(\frac{\pi\delta\lambda}{p+q})}{\frac{\pi\delta\lambda}{p+q}} \right)^{2p} \right| \leq \frac{1}{3} \left(\frac{p+q}{\pi} \right)^{p+q} \delta^{-(p+q)} \frac{1}{|\lambda|^{p+2}} \frac{1}{|\lambda|^q} \leq \frac{\nu}{|\lambda|^q},$$

since $|\lambda| \geq \frac{1}{3^{\frac{1}{p+2}}} \left(\frac{p+q}{\pi} \right)^{\frac{p+q}{p+2}} \delta^{-\frac{p+q}{p+2}} \nu^{\frac{1}{p+2}}$. Thus this fact is established with $c_0 = \frac{1}{3^{\frac{1}{p+2}}} \left(\frac{p+q}{\pi} \right)^{\frac{p+q}{p+2}}$.

For the last one, we take δ_0 small enough to have $\operatorname{sinc} \frac{\pi\delta_0}{p+q} = \sup_{t \geq \delta_0} |\operatorname{sinc} t|$ and then $|\widehat{g}_\delta(\lambda)| \leq \gamma_\delta$ when $|\lambda| \geq 1$. The first identity allows to conclude. \square

We can now state the first crucial result in this proof.

Lemma 4.4. *There exist $\delta_1 > 0$, $c_1 > 0$ such that, if $0 < \delta < \delta_1$, $0 < c_1 \delta^2 < 1 - \sqrt{\gamma_\delta}$.*

Moreover, let

$$\alpha_\delta = 1 - \left(c_1 \delta^2 + \frac{\gamma_\delta}{1 - c_1 \delta^2} \right)$$

and let m_δ be such that

$$N_\delta = 2^{m_\delta} \geq \delta^{-7/2}.$$

Then, for $0 \leq k \leq N$,

$$\sum_{\substack{0 \leq j \leq N \\ j \neq k}} \frac{|\widehat{\varphi}(\lambda_j - \lambda_k)|}{j + N_\delta} \leq \frac{1 - \alpha_\delta}{k + N_\delta}.$$

Proof. The Taylor expansion when $\delta \rightarrow 0$ of $1 - \sqrt{\gamma_\delta}$ is of the form

$$1 - \sqrt{\gamma_\delta} = A\delta^2 + O(\delta^4),$$

with $A > 0$. Thus, if $c_1 < A$, for δ small enough, $0 < c_1 \delta^2 < 1 - \sqrt{\gamma_\delta}$. Next, notice that $0 < 1 - \left(\beta + \frac{\gamma_\delta}{1 - \beta} \right) < 1$ if $0 < \beta < 1 - \sqrt{\gamma_\delta}$ which shows that $0 < \alpha_\delta < 1$. For future use, note that there is a $\kappa > 0$ such that

$$(4.8) \quad \alpha_\delta = \kappa \delta^2 + O(\delta^4).$$

We will further assume that δ is small enough for

$$\delta^{-7/2} \geq \max \left(c_1^{-\frac{q+1}{q-2}} \delta^{-2\frac{q+1}{q-2}}, c_2 \delta^{-2-\frac{p+q}{p+2}} \right) = \max \left(c_1^{-\frac{q+1}{q-2}} \delta^{-3}, \frac{c_0}{c_1} \delta^{-7/2} \right)$$

since we chose $q = 8$ and $p = 10$.

Note that, for every $\varepsilon > 0$, the power $7/2$ could be reduced to $3 + \varepsilon$ by taking p large enough, but could not be reduced below 3 with this construction.

We can now turn to the estimate itself. Set $\beta = c_1 \delta^2$ and split the sum in the left hand side of the main inequality into two sums

$$E := \sum_{\substack{0 \leq j \leq N \\ j \neq k}} \frac{|\widehat{\varphi}(\lambda_j - \lambda_k)|}{j + N_\delta} = E_1 + E_2$$

where

$$E_1 = \sum_{j+N_\delta < (1-\beta)(k+N_\delta)} \frac{|\widehat{\varphi}(\lambda_j - \lambda_k)|}{j + N_\delta}$$

and

$$E_2 = \sum_{\substack{j+N_\delta \geq (1-\beta)(k+N_\delta) \\ j \neq k}} \frac{|\widehat{\varphi}(\lambda_j - \lambda_k)|}{j + N_\delta}.$$

The result is obtained if we prove the two estimates

$$E_1 \leq \frac{\beta}{k + N_\delta} \quad \text{et} \quad E_2 \leq \frac{\gamma_\delta/(1-\beta)}{k + N_\delta}.$$

Now, as $N_\delta \geq \frac{c_0}{c_1} \delta^{-2-\frac{p+q}{p+2}}$, $\beta N_\delta \geq D_\delta$. Then, if j is an index corresponding to E_1 , then

$$|\lambda_k - \lambda_j| \geq |k - j| = (k + N_\delta) - (j + N_\delta) \geq \beta(k + N_\delta) \geq \beta N_\delta \geq D_\delta$$

hence, from Lemma 4.33 with $\nu = 1$,

$$\begin{aligned} E_1 &\leq \sum_{j+N_\delta < (1-\beta)(k+N_\delta)} |\widehat{\varphi}(\lambda_j - \lambda_k)| \leq \sum_{j+N_\delta < (1-\beta)(k+N_\delta)} \frac{1}{|\lambda_j - \lambda_k|^q} \\ &\leq \sum_{j+N_\delta < (1-\beta)(k+N_\delta)} \frac{1}{(\beta(k + N_\delta))^q}. \end{aligned}$$

But, E_1 contains less than k terms so

$$(4.9) \quad E_1 \leq \frac{k}{k + N_\delta} \frac{\beta^{-(q+1)}}{(k + N_\delta)^{q-2}} \frac{\beta}{k + N_\delta} \leq \frac{\beta}{k + N_\delta}$$

since $N_\delta \geq \beta^{-\frac{q+1}{q-2}} = c_1^{-\frac{q+1}{q-2}} \delta^{-2\frac{q+1}{q-2}}$.

We shall now bound E_2 . In this sum,

$$j + N_\delta \geq (1 - \beta)(k + N_\delta) \quad \text{and} \quad |\lambda_j - \lambda_k| \geq 1$$

then

$$\begin{aligned} E_2 &\leq \frac{1}{(1-\beta)(k + N_\delta)} \sum_{\substack{1 \leq j \leq N \\ j \neq k}} \frac{\gamma_\delta}{4(\lambda_j - \lambda_k)^2 - 1} \\ &\leq \frac{\gamma_\delta}{(1-\beta)(k + N_\delta)} \sum_{\substack{1 \leq j \leq N \\ j \neq k}} \frac{1}{4(j-k)^2 - 1} \\ &\leq \frac{\gamma_\delta}{(1-\beta)(k + N_\delta)} \sum_{\ell=1}^{\infty} \frac{2}{4\ell^2 - 1}. \end{aligned}$$

Since $\frac{2}{4\ell^2 - 1} = \frac{1}{2\ell - 1} - \frac{1}{2\ell + 1}$, we obtain the expected bound $E_2 \leq \frac{\gamma_\delta/(1-\beta)}{k + N_\delta}$. \square

The following lemma is a first step towards proving Theorem 4.2 and is a consequence of Lemma 4.4.

Lemma 4.5. *Let us use the notations of the lemma 4.4. Then,*

$$(4.10) \quad \left| \int_{I_\delta} T_\delta(t) \left(\sum_{k=0}^N a_k e^{2i\pi\lambda_k t} \right) \varphi(t) dt \right| \geq \alpha_\delta \sum_{k=0}^N \frac{|a_k|}{k + N_\delta}.$$

Proof. By definition of T_δ ,

$$\begin{aligned} \int_{I_\delta} T_\delta(t) e^{2i\pi\lambda_k t} \varphi(t) dt &= \sum_{j=0}^M \frac{u_j}{j + N_\delta} \int_{I_\delta} e^{-2i\pi\lambda_j t} e^{2i\pi\lambda_k t} \varphi(t) dt \\ &= \sum_{j=0}^M \frac{u_j}{j + N_\delta} \widehat{\varphi}(\lambda_j - \lambda_k) \\ &= \frac{u_k}{k + N_\delta} + \sum_{\substack{0 \leq j \leq M \\ j \neq k}} \frac{u_j}{j + N_\delta} \widehat{\varphi}(\lambda_j - \lambda_k) \end{aligned}$$

thus

$$\left| \int_{I_\delta} T_\delta(t) e^{2i\pi\lambda_k t} \varphi(t) dt - \frac{u_k}{k + N_\delta} \right| \leq \sum_{\substack{0 \leq j \leq M \\ j \neq k}} \frac{1}{j + N_\delta} \widehat{\varphi}(\lambda_j - \lambda_k).$$

By applying the lemma 4.4, that

$$\left| \int_{I_\delta} T_\delta(t) e^{2i\pi\lambda_k t} \varphi(t) dt - \frac{u_k}{k + N_\delta} \right| \leq \frac{1 - \alpha_\delta}{k + N_\delta}.$$

It follows that

$$\left| \int_{I_\delta} T_\delta(t) a_k e^{2i\pi\lambda_k t} \varphi(t) dt - \frac{u_k a_k}{k + N_\delta} \right| \leq \frac{1 - \alpha_\delta}{k + N_\delta} |a_k|.$$

Using the fact that $u_k a_k = |a_k|$ and the triangular inequality, we obtain

$$\left| \int_{I_\delta} T_\delta(t) \left(\sum_{k=0}^N a_k e^{2i\pi\lambda_k t} \right) \varphi(t) dt - \sum_{k=0}^N \frac{|a_k|}{k + N_\delta} \right| \leq (1 - \alpha_\delta) \sum_{k=0}^N \frac{|a_k|}{k + N_\delta}$$

from which the lemma follows immediately. \square

Construction of \tilde{T}_δ

Recall that, from our assumption on N_δ and N , we can write

$$T_\delta(t) = \sum_{k=0}^M \frac{u_k}{k + N_\delta} e^{-2i\pi\lambda_k t} = \sum_{j=m_\delta}^{n_\delta} f_j(t)$$

where we set $D_j = \{k \in \mathbb{N} : 2^j \leq k < 2^{j+1}\}$ and

$$f_j(t) = \sum_{r+N_\delta \in D_j} \frac{u_r}{r + N_\delta} e^{-2i\pi\lambda_r t}.$$

Before we estimate the norms of the f_j 's, let us recall Hilbert's inequality (see e.g. [4, Chapter 10]).

Lemma 4.6 (Hilbert's inequality). *Let $\lambda_1, \dots, \lambda_N$ be real numbers with $|\lambda_k - \lambda_\ell| \geq 1$ when $k \neq \ell$, and let z_1, \dots, z_N be complex numbers. We have*

$$\left| \sum_{\substack{1 \leq k, \ell \leq N \\ k \neq \ell}} \frac{z_k \overline{z_\ell}}{\lambda_k - \lambda_\ell} \right| \leq \pi \sum_{k=1}^N |z_k|^2.$$

We can now prove the following:

Lemma 4.7. *For $m_\delta \leq j \leq n_\delta$,*

$$(1) \quad \|f_j\|_\infty \leq 1$$

$$(2) \|f_j\|_{L^2(I_\delta)} \leq 2^{-\frac{j}{2}} \sqrt{|I_\delta| + 1}.$$

Proof. The first estimate is obtained by straightforward computations: for all t ,

$$|f_j(t)| \leq \sum_{r+N_\delta \in D_j} \frac{1}{r+N_\delta} \leq \frac{|\mathbb{D}_j|}{2^j} = 1$$

hence $\|f_j\|_\infty \leq 1$.

For the second one, set $v_r = \frac{u_r}{r+N_\delta}$. Then

$$\begin{aligned} \|f_j\|_{L^2(I_\delta)}^2 &= \int_{I_\delta} f_j(t) \overline{f_j(t)} dt = \int_{I_\delta} \sum_{r+N_\delta, s+N_\delta \in D_j} v_r \overline{v_s} e^{-2i\pi(\lambda_r - \lambda_s)t} dt \\ &= |I_\delta| \sum_{r+N_\delta \in D_j} |v_r|^2 + \sum_{\substack{r+N_\delta, s+N_\delta \in D_j \\ r \neq s}} v_r \overline{v_s} \int_{-|I_\delta|/2}^{|I_\delta|/2} e^{-2i\pi(\lambda_r - \lambda_s)t} dt \\ &= |I_\delta| \sum_{r+N_\delta \in D_j} |v_r|^2 + \sum_{\substack{r+N_\delta, s+N_\delta \in D_j \\ r \neq s}} v_r \overline{v_s} \left(\frac{e^{i|I_\delta|\pi(\lambda_r - \lambda_s)} - e^{-i|I_\delta|\pi(\lambda_r - \lambda_s)}}{2i\pi(\lambda_r - \lambda_s)} \right). \end{aligned}$$

It follows that

$$\|f_j\|_{L^2(I)}^2 \leq |I_\delta| \sum_{r+N_\delta} |v_r|^2 + \frac{1}{2\pi} \left| \sum_{\substack{r,s \\ r \neq s}} \frac{v_r e^{-i\pi\lambda_r(1+\delta)} \overline{v_s e^{-i\pi\lambda_s(1+\delta)}}}{\lambda_s - \lambda_r} \right| + \frac{1}{2\pi} \left| \sum_{\substack{r,s \\ r \neq s}} \frac{v_r e^{i\pi\lambda_r(1+\delta)} \overline{v_s e^{i\pi\lambda_s(1+\delta)}}}{\lambda_s - \lambda_r} \right|$$

We then apply Hilbert's inequality 4.6 to the last two sums to obtain

$$\begin{aligned} \|f_j\|_2^2 &\leq |I_\delta| \sum_{r+N_\delta} |v_r|^2 + \frac{1}{2} \sum_{r+N_\delta, \beta \in D_j} |v_r|^2 + \frac{1}{2} \sum_{r+N_\delta \in D_j} |v_r|^2 \\ &= (|I_\delta| + 1) \sum_{r+N_\delta} |v_r|^2. \end{aligned}$$

Since

$$\begin{aligned} \sum_{r+N_\delta \in D_j} |v_r|^2 &= \sum_{r+N_\delta \in D_j} \frac{|u_r|^2}{(r+N_\delta)^2} = \sum_{r+N_\delta \in D_j} \frac{1}{(r+N_\delta)^2} \\ &\leq \sum_{r+N_\delta \in D_j} \frac{1}{2^{2j}} \leq \frac{2^j}{2^{2j}} = 2^{-j}, \end{aligned}$$

we get $\|f_j\|_{L^2(I)}^2 \leq (|I_\delta| + 1)2^{-j}$ as claimed. \square

We then write the Fourier series expansion of $|f_j| \in L^2(I_\delta)$ as

$$|f_j(t)| = \sum_{s \in \mathbb{Z}} a_{s,j} e^{\frac{2i\pi}{|I_\delta|} st},$$

and define $h_j \in L^2(I_\delta)$ via its Fourier series expansion

$$h_j(t) = a_{0,j} + 2 \sum_{s=1}^{\infty} a_{s,j} e^{\frac{2is\pi}{|I_\delta|} t}.$$

Lemma 4.8. *For $m_\delta \leq j \leq n_\delta$, the following properties hold*

- (1) $\operatorname{Re}(h_j) = |f_j|$;
- (2) $\|h_j\|_2 \leq \sqrt{2} \|f_j\|_2$;
- (3) $h_j \in H^\infty(I_\delta)$.

Proof. First, as $|f_j|$ is real valued, its zero Fourier coefficient $a_{0,j}$ is also real, and for the remaining ones $\overline{a_{s,j}} = a_{-s,j}$ for every $s \geq 1$. A direct computation then shows that $\operatorname{Re}(h_j) = |f_j|$ which is less than 1 by lemma 4.7 while Parseval shows that $\|h_j\|_2 \leq \sqrt{2}\|f_j\|_2$. \square

We now define a sequence $(F_j)_{j \geq m_\delta}$ inductively through

$$F_{m_\delta} = f_{m_\delta} \quad \text{and} \quad F_{j+1} = F_j e^{-\varepsilon h_{j+1}} + f_{j+1}$$

where $0 < \varepsilon < 1$ is a small parameter that we will adjust later. Further set

$$E_\varepsilon := \sup_{0 < x \leq 1} \frac{x}{1 - e^{-\varepsilon x}} = \frac{1}{\varepsilon} \sup_{0 < x \leq \varepsilon} \frac{x}{1 - e^{-x}} = \frac{1}{1 - e^{-\varepsilon}}.$$

From the next to last identity, it is easy to obtain the following simple bound:

$$\frac{1}{\varepsilon} \leq E_\varepsilon \leq \frac{2}{\varepsilon}.$$

Lemma 4.9. *For all $m_\delta \leq j \leq n_\delta$,*

$$\|F_j\|_\infty \leq \frac{2}{\varepsilon}.$$

Proof. By definition of E_ε , if $C \leq E_\varepsilon$ and $0 \leq x \leq 1$, then $Ce^{-\varepsilon x} + x \leq E_\varepsilon e^{-\varepsilon x} + x \leq E_\varepsilon$.

We can now prove by induction over j that $|F_j| \leq E_\varepsilon$ from which the lemma follows. First, when $j = 0$, from Lemma 4.7 we get

$$\|F_0\|_\infty = \|f_0\|_\infty \leq 1 \leq E_\varepsilon.$$

Assume now that $\|F_j\|_\infty \leq E_\varepsilon$, then

$$\begin{aligned} |F_{j+1}(t)| &= |F_j(t)e^{-\varepsilon h_{j+1}(t)} + f_{j+1}(t)| \leq |F_j(t)|e^{-\varepsilon \Re(h_{j+1}(t))} + |f_{j+1}(t)| \\ &= |F_j(t)|e^{-\varepsilon |f_{j+1}(t)|} + |f_{j+1}(t)|. \end{aligned}$$

As $|f_{j+1}(t)| \leq 1$ and $|F_j(t)| \leq E_\varepsilon$, we get $|F_{j+1}(t)| \leq E_\varepsilon$ as claimed. \square

Lemma 4.10. *Let $m_\delta \leq n \leq n_\delta$. For $j = m_\delta, \dots, n$ we define $g_{j,n} = e^{-\varepsilon H_{j,n}}$ with*

$$H_{j,n} = \begin{cases} h_{j+1} + \dots + h_n & \text{if } j < n \\ 0 & \text{if } j = n \end{cases}.$$

Then $F_n = \sum_{j=m_\delta}^n f_j g_{j,n}$. Moreover $\|H_{j,n}\|_2 \leq \frac{\sqrt{2(|I_\delta| + 1)}}{\sqrt{2} - 1} 2^{-\frac{j}{2}}$.

Proof. The first part is obtained by a simple induction on n . Moreover, the lemma 4.8 implies

$$\begin{aligned} \|H_{j,n}\|_2 &= \left\| \sum_{r=j+1}^n h_r \right\|_2 \leq \sum_{r=j+1}^n \|h_r\|_2 \leq \sqrt{2} \sum_{r=j+1}^n \|f_r\|_2 \\ &\leq \sqrt{2(|I_\delta| + 1)} \sum_{r=j+1}^{\infty} 2^{-\frac{r}{2}} = \frac{\sqrt{2(|I_\delta| + 1)}}{\sqrt{2} - 1} 2^{-\frac{j}{2}}, \end{aligned}$$

as stated. \square

Next, we will need the following simple well-known lemma:

Lemma 4.11. *If $H \in H^\infty$ and $\operatorname{Re}(H) \geq 0$, then $e^{-H} \in H^\infty$ and*

$$\|e^{-H} - 1\|_2 \leq \|H\|_2.$$

Proof. Since H^∞ is a Banach algebra, the partial sums $\sum_{k=0}^n (-1)^k \frac{H^k}{k!}$ of e^{-H} are elements of H^∞ . Moreover, since H is bounded, these sums converge uniformly toward e^{-H} , with $e^{-H} \in H^\infty$. Finally, if $z \in \mathbb{C}$ and $\Re(z) \geq 0$,

$$|e^{-z} - 1| = \left| \int_0^1 z e^{-tz} dt \right| \leq \int_0^1 |z| e^{-t\Re(z)} dt \leq |z|.$$

In our case $z = H(t)$, and we have

$$|e^{-H(t)} - 1| \leq |H(t)|$$

and by integration we have the desired inequality. \square

Lemma 4.12. *Assume that $0 < \varepsilon \leq \frac{\sqrt{|I_\delta|}(\sqrt{2}-1)}{\sqrt{2(|I_\delta|+1)}}$. Then, for $m_\delta \leq j \leq n \leq n_\delta$,*

$$(1) \|g_{j,n} - 1\|_2 \leq \varepsilon \|H_{j,n}\|_2 \leq \frac{\sqrt{2(|I_\delta|+1)}}{\sqrt{2}-1} \varepsilon 2^{-\frac{j}{2}}$$

$$(2) \text{ The Fourier series of } g_{j,n}(t) - 1 \text{ writes } g_{j,n}(t) - 1 = \sum_{s \geq 0} c_{s,j} e^{\frac{2is\pi}{|I|}t}, \text{ with } \sum_{s=0}^{+\infty} |c_{s,j}|^2 \leq 1.$$

Proof. By lemma 4.11, $\|g_j - 1\|_2 \leq \varepsilon \|H_j\|_2$. Then, since $g_{j,n}$ is analytic, its Fourier series writes

$$g_j(t) - 1 = \sum_{s \geq 0} c_{s,j} e^{\frac{2is\pi}{|I_\delta|}st}.$$

But then, with Parseval

$$\left(\sum_{s \geq 0} |c_{s,j}|^2 \right)^{\frac{1}{2}} = \frac{1}{\sqrt{|I_\delta|}} \|g_j - 1\|_2 \leq \frac{\varepsilon}{\sqrt{|I_\delta|}} \|H_j\|_2 \leq \frac{\varepsilon}{\sqrt{|I_\delta|}} \frac{\sqrt{2(|I_\delta|+1)}}{\sqrt{2}-1} 2^{-\frac{j}{2}} \leq 2^{-\frac{j}{2}} < 1$$

which implies the claimed bound. \square

Now recall that

$$T_\delta = \sum_{m_\delta \leq j \leq n_\delta} f_j$$

and define

$$\tilde{T}_\delta = F_{n_\delta}.$$

In particular, from Lemma 4.9, we have

$$(4.11) \quad \|\tilde{T}_\delta\|_\infty \leq \frac{2}{\varepsilon}.$$

The key estimates here is the following;

Lemma 4.13. *Once again, we use the notations of the lemma 4.4. There exists $\delta_2 > 0$ such that, if $0 < \delta < \delta_2$ and $N_\delta \geq \delta^{-7/2}$ then*

$$(4.12) \quad \left| \int_{I_\delta} (\tilde{T}_\delta - T_\delta)(t) \left(\sum_{k=0}^N a_k e^{2i\pi\lambda_k t} \right) \varphi(t) dt \right| \leq \frac{2}{3} \alpha_\delta \sum_{k=0}^N \frac{|a_k|}{k + N_\delta}.$$

where φ is the function defined in (4.7).

Proof. It is enough to prove that, under the conditions of the lemma, for $0 \leq k \leq N$ we have

$$(4.13) \quad \left| \int_{I_\delta} (\tilde{T}_\delta - T_\delta)(t) e^{2i\pi\lambda_k t} \varphi(t) dt \right| \leq \frac{2}{3} \frac{\alpha_\delta}{k + N_\delta}.$$

Once (4.13) is established, it will then be enough to multiply the left hand side by a_k and to use the triangular inequality

We fix $k \in [0, N]$ and let ℓ the index such that $k + N_\delta \in D_\ell$. We define R, R_1 and R_2 as follows

$$\begin{aligned} R &= \int_{I_\delta} (\tilde{T}_{\delta, \beta} - T_{\delta, \beta})(t) e^{2i\pi\lambda_k t} \varphi(t) dt \\ &= \int_{I_\delta} \sum_{m_\delta \leq j \leq \ell-2} f_j(t) (g_j(t) - 1) e^{2i\pi\lambda_k t} \varphi(t) dt + \int_{I_\delta} \sum_{\ell-1 \leq j \leq n_\delta} f_j(t) (g_j(t) - 1) e^{2i\pi\lambda_k t} \varphi(t) dt \\ &:= R_1 + R_2 \end{aligned}$$

We will first bound R_1 . Note that if $s \in \mathbb{Z}$,

$$\begin{aligned} \int_{I_\delta} f_j(t) \varphi(t) e^{2i\pi\lambda_k t} e^{\frac{2i\pi}{|\delta|} s t} dt &= \int_{I_\delta} \sum_{r+N_\delta \in D_j} \frac{u_r}{r+N_\delta} \varphi(t) e^{2i\pi(-\lambda_r + \lambda_k + \frac{s}{|\delta|}) t} dt \\ &= \sum_{r+N_\delta \in D_j} \frac{u_r}{r+N_\delta} \hat{\varphi}(\lambda_r - \lambda_k - \frac{s}{|\delta|}). \end{aligned}$$

From there, we obtain

$$\begin{aligned} \int_{I_\delta} f_j(t) (g_j(t) - 1) e^{2i\pi\lambda_k t} \varphi(t) dt &= \int_{I_\delta} f_j(t) \varphi(t) e^{2i\pi\lambda_k t} \sum_{s \geq 0} c_{s,j} e^{\frac{2i\pi}{|\delta|} s t} dt \\ &= \sum_{s=0}^{+\infty} c_{s,j} \int_{I_\delta} f_j(t) \varphi(t) e^{2i\pi\lambda_k t} e^{\frac{2i\pi}{|\delta|} s t} dt \\ &= \sum_{s=0}^{+\infty} c_{s,j} \sum_{r+N_\delta \in D_j} \frac{u_r}{r+N_\delta} \hat{\varphi}(\lambda_r - \lambda_k - \frac{s}{|\delta|}) \\ &= \sum_{r+N_\delta \in D_j} \frac{u_r}{r+N_\delta} \sum_{s=0}^{\infty} c_{s,j} \hat{\varphi}(\lambda_r - \lambda_k - \frac{s}{|\delta|}). \end{aligned}$$

So finally we get

$$R_1 = \sum_{m_\delta \leq j \leq \ell-2} \sum_{r+N_\delta \in D_j} \frac{u_r}{r+N_\delta} \sum_{s=0}^{\infty} c_{s,j} \hat{\varphi}(\lambda_r - \lambda_k - \frac{s}{|\delta|}).$$

Let $c_s(r) = c_{s,j}$ if $r + N_\delta \in I_j$. Since $\hat{\varphi}$ is an even function, we can write

$$R_1 = \sum_{2^{m_\delta} \leq r+N_\delta < 2^{\ell-1}} \frac{u_r}{r+N_\delta} \sum_{s=0}^{\infty} c_s(r) \hat{\varphi}(\lambda_k - \lambda_r + \frac{s}{|\delta|}) := \sum_{2^{m_\delta} \leq r+N_\delta < 2^{\ell-1}} \frac{u_r}{r+N_\delta} E_r.$$

From Lemma 4.3, recall that, with $D_\delta = c_0 \delta^{-\frac{p+q}{p+2}}$ and $\nu = \frac{\alpha_\delta}{|I_\delta|^{1/q}}$ then, for

$$|\lambda| \geq \max(1, D_\delta \nu^{-\frac{1}{p+2}}),$$

we have

$$(4.14) \quad |\hat{\varphi}(\lambda)| \leq \frac{\alpha_\delta}{(|I_\delta| |\lambda|)^q}.$$

Now, $s \geq 0$, $|I_\delta| \geq 1$ and, as $r + N_\delta < 2^{\ell-1}$, $2^\ell \leq k + N_\delta < 2^{\ell+1}$, $\lambda_k > \lambda_r$ thus

$$(4.15) \quad \begin{aligned} |I_\delta| \left| \lambda_k - \lambda_r + \frac{s}{|\delta|} \right| &= |I_\delta| (\lambda_k - \lambda_r) + s \geq \lambda_k - \lambda_r + s \geq k - r + s \\ &= (k + N_\delta) - (r + N_\delta) > 2^\ell - 2^{\ell-1} = 2^{\ell-1} \geq 2^{m_\delta-1}. \end{aligned}$$

Further, from (4.8) and $1 \leq |I_\delta| \leq 2$, $D_\delta \nu^{-\frac{1}{p+2}} \leq c_3 \delta^{-\frac{p+q+2}{p+2}} = c_3 \delta^{-\frac{20}{12}}$. Thus, choosing m_δ sufficiently large for $N_\delta = 2^{m_\delta} \delta^{-7/2}$ and $\delta \leq \delta_2$ for some $\delta_2 > 0$ small enough, we are able to apply (4.14) and obtain, with (4.15)

$$|\hat{\varphi}(\lambda_k - \lambda_r + \frac{s}{|\delta|})| \leq \frac{\alpha_\delta}{(k - r + s)^q}.$$

We can now bound E_r . Since $\sum_{s=0}^{\infty} |c_s(r)|^2 \leq 1$, the previous bound and Cauchy-Schwarz give us

$$\begin{aligned} |E_r| &= \sum_{s=0}^{\infty} |c_s(r)| \left| \widehat{\varphi} \left(\lambda_k - \lambda_r + \frac{s}{1+\delta} \right) \right| \leq \alpha_{\delta} \left(\sum_{s=0}^{\infty} \frac{1}{(k-r+s)^{2q}} \right)^{1/2} \\ &\leq \alpha_{\delta} \left(\sum_{n=k-r}^{\infty} \sum_{n \geq k-r} \frac{1}{n^{2q}} \right)^{1/2} \\ &\leq \alpha_{\delta} \left(\int_{k-r-1}^{\infty} \frac{dt}{t^{2q}} \right)^{1/2} = \frac{\sqrt{2q-1} \alpha_{\delta}}{(k-r-1)^{q-1/2}}. \end{aligned}$$

But $k-r > 2^{\ell-1}$ then $k-r-1 \geq 2^{\ell-1}$. Since $k+N_{\delta} \in \mathcal{D}_{\ell}$ i.e. $2^{\ell} \leq k+N_{\delta} \leq 2^{\ell+1}$ we get $\frac{1}{k-r-1} \leq \frac{4}{k+N_{\delta}}$ and then

$$|E_r| \leq \frac{4^{q-1/2} \sqrt{2q-1} \alpha_{\delta}}{25(k+N_{\delta})^{q-1/2}}$$

Finally, we deduce that

$$\begin{aligned} |R_1| &= \left| \sum_{2^{m_{\delta}} \leq r+N_{\delta} < 2^{\ell-1}} \frac{u_r}{r+N_{\delta}} E_r \right| \leq \sum_{2^{m_{\delta}} \leq r+N_{\delta} < 2^{\ell-1}} \frac{|E_r|}{r+N_{\delta}} \\ &\leq \frac{4^{q-1/2} \sqrt{2q-1} \alpha_{\delta}}{(k+N_{\delta})^{q-1/2}} \end{aligned}$$

since last sum has at most $2^{\ell-1} \leq k+N_{\delta}$.

We will now bound R_2 .

$$\begin{aligned} |R_2| &\leq \sum_{\ell-1 \leq j \leq n_{\delta}} \int_I |f_j(t)| |g_j(t) - 1| |\varphi(t)| dt \\ &\leq \|\varphi\|_{\infty} \sum_{\ell-1 \leq j \leq n_{\delta}} \|f_j\|_2 \|g_j - 1\|_2. \end{aligned}$$

According to the lemmas 4.7 (1) and 4.12 (1) we get

$$|R_2| \leq \|\varphi\|_{\infty} \varepsilon \frac{\sqrt{2}(2+\delta)}{\sqrt{2}-1} \sum_{l-1 \leq j \leq m} 2^{-j}.$$

since

$$\sum_{\ell-1 \leq j \leq n_{\delta}} 2^{-j} \leq \sum_{j=\ell-1}^{\infty} 2^{-j} = 2^{-\ell+2} = 8 \cdot 2^{-(\ell+1)}$$

and $k+N_{\delta} \in I_l$ then $\frac{1}{2^{\ell+1}} \leq \frac{1}{k+N_{\delta}}$. Consequently

$$\sum_{\ell-1 \leq j \leq n_{\delta}} 2^{-j} \leq \frac{8}{k+N_{\delta}}.$$

Finally, we deduce that

$$R_2 \leq \varepsilon \|\varphi\|_{\infty} \frac{\sqrt{2}(|I_{\delta}|+1)}{\sqrt{2}-1} \frac{8}{k+N_{\delta}}$$

and we obtain $R_2 \leq \frac{\alpha_{\delta}}{3} \frac{1}{k+N_{\delta}}$ when $\varepsilon \leq \frac{(\sqrt{2}-1)}{24\|\varphi\|_{\infty}(|I_{\delta}|+1)\sqrt{2}} \alpha_{\delta}$. □

Note that, from (4.8) and $|I_\delta| \leq 2$, we can take $\varepsilon = c_4 \delta^2$ for some $c_4 > 0$.

It is now easy to deduce the theorem 4.2 using the 2 inequalities (4.10) and (4.12).

Proof of theorem 4.2. Let $S_\delta = \sum_{k=0}^N \frac{|a_k|}{k + N_\delta}$ and $\Phi(t) = \sum_{k=0}^N a_k e^{2i\pi\lambda_k t}$ as previously defined. Recall that in (4.10), we have shown that

$$\begin{aligned} \alpha_\delta S_\delta &\leq \left| \int_{I_\delta} T_\delta(t) \phi(t) \varphi(t) dt \right| \\ &\leq \left| \int_{I_\delta} \tilde{T}_\delta(t) \phi(t) \varphi(t) dt \right| + \left| \int_{I_\delta} (\tilde{T}_\delta(t) - T_\delta(t)) \phi(t) \varphi(t) dt \right| \\ &\leq \left| \int_{I_\delta} \tilde{T}_\delta(t) \phi(t) \varphi(t) dt \right| + \frac{2}{3} \alpha_\delta S_\delta \end{aligned}$$

with (4.12).

It follows that

$$\begin{aligned} S_\delta &\leq \frac{3}{\alpha_\delta} \left| \int_{I_\delta} \tilde{T}_\delta(t) \phi(t) \varphi(t) dt \right| \\ &\leq \frac{3 \|\tilde{T}\|_\infty \|\varphi\|_\infty}{\alpha_\delta} \int_{I_\delta} |\phi(t)| dt. \end{aligned}$$

But, if δ is small enough and $N_\delta \geq \delta^{-7/2}$,

– from Lemma 4.3, $\|\varphi\|_\infty \leq \frac{\pi}{2}$;

– from (4.8), $\alpha_\delta = \kappa \delta^2 + O(\delta^4)$

– as $\varepsilon = c_4 \delta^2$, from (4.11), $\|\tilde{T}\|_\infty = \frac{2}{c_4 \delta^2}$.

Therefore, there are two absolute constants δ_* and c_* such that, if $\delta \leq \delta_*$,

$$S_\delta \leq \frac{c_*}{\delta^4} \int_{I_\delta} |\phi(t)| dt.$$

As noticed at the start of the proof, this implies that

$$\sum_{n=0}^N \frac{|a_n|}{k+1} \leq \frac{c_*}{\delta^{\frac{15}{2}}} \int_{-\frac{1+\delta}{2}}^{\frac{1+\delta}{2}} \left| \sum_{k=0}^N a_k e^{2i\pi\lambda_k t} \right| dt$$

for every N , every sequence of real numbers $(\lambda_j)_{j \geq 0}$ with $\lambda_{j+1} - \lambda_j \geq 1$ and every complex sequence $(a_j)_{j \geq 0}$. \square

We have not fully optimised the proof, by taking q sufficiently large and p/q sufficiently large, one can replace $\delta^{15/2}$ by $\delta^{7+\eta}$ for any fixed η .

5. SOME OPEN PROBLEMS

The L^2 theory of exponential sums is rather well understood. This is not the case for L^1 theory for which there are still many open questions. Let us mention a few of them.

- (1) There is a major difference between the sums that appear in Ingham's Theorem and those that appear in Mc Geehe, Pigno, Smith and Nazarov's Theorems. In the L^2 case, the sums can be two sided and not in the L^1 case. The proof given here does not work in this case (this is due to the construction of \tilde{T}) and we are tempted to conjecture the following

Question 1. Let $T > 1$ and $(\lambda_k)_{k \in \mathbb{Z}}$ a real sequence such that $\lambda_{k+1} - \lambda_k \geq 1$ for every k and λ_k has same sign as k .

Does there exist a constant C such that, for every N and every sequence $(c_j)_{j=-N, \dots, N}$ of complex numbers,

$$\sum_{k=-N}^N \frac{|c_k|}{|k|+1} \leq C \int_{-T/2}^{T/2} \left| \sum_{k=-N}^N c_k e^{2i\pi\lambda_k t} \right| dt.$$

- (2) The second question is the optimality of the condition $T = 1$ in Nazarov's Theorem. In view of Ingham's counterexample, it is tempting to conjecture that $T > 1$ is requested (as observed by Nazarov, his proof requires this).

Question 2. *Is the condition $T > 1$ necessary in Nazarov's Theorem. If yes, what is the right behaviour of the constants.*

We think the estimate shown here is not optimal.

- (3) Another question related to the previous one comes from Haraux's Theorem. When $\lambda_{k+1} - \lambda_k \rightarrow +\infty$, T can be chosen arbitrarily small. A key element of Haraux's strategy is that the lower and upper bounds in Ingham's inequality are both multiples of the ℓ^2 -norm of the coefficients. This is no longer the case in the L^1 setting.

In forthcoming work, we have been able to use a compactness argument to show that one may take T arbitrarily small in Nazarov's Theorem when $\lambda_{k+1} - \lambda_k \rightarrow +\infty$. However, in doing so, we loose control of constants. This leads to the following:

Question 3. *Prove a quantitative version Nazarov's Theorem with T arbitrarily small when $\lambda_{k+1} - \lambda_k \rightarrow +\infty$.*

- (4) Are $L^1([-T, T])$ -norms of lacunary non-harmonic Fourier series comparable to the ℓ^2 -norms of the coefficients? This is the case for Besikovich norms.

Note that

$$\sum_{j=0}^{\infty} \frac{|a_j|}{j+1} \leq \left(\sum_{j=0}^{\infty} \frac{1}{(j+1)^2} \right)^{1/2} \left(\sum_{j=0}^{\infty} |a_j|^2 \right)^{1/2} = \frac{\pi}{\sqrt{6}} \left(\sum_{j=0}^{\infty} |a_j|^2 \right)^{1/2}.$$

The question is then

Question 4. *Find a (gap) condition on (λ_k) and on T that implies that there is a constant $C > 0$ such that, for every (a_k) with finite support*

$$\frac{1}{T} \int_{-T/2}^{T/2} \left| \sum_{j=0}^{+\infty} a_j e^{2i\pi\lambda_j t} \right| dt \geq C \left(\sum_{j=0}^{\infty} |a_j|^2 \right)^{1/2}.$$

One may also go the other way and ask whether one can still obtain a result like Theorem B when $\lambda_{k+1} - \lambda_k \rightarrow 0$ but with a smaller power on the denominator. For instance, when $\lambda_k = \ln k$ then the Fourier series become Dirichlet series

$$\sum_{k=1}^{+\infty} \frac{a_k}{k^{2i\pi t}} = \sum_{k=1}^{+\infty} a_k e^{2i\pi(\ln k)t}.$$

The following question was asked in [3]:

Question 5. *Is it true that, for every sequence $(a_k)_{k \geq 1}$ with finite support,*

$$C \left(|a_1| + \sum_{k \geq 2} \frac{|a_k|}{\sqrt{k} \ln k} \right) \leq \lim_{T \rightarrow +\infty} \int_{-T/2}^{T/2} \left| \sum_{k=1}^{+\infty} \frac{a_k}{k^{2i\pi t}} \right| dt.$$

REFERENCES

- [1] A. S. BELOV & S. V. KONYAGIN *On the conjecture of Littlewood and minima of even trigonometric polynomials*. Harmonic analysis from the Pichorides viewpoint (Anogia, 1995), 1–11, Publ. Math. Orsay, **96-01**, Univ. Paris XI, Orsay, 1996.
 [2] E. BELLER & D. J. NEWMAN *An l^1 extremal problem for polynomials*. Proc. Amer. Math. Soc. **29** (1971), 474–481.
 [3] O. F. BREVIK, K.-M. PERFEKT, K. SEIP, A. G. SISKAKIS & D. VUKOTIĆ *The multiplicative Hilbert matrix*. Adv. Math. **302** (2016), 410–432.

- [4] D. CHOIMET & H. QUEFFÉLEC *Twelve landmarks of twentieth-century analysis*. Cambridge University Press, New York, 2015
- [5] P. J. COHEN *On a conjecture of Littlewood and idempotent measures*. Amer. J. Math. **82** (1960), 191–212.
- [6] H. DAVENPORT *On a theorem of P. J. Cohen*. Mathematika **7** (1960), 93–97.
- [7] R. A. DEVORE & G. G. LORENTZ *Constructive Approximation*. Grundlehren der mathematischen Wissenschaften, 303. Springer-Verlag, Berlin, 1993.
- [8] J. J. F. FOURNIER *On a theorem of Paley and the Littlewood conjecture*. Ark. Mat. **17** (1979), 199–216.
- [9] L. GRAFAKOS *Classical Fourier analysis*. Third edition. Graduate Texts in Mathematics, **249**. Springer, New York, 2014
- [10] B. HANSON *Littlewood’s Problem for Sets with Multidimensional Structure.*, I.M.R.N. **21** (2021), 16736–16750.
- [11] S. HUDSON & M. LECKBAND *Hardy’s inequality and fractal measures*. J. Funct. Anal. **108** (1992), 133–160.
- [12] A. E. INGHAM *Some trigonometrical inequalities with applications to the theory of series*. Math. Z. **41** (1936), 367–379.
- [13] PH. JAMING, K. KELLAY & C. SABA *The Littlewood problem and non-harmonic Fourier series*. Math. Ann. (2023). <https://doi.org/10.1007/s00208-023-02692-w>.
- [14] V. KOMORNIK & P. LORETI *Fourier Series in Control Theory*. Springer-Verlag, New York, 2005.
- [15] S. V. KONYAGIN *On the Littlewood problem*. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. **45** (1981), no. 2, 243–265, 463.
- [16] O. C. MCGEHEE, L. PIGNO & B. SMITH *Hardy’s inequality and the L^1 norm of exponential sums*. Ann. of Math. (2) **113** (1981), 613–618.
- [17] F. L. NAZAROV *On a proof of the Littlewood conjecture by McGehee, Pigno and Smith*. St. Petersburg Math. J. **7** (1996), no. 2, 265–275
- [18] D. J. NEWMAN *An L^1 Extremal Problem for Polynomials*. Proc. Amer. Math. Soc. **16** (1965), no. 6, 1287–1290
- [19] S. K. PICHORIDES *A lower bound for the L^1 norm of exponential sums*. Mathematika **21** (1974), 155–159.
- [20] S. K. PICHORIDES *On a conjecture of Littlewood concerning exponential sums. I*. Bull. Soc. Math. Grèce (N.S.) **18** (1977), 8–16.
- [21] S. K. PICHORIDES *On a conjecture of Littlewood concerning exponential sums. II*. Bull. Soc. Math. Grèce (N.S.) **19** (1978), 274–277.
- [22] S. K. PICHORIDES *On the L^1 norm of exponential sums*. Ann. Inst. Fourier (Grenoble) **30** (1980), 79–89.
- [23] X. SHAO *On character sums and exponential sums over generalized arithmetic progressions*. Bull. London Math. Soc. **45** (2013), 541–550.
- [24] R. M. TRIGUB & E. S. BELINSKY *Fourier Analysis and Approximation of Functions*, Kluwer, 2004.
- [25] A. ZYGMUND *Trigonometric Series, vol. I*. Cambridge 1959.