



HAL
open science

Transient imine as a directing group for the Pd-catalyzed anomeric C(sp³)-H arylation of 3-aminosugars

Juba Ghouillem, Sokna Bazzi, Nicolas Grimblat, Pascal Retailleau, Vincent Gandon, Samir Messaoudi

► To cite this version:

Juba Ghouillem, Sokna Bazzi, Nicolas Grimblat, Pascal Retailleau, Vincent Gandon, et al.. Transient imine as a directing group for the Pd-catalyzed anomeric C(sp³)-H arylation of 3-aminosugars. *Chemical Communications*, 2023, 59 (17), pp.2497-2500. 10.1039/D3CC00046J . hal-04315221

HAL Id: hal-04315221

<https://hal.science/hal-04315221>

Submitted on 30 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Transient imine as a directing group for the Pd-catalyzed anomeric C(sp³)-H arylation of 3-aminosugars

Juba Ghouillem,^a Sokna Bazzi,^a Nicolas Grimblat,^b Pascal Retailleau,^c Vincent Gandon^{d,b} and Samir Messaoudi^{a*}

⁵ Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

The first example of Pd(II)-catalyzed anomeric arylation of 3-aminosugars is reported by using an L,X-type transient directing groups (TDG) approach combined with an external 2-pyridone ligand. The released free amine was *in situ* transformed into an azide function, which was then exploited in a CuAAC to increase the molecular complexity and prepare a variety of complex substituted C3-triazolo C-glycosides in good yields.

The aryl C-glycoside motif, which is found in many natural products and drug candidates, has attracted tremendous attention in medicinal chemistry and drug discovery.¹ These glycomimetics often feature enhanced resilience to enzymatic and hydrolytic cleavage under biological conditions compared to their O/N-glycoside congeners.² In particular, aryl C-glycosides bearing a nitrogen atom at the C3 position of the sugar moiety (e.g. 3-amino or 3-triazolo C-glycosides) are one of the most important families of C-glycosides found in natural bioactive compounds.³ These glycosides emerged as a privileged class with diverse potential applications (Figure 1).

Classical methods for the synthesis of aryl C-glycoside have largely relied on the use of metal-catalyzed cross-coupling strategies involving the use of stoichiometric organometallic coupling partners.^{3f-h} The persistent need from drug discovery programs to elaborate complex aryl C-glycosides has led to pioneering developments in the field of C-H functionalization.⁴ This approach, which avoids the preactivation of the coupling partners, has led to new paradigms for the synthesis of aryl C-glycosides. In this context, several groups have reported elegant methods by applying metal-catalyzed activation of C(sp²)-H bonds. However, methods involving a direct activation of the less reactive C(sp³)-H bonds of the sugar partners are rare.⁵ Thus far, precise structural design and selective modification of glycosides remain challenging tasks. Recently, Ackermann's group^{5e} and our

⁴⁰ ^a Université Paris-Saclay, CNRS, BioCIS, 92290, Châtenay-Malabry, France, Tel: + (33) 0146835887; E-mail: samir.messaoudi@universite-paris-saclay.fr

^b Laboratoire de Chimie Moléculaire (LCM), CNRS UMR 9168, Ecole Polytechnique, Institut Polytechnique de Paris, route de Saclay, 91120 Palaiseau France.

^c Institut de Chimie des Substances Naturelles, CNRS UPR 2301, Université Paris-

⁴⁵ Saclay, avenue de la terrasse, 91198 Gif-sur-Yvette, France

^d Université Paris-Saclay, CNRS, ICMMO, 91405, Orsay cedex, France

[†] Electronic Supplementary Information (ESI) available: General, experimental procedures for starting materials and ¹H and ¹³C spectra for all new compounds. See DOI: 10.1039/b000000x/





Figure 1. Aryl C-glycosides-based bioactive molecules



Scheme 1. Strategies to access aryl C-glycosides.

group^{5d} reported independently two studies in which the anomeric, as well as C-3 position of the sugar moieties, were functionalized selectively through a Pd-catalyzed C(sp³)-H activation reaction (Scheme 1-A-B). Nevertheless, these strategies require synthetic steps to prepare the activated sugar substrate and remove the directing group. To address this issue, we sought to develop a transient imine directing group approach⁶ for the Pd-catalyzed anomeric C(sp³)-H arylation of

3-aminosugars to access aryl C-glycosides in an expedient way (Scheme 1-C).

We hypothesized that the stereoselective introduction of the aryl moiety could be accomplished through a C(sp³) arylation of the *in situ* generated axial imine intermediate (A, Figure 2) as an L,X-type transient directing groups, to direct the palladium (II) catalyst selectively to the α-anomeric C(sp³)–H bond through a bidentate coordinating mechanism (intermediates B and C). Before testing the coupling experimentally, a computational study of the key CMD step was performed at the M06//6-311++G(d,p)/SDD level of theory in order to evaluate the free energy demand. It was found that when the axial imine pyridol directing group was used (R = Ac), the CMD transition state (CMD-TS) that gives rise to the α-configuration necessitates a barrier of 19.71 kcal/mol (Figure 2), which is comparable to our previous reported study with the use of a classical directing group strategy.^{5d}




Figure 2. Proposed mechanism and DFT calculations

We anticipated at this stage that the final C-aryl glycosides bearing a free amine will be extremely polar and tricky to purify, so to ease their isolation and analysis, we envisioned transforming the released free amine products without purification into azides. This later could then undergo a CuAAC process to increase the molecular complexity and prepare a variety of complex substituted C3-triazolo C-glycosides. This approach is conceptually attractive in terms of diversifying the C-aryl glycoside pharmacophore, aiming to identify novel scaffolds of biological interest.

To initiate this study, we selected *O*-protected 3-amino sugars **1a-c** and 4-iodotoluene **2a** as models. To facilitate isolation and purification of the desired C3-azido C-glycosides and reduce the generation of wastes, we decided to examine both transformations in a pot manner as depicted in Table 1: (i) the Pd-catalyzed C(sp³) activation directed by the *in situ* generated transient imine group and (ii) the Cu-catalyzed azidation of the

released free amine. Representative results are summarized in Table 1.

Table 1 Survey of the conditions for the anomeric arylation of **1a-c** with **2a**

4-Iodotoluene **2a** (3 equiv)

Pd(OAc)₂ (10 mol%), TDG (50 mol%), L (50 mol%), [Ag] (2 equiv), H₂O (10 equiv), HFIP/AcOH (19:1) 0.1M, 16h, 90°C

CuSO₄·H₂O (xx mol%), K₂CO₃ (xx equiv), MeOH, RT, 4 h

1a (R = Bn), 1b (R = Ac), 1c (R = benzylidene) (1 equiv)

1. C-H arylation; 2. azidation

Entry	TDG	Ligand	[Pd]	[Ox]	Yield (%) ^b
1	TDG1	L1	Pd(OAc) ₂	AgTFA	27
2	TDG2	L1	Pd(OAc) ₂	AgTFA	0
3	TDG3	L1	Pd(OAc) ₂	AgTFA	23
4	TDG4	L1	Pd(OAc) ₂	AgTFA	32
5	TDG5	L1	Pd(OAc) ₂	AgTFA	0
6	TDG4	L2	Pd(OAc) ₂	AgTFA	34
7	TDG4	L3	Pd(OAc) ₂	AgTFA	41
8	TDG4	L4	Pd(OAc) ₂	AgTFA	43
9	TDG4	L5	Pd(OAc) ₂	AgTFA	42
10	TDG4	L6	Pd(OAc) ₂	AgTFA	37
11	TDG4	L7	Pd(OAc) ₂	AgTFA	51
12	TDG4	L8	Pd(OAc) ₂	AgTFA	40
13	TDG4	-	Pd(OAc) ₂	AgTFA	27
14	TDG4	L7	Pd(OAc) ₂	CuTFA	28
15	TDG4	L7	Pd(OAc) ₂	AgOAc	0
16	TDG4	L7	Pd(OAc) ₂	Ag ₂ O	0
17	TDG4	L7	Pd(OAc) ₂	AgOMe	0
18	TDG4	L7	PdCl ₂	AgTFA	20
19	TDG4	L7	Pd(OAc) ₂ / Pd(TFA)	AgTFA	42
20	TDG4	L7	Pd(OAc) ₂	AgTFA	51 ^c

TDGs

Ligands

Reactions were conducted with substrate **1a-c** (0.4 mmol), **2a** (1.2 mmol), Pd. cat. (10 mol%), TDG (50 mol%), L (50 mol%), Ag-source (2 equiv), H₂O (10 equiv) in a mixture of HFIP:AcOH (19:1) [0.1 M] at 90 °C for 16 h. ^b Yield of isolated product **3a**. ^c 30 min heating under microwave irradiation.


In preliminary experiments, the C(sp³)-arylation was examined first by using Pd(OAc)₂ as a catalyst (10 mol%), TDG1 (50 mol%), L1 (50 mol%) as a ligand, AgTFA (2 equiv) and H₂O (10 equiv) in a mixture of HFIP/AcOH (19:1 ratio, 0.1 M) at 110 °C for 16 h. The reaction mixture was monitored by LCMS and once the disappearance of the starting material was observed and the formation of the 3-amino C-tolyl glycoside intermediate was detected, the diazo-transfer reagent imidazole-1-sulfonyl azide⁷ (3 equiv), CuSO₄·H₂O (10 mol%)

and K_2CO_3 (3 equiv), was added to the mixture which was stirred two additional hours at RT. Under these conditions, no desired product was detected by LCMS analysis when both starting substrates **1b** and **1c** in which the hydroxyl functions were protected by acetyl or benzylidene groups, were used (Table 1). However, the use of *O*-benzylated sugar **1a** led to the formation of the desired C3-azido *C*-aryl glycoside in a modest 27% yield for the two steps (entry 1). X-ray analysis clearly showed the α -configuration of the aryl group and the 1C_4 conformation of the sugar nucleus, demonstrating the high diastereoselectivity of this reaction. This first set of results clearly demonstrated that the choice of the protecting group of the sugar moiety plays a pivotal role in this reaction.

It is worth noting that all our attempts to isolate the 3-amino *C*-tolyl glycoside intermediate [**3a-NH₂**] were not an easy task and revealed that this aryl *C*-glycoside displays unexpected photophysical properties. Precisely, no any absorbance at ~240 nm and at ~360 nm wavelength was detected despite the presence of three aromatic nucleus (SI). This issue rends the monitoring of the reaction by LCMS or TLC much more complicate, thus prompting us to continue our study by using the initial approach involving the *in situ* transformation of the amino group to an azide.

Knowing that the azidation step works quantitatively, we screened several parameters (Pd-catalyst, TDG, ligand, base, solvent, temperature, and reaction time, Table 1) to increase the yield of the first steps. The screening of various transient directing groups revealed that TDG3 and TDG4 display the same reactivity as TDG1 (23% and 32% yields, respectively) while quinoline benzaldehyde TDG2 or TDG5 were completely unreactive. A characteristic aspect of the developed method lies in the pyridone ligands used for this coupling. 5-Nitro-pyridone L2 was firstly examined, resulting in a similar yield than with 5-CF₃ pyridone L1 (34% vs 32% yields). Substitution of the C3 position of the pyridine with electron-withdrawing groups further improved the yield up to 51% (L3–L7), except for L6, which produced **3a** in a yield similar to L1. Not surprisingly, only a 25% yield of **3a** was obtained when the reaction was performed without ligand. Various other parameters were examined (Pd-catalyst, source of Ag, other oxidants, reaction temperature) but without success to improve the yield. Finally, we found that the arylation/azidation occurred smoothly with 51% yield when using Pd(OAc)₂ as a catalyst (10 mol%), TDG4 (50 mol%), L7 as the ligand (50 mol%), AgTFA (2 equiv), H₂O (10 equiv) as an additive in a mixture of HFIP:AcOH (19:1) at 90 °C for 16 h (entry 11). Interestingly, the reaction time was reduced to only 30 min when the coupling was performed under microwave irradiation (entry 20). Of note, the use of Pd-catalyst and TDG was necessary to achieve this transformation since no reaction took place when the coupling was conducted in their absence.

Table 2 Scope of the coupling of **1a** with various aryl iodides **2** followed by the azidation^a




^a Reactions were performed in a flame dried re-sealable tube using sugar **1a** (0.4 mmol), ArI (1.2 mmol), Pd(OAc)₂ (10 mol%), TDG (50 mol%), L (50 mol%), AgTFA (2 equiv), H₂O (10 equiv) in a mixture of HFIP:AcOH (19:1) [0.1 M] at 90 °C for 16 h. ^a Yield of isolated product.

Motivated by these results, we next explored the scope of this new anomeric arylation procedure of **1a** with aryl iodides. At first, we were pleased to see that various aryl iodides bearing diverse functions such as -OMe, -Me, -SMe, -Cl, -OCF₃, and -alkyl chain (butyl or pentyl) reacted smoothly with **1a** to afford the desired aryl *C*-glycosides **3a-w** in moderate to good yields and exclusive α -anomeric stereoselectivity (Table 2). Surprisingly, the presence of an *ortho* substituent at the aromatic ring of the coupling partner did not affect the reaction process, as compound **3c** was obtained in satisfactory yield. Moreover, bulky polycyclic aromatic partners such as fluorene and phenanthrene were also tolerated under these conditions (compounds **3n-p**). Finally, the synthetic utility of this methodology was demonstrated by the synthesis of the *C*-glycoside **3q** as an analog of the dapagliflozin drug (FORXIGAs) used to treat type 2 diabetes (Figure 1). Although

the yields remained modest due to the unreacted anomeric C–H bond of sugars of type **1a**, this constitutes the first example of intermolecular $C(sp^3)$ –H functionalization using a transient directing group.

It is worth noting that the role of protecting groups turned out to be crucial for this reaction particularly the primary alcohol at C6 position of the sugar. Low yields were obtained when acetyl and naphthyl groups were used instead to benzyl one (compounds **3r** and **3s**), however the pivalyl group seems to be better (compound **3t**, 28% yield). Furthermore, switching from benzyl- to methoxyl- protecting groups led to the desired compound **3u** in only a 15% yield.

Another important application of this method is the orthogonal preparation of a series of complex 3-triazolo-aryl *C*-glycosides through the simple copper-catalyzed azide-alkyne cycloaddition (CuAAC reaction).⁸ Different motifs were introduced at the azide function such as the trifluorinated benzene **4a** which can be considered as a direct analog of GB1107 (Figure 1).



Scheme 2. Reactions were performed in a flame dried re-sealable tube using **3a** (0.4 mmol), alkyne (0.4 mmol), $\text{CuSO}_4 \cdot \text{H}_2\text{O}$ (0.04 mmol, 10 mol%), ascorbic acid (40 mol%) in *t*-BuOH: H_2O (2:1) 0.1M at RT. ^a Yield of isolated product.

In addition, other triazolo aryl *C*-glycosides bearing biologically valuable functions such as a fluorescein glycoconjugate **4b**, as well as the uridine- glycoconjugate **4c** were synthesized in 59% and 65% yields, respectively.

In conclusion, we report the first example of intermolecular anomeric $C(sp^3)$ –H functionalization using a transient directing group. This approach obviates the need for installation and removal of the directing group and furnishes in one pot manner a diverse collection of synthetically valuable 3-azido aryl *C*-glycosides with excellent α -selectivity. Moreover, the azide function allowed us to increase the structural complexity of this series of glycosides through a CuAAC reaction. Notably, the facile nature of this reaction has successfully been applied to the synthesis of various substituted triazolo aryl *C*-glycosides.

Acknowledgements

Authors acknowledge support of this project by CNRS, Universite Paris-Saclay, ANR SeIFSuCHi (ANR-18-CE07-0012).

Notes and references

- (a) J. Štambaský, M. Hocek, P. Kočovský, *Chem. Rev.* 2009, **109**, 6729 (b) E. De Clercq, *J. Med. Chem.* **2016**, *59*, 2301 (c) Y. Yang, B. Yu, *Chem. Rev.* 2017, **117**, 12281; (d) K. Kitamura, Y. Ando, T. Matsumoto, K. Suzuki, *Chem. Rev.* 2018, **118**, 1495 (e) É. Bokor, S. Kun, D. Goyard, M. Tóth, J.-P. Praly, S. Vidal, L. Somsák *Chem. Rev.* 2017, **117**, 3, 1687 (f) C.-F. Liu *Molecules* 2022, **27**, 7439.
- (a) R. Duval, C. Duplais, *Nat. Prod. Rep.*, 2017, **34**, 161 (b) Jaramillo, C.; Knapp, S. *Synthesis*, 1994, *1*, 1. (c) K. Kitamura, U. Ando, T. Matsumoto, K. Suzuki, *Angew. Chem. Int. Ed.*, 2014, **53**, 1258.
- T. Cañeque, F. Gomes, T. Thi Mai, G. Maestri, M. Malacria, R. Rodriguez, *Nature Chem.* 2015, **7**, 744 (b) Y. Chang, L. Xing, Q. Sun, S. Liang, T. Liu, X. Zhang, T. Zhu, B. A., Pfeifer, Q. Che, G. Zhang, *J. Nat. Prod.* 2020, **83**, 2749 (c) G. D. A. Martin, L.T. Tan, P. R. Jensen, R. E. Dimayuga, C. R. Fairchild, C. Raventos-Suarez, W. Fenical, *J. Nat. Prod.* 2007, **70**, 1406. (d) H. Huang, T. Yang, X. Ren, J. Liu, Y. Song, A. Sun, J. Ma, B. Wang, Y. Zhang, C. Huang, *J. Nat. Prod.* 2012, **75**, 202. Y. Song, G. Liu, J. Li, H. Huang, X. Zhang, H. Zhang, J. Ju, *Mar. Drugs* 2015, **13**, 1304. (e) K. Toshima, H. Ouchi, Y. Okazaki, T. Kano, M. Moriguchi, A. Asai, S. Matsumura, *Angew. Chem., Int. Ed.*, 1997, **36**, 2748 (f) K. Toshima, Y. Maeda, H. Ouchi, A. Asaib, S. Matsumura, *Bioorg. Med. Chem. Lett.* 2000, **10**, 2163. (g) C. Liu, W. Wang, B. J. A. Elisworth, A. REGUEIRO-REN, WO/2021/242982. (h) C. Liu, P. Rao Jalagam, J. Feng, W. Wang, T. Raja, M. Reddy Sura, R. K. V. L. P. Manepalli, B. Reddy Aliphedi, S. Medavarapu, S. K. Nair, V. Muthalagu, R. Natesan, A. Gupta, B. Beno, M. Panda, K. Ghosh, J. K. Shukla, H. Sale, P. Haldar, N. Kalidindi, D. Shah, D. Patel, A. Mathur, B. A. Ellsworth, D. Cheng, A. Regueiro-Ren. *J. Med. Chem.* 2022, **65**, 11084.
- J. Ghouilem, M. de Robichon, F. Le Bideau, A. Ferry, S. Messaoudi, *Chem. Eur. J.* 2021, **27**, 491.
- (a) N. Probst, G. Grelier, N. Ghermani, V. Gandon, M. Alami, S. Messaoudi, *Org. Lett.* 2017, **19**, 5038 (b) N. Probst, G. Grelier, S. Dahaoui, M. Alami, V. Gandon and S. Messaoudi, *ACS Catal.*, 2018, **8**, 7781. (c) J. Ghouilem, R. Franco, P. Retailleau, M. Alami, V. Gandon, S. Messaoudi, *Chem. Commun.* 2020, **56**, 7175. (d) J. Ghouilem, C. Tran, N. Grimblat, P. Retailleau, M. Alami, V. Gandon, S. Messaoudi, *ACS Catal.* 2021, **11**, 1818, (e) J. Wu, A. Kopp, L. Ackermann, *Angew. Chem. Int. Ed.* 2022, **61**, e202114993.
- For recent reviews, see: J. I. Higham, J. A. Bull, *Org. Biomol. Chem.*, 2020, **18**, 7291, (b) M. I. Lapuh, S. Mazeh, T. Besset, *ACS Catal.* 2020, **10**, 12898, (c) C. Jacob, B. U. W. Maes, G. Evano, *Chem. Eur. J.* 2021, **27**, 13899, (d) P. Gandeepan, L. Ackermann, *Chem.*, **4**, 199.
- (a) S. P. Green, K. M. Wheelhouse, A. D. Payne, J. P. Hallett, P. W. Miller, J. A. Bull, *Org. Process Res. Dev.* 2020, **24**, 67. (b) A. Vasella, C. Witzig, J.-L. Chiara, M. Martin-Lomas, *Helv. Chim. Acta* 1991, **74**, 2073 (c) P. B. Alper, S.-C. Hung, C.-H. Wong, *Tetrahedron Lett.* 1996, **37**, 6029 (d) P. T. Nyffeler, C.-H. Liang, K. M. Koeller, C.-H. Wong, *J. Am. Chem. Soc.* 2002, **124**, 10773.
- (a) N. Nebra, J. García-Álvarez *Molecules* 2020, **25**, 2015, (b) E. Haldon, M. C. Nicasio, P. J. Pérez, *Org. Biomol. Chem.* 2015, **13**, 9528-9550 (c) D. H. Hilko, L. F. Bornaghi, S.-A. Poulsen, *J. Org. Chem.* 2018, **83**, 11944; (d) G. G. Crisp, B. L. Flynn, *J. Org. Chem.* 1993, **58**, 6614. (e) K. Peterson, R. Kumar, O. Stenström, P. Verma, P. R. Verma, M. Hakansson, B. Kahl-Knutsson, F. Zetterberg, H. Leffler, M. Akke, D. T. Logan, U. J. Nilsson. *J. Med. Chem.* 2018, **61**, 3, 1164.