



HAL
open science

Perspective: Plant-Based Meat Alternatives Can Help Facilitate and Maintain a Lower Animal to Plant Protein Intake Ratio

Mark Messina, Alison Duncan, Andrea Glenn, François Mariotti

► **To cite this version:**

Mark Messina, Alison Duncan, Andrea Glenn, François Mariotti. Perspective: Plant-Based Meat Alternatives Can Help Facilitate and Maintain a Lower Animal to Plant Protein Intake Ratio. *Advances in Nutrition*, 2023, 14 (3), pp. 392-405. 10.1016/j.advnut.2023.03.003 . hal-04319571

HAL Id: hal-04319571

<https://hal.science/hal-04319571v1>

Submitted on 3 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Title: Plant-based meat alternatives can help facilitate and maintain a lower animal to plant protein intake ratio

Mark Messina¹

Alison M. Duncan²

Andrea J. Glenn³⁻⁵

Francois Mariotti⁶

¹Soy Nutrition Institute Global, 1747 Pennsylvania Ave., NW, Suite 1000, Washington, D.C. 20006

²Department of Human Health and Nutritional Sciences, University of Guelph, Guelph, Ontario, Canada N1G 2W1

³Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA

⁴Department of Nutritional Sciences, University of Toronto, Toronto, ON, Canada

⁵Toronto 3D Knowledge Synthesis and Clinical Trials Unit, Clinical Nutrition and Risk Factor Modification Centre, St. Michael's Hospital, Toronto, ON, Canada

⁶Université Paris-Saclay, AgroParisTech, INRAE, UMR PNCA, 91120, Palaiseau, France

Corresponding author: Mark Messina, 26 Spadina Parkway, Pittsfield, MA 01201; Ph: 1-360-531-1405; email: mark.messina@sniglobal.org

Word count: 6578

Figures: 0

Tables: 3

Running title: Plant-based meat alternatives and the protein transition

Source of financial support: None.

Conflicts of interest: MM is employed by the Soy Nutrition Institute Global, an organization that receives funding from the United Soybean Board and industry members who are involved in the manufacture and/or sale of soyfoods and/or soybean components. AD is a scientific advisory to the Soy Nutrition Institute Global. AJG has received an honorarium from the Soy Nutrition Institute and Academy of Nutrition and Dietetics. FM has received research grants for PhD fellowships under his direction by AgroParisTech and INRAE from Terres Univia and Ecotone Foundation, under the aegis of the Foundation de France. Terres Univia is the professional association for oilseeds and protein-rich plants in France, which are mainly used as livestock feeds. The Ecotone Foundation for biodiversity is founded by the Ecotone Group that sells mainly organic and plant-based foods.

Abstract

The health and environmental advantages of plant-predominant diets will likely lead to increasing numbers of consumers reducing their reliance on animal products. Consequently, health organizations and professionals will need to provide guidance on how best to make this change. In many developed countries, nearly twice as much protein is derived from animal versus plant sources. Potential benefits could result from consuming a higher share of plant protein. Advice to consume equal amounts from each source is more likely to be embraced than advice to eschew all or most animal products. However, much of the plant protein currently consumed comes from refined grains, which is unlikely to provide the benefits associated with plant-predominant diets. In contrast, legumes provide ample amounts of protein as well other components such as fiber, resistant starch, and polyphenolics, which are collectively thought to exert health benefits. But despite their many accolades and endorsement by the nutrition community, legumes make a negligible contribution to global protein intake, especially in developed countries. Furthermore, evidence suggests the consumption of cooked legumes will not substantially increase over the next several decades. We argue here that plant-based meat alternatives (PBMA) made from legumes are a viable alternative, or a complement, to consuming legumes in the traditional manner. These products may be accepted by meat eaters because they can emulate the orosensory properties and functionality of the foods they are intended to replace. PBMA can be both transition foods and maintenance foods in that they can facilitate the transition to a plant-predominant diet and make it easier to maintain that diet by satiating cravings. PBMA also have a distinct advantage of being able to be fortified with shortfall nutrients in plant-predominant diets. Whether existing PBMA provide similar health benefits as whole legumes, or can be formulated to do so, remains to be established.

Key words: Plant-based meat alternatives, animal protein, plant protein, legumes, sustainability, nutrient fortification, shortfall nutrients

Introduction

For the past several decades health organizations have recommended increasing the intake of whole plant foods (1). These recommendations include consuming more fruits and vegetables (2, 3), whole grains (4) and fiber-rich foods (5). To varying degrees, following this dietary advice is associated with reduced risks of cancer, cardiovascular disease (CVD), diabetes and overall mortality (6-9). More recently, the health benefits of replacing animal protein with plant protein have been emphasized (10-13). The environmental advantages of this exchange are also highlighted (14-16).

In developed countries, considerably more protein is consumed from animal sources than from plants. For example, National Health and Nutrition Examination Survey (NHANES) (2015-2018) data indicate the US animal to plant protein intake ratio is approximately 2.1:1 (54.8 g/d animal protein, 25.8 g/d plant protein) (17). Similar ratios have been reported in many developed countries (18, 19).

Although plant protein composition (i.e. amino acids) may play a role (13, 20), the health benefits associated with consuming diets higher in plant protein likely rather result, at least partly, from the non-protein components of whole food sources of plant protein. It is therefore noteworthy that in the US, 46% of the plant protein intake is from refined grains (21). Similarly in France, one-third of dietary protein intake is from plant sources and of that, 55% is from refined grains (18). As noted by Perraud et al. (22) and Mariotti and Huneau (23), protein intake profiles largely define diets since protein food sources contribute other nutrients that tend to cluster together as a “protein package”, e.g. red meat tends to contribute saturated fatty acids and bioavailable iron, and foods high in plant protein tend to contribute fiber and polyunsaturated fatty acids.

The optimal dietary intake ratio of animal to plant protein has not been established and may differ according to food availability. The EAT–Lancet Commission recommends obtaining *most* protein from plant sources (14), whereas Canada’s Food Guide states “Choose protein foods that come from plants more often” when referring to the “protein foods” (24), and the Health Council of the Netherlands recommends a 1:1 animal to plant protein ratio (25). The latter two recommendations align with efforts to reduce, but not eliminate, the dietary intake of animal products, and especially meat (26, 27). These recommendations can still result in a marked reduction in diet-related environmental pressures, such as reducing greenhouse gas emissions, and land and water use (22) and are more likely to succeed because relatively few people will ever completely eliminate animal protein from their diet (28-30), whereas many may be willing to substantially reduce their intake (31)(32).

Effective reduction in meat consumption (flexitarianism) or intention to do so (pro-flexitarianism) is largely explained by current attitudinal changes related to the impact of meat intake on human health, climate change and animal welfare (33). However, meat has an important culinary and cultural role in most societies (34, 35) and is viewed as an essential part of a meal in many cultures (36, 37). Consequently, even complying with the Dutch recommendation to consume similar amounts of protein from animal and plant sources will be a challenge for populations in developed countries, despite there being many readily available plant sources of protein.

The premise of this paper is that the new generation of plant-based meat alternatives (PBMA) can play a pivotal role in this protein intake transition from animal to plant protein sources. Not only can PBMA aid in the transition to a diet higher in plant protein, they can also help to maintain that diet once the transition has been made. The role of PBMA as foods facilitating the transition from an animal-based to a more plant-predominant diet is not a new concept (38-40). Less appreciated is the utility of PBMA in maintaining that newly transformed plant-predominant diet over the long term (41). Maintenance of a plant-predominant diet is an important consideration because a US study found that 10% of adults (≥ 17 years old) are former vegetarians/vegans whereas only 2% are currently vegetarian or vegan (42). Thus, ensuring continued adherence to a plant-predominant diet may be a seminal advantage of PBMA. Newly developed PBMA are designed to provide the consumer with the same orosensory properties as and functionality of the meat products they are intended to replace. For these reasons, these products are more likely to be embraced by populations in developed countries where meat has come to play a central role in the culture (43).

Given the potential role of PBMA in the transition to and maintenance of a diet that contains more plant protein, it is important to consider their nutritional and health attributes in comparison to both the meat they replace and the whole plant foods (legumes, grains, nuts, seeds) for which they serve as an alternative (16, 44, 45). Globally, dietary guidelines provide relatively little counsel on the use of PBMA (46). An exception is Canada's Food Guide which notes in their guidance on how to eat more protein foods that come from plants, that many simulated meat products are highly processed and can add excess sodium and saturated fat to the diet, and therefore it is important to use food labels to make a healthy choice (47).

Thus, the ramifications of reducing meat consumption on nutrient intake must be considered since meat contributes significantly to nutrient intake globally (48), including in Europe (49) and the US (50). However, there is limited evidence of the effect of reducing meat intake on nutrient adequacy in the context of the overall diet. That is, the impact on nutrient status when meat is replaced by other sources of calories. Recently, Dussiot et al. (51) ran gradual meat intake reduction scenarios in successive 10% steps for diets that were both nutritionally adequate and as close as possible to a reference healthy dietary pattern. They found that some food groups were able to replace the contribution of meat to nutrient adequacy while also leading to overall healthier dietary patterns (i.e., whole grains, fruit, vegetables, seafood). However, these healthy dietary patterns require profound reorganization of the diet.

Vieux et al. (52) estimated that approximately half of total protein intake by French adults must be animal-based to meet non-protein nutrient-based recommendations, which would be consistent with the 1:1 dietary animal:plant intake ratio promoted by the Health Council of the Netherlands. However, the optimization model and the research strategy that the authors used have been strongly criticized, and the results appeared to be explained by a marked over-restriction of the possible solutions.(53) Also, this estimate excluded the use of dietary supplements or fortified foods and did not consider the types of more novel foods that may be consumed when meat intake is consciously reduced. Furthermore, Fouillet et al. (54) concluded that among French adults, only when plant protein accounts for more than 80% of total protein, are nutrient fortification/supplementation and/or new foods required to meet nutrient

requirements. In any case, all modelled diets greatly departed from current observed diets, which again indicates the major change needed to most diets.

Finally, it is acknowledged that in the near term the dietary contribution of PBMA will be a consideration primarily relevant only to developed countries due to their high meat intake and ability to afford the relatively high cost of many PBMA relative to more conventional meat alternatives such as legumes (55). However, projections that the global demand for animal products will rise by 60–70% by 2050 stem primarily from the anticipated increased consumption in low- and middle-income countries (56). As the price of PBMA decreases, as is expected, and consumer awareness of the link between diet and climate change and the relatively low environmental footprint of PBMA increases, these products may be embraced in both developed and developing countries (57-59).

The general topics covered in this manuscript include background information on PBMA; current animal to plant protein intake ratio; protein and chronic disease risk; protein quality and global protein needs; using legumes and PBMA to lower the animal to plant protein ratio and current understanding of the health effects of PBMA.

Historical perspective on plant-based meat alternatives

Meat substitutes have existed in the Western world for more than a century and considerably longer in some Asian countries. Notable in this regard is the creation of Nuttose in 1896 by Dr. John Harvey Kellogg, which was a canned product made primarily from peanuts (60). In 1911, in France, cold cuts made from soy became available and in 1922, the first soy-based meat alternative was developed by Madison Foods in Tennessee (61). However, in part because of improvements in technology and increased consumer demand, within the past two decades the PBMA industry has been especially active and innovative (62).

Traditionally, adherents of plant-predominant diets met their protein needs by consuming a mix of grains and legumes, a common culinary practice among cultures throughout the world. These food groups have indispensable amino acid (IAA) profiles that complement one another such that the combination produces a protein with an IAA pattern that more closely matches biological requirements (63-66). Traditional soyfoods such as tofu and tempeh, which have been consumed in many parts of Asia for centuries, have been embraced by flexitarians (67) and vegetarians over the past several decades because of the abundant amounts of high-quality protein they provide (68, 69). There are also “veggie” burgers, such as a black bean burger, made using beans often in combination with grains and vegetables that may or may not be vegan (70). This type of product is especially attractive to those wanting a food that can be consumed in the same manner as a meat patty. The wheat (gluten)-derived product seitan also fills a role as a meat substitute (71).

However, none of the aforementioned products truly emulate the orosensory properties of meat and for this reason, their entrance and acceptance by the mainstream consumer has been somewhat limited (72, 73). Research shows that for PBMA to successfully replace meat, they need to emulate the taste, texture, visual appearance, and cooking method of meat (74, 75). The new generation of PBMA, which are typically comprised of protein extracted from legumes such

as pea, soy and/or wheat gluten and require new technology for their production, meets these criteria (45, 76, 77). These products, while embraced by vegetarians, are designed to appeal to a much broader demographic – basically anyone wanting to reduce meat intake (78). In fact, US household data indicate that over a recent two-year period, 86% of purchasers of PBMA were consumers of beef (79).

Several research groups have published comprehensive reviews on the nutrient content of a wide range of PBMA (76, 80-83) so this information will not be discussed in detail here. The nutrient content of PBMA varies widely and differs according to the primary sources of protein, secondary ingredients (e.g., type of fat added), and the addition of micronutrients. It is also recognized that meat alternatives not made from legumes, but from algae, fungi, and insects, are gaining relevance, and that animal protein is being produced by in vitro methods and precision fermentation (84). However, the focus of this manuscript is on legume-based PBMA. That is, products such as burgers that are made primarily from extracts of legume protein. Surveys to date indicate consumers are more accepting of PBMA than other types of alternatives (85, 86). It is notable that a recent analysis of the French diet found that plant-based substitutes that include legumes appear more nutritionally adequate to substitute for animal products than other alternatives (87).

Animal to plant protein intake ratios

As highlighted above, in developed countries, approximately twice as much protein is consumed from animal sources compared to plant sources (18, 19, 21, 88) whereas globally, this ratio is reversed – in many developing countries, much less than one-third of dietary protein is derived from animal sources (89, 90). In Bangladesh for example, 80% of the protein consumed is from plant sources (90). Interestingly, according to crude estimates based on disappearance data, animal and plant sources provided similar amounts of protein in the US in 1909 (91). However, over the past century, the animal to plant protein intake ratio has steadily increased as a result of an increase in animal protein intake and a decrease in plant protein intake (21, 91).

Differences between animal and plant protein

Chronic disease risk

As noted previously, most evidence suggests that the health benefits of plant-predominant diets likely stem from the non-protein components rather than the protein per se (92, 93), and likely also from the parallel reduction in animal product intake. To this point, eliminating red and processed meat from the diet in and of itself may reduce risk of coronary artery disease (CAD) (94), although there is disagreement on this point (95). In any event, that which replaces meat in the diet likely determines the extent to which risk is reduced. For example, data from the combined analysis of the Nurses' Health Study (n=84,628 women) and the Health Professionals Follow-up Study (n= 42,908 men) showed that replacing 5% of energy intake from saturated fats with equivalent energy intake from polyunsaturated fat, monounsaturated fat, or carbohydrates from whole grains was associated with a 25%, 15%, and 9% lower risk of CAD, respectively, whereas replacement with carbohydrates from refined starches/added sugars was not significantly associated with risk reduction (96, 97).

However, it is also possible that differences between animal and plant proteins may affect chronic disease risk as a result of the differences in their amino acid profiles (13, 98-100). For example, there is mounting evidence that the concentration of certain amino acids that is higher in animal protein than plant protein, such as the branched-chain amino acids and methionine, may exert adverse effects on metabolism, whereas the concentration of certain amino acids that is higher in plant protein than animal protein, such as glutamine, cysteine and arginine, may exert beneficial effects (20, 100, 101). Thus, diets differing in the amounts of animal and plant protein content may exert differing metabolic effects due to the differences in amino acid profiles as well as the differences in nutrients and non-nutrients found in their respective protein packages. For example, there are phytochemicals such as isoflavones and glucosinolates, which theoretically could exert both favorable and unfavorable effects, and phytate, which inhibits mineral absorption but is also an anti-oxidant (102, 103). In general, over the past several years, systematic reviews and meta-analyses of observational data have generally found that animal protein and plant protein are associated with an increased and decreased risk, respectively, of type-2 diabetes, cardiovascular mortality and all-cause mortality (12, 104, 105).

Finally, it would be remiss not to mention that there are short-fall nutrients in vegetarian and especially, vegan diets. Notable in this regard are vitamin B12, vitamin D, zinc, choline, iodine and calcium (106). However, deficiencies are much less likely to occur when adhering to a plant-predominant diet as is discussed here, than a vegan diet. Furthermore, deficiencies are not necessarily more likely to occur because a serving of legumes is replaced by a PBMA. In fact, as noted later, because PBMA's have the potential to be fortified, they may be less likely to occur.

Protein quality

Differences between the amino acid profiles of animal and plant protein are associated with differences in protein quality. Animal proteins tend to be higher in IAA (99, 107) and as such have higher protein quality scores as determined by the protein digestibility corrected amino acid score (PDCAAS) and the digestible indispensable amino acid score (DIAAS) (108-110). However, in populations consuming a mix of animal and plant protein these differences are not likely to be clinically relevant. In fact, it is not conclusive whether the lower quality of some plant proteins necessitates that adherents of plant-predominant diets consume higher amounts of protein to meet their biological requirements for nitrogen and amino acids (111). To this point, there is no vegan US protein recommended dietary allowance (RDA) as there is for iron (1.8 vs 1.0 mg). On the other hand, some (111-114), but not all (115), experts have called for vegans to consume more protein to account for the lower digestibility of protein from plants (116). The Dutch Ministry of Health recommends that vegans consume 30% more protein than omnivores to compensate for the lower quality of plant protein (117); however, this recommendation greatly exceeds recommendations made by most experts (29).

Global protein needs

The world population is predicted to reach close to 10 billion people three decades from now (118). Determining how best to meet the global protein and caloric needs of the 2050 population is a matter of some urgency. Protein is by far the more important consideration because as was noted in a recent report from the Stockholm Resilience Centre, "Meeting the demand for protein,

within environmental limits, is one of the biggest challenges for the global food system in the 21st century” (119).

Precisely how much protein will be needed is unclear, but in an attempt to answer this question, Irish researchers proposed five scenarios (120). At the high end of the spectrum, it was estimated that protein production will need to increase by 78% to meet global protein needs based on the premise that the entire population will consume ~100 g/d protein. On the low end, which is based on the premise the population will consume only 50 g/d protein, production could decrease by 13% without compromising protein availability. The high estimate may be the more probable scenario because although the types of protein consumed vary markedly among cultures and societies, if availability allows, populations tend to consume approximately 16% of their calories in the form of protein (121), which is much greater than the approximate 10% of calories needed to meet the RDA (122).

Finally, for an assortment of reasons, including concern over the environment and animal welfare as well as personal health considerations, evidence suggests that over the next 30 years in developed countries, there is likely to be a shift in the direction of a plant-predominant diet (123). Therefore, the question becomes how the world is going to produce the calories and *plant* protein needed by 2050. If the plant protein is to come from non-soy legumes, global production would need to substantially increase to meet demand whereas worldwide soybean production is sufficient to meet that need if some of the soy protein currently used for animal feed is instead used for direct human consumption (124).

Shifting the animal to plant protein intake ratio

Consuming more legumes is an obvious choice for increasing plant protein intake (125, 126). In 2013, the United Nations declared 2016 as the International Year of Pulses (127), and the International Lipid Expert Panel places soy, legumes and nuts at the top (most desirable) of the protein source pyramid for promoting cardiometabolic health (128). Legumes (including beans and other pulses) are beneficial to long-term health (129, 130), good sources of protein and fiber (52), affordable (131), and have a low environmental footprint (132). However, despite their many accolades, pulses are vastly underutilized sources of nutrition in most regions in the world (Table 1) (133). Worldwide, pulses account for only 6% of total protein intake.

In the US, dry bean intake accounts for only about 2% of the total protein intake and 6% of plant protein (21). A recent French study found that even knowledge of the health benefits of pulses did not lead to greater intake of these foods (134). Reuzé et al. (135) recently summarized motives associated with greater legume consumption among French adults. Furthermore, only very recently (2017) did French public health authorities include a specific guideline on legumes in the dietary guidelines (136, 137).

Although the terms legumes, pulses and beans are often used interchangeably, pulses are the edible seed part of the legume plant which include beans, lentils, dry peas, chickpeas and cow peas. Legumes not only include pulses, but soybeans, peanuts, snap beans and snap peas (138). Soybeans and peanuts are referred to as oilseed legumes because of their high fat content. Legume consumption has traditionally been a larger part of the cuisines in countries and regions

such as Mexico (refried kidney beans), India (dhal and pappadums), the Mediterranean (navy bean soup and Greek fava), and the Middle East (falafel and hummus).

As can be seen from Table 2, on a caloric basis, legumes are approximately 29% protein, which is approximately twice the percentage (13%) found in grains, and legumes on average are also higher in fiber than grains (139). To lower the US animal to plant protein intake ratio from approximately 2.1:1 to 1.3:1, one would need to substitute the protein from one serving of cooked beans daily for an equivalent amount of animal protein. This exchange would result in an additional 8.6 g of plant protein being consumed (140). Adding that amount to the 25.8 g/d of plant protein per capita consumed in the US would result in a total of 34.4 g/d (17). Subtracting 8.6 g/d from the 54.8 g/d of animal protein per capita consumed in the US would result in a total of 46.2 g/d. Thus, the dietary animal to plant protein ratio would be 1.3:1 (46.2:34.4). On the other hand, meeting the cooked beans, peas, and lentils intake recommendation of the current US Dietary Guidelines – 3 servings weekly (1.5 cups cooked) – would clearly have much less effect (141). If the approximate 3.7 g/d protein from these plant foods replaced an equal amount of animal protein, the ratio would decrease only from 2.1:1 to about 1.7:1 (51.1:29.5). (Note that the USDA designates one serving of beans as ½ cup, which weighs approximately 90 g, whereas in this manuscript, 100 g is considered to be one serving based on work by Marinangeli et al.(140))

Is the direct consumption of pulses likely to increase?

How likely is it that Americans and other Western populations will consume one serving of cooked pulses daily? Despite calls for greater public health efforts promoting pulse consumption (142, 143), there is little evidence that these efforts, if undertaken, will be successful. In NHANES (2013–2014; n = 6,048), only 4% of participants consumed legumes on both days of the survey (144). Furthermore, that figure was lower than the 5.6% of participants who consumed legumes on both days of the survey in the previous two years. More recently, Tao et al. (145) reported that among the 4,058 participants in NHANES 2017-18, daily legume intake averaged only 0.11 servings, which was not higher than reported in NHANES 2011-12 (n=4313, 0.12 servings/d). This downward trend in legume intake is also evident from data from the Continuing Survey for Individual Intakes (1994–1996) wherein 14% of the US adults consumed dry beans on both days of the survey (146). In contrast, in the 2017 Beans, Lentils, Peas Survey, only 4.9% of participants consumed beans on both days of the survey (144). Even more striking in some ways is that based on grocery purchases, US mean annual per capita expenditure on legumes was only \$4.76 during 2017–2019 (147).

In NHANES, because of their higher fat content, neither soybeans nor peanuts are included in the legume category. As seen in Table 3, one serving (2 tbsp) of peanut butter provides almost as much protein as a serving of pulses (7.27 g vs 8.62 g) although it contains considerably more kcal (~189 vs ~127). Furthermore, according to the Adventist Health Study 2, reported legume intake (minus soy) of male vegans was only about 65 g/d legumes, about ¾ serving, although they also consumed on average 207 g/d soyfoods (148).

Legume intake is similarly low in the UK. In the Oxford component of the European Prospective Investigation into Cancer and Nutrition, when standardized for an intake of 2,000 kcal, regular meat eaters (n= 2,852) reported consuming only 26.7 g/d of pulses/legumes. Vegans (n=269) consumed 68.4 g/d, but that is still less than one serving.(149) However, vegans also consumed

59.6 g/d of vegetarian protein alternatives (e.g., tofu, soymilk, soy burgers, Quorn), much higher than the 4.9 g/d consumed by regular meat eaters.

Barriers to bean consumption in Western countries include beans not being part of the traditional diet, concerns about flatulence/abdominal discomfort, lack of knowledge about preparation/cooking (150), objectionable taste and texture and high carbohydrate content (144), the perception that beans are “poor man’s food” or poor man’s meat,” (151) and concerns about antinutrients. Considerable efforts will be needed to increase intake of whole legumes in the general population, including additional public education campaigns, changing the food environment, and more legume-based dishes being served at schools, hospitals, and restaurants. However, if protein intake from the consumption of cooked legumes served in the traditional manner is unlikely to substantially increase, a viable alternative is to consume PBMA made from legumes. Importantly, and as noted previously, PBMA should not be viewed as a replacement for legumes, but as a complement, that is, it is not one or the other. Both foods can be used as a means of increasing legume and plant protein intake.

Perspectives on PBMA

Introduction

The potential role of meat substitutes in reducing reliance on and changing attitudes toward meat was recently demonstrated by Bianchi et al. (152) These authors found that a 4-week behavioral program in which British study participants were provided plant-based meat substitutes along with recipes for their use and information about the benefits of eating less meat and told of success stories of people who had reduced their meat intake, resulted in a reduction in meat intake and changes in psychosocial constructs consistent with a sustained reduction in meat intake.

Evidence that PBMA can substantially contribute to reducing the animal to plant protein intake ratio is readily apparent. According to Messina et al. (153), the protein content of 5 soy-based patties ranged from 14 to 27 g per patty, considerably more than an average serving of pulses (~8.6 g)(Table 3). Even a conservative estimate indicates that the consumption of only 4 servings (4 patties) of soy-based patties per week or about 75 g protein, if replacing animal protein in the diet, would bring the US animal to plant protein intake ratio down to approximately 1.2:1. Arguably, the higher protein content of the PBMA vs legumes may allow for the former to more effectively facilitate the transition to a diet containing similar amounts of animal and plant protein.

Skepticism about the utility of PBMA

However, the increased popularity of PBMA has led to greater scrutiny of their nutritional attributes and questions about their role in the diet. For example, Toh et al. (154) recently commented that “There is a gap in our understanding of the long-term impacts of dietary patterns that characteristically feature PBMA compared with PBDs [plant-based diets].” The PBDs in this case refers to a balanced consumption of grains, legumes, nuts, seeds, fruits and vegetables. Fardet et al. (155, 156) maintain that as a rule, degrading the food matrix, as occurs

for example in the extraction of protein from beans, adversely impacts the healthfulness of that food, regardless of its nutrient and calorie content. Furthermore, based on associations between ultra-processed food (UPF) intake and adverse health outcomes, Gehring et al. (157) hypothesized that a higher intake of the processed form of plant-based foods might reduce or cancel their potential health benefits.

Concerns about the processing involved in the production of PBMAs

The processed nature of PBMA

s was also highlighted by Macdiarmid (158), when this author concluded that the trend towards consuming more highly processed plant-based convenience foods is a concern regarding both public health and achieving the targets of reducing greenhouse gas emissions. If based on the NOVA classification, it was reported that swapping meat for PBMAs in France would result in an increase in the percentage of energy from UPF from 29% in observed diets to 34% on average and up to 40% according to type of substitution in modelled diets.(87) As an aside, in reference to the comment by Macdiarmid (158), while it may be true that a soy burger is more convenient than having to boil dry beans, it may not be more convenient than eating canned beans or ground beef. Furthermore, research indicates that plant-based burgers have a low environmental footprint (57, 58, 159, 160).

The considerable discussion about the NOVA food classification and the harms of consuming foods classified by NOVA as ultra-processed shines a negative light on the entire new generation of PBMA

s because they are classified by NOVA as UPFs (161, 162), foods whose intake is to be discouraged (163). However, a recent analysis found that of the many common criticisms of UPFs, none of those examined apply to soy-based meat alternatives (or soy-based dairy alternatives), all of which were NOVA-classified as UPFs, more so than they apply to their animal-based counterparts, beef and cow's milk, which are classified as unprocessed/minimally processed foods. The criticisms addressed were 1) hyperpalatability 2) high energy density 3) increased energy intake rate 4) low satiation 5) low cost/high snackability and 6) high glycemic index (milks only) (153). One concern not addressed by the authors of this analysis is that UPFs cause adverse changes in the microbiome (164); however, this does not appear to be the case for PBMAs (165).

Furthermore, many PBMA

s are rated highly by nutrient profiling models other than NOVA (166). For example, Australian researchers recently found the average Health Star rating for 50 plant burgers was 3.9 (5-point scale) versus a rating of 2.9 for a meat burger (167). A French analysis found that plant-based meat and dairy substitutes had a small effect on overall quality of the French diet and heterogeneous impacts on nutrient adequacy and security. Importantly, as mentioned previously, it was found that plant-based substitutes that include legumes appear more nutritionally adequate to substitute animal products than other alternatives such as cereal-based substitutes (87).

Clinical data supportive of PBMAs

Evidence offering the most direct insight regarding the comparison between meat and PBMA

s comes from the clinical trial Study With Appetizing Plantfood-Meat Eating Alternative Trial (SWAP-MEAT) (168). Participants (n=36) consumed about 2½ servings daily of PBMAs or analogous meat products, providing 25% of total calories and 50% of total protein, for 8 weeks each in a randomized, crossover design. Results showed that consumption of PBMAs significantly

decreased circulating levels of trimethylamine oxide and low-density-lipoprotein cholesterol as well as body weight when compared to the meat products (168). There were also no differences in inflammatory markers between the groups (169), nor were adverse effects observed on any of the other endpoints analyzed.

Also of relevance are the results of the SWAP-MEAT Athlete study, which involved 24 athletes (12 recreational runners and 12 resistance trainers) who were randomly assigned to three diets for 4 weeks each in a crossover design: whole foods plant-based, plant-based with PBMA, and an omnivorous diet (170). At study termination, there were no differences in running outcomes for the runners or lifting outcomes for the resistance trainers, suggesting that inclusion of PBMA in a plant-predominant diet may not impact fitness outcomes in athletes compared to a whole foods plant-predominant diet and an omnivorous diet.

Additionally, although not a direct comparison of meat and PBMA, a recent 3-month parallel-arm designed clinical trial compared the effects of a low-carbohydrate vegan diet with a moderate-carbohydrate vegetarian diet in 164 participants with type 2 diabetes (171). To achieve a lower carbohydrate content in the vegan diet, the diet was higher in fat (from canola oil) and protein. Approximately 25% of total caloric intake was derived from protein, nearly all of which came from plant sources and much of that was derived from soy-based meat alternatives. At study termination, both diets caused significant reductions in body weight, glycated hemoglobin, blood pressure and blood lipids, with no adverse effects reported (171).

Legumes compared with legume-based PBMA

As noted previously, PBMA may be regarded as transition foods or gateway foods to facilitate the conversion to a plant-predominant diet. Or, in the words of Alae-Carews et al. (172), PBMA are an important “stepping stone” for dietary change. Implicit in this perspective is that PBMA are more healthful than the meat they replace but perhaps less so than less processed forms of plant protein, such as legumes and whole grains. Several authors have pondered whether PBMA derived from legumes offer similar nutritional benefits or chronic disease reductions as whole legumes (44, 45). Van Vliet et al. (173) went a step further by stating that “The mimicking of animal foods using isolated plant proteins, fats, vitamins, and minerals likely underestimates the true nutritional complexity of whole foods in their natural state, which contain hundreds to thousands of nutrients that impact human health.”

It is true that in addition to providing nutrients and fiber, legumes contain an assortment of phenolic compounds that may contribute to their health effects (174-176). Observational data indicate legume intake is associated with reductions in the risk of colorectal cancer (177) and all-cause mortality (178), and clinical data suggest pulses may help to manage body weight (179) and improve glycemic control (180). However, the extent to which bean phenolics contribute to these associations is not known.

The new generation of PBMA have not been commercially available long enough for their intake to be examined in observational studies, and limited clinical data are available. On the other hand, the primary proteins used in these products, such as soy and to a much lesser extent pea, have been rigorously investigated in the same processed form (e.g., soy protein isolate and soy protein concentrate) as that used in PBMA. For example, soy protein is a high quality protein

(110, 181), lowers blood cholesterol levels (182, 183) and promotes gains in muscle mass and strength to a similar extent as animal proteins, including whey (184). Furthermore, one aspect of PBMA that is cited above by van Vliet et al. (173) as a disadvantage – the use of isolated components – may actually be an advantage in comparison to consuming legumes in their unprocessed (raw) state, as the manufacturing process allows the fortification of shortfall nutrients in both plant-predominant and animal-based diets which, if need be, can be tailored to specific populations.

For example, one can envision that in populations where intake of a particular nutrient, such as iron, is marginal, PBMA could serve as a vehicle for fortification of that nutrient. Industry could be encouraged to fortify their products with nutrients that may be of concern when replacing animal protein with plant protein. Interestingly, in Canada, for a product to be considered a “simulated meat product” the nutritional content must be equal to that of the meat product it is intended to substitute (185). Of relevance to this discussion is a recent study which optimized the recipe for creating a legume-based patty aimed at improving nutritional content while taking account of technological constraints and applying nutritional constraints to limit the risk of overt deficiency in 12 key nutrients. Even in this case, there remained the need to fortify with vitamin B₁₂, zinc and iron (186).

Another possible advantage of PBMA is that components of legumes that may be objectionable to some consumers such as oligosaccharides because they can cause flatulence are greatly reduced when the protein is extracted from the beans (187). The digestibility of the extracted protein may also be greater because of the elimination of compounds that can inhibit protein digestion (108, 116). Additionally, processing can eliminate or reduce compounds traditionally classified as anti-nutrients, such as phytate (188). The nutrient content of PBMA varies widely, but some tend to be relatively high in sodium although they do not meet the US Food and Drug definition of a high-sodium food (≥ 400 mg/serving) (81, 153, 167). Also, some are high in saturated fat (but lower than beef), a result of the attempt to emulate the orosensory properties of meat (153).

The addition of sodium can improve the taste and flavor of plant-based ingredients and mask unpleasant flavors typical in plant-based proteins, such as beany and chalky (189). It is important to note however, that because salt is commonly added to ground beef when cooking to improve texture and taste, there may be little difference in the sodium content between PBMA and beef as they are typically consumed (190). Given the improvements in taste and texture of PBMA made in recent years, it is not unreasonable to speculate that future iterations of PBMA will be able to maintain or improve current orosensory properties while reducing sodium and saturated fat content. In fact, the newest version of one popular soy-based burger contains less saturated fat (8 vs 6 g per serving) than the original version (191).

One concern about PBMA is their potential to be consumed similarly to the traditional way in which hamburgers are consumed – with a refined bun and perhaps fries or chips (159). While this concern is understandable, PBMA are now available in ground form in addition to patties and many other forms. For example, products that emulate steak, chicken and seafood are either already available or soon expected. Thus, the way PBMA can be consumed is much broader than was initially the case.

Hybrid meat products

Hybrid meat products, which refers to the combination of animal and plant protein, may be an especially appealing and efficacious approach to decreasing meat intake as research shows that to create an effective dietary change, new practices should not diverge too much from consumers' previous behavior (192-194). These products preserve the animal flavor while reducing the amount of meat in the formulation. Such products offer convenience and a more characteristic taste to flexitarian consumers (195, 196). The purpose of hybrid meat products is to substitute a portion of the meat with a more sustainable source, while plant-based ingredients in meat products are often added for their functionality, such as extenders, fillers, and binders (84, 197). If hybrid meat products increase in popularity, soy protein ingredients can contribute to their success as manufacturers have considerable experience using these combinations (198, 199). This is also likely the case for other legume proteins.

Conclusion and perspective on the protein intake transition

Evidence indicates that PBMA's represent a convenient, nutritious, and sustainable way to lower the animal to plant protein intake ratio in developed countries. The occasional replacement of meat, such as beef, with a PBMA is unlikely to result in nutrient intake being compromised, especially if the PBMA's are fortified with shortfall micronutrients. A conservative estimate indicates that replacing just 4 servings per week of meat with 4 servings of soy-based PBMA's is sufficient to lower the animal to plant protein intake ratio from about 2.1:1 to 1.2:1, which is a reasonable goal given the glacial pace at which changes in dietary habits typically occur. As new PBMA's are developed, additional research on these products will be needed to help further understand their role in a healthy diet.

Future iterations of PBMA's should reflect a continued emphasis on improving nutrient content while maintaining the orosensory properties that make them effective options for individuals moving away from an animal-based to a more plant-predominant diet. In some sense, the comparison between the health attributes of legumes and PBMA's is irrelevant if in fact, legume intake is unlikely to substantially increase, although continued efforts aimed at increasing intake should be undertaken.

Health professionals need to be well informed about PBMA's so they can competently advise their patients and clients about the role these foods can play in the diet. Both health professionals and consumers need to recognize that the nutrient content of PBMA's varies markedly. Such variation also exists for plant-based milks, for example, some are low in protein and not all are fortified with key nutrients found in cow's milk, such as calcium and vitamin D (200). Recently, Klapp et al. (46) recommended that dietary guidelines "should differentiate between plant-based alternatives that can be consumed frequently and those that should be consumed in moderation or merely for enjoyment."

In a recently published textbook on vegetarian diets for dietitians, intake recommendations for vegans included legumes at least three times daily (201). Although both pulses/dry beans and soybeans are legumes, because of the high fat content of the latter and because of the many forms in which soybeans are consumed, soyfoods can be viewed as a separate category from

pulses. Given their high fat content, peanuts (and peanut butter) can also be viewed similarly to soy (108). The pulses/dry beans and soy/peanut categories can be part of legume-based PBMA. Thus, there are three distinct ways to incorporate legumes into the diet, i.e., dry beans/pulses, peanuts and foods derived from soybeans, and PBMA.

In the interest of diet/nutritional diversity and convenience, consumers may consider choosing foods from each of these three legume categories because doing so may enhance adherence – by increasing variety – to plant-predominant diets, and it may facilitate sufficient nutrient intake. On a per serving basis, many PBMA are considerably higher in calories than pulses (140) and traditional soyfoods such as tofu (Table 3), so when caloric intake is restricted, this difference needs to be appreciated. Meeting the legume intake recommendation by consuming foods from the three legume-based food categories needs to be individualized based on health and personal factors including cultural considerations, economic status and overall lifestyle.

In conclusion, PBMA provide a convenient option for omnivores to transition to a diet that has a lower animal to plant protein ratio. As more such products enter the market, and their cost decreases as is expected, PBMA are likely to become increasingly mainstream. For adherents of plant-predominant diets, well-designed, nutrient-dense formulations of PBMA should be able to provide shortfall nutrients and for many individuals, they will likely enhance compliance to their diet by increasing the variety of protein-rich plant foods available. Finally, whereas a deeper redesign of the diet to include more traditional, minimally processed plant protein sources that have proven nutritional/health benefits should remain a primary goal, PBMA are useful and indeed practically indispensable to facilitate and maintain the transition toward a more plant-predominant diet for a large part of the population and are apt to make continued adherence to a plant-predominant diet easier.

References

1. Rock CL, Thomson C, Gansler T, Gapstur SM, McCullough ML, Patel AV, Andrews KS, Bandera EV, Spees CK, Robien K, et al. American Cancer Society guideline for diet and physical activity for cancer prevention. *CA Cancer J Clin* 2020;70(4):245-271.
2. WHO Fruit and Vegetable Promotion Initiative – report of the meeting, Geneva, 25–27 August 2003. 2003;
3. Deckelbaum RJ, Fisher EA, Winston M, Kumanyika S, Lauer RM, Pi-Sunyer FX, St Jeor S, Schaefer EJ, and Weinstein IB. Summary of a scientific conference on preventive nutrition: pediatrics to geriatrics. *Circulation* 1999;100(4):450-6.
4. Curtain F and Grafenauer S. Historical and global perspective on grains and whole grains within Dietary Guidelines. *Cereal Foods World* 2020;65(3):
5. Slavin JL. Position of the American Dietetic Association: health implications of dietary fiber. *J Am Diet Assoc* 2008;108(10):1716-31.
6. Aune D, Giovannucci E, Boffetta P, Fadnes LT, Keum N, Norat T, Greenwood DC, Riboli E, Vatten LJ, and Tonstad S. Fruit and vegetable intake and the risk of cardiovascular disease, total cancer and all-cause mortality-a systematic review and dose-response meta-analysis of prospective studies. *Int J Epidemiol* 2017;46(3):1029-1056.
7. Gaesser GA. Whole grains, refined grains, and cancer risk: A systematic review of meta-analyses of observational studies. *Nutrients* 2020;12(12):
8. Aune D, Keum N, Giovannucci E, Fadnes LT, Boffetta P, Greenwood DC, Tonstad S, Vatten LJ, Riboli E, and Norat T. Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: systematic review and dose-response meta-analysis of prospective studies. *BMJ* 2016;353(i2716).
9. Veronese N, Solmi M, Caruso MG, Giannelli G, Osella AR, Evangelou E, Maggi S, Fontana L, Stubbs B, and Tzoulaki I. Dietary fiber and health outcomes: an umbrella review of systematic reviews and meta-analyses. *Am J Clin Nutr* 2018;107(3):436-444.
10. Abdelhamid AS, Brown TJ, Brainard JS, Biswas P, Thorpe GC, Moore HJ, Deane KH, AlAbdulghafoor FK, Summerbell CD, Worthington HV, et al. Omega-3 fatty acids for the primary and secondary prevention of cardiovascular disease. *Cochrane Database Syst Rev* 2018;7(CD003177).
11. Al-Shaar L, Satija A, Wang DD, Rimm EB, Smith-Warner SA, Stampfer MJ, Hu FB, and Willett WC. Red meat intake and risk of coronary heart disease among US men: prospective cohort study. *BMJ* 2020;371(m4141).
12. Naghshi S, Sadeghi O, Willett WC, and Esmailzadeh A. Dietary intake of total, animal, and plant proteins and risk of all cause, cardiovascular, and cancer mortality: systematic review and dose-response meta-analysis of prospective cohort studies. *BMJ* 2020;370(m2412).
13. Gardner CD, Hartle JC, Garrett RD, Offringa LC, and Wasserman AS. Maximizing the intersection of human health and the health of the environment with regard to the amount and type of protein produced and consumed in the United States. *Nutr Rev* 2019;77(4):197-215.

14. Willett W, Rockstrom J, Loken B, Springmann M, Lang T, Vermeulen S, Garnett T, Tilman D, DeClerck F, Wood A, et al. Food in the Anthropocene: the EAT-Lancet Commission on healthy diets from sustainable food systems. *Lancet* 2019;393(10170):447-492.
15. Gibbs J and Cappuccio FP. Plant-based dietary patterns for human and planetary health. *Nutrients* 2022;14(8):
16. Nelson ME, Hamm MW, Hu FB, Abrams SA, and Griffin TS. Alignment of healthy dietary patterns and environmental sustainability: A systematic review. *Adv Nutr* 2016;7(6):1005-1025.
17. Hoy MK, Murayi T, and Moshfegh AJ. Diet quality and food intakes among US adults by level of animal protein intake, What We Eat in America, NHANES 2015-2018. *Curr Dev Nutr* 2022;6(5):nzac035.
18. Salome M, de Gavelle E, Dufour A, Dubuisson C, Volatier JL, Fouillet H, Huneau JF, and Mariotti F. Plant-protein diversity is critical to ensuring the nutritional adequacy of diets when replacing animal with plant protein: Observed and modeled diets of French adults (INCA3). *J Nutr* 2020;150(3):536-545.
19. Halkjaer J, Olsen A, Bjerregaard LJ, Deharveng G, Tjonneland A, Welch AA, Crowe FL, Wirfalt E, Hellstrom V, Niravong M, et al. Intake of total, animal and plant proteins, and their food sources in 10 countries in the European Prospective Investigation into Cancer and Nutrition. *Eur J Clin Nutr* 2009;63 Suppl 4(S16-36).
20. Tharrey M, Mariotti F, Mashchak A, Barbillon P, Delattre M, Huneau JF, and Fraser GE. Patterns of amino acid intake are strongly associated with cardiovascular mortality, independently of the sources of protein. *Int J Epidemiol* 2020;49(1):312-321.
21. Shan Z, Rehm CD, Rogers G, Ruan M, Wang DD, Hu FB, Mozaffarian D, Zhang FF, and Bhupathiraju SN. Trends in dietary carbohydrate, protein, and fat intake and diet quality among US adults, 1999-2016. *JAMA* 2019;322(12):1178-1187.
22. Perraud E, Wang J, Salome M, Mariotti F, and Kesse-Guyot E. Dietary protein consumption profiles show contrasting impacts on environmental and health indicators. *Sci Total Environ* 2022;856(159052).
23. Mariotti F and Huneau JF. Plant and animal protein intakes are differentially associated with large clusters of nutrient intake that may explain part of their complex relation with CVD risk. *Adv Nutr* 2016;7(3):559-60.
24. Health Canada [Internet]. Ottawa: Canada's Food Guide. Healthy eating recommendations. [accessed 2022 Dec 04]. Available from: <https://food-guide.canada.ca/en/healthy-eating-recommendations/>.
25. Grasso AC, Olthof MR, van Dooren C, Broekema R, Visser M, and Brouwer IA. Protein for a healthy future: How to increase protein intake in an environmentally sustainable way in older adults in the Netherlands. *J Nutr* 2021;151(1):109-119.
26. Kwasny T, Dobernig K, and Riefler P. Towards reduced meat consumption: A systematic literature review of intervention effectiveness, 2001-2019. *Appetite* 2022;168(105739).
27. Leroy F, Abraini F, Beal T, Dominguez-Salas P, Gregorini P, Manzano P, Rowntree J, and van Vliet S. Animal board invited review: Animal source foods in healthy, sustainable, and ethical diets - An argument against drastic limitation of livestock in the food system. *Animal* 2022;16(3):100457.

28. Paslakis G, Richardson C, Nohre M, Brahler E, Holzapfel C, Hilbert A, and de Zwaan M. Prevalence and psychopathology of vegetarians and vegans - Results from a representative survey in Germany. *Sci Rep* 2020;10(1):6840.
29. Melina V, Craig W, and Levin S. Position of the Academy of Nutrition and Dietetics: Vegetarian diets. *J Acad Nutr Diet* 2016;116(12):1970-1980.
30. Bryant CJ. We can't keep meating like this: Attitudes towards vegetarian and vegan diets in the United Kingdom. *Sustainability* 2019;11(6844).
31. Neff RA, Edwards D, Palmer A, Ramsing R, Righter A, and Wolfson J. Reducing meat consumption in the USA: a nationally representative survey of attitudes and behaviours. *Public Health Nutr* 2018;21(10):1835-1844.
32. OpinionWay - Good Food Institute Meat consumption and attitudes towards sustainable proteins in Europe. 2022.
33. de Gavelle E, Davidenko O, Fouillet H, Delarue J, Darcel N, Huneau JF, and Mariotti F. Self-declared attitudes and beliefs regarding protein sources are a good prediction of the degree of transition to a low-meat diet in France. *Appetite* 2019;142(104345).
34. Stoll-Kleemann S and Schmidt UJ. Reducing meat consumption in developed and transition countries to counter climate change and biodiversity loss: A review of influence factors. *Regional Environmental Change* 2017;17(1261-1277).
35. Nungesser F and Winter M. Meat and social change: Sociological perspectives on the consumption and production of animals. *OZS Osterr Z Soziol* 2021;46(2):109-124.
36. Hargreaves SM, Raposo A, Saraiva A, and Zandonadi RP. Vegetarian diet: An overview through the perspective of quality of life domains. *Int J Environ Res Public Health* 2021;18(8):
37. Sobal J. Men, meat, and marriage: Models of masculinity. *Food Foodways* 2005;13(135-158).
38. Gastaldello A, Giampieri F, de Giuseppe R, Grosso G, Baroni L, and Battino M. The rise of processed meat alternatives: A narrative review of the manufacturing, composition, nutritional profile and health effects of newer sources of protein, and their place in healthier diets. *Trends Food Science Technol* 2022;
39. O'Connor A. Fake meat vs real meat. *New York Times* 2020, December 2;
40. Pointing C. Meat substitutes help people transition to a vegan diet, says PCRM researcher. *LIVEKINDLY*
41. Romao B, Botelho RBA, Nakano EY, Raposo A, Han H, Vega-Munoz A, Ariza-Montes A, and Zandonadi RP. Are vegan alternatives to meat products healthy? A study on nutrients and main ingredients of products commercialized in Brazil. *Front Public Health* 2022;10(900598).
42. Faunalytics. Faunalytics. How many former vegetarians and vegans are there? 2014;
43. Chiles RM and Fitzgerald AJ. Why is meat so important in Western history and culture? A genealogical critique of biophysical and political-economic explanations. *Agric Hum Values* 2018;35):1-17.
44. Tso R and Forde CG. Unintended consequences: Nutritional impact and potential pitfalls of switching from animal- to plant-based foods. *Nutrients* 2021;13(8):
45. Hu FB, Otis BO, and McCarthy G. Can plant-based meat alternatives be part of a healthy and sustainable diet? *JAMA* 2019;322(16):1547-8.

46. Klapp A-L, Feil N, and Risius A. A global analysis of national dietary guidelines on plant-based diets and substitutions for animal-based foods border. *Current Developments in Nutr* 2022;
47. Canada's Food Guide. Eat Protein Foods. Accessed December 3, 2022. <https://food-guide.canada.ca/en/healthy-eating-recommendations/make-it-a-habit-to-eat-vegetables-fruit-whole-grains-and-protein-foods/eat-protein-foods/>.
48. Smith NW, Fletcher AJ, Hill JP, and McNabb WC. Modeling the contribution of meat to global nutrient availability. *Front Nutr* 2022;9(766796).
49. Cocking C, Walton J, Kehoe L, Cashman KD, and Flynn A. The role of meat in the European diet: current state of knowledge on dietary recommendations, intakes and contribution to energy and nutrient intakes and status. *Nutrition Res Reviews* 2020;33(181-9).
50. Agarwal S and Fulgoni VL, 3rd. Contribution of beef to key nutrient intakes in American adults: an updated analysis with NHANES 2011-2018. *Nutr Res* 2022;105(105-112).
51. Dussiot A, Fouillet H, Perraud E, Salome M, Huneau JF, Kesse-Guyot E, and Mariotti F. Nutritional issues and dietary levers during gradual meat reduction - A sequential diet optimization study to achieve progressively healthier diets. *Clin Nutr* 2022;41(12):2597-2606.
52. Vieux F, Remond D, Peyraud JL, and Darmon N. Approximately Half of Total Protein Intake by Adults must be Animal-Based to Meet Non-Protein Nutrient-Based Recommendations with Variation Due to Age and Sex. *J Nutr* 2022;
53. Mariotti F, Huneau JF, and Fouillet H. No nutritional lessons can be learned from a misspecified and over-restricted model with no sensitivity analysis. *J Nutr* In press;
54. Fouillet H, Dussiot A, Perraud E, Wang J, Kesse-Guyot E, and Mariotti F. Plant to animal protein ratio in the diet: nutrient adequacy, long-term health and environmental pressure. *medRxiv* 2022;
55. Lusk JL, Blaustein-Rejto D, Shah S, and Tonsor GT. Impact of plant-based meat alternatives on cattle inventories and greenhouse gas emissions. *Environ Res Lett* 2022;17(024035).
56. Makkar HPS. Feed demand landscape and implications of food-not feed strategy for food security and climate change. *Animal* 2018;12:1744-54. *Animal* 2018;12(1744-54).
57. Khan S, Loyola C, Dettling J, Hester J, and Moses R Comparative environmental LCA of the Impossible burger with conventional ground beef burger. (accessed January 15, 2021). 2019.
58. Quantis USA. A comparative life cycle assessment of plant-based foods and meat foods. Assessing the environmental benefits of plant-based dietary choices through: a comparison of meal choices, and a comparison of meat products and MorningStar Farms® veggie products. https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&cad=rja&uact=8&ved=2ahUKEwiEot-ZiYb1AhWSjokEHR4AQ4QFnoECAQQAQ&url=https%3A%2F%2Fwww.morningstarfarms.com%2Fcontent%2Fdam%2FNorthAmerica%2Fmorningstarfarms%2Fpdf%2FMSFPlantBasedLCARreport_2016-04-10_Final.pdf&usg=AOvVaw2LHWffjW87k84IKhEWTwfp. 2016;
59. Detzel A, Kruger M, Busch M, Blanco-Gutierrez I, Varela C, Manners R, Bez J, and Zannini E. Life cycle assessment of animal-based foods and plant-based protein-rich alternatives: an environmental perspective. *J Sci Food Agric* 2022;102(12):5098-5110.

60. Markel H. John Harvey Kellogg and the pursuit of wellness. *JAMA* 2011;305(17):1814-5.
61. Pantry N. The fascinating history of meat alternatives—And why they're not "unnatural" *Peaceful Dumpling* 2020;
62. Tziva M, Negro SO, Kalfagianni A, and Hekkert MP. Understanding the protein transition: The rise of plant-based meat substitutes. *Environmental Innovation Societal Transitions* 2020;35(217-231).
63. Woolf PJ, Fu LL, and Basu A. vProtein: identifying optimal amino acid complements from plant-based foods. *PLoS One* 2011;6(4):e18836.
64. Rafii M, Pencharz PB, Ball RO, Tomlinson C, Elango R, and Courtney-Martin G. Bioavailable methionine assessed using the indicator amino acid oxidation method is greater when cooked chickpeas and steamed rice are combined in healthy young men. *J Nutr* 2020;
65. Muoki PN, de Kock HL, and Emmambux MN. Effect of soy flour addition and heat-processing method on nutritional quality and consumer acceptability of cassava complementary porridges. *J Sci Food Agric* 2012;92(8):1771-9.
66. Dimina L, Remond D, Huneau JF, and Mariotti F. Combining Plant Proteins to Achieve Amino Acid Profiles Adapted to Various Nutritional Objectives-An Exploratory Analysis Using Linear Programming. *Front Nutr* 2021;8(809685).
67. Derbyshire EJ. Flexitarian diets and health: A review of the evidence-based literature. *Front Nutr* 2016;3(55).
68. Rizzo G and Baroni L. Soy, soy foods and their role in vegetarian diets. *Nutrients* 2018;10(1):
69. Messina M and Messina V. The role of soy in vegetarian diets. *Nutrients* 2010;2(855-888).
70. Hull MA. Veggie burgers. *J Renal Nutr* 2013;23(e99-e100).
71. Lima M, Costa R, Rodrigues I, Lameiras J, and Botelho G. A narrative review of alternative protein sources: Highlights on meat, fish, egg and dairy analogues. *Foods* 2022;11(
72. Dekkers BL, Boom RM, and van der Goot AJ. Structuring processes for meat analogues. *Trends Food Sci Technol* 2018;81(25-36).
73. Pohjolainen P, Vinnari M, and Jokinen P. Consumers' perceived barriers to following a plant-based diet. *British Food J* 2015;117(1150–67).
74. Szenderak J, Frona D, and Rakos M. Consumer acceptance of plant-based meat substitutes: A narrative review. *Foods* 2022;11(
75. Michel F, Hartman C, and Siegrist M. Consumers' associations, perceptions and acceptance of meat and plant-based meat alternatives. *Food Quality Preference* 2021;87(104063).
76. Cole E, Goeler-Slough N, Cox A, and Nolden A. Examination of the nutritional composition of alternative beef burgers available in the United States. *Int J Food Sci Nutr* 2022;73(4):425-432.
77. He J, Evans NM, Liu H, and Shao S. A review of research on plant-based meat alternatives: Driving forces, history, manufacturing, and consumer attitudes. *Compr Rev Food Sci Food Saf* 2020;19(5):2639-2656.
78. Szejda KUT and Matti W. Accelerating consumer adoption of plant-based meat: An evidence-based guide for effective practice. *The Good Food Institute* 2020;

79. Neuhofer ZT and Lusk JL. Most plant-based meat alternative buyers also buy meat: an analysis of household demographics, habit formation, and buying behavior among meat alternative buyers. *Sci Rep* 2022;12(1):13062.
80. Alessandrini R, Brown MK, Pombo-Rodrigues S, Bhageerutty S, He FJ, and MacGregor GA. Nutritional quality of plant-based meat products available in the UK: A cross-sectional survey. *Nutrients* 2021;13(12):
81. Pointke M and Pawelzik E. Plant-based alternative products: Are they healthy alternatives? Micro- and macronutrients and nutritional scoring. *Nutrients* 2022;14(3):
82. Harnack L, Mork S, Valluri S, Weber C, Schmitz K, Stevenson J, and Pettit J. Nutrient composition of a selection of plant-based ground beef alternative products available in the United States. *J Acad Nutr Diet* 2021;21(12):2401-2408 e12.
83. Bryngelsson S, Moshtaghian H, Bianchi M, and Hallstrom E. Nutritional assessment of plant-based meat analogues on the Swedish market. *Int J Food Sci Nutr* 2022;1-13.
84. Boukid F. Plant-based meat analogues: from niche to mainstream. *Eur Food Res Technol* 2021;247(297-308).
85. Grasso AC, Hung Y, Olthof MR, Verbeke W, and Brouwer IA. Older consumers' readiness to accept alternative, more sustainable protein sources in the European Union. *Nutrients* 2019;11(8):
86. Parodi A, Leip A, De Boer IJM, Slegers PM, Ziegler F, Temme EHM, Herrero M, Tuomisto H, Valin H, Van Middelaar CE, et al. The potential of future foods for sustainable and healthy diets. *Nature Sustainability* 2018;1(782-789).
87. Salome M, Huneau JF, Le Baron C, Kesse-Guyot E, Fouillet H, and Mariotti F. Substituting meat or dairy products with plant-based substitutes has small and heterogeneous effects on diet quality and nutrient security: a simulation study in French adults (INCA3). *J Nutr* 2021;151(8):2435-2445.
88. Pasiakos SM, Agarwal S, Lieberman HR, and Fulgoni VL, 3rd. Sources and Amounts of Animal, Dairy, and Plant Protein Intake of US Adults in 2007-2010. *Nutrients* 2015;7(8):7058-69.
89. Grigg D. The pattern of world protein consumption. *Geoforum* 1995;26(1):1-17.
90. Daily per capita protein supply -. *Our World in Data*, 2017.
91. Gortner WA. Nutrition in the United States, 1900 to 1974. *Cancer Res* 1975;35(11 Pt. 2):3246-53.
92. Samtiya M, Aluko RE, Dhewa T, and Moreno-Rojas JM. Potential health benefits of plant food-derived bioactive components: An overview. *foods* 2021;10(
93. Dhingra D, Michael M, Rajput H, and Patil RT. Dietary fibre in foods: A review. *J Food Science Technology* 2012;255-266.
94. Papier K, Knuppel A, Syam N, Jebb SA, and Key TJ. Meat consumption and risk of ischemic heart disease: A systematic review and meta-analysis. *Crit Rev Food Sci Nutr* 2021;1-12.
95. Johnston BC, Zeraatkar D, Han MA, Vernooij RWM, Valli C, El Dib R, Marshall C, Stover PJ, Fairweather-Tait S, Wojcik G, et al. Unprocessed Red Meat and Processed Meat Consumption: Dietary Guideline Recommendations From the Nutritional Recommendations (NutriRECS) Consortium. *Ann Intern Med* 2019;171(10):756-764.
96. Li Y, Hruby A, Bernstein AM, Ley SH, Wang DD, Chiuve SE, Sampson L, Rexrode KM, Rimm EB, Willett WC, et al. Saturated fats compared with unsaturated fats and sources of

- carbohydrates in relation to risk of coronary heart disease: A prospective cohort study. *J Am Coll Cardiol* 2015;66(14):1538-48.
97. Astrup A, Teicholz N, Magkos F, Bier DM, Brenna JT, King JC, Mente A, Ordovas JM, Volek JS, Yusuf S, et al. Dietary saturated fats and health: Are the U.S. guidelines evidence-based? *Nutrients* 2021;13(10):
 98. Day L, Cakebread JA, and Loveday SM. Food proteins from animals and plants: Differences in the nutritional and functional properties. *Trends Food Sci Technol* 2022;119(428-442).
 99. Gorissen SHM, Crombag JJR, Senden JMG, Waterval WAH, Bierau J, Verdijk LB, and van Loon LJC. Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids* 2018;50(12):1685-1695.
 100. Mariotti F. Animal and plant protein sources and cardiometabolic health. *Adv Nutr* 2019;10(Suppl_4):S351-S366.
 101. Green CL, Lamming DW, and Fontana L. Molecular mechanisms of dietary restriction promoting health and longevity. *Nat Rev Mol Cell Biol* 2022;23(1):56-73.
 102. Lopez-Moreno M, M G-R, and Miguel M. Antinutrients: Lectins, goitrogens, phytates and oxalates, friends or foe? *J Functional Foods* 2022;89(104939).
 103. Petroski W and Minich DM. Is there such a thing as "anti-nutrients"? A narrative review of perceived problematic plant compounds. *Nutrients* 2020;12(10):
 104. Lv JL, Wu QJ, Li XY, Gao C, Xu MZ, Yang J, Zang ST, Luan J, Cai DZ, Chang Q, et al. Dietary protein and multiple health outcomes: An umbrella review of systematic reviews and meta-analyses of observational studies. *Clin Nutr* 2022;41(8):1759-1769.
 105. Qi XX and Shen P. Associations of dietary protein intake with all-cause, cardiovascular disease, and cancer mortality: A systematic review and meta-analysis of cohort studies. *Nutr Metab Cardiovasc Dis* 2020;30(7):1094-1105.
 106. Bakaloudi DR, Halloran A, Rippin HL, Oikonomidou AC, Dardavesis TI, Williams J, Wickramasinghe K, Breda J, and Chourdakis M. Intake and adequacy of the vegan diet. A systematic review of the evidence. *Clin Nutr* 2021;40(5):3503-3521.
 107. Pinckaers PJM, Trommelen J, Snijders T, and van Loon LJC. The anabolic response to plant-based protein ingestion. *Sports Med* 2021;51(Suppl 1):59-74.
 108. Rutherford SM, Fanning AC, Miller BJ, and Moughan PJ. Protein digestibility-corrected amino acid scores and digestible indispensable amino acid scores differentially describe protein quality in growing male rats. *J Nutr* 2015;145(2):372-9.
 109. Mathai JK, Liu Y, and Stein HH. Values for digestible indispensable amino acid scores (DIAAS) for some dairy and plant proteins may better describe protein quality than values calculated using the concept for protein digestibility-corrected amino acid scores (PDCAAS). *Br J Nutr* 2017;117(4):490-499.
 110. Hughes GJ, Ryan DJ, Mukherjea R, and Schasteen CS. Protein digestibility-corrected amino acid scores (PDCAAS) for soy protein isolates and concentrate: Criteria for evaluation. *J Agric Food Chemistry* 2011;59(23):12707-12.
 111. Mariotti F, *Plant protein, animal protein, and protein quality*, in *Vegetarian and plant-based diets in health and disease prevention*, F. Mariotti, Editor 2017, Academic Press: Cambridge, MA. p. 621-642.

112. Domic J, Grootswagers P, van Loon LJC, and de Groot L. Perspective: Vegan diets for older adults? A perspective on the potential impact on muscle mass and strength. *Adv Nutr* 2022;
113. Mangels R, Messina V, and Messina M, *The Dietitian's Guide to Vegetarian Diets*. 3rd ed 2011, Burlington, Massachusetts: Jones and Bartlett.
114. Ciuris C, Lynch HM, Wharton C, and Johnston CS. A comparison of dietary protein digestibility, based on DIAAS scoring, in vegetarian and non-vegetarian athletes. *Nutrients* 2019;11(12):
115. Mariotti F and Gardner CD. Dietary protein and amino acids in vegetarian diets-A review. *Nutrients* 2019;11(11):
116. Gilani GS, Cockell KA, and Sepehr E. Effects of antinutritional factors on protein digestibility and amino acid availability in foods. *J AOAC Int* 2005;88(3):967-87.
117. Health Council of the Netherlands. Dietary Reference Intakes: energy, proteins, fats and digestible carbohydrates. The Hague (Netherlands): Health Council of the Netherlands; 2001.
118. Adam D. How far will global population rise? Researchers can't agree. *Nature* 2021;597(7877):462-465.
119. Porritt J and McCarthy M The global protein challenge. 2017.
120. Henchion M, Hayes M, Mullen AM, Fenelon M, and Tiwari B. Future protein supply and demand: Strategies and factors influencing a sustainable equilibrium. *Foods* 2017;6(7):
121. Lieberman HR, Fulgoni VL, Agarwal S, Pasiakos SM, and Berryman CE. Protein intake is more stable than carbohydrate or fat intake across various US demographic groups and international populations. *Am J Clin Nutr* 2020;112(1):180-186.
122. Wolfe RR, Cifelli AM, Kostas G, and Kim IY. Optimizing protein intake in adults: Interpretation and application of the recommended dietary allowance compared with the acceptable macronutrient distribution range. *Adv Nutr* 2017;8(2):266-275.
123. Paris JMG, Falkenberg T, Nothlings U, Heinzl C, Borgemeister C, and Escobar N. Changing dietary patterns is necessary to improve the sustainability of Western diets from a One Health perspective. *Sci Total Environ* 2022;811(151437).
124. Messina M. Perspective: Soybeans can help address the caloric and protein needs of a growing global population. *Front Nutr* 2022;9(909464).
125. van der Weele C, Feindt P, van der Goot AJ, van Mierlo B, and van Boekel M. Meat alternatives: An integrative comparison. *Trends Food Sci Technol* 2019;88(505-512).
126. Onwezen MC, Bouwman EP, Reinders MJ, and Dagevos H. A systematic review on consumer acceptance of alternative proteins: Pulses, algae, insects, plant-based meat alternatives, and cultured meat. *Appetite* 2021;159(105058).
127. Calles T, Xipsiti M, and del Castell R. Legacy of the international year of pulses. *Environmental Earth* 2022;78(124).
128. Zhubi-Bakija F, Bajraktari G, Bytyci I, Mikhailidis DP, Henein MY, Latkovskis G, Rexhaj Z, Zhubi E, Banach M, and International Lipid Expert P. The impact of type of dietary protein, animal versus vegetable, in modifying cardiometabolic risk factors: A position paper from the International Lipid Expert Panel (ILEP). *Clin Nutr* 2021;40(1):255-276.
129. Mullins AP and Arjmandi BH. Health Benefits of Plant-Based Nutrition: Focus on Beans in Cardiometabolic Diseases. *Nutrients* 2021;13(2):

130. Becerra-Tomas N, Papandreou C, and Salas-Salvado J. Legume Consumption and Cardiometabolic Health. *Adv Nutr* 2019;10(Suppl_4):S437-S450.
131. Drewnowski A and Rehm CD. Vegetable cost metrics show that potatoes and beans provide most nutrients per penny. *PLoS One* 2013;8(5):e63277.
132. Harwatt HS, J, Eshel G, Soret S, and Ripple W. Substituting beans for beef as a contribution toward US climate change targets. *Clim Change* 2017;143(262-270).
133. Rawal V and Navarro DK. *The Global Economy of Pulses*. FAO 2019;
134. Melendrez-Ruiz J, Buatois Q, Chambaron S, Monnery-Patris S, and Arvisenet G. French consumers know the benefits of pulses, but do not choose them: An exploratory study combining indirect and direct approaches. *Appetite* 2019;141(104311).
135. Reuze A, Mejean C, Carrere M, Sirieix L, Druésne-Pecollo N, Peneau S, Touvier M, Hercberg S, Kesse-Guyot E, and Alles B. Rebalancing meat and legume consumption: change-inducing food choice motives and associated individual characteristics in non-vegetarian adults. *Int J Behav Nutr Phys Act* 2022;19(1):112.
136. HCSP. Avis relatif aux objectifs de santé publique quantifiés pour la politique nutritionnelle de santé publique (PNNS) 2018–2022. Paris: Haut Conseil de la santé publique (HCSP); 2018. Available from: https://www.hcsp.fr/Explo_re.cgi/Telecharger?NomFichier=hcspa20180209_avisrelauxobjequantpourlapolitique. pdf.
137. ANSES OPINION of the French Agency for Food, Environmental and Occupational Health & Safety (ANSES Opinion Request No 2017-SA-014). 2019.
138. Didinger C and Thompson HJ. Defining nutritional and functional niches of legumes: A call for clarity to distinguish a future role for pulses in the Dietary Guidelines for Americans. *Nutrients* 2021;13(4):
139. Jonnalagadda SS, Harnack L, Liu RH, McKeown N, Seal C, Liu S, and Fahey GC. Putting the whole grain puzzle together: health benefits associated with whole grains--summary of American Society for Nutrition 2010 Satellite Symposium. *J Nutr* 2011;141(5):1011S-22S.
140. Marinangeli CPF, Curran J, Barr SI, Slavin J, Puri S, Swaminathan S, Tapsell L, and Patterson CA. Enhancing nutrition with pulses: defining a recommended serving size for adults. *Nutr Rev* 2017;75(12):990-1006.
141. U.S. Department of Agriculture and U.S. Department of Health and Human Services. *Dietary Guidelines for Americans, 2020-2025*. 9th Edition. December 2020. Available at [DietaryGuidelines.gov](https://www.dietaryguidelines.gov).
142. Foyer CH, Lam HM, Nguyen HT, Siddique KH, Varshney RK, Colmer TD, Cowling W, Bramley H, Mori TA, Hodgson JM, et al. Neglecting legumes has compromised human health and sustainable food production. *Nat Plants* 2016;2(16112).
143. Didinger C and Thompson H. Motivating pulse-centric eating patterns to benefit human and environmental well-being. *Nutrients* 2020;12(11):
144. Perera T, Russo C, Takata Y, and Bobe G. Legume consumption patterns in US adults: National Health and Nutrition Examination Survey (NHANES) 2011-2014 and Beans, Lentils, Peas (BLP) 2017 Survey. *Nutrients* 2020;12(5):
145. Tao MH, Liu JL, and Nguyen UDT. Trends in Diet Quality by Race/Ethnicity among Adults in the United States for 2011-2018. *Nutrients* 2022;14(19):

146. Lucier G, Lin B-H, Allshouse J, and Kanto LS, *Factors Affecting Dry Bean Consumption in the United States. Economic Research Service USDA. Vegetables and Specialties S&O/VGS-280/April*, 2000.
147. Semba RD, Rahman N, Du S, Ramsing R, Sullivan V, Nussbaumer E, Love D, and Bloem MW. Patterns of legume purchases and consumption in the United States. *Front Nutr* 2021;8(732237).
148. Orlich MJ, Mashchak AD, Jaceldo-Siegl K, Utt JT, Knutsen SF, Sveen LE, and Fraser GE. Dairy foods, calcium intakes, and risk of incident prostate cancer in Adventist Health Study-2. *Am J Clin Nutr* 2022;
149. Papier K, Tong TY, Appleby PN, Bradbury KE, Fensom GK, Knuppel A, Perez-Cornago A, Schmidt JA, Travis RC, and Key TJ. Comparison of Major Protein-Source Foods and Other Food Groups in Meat-Eaters and Non-Meat-Eaters in the EPIC-Oxford Cohort. *Nutrients* 2019;11(4):
150. Doma KM, Farrell EL, Leith-Bailey ER, Soucier VD, and Duncan AM. Motivators, barriers and other factors related to bean consumption in older adults. *J Nutr Gerontol Geriatr* 2019;38(4):397-413.
151. Leterme P and Carmenza Munoz L. Factors influencing pulse consumption in Latin America. *Br J Nutr* 2002;88 Suppl 3(S251-5).
152. Bianchi F, Stewart C, Astbury NM, Cook B, Aveyard P, and Jebb SA. Replacing meat with alternative plant-based products (RE-MAP): a randomized controlled trial of a multicomponent behavioral intervention to reduce meat consumption. *Am J Clin Nutr* 2022;115(5):1357-1366.
153. Messina M, Sievenpiper JL, Williamson P, Kiel J, and Erdman JW. Perspective: Soy-based meat and dairy alternatives, despite classification as ultra-processed foods, deliver high-quality nutrition on par with unprocessed or minimally processed animal-based counterparts. *Adv Nutr* 2022;13(3):726-738.
154. Toh DWK, Akila SRV, and Henry CJ. Unknown impacts of plant-based meat alternatives on long-term health. *Nature Food* 2022;
155. Fardet A and Rock E. Chronic diseases are first associated with the degradation and artificialization of food matrices rather than with food composition: calorie quality matters more than calorie quantity. *Eur J Nutr* 2022;
156. Fardet A and Rock E. Perspective: Reductionist nutrition research has meaning only within the framework of holistic and ethical thinking. *Adv Nutr* 2018;9(6):655-670.
157. Gehring J, Touvier M, Baudry J, Julia C, Buscail C, Srour B, Hercberg S, Peneau S, Kesse-Guyot E, and Alles B. Consumption of ultra-processed foods by pesco-vegetarians, vegetarians, and vegans: Associations with duration and age at diet initiation. *J Nutr* 2021;151(1):120-131.
158. Macdiarmid JI. The food system and climate change: are plant-based diets becoming unhealthy and less environmentally sustainable? *Proc Nutr Soc* 2021;1-6.
159. Santo RE, Kim BF, Goldman SE, Dutkiewicz J, Biehl EMB, Bloem MW, Neff RA, and Nachman KE, 2,4,5*. Considering plant-based meat substitutes and cell-based meats: A public health and food systems perspective. *Frontiers Sustainable Food Systems* 2020;4(134).

160. Clark M, Springmann M, Rayner M, Scarborough P, Hill J, Tilman D, Macdiarmid JI, Fanzo J, Bandy L, and Harrington RA. Estimating the environmental impacts of 57,000 food products. *Proc Natl Acad Sci U S A* 2022;119(33):e2120584119.
161. Monteiro CA. Nutrition and health. The issue is not food, nor nutrients, so much as processing. *Public Health Nutr* 2009;12(5):729-31.
162. Monteiro CA, Cannon G, Levy RB, Moubarac JC, Louzada ML, Rauber F, Khandpur N, Cediel G, Neri D, Martinez-Steele E, et al. Ultra-processed foods: what they are and how to identify them. *Public Health Nutr* 2019;22(5):936-941.
163. Lane MM, Davis JA, Beattie S, Gomez-Donoso C, Loughman A, O'Neil A, Jacka F, Berk M, Page R, Marx W, et al. Ultraprocessed food and chronic noncommunicable diseases: A systematic review and meta-analysis of 43 observational studies. *Obes Rev* 2021;22(3):e13146.
164. Zinocker MK and Lindseth IA. The Western diet-microbiome-host interaction and its role in metabolic disease. *Nutrients* 2018;10(3):
165. Toribio-Mateas MA, Bester A, and Klimenko N. Impact of plant-based meat alternatives on the gut microbiota of consumers: A real-world study. *Foods* 2021;10(2040).
166. Mozaffarian D, El-Abbadi NH, O'Hearn M, J E-M, Masters WA, Jacques P, Shi P, J.B B, and Micha R. Food Compass is a nutrient profiling system using expanded characteristics for assessing healthfulness of foods. *Nature Food* 2021;2(809-818).
167. Curtain F and Grafenauer S. Plant-based meat substitutes in the flexitarian age: An audit of products on supermarket shelves. *Nutrients* 2019;11(11):
168. Crimarco A, Springfield S, Petlura C, Streaty T, Cunanan K, Lee J, Fielding-Singh P, Carter MM, Topf MA, Wastyk HC, et al. A randomized crossover trial on the effect of plant-based compared with animal-based meat on trimethylamine-N-oxide and cardiovascular disease risk factors in generally healthy adults: Study With Appetizing Plantfood-Meat Eating Alternative Trial (SWAP-MEAT). *Am J Clin Nutr* 2020;112(5):1188-1199.
169. Crimarco A, Landry MJ, Carter MM, and Gardner CD. Assessing the effects of alternative plant-based meats v. animal meats on biomarkers of inflammation: a secondary analysis of the SWAP-MEAT randomized crossover trial. *J Nutr Sci* 2022;11(e82).
170. Roberts AK, Busque V, Robinson JL, Landry MJ, and Gardner CD. SWAP-MEAT Athlete (study with appetizing plant-food, meat eating alternatives trial) - investigating the impact of three different diets on recreational athletic performance: a randomized crossover trial. *Nutr J* 2022;21(1):69.
171. Jenkins DJA, Jones PJH, Abdullah MMH, Lamarche B, Faulkner D, Patel D, Sahye-Pudaruth S, Paquette M, Bashyam B, Pichika SC, et al. Low-carbohydrate vegan diets in diabetes for weight loss and sustainability: a randomized controlled trial. *Am J Clin Nutr* 2022;
172. Alae-Carew C, Green R, Stewart C, Cook B, Dangour AD, and Scheelbeek PFD. The role of plant-based alternative foods in sustainable and healthy food systems: Consumption trends in the UK. *Sci Total Environ* 2022;807(Pt 3):151041.
173. van Vliet S, Kronberg SL, and Provenza FD. Plant-based meats, human health, and climate change. *Frontiers Sustainable Food Systems* 2020;4(
174. Rodriguez Madrera R, Campa Negrillo A, Suarez Valles B, and Ferreira Fernandez JJ. Phenolic content and antioxidant activity in seeds of common bean (*Phaseolus vulgaris* L.). *Foods* 2021;

175. Carbas B, Machado N, Oppolzer D, Ferreira L, Queiroz M, Brites C, Rosa EA, and Barros AI. Nutrients, antinutrients, phenolic composition, and antioxidant activity of common bean cultivars and their potential for food applications. *Nutrients* 2020;9(
176. Yang QQ, Gan RY, Ge YY, Zhang D, and Corke H. Polyphenols in common beans (*Phaseolus vulgaris* L.): Chemistry, analysis, and factors affecting composition. *Compr Rev Food Sci Food Saf* 2018;17(6):1518-1539.
177. Zhu B, Sun Y, Qi L, Zhong R, and Miao X. Dietary legume consumption reduces risk of colorectal cancer: evidence from a meta-analysis of cohort studies. *Sci Rep* 2015;5(8797).
178. Li H, Li J, Shen Y, Wang J, and Zhou D. Legume consumption and all-cause and cardiovascular disease mortality. *Biomed Res Int* 2017;2017(8450618).
179. Kim SJ, de Souza RJ, Choo VL, Ha V, Cozma AI, Chiavaroli L, Mirrahimi A, Blanco Mejia S, Di Buono M, Bernstein AM, et al. Effects of dietary pulse consumption on body weight: a systematic review and meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2016;103(5):1213-23.
180. Hafiz MS, Campbell MD, O'Mahoney LL, Holmes M, Orfila C, and Boesch C. Pulse consumption improves indices of glycemic control in adults with and without type 2 diabetes: a systematic review and meta-analysis of acute and long-term randomized controlled trials. *Eur J Nutr* 2022;61(2):809-824.
181. van den Berg LA, Mes JJ, Mensink M, and Wanders AJ. Protein quality of soy and the effect of processing: A quantitative review. *Frontiers Nutr* 2022;
182. Blanco Mejia S, Messina M, Li SS, Vigiouliouk E, Chiavaroli L, Khan TA, Srichaikul K, Mirrahimi A, Sievenpiper JL, Kris-Etherton P, et al. A meta-analysis of 46 studies identified by the FDA demonstrates that soy protein decreases circulating LDL and total cholesterol concentrations in adults. *J Nutr* 2019;149(6):968-981.
183. Jenkins DJA, Blanco Mejia S, Chiavaroli L, Vigiouliouk E, Li SS, Kendall CWC, Vuksan V, and Sievenpiper JL. Cumulative meta-analysis of the soy effect over time. *J Am Heart Assoc* 2019;8(13):e012458.
184. Messina M, Lynch H, Dickinson JM, and Reed KE. No difference between the effects of supplementing with soy protein versus animal protein on gains in muscle mass and strength in response to resistance exercise. *Int J Sport Nutr Exerc Metab* 2018;28(6):674-685.
185. Musa-Veloso K and Juana J. Regulation and labeling of plant-based beverages and simulated meat, poultry, and egg products in Canada and the United States. *Cereal Foods World* 2020;65(4):July-August.
186. Salome M, Mariotti F, Nicaud MC, Dussiot A, Kesse-Guyot E, Maillard MN, Huneau JF, and Fouillet H. The potential effects of meat substitution on diet quality could be high if meat substitutes are optimized for nutritional composition—a modeling study in French adults (INCA3). *Eur J Nutr* 2022;61(4):1991-2002.
187. Winham DM and Hutchins AM. Perceptions of flatulence from bean consumption among adults in 3 feeding studies. *Nutr J* 2011;10(128).
188. Shi L, Arntfield SD, and Nickerson M. Changes in levels of phytic acid, lectins and oxalates during soaking and cooking of Canadian pulses. *Food Res Int* 2018;107(660-668).
189. Mittermeier-Klessinger VK, Hofmann T, and Dawid C. Mitigating off-flavors of plant-based proteins. *J Agric Food Chem* 2021;69(32):9202-9207.

190. Tobin BD, O'Sullivan MG, Hamill RM, and Kerry JP. Effect of varying salt and fat levels on the sensory quality of beef patties. *Meat Sci* 2012;91(4):460-5.
191. Watson E Impossible Foods tweaks formulation, slashes saturated fat: 'The category has done a lousy job of explaining itself,' says CEO. 2022.
192. Ryan RM and Deci EL. Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *Am Psychol* 2000;55(1):68-78.
193. Grasso S and Jaworska S. Part meat and part plant: Are hybrid meat products fad or future? *Foods* 2020;9(
194. Askew K Making 'healthier' meat: Consumer attitudes towards to better-for-you-burgers and bangers. 2020.
195. Alcorta A, Porta A, Tárrega A, Alvarez MD, and Pilar Vaquero M. Foods for plant-based diets: challenges and innovations. *Foods* 2021;10(293).
196. Hicks TM, Knowles SO, and Farouk MM. Global provisioning of red meat for flexitarian diets. *Front Nutr* 2018;5(50).
197. Boukid F and Castellari M. Veggie burgers in the EU market: a nutritional challenge? *Eur Food Res Technol* 2021;1-9.
198. Thrane M, Paulsen PV, Orcutt MW, and Krieger TM, *Soy protein: Impacts, production, and applications*, in *Sustainable Protein Sources*, S.R. Nadathur, J.P.D. Wanasundara, and L. Scanlin, Editors. 2017, Academic Press: United Kingdom. p. 23-46.
199. Grasso S, Smith G, Bowers S, Ajayi OM, and Swainson M. Effect of texturised soy protein and yeast on the instrumental and sensory quality of hybrid beef meatballs. *J Food Sci Technol* 2019;56(6):3126-3135.
200. Drewnowski A, Henry CJ, and Dwyer JT. Proposed nutrient standards for plant-based beverages intended as milk alternatives. *Front Nutr* 2021;8(761442).
201. Mangels R, Messina V, and Messina M, *The Dietitian's Guide to Vegetarian Diets: Issues and Applications* 2022, Burlington, MA: Jones and Bartlett Learning. 600.

Table 1. World and selected region pulse intake

Region	g/d	% total intake	
		protein	kcal
World	21	6	3
Latin America & Caribbean	34	9	4
Sub-Saharan Africa	33	12	5
South Asia	33	11	5
North Africa	19	5	2
West Asia	19	6	3
Oceania	12	2	1
North America	11	2	1
Southeast Asia	9	3	1
Europe	7	2	1
East Asia	4	1	0
Caucasus & Central Asia	1	0	0

Source: Rawal V, Navarro DK. The Global Economy of Pulses. FAO 2019

Table 2. Protein and fiber content of selected legumes (per 100 g cooked) from the USDA's FoodData Central*

Legume	USDA FDC ID	Protein		Fiber
		(g)	% kcal	(g)
Soybeans	174271	18.21	42.35	6.0
Lupin	173804	15.57	53.69	2.8
Lentils	175254	9.02	31.65	7.9
Pinto beans	175200	9.01	25.20	9.0
Great Northern	173790	8.33	28.24	7.0
Kidney beans (red)	175242	8.67	27.31	7.4
Black beans	175237	8.86	26.85	8.7
Mung beans	175255	7.02	26.74	7.6
Peas (green)	170102	5.36	25.52	5.5
Navy beans	173794	8.23	23.51	10.5
Adzuki beans	173789	7.52	23.50	7.3
Lima beans	169316	6.81	22.15	5.3
Garbanzo beans	173799	8.86	21.61	7.6

* U.S. Department of Agriculture, Agricultural Research Service. FoodData Central, 2019. fdc.nal.usda.gov.

Table 3. Energy, macronutrient and fiber content of legumes and foods made from legumes*

	Pulses (N=29)	Tofu (N=10)	Peanut butter (N=3)	Peanuts (N=3)	Soy-PBMAs (N=5)
Energy (kcal)	127 ± 14	100.10 ± 24.86	189.00 ± 1.73	161.33 ± 0.94	198 ± 83.28
Protein (g)	8.62 ± 1.54	10.95 ± 2.53	7.27 ± 0.38	7.23 ± 0.14	18.8 ± 5.08
% kcal protein	27.1	43.8	15.4	17.9	38.0
Fat (g)	0.66 ± 0.62	5.76 ± 1.91	16.07 ± 0.31	13.8 0.24	11.2 ± 6.97
Carbohydrate (g)	22.55 ± 3.49	2.20 ± 0.98	7.24 ± 0.40	5.03 ± 0.64	8.2 ± 2.23
Fiber (g)	7.4 ± 2.0	1.07 ± 0.60	1.99 ± 0.50	2.52 ± 0.12	4.2 ± 2.04

*Values = mean ± standard deviation. Serving sizes: pulses, 100 g cooked tofu, 100 g raw; peanut butter, 2 tablespoons (32 g); peanuts, 1 oz (28 g); soy-PBMA (burgers, 1 patty). Sources: Values for tofu, peanut butter, peanuts from U.S. Department of Agriculture, Agricultural Research Service. FoodData Central, 2019. fdc.nal.usda.gov. All tofus except one either hard or firm. Values for pulses and soy-PBMAs come from references 133 and 145, respectively.

Nutrient database number for tofus: 16160, 16281, 16159, 16277, 16276, 16212, 16211, 16426, 16427, 16213; Peanut butter: 16097, 16398, 16167 and peanuts; 16091, 16093, 16095