



HAL
open science

Editorial: Sustainable production of bioactive pigments, volume II

Wee Sim Choo, Laurent Dufossé, Lourdes Morales-Oyervides

► To cite this version:

Wee Sim Choo, Laurent Dufossé, Lourdes Morales-Oyervides. Editorial: Sustainable production of bioactive pigments, volume II. *Frontiers in Sustainable Food Systems*, 2023, 7, 10.3389/fsufs.2023.1300072 . hal-04320165

HAL Id: hal-04320165

<https://hal.science/hal-04320165v1>

Submitted on 4 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN ACCESS

EDITED BY

Ashish Rawson,
National Institute of Food Technology
Entrepreneurship and Management, India

REVIEWED BY

Priyanka Kajla,
Guru Jambheshwar University of Science and
Technology, India
Rebeca Garcia,
Monterrey Institute of Technology and Higher
Education (ITESM), Mexico

*CORRESPONDENCE

Wee Sim Choo
✉ choo.wee.sim@monash.edu
Laurent Dufossé
✉ laurent.dufosse@univ-reunion.fr
Lourdes Morales-Oyervides
✉ lourdesmorales@uadec.edu.mx

RECEIVED 23 September 2023

ACCEPTED 12 October 2023

PUBLISHED 23 October 2023

CITATION

Choo WS, Dufossé L and Morales-Oyervides L
(2023) Editorial: Sustainable production of
bioactive pigments, volume II.
Front. Sustain. Food Syst. 7:1300072.
doi: 10.3389/fsufs.2023.1300072

COPYRIGHT

© 2023 Choo, Dufossé and Morales-Oyervides.
This is an open-access article distributed under
the terms of the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Editorial: Sustainable production of bioactive pigments, volume II

Wee Sim Choo^{1*}, Laurent Dufossé^{2*} and
Lourdes Morales-Oyervides^{3*}

¹School of Science, Monash University Malaysia, Bandar Sunway, Malaysia, ²ESIROI Agroalimentaire, Université de la Réunion, Saint-Denis, France, ³Facultad de Ciencias Químicas, Universidad Autónoma de Coahuila, Saltillo, Coahuila, Mexico

KEYWORDS

anthocyanin, bacterial pigments, fungal pigments, biological activities, ultrasonic extraction

Editorial on the Research Topic

Sustainable production of bioactive pigments, volume II

Pigments are compounds that humans perceive to have color. In nature, the most abundant natural pigments are chlorophylls, which are the source of green in all plants. Chlorophylls are the primary pigments for photosynthesis to produce energy for the development and growth of plants. Natural plant pigments that contribute to blue, purple, red, yellow and orange in nature are accessory pigments and secondary metabolites that possess diverse structures and functions in plants and offer potential health benefits to humans (Choo, 2019). Anthocyanins, betalains and carotenoids are the common natural plant pigments besides chlorophylls (Choo, 2018). Besides plant pigments, natural pigments can be obtained to a lesser extent from animals [e.g., astaxanthin from shrimp processing waste (Cahú et al., 2012)] or insects [e.g., cochineal colorant or carminic acid from *Dactylopius coccus* Costa (Dactylopiidae, Hemiptera) (Madsen et al., 1993)]. Microalgae and microorganisms are emerging sources of natural pigments (Morales-Oyervides et al., 2020). Bioactive pigments are pigments that interact with or affect cell tissues in the human body. Bioactive pigments that provide health benefits to humans have generated lots of interest among researchers, industries, and consumers. The continual prospecting of terrestrial and aquatic natural resources for bioactive pigments is unsustainable. Therefore, to ensure the continuity of utilization of bioactive pigments for future generations and feasible increases in their usage, sustainable production of bioactive pigments is needed (Choo et al., 2021).

This Research Topic is the second volume of the earlier Research Topic on “Sustainable production of bioactive pigments”. This Research Topic provides 1 review, 1 mini review and 3 original research articles on the production of pigments, colorants and dyes by microbial and biotechnological means. Joshi et al. summarizes the recent biotechnological innovations in microbial pigment production. Fermentation strategies, co-cultures, and genetic modification of microbial strains (mutagenesis, whole genome shuffling, metabolic engineering, use of CRISPR system) are among the most promising avenues for improvement. Microbial production is not a novel idea from researchers but a reality in many countries worldwide (e.g., astaxanthin, β -carotene, lycopene in Europe).

Filamentous fungi of the *Monascus* genus have been used for centuries in Asia for the production of mainly red pigments and, as such, they remain a subject of ongoing research from various angles. In a mini-review from Buranelo Egea et al., the potential, strategies, and challenges of *Monascus* pigment for food application are summarized. Regulatory aspects are still quite divergent around the world—as with any other product still in the

implementation phase—and are mainly related to the (hepatotoxic mycotoxin) citrinin content. All bioactive properties of *Monascus* pigments (over 60 different structures up to now) are also emphasized in this work.

In a research paper, Dias Oliveira et al. demonstrated that maltose production residues can be used to obtain natural pigments by *Monascus ruber* strain CCT 3802 in solid and submerged cultures. Using submerged fermentation, the culture medium containing 10 g L⁻¹ of maltose syrup provided the highest concentrations of red pigments (14.54 AU_{510nm} g⁻¹ dry biomass) with an intense dark red color. Using by-products, leftovers, food or feed residues is paramount in reducing the costs of microbial pigment production.

Another research paper dealing with *Monascus* focuses on the inhibitory effect of *Monascus purpureus* pigment extracts against fungi and its mechanism of action. In this study by Majhi et al., *Monascus purpureus* pigment extracts were tested (*in vitro*) against *Penicillium expansum* MTCC 4900, *Rhizopus stolonifer* MTCC 10595, and *Aspergillus niger* MTCC 8652 for antifungal activity. Numerous studies mention the antimicrobial properties of microbial pigments. Some cases are clearly established, for pure compounds (e.g., prodigiosin). Others require more experimentation, as they are still based on pigment extracts, whether crude or semi-purified.

As mentioned earlier in this editorial, plants are still playing a big role in the production of pigment, dye and color. Researchers continue investigating new genera and species, discovering new molecules (among the carotenoids, betalains, chlorophylls) and sources (such as wild plants, Amazonian plants, barks from Madagascar). Elgudayem et al. investigated the ultrasonic aqueous extraction of phenolic compounds from *Polygonum equisetifome* roots. The process was optimized using a Box-Behnken experimental design where the factors investigated were the extraction temperature, ultrasonic-assisted extraction time and liquid-solid ratio. The roots of *P. equisetifome* contained high anthocyanin content, water-soluble phenols and condensed tannins. The extracts exhibited promising

antioxidant potential and antibacterial activities against several pathogenic bacteria.

This editorial is a summary of the articles in this Research Topic. We would like to thank all the authors, reviewers, and Frontiers editorial team for their contributions to this Research Topic.

Author contributions

WC: Writing—review & editing. LD: Writing—original draft, Writing—review & editing. LM-O: Writing—review & editing.

Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The author(s) declared that they were an editorial board member of Frontiers, at the time of submission. This had no impact on the peer review process and the final decision.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Cahú, T. B., Santos, S. D., Mendes, A., Nader, H. B., and Bezerra, R. S. (2012). Recovery of protein, chitin, carotenoids and glycosaminoglycans from Pacific white shrimp (*Litopenaeus vannamei*) processing waste. *Proc. Biochem.* 47, 570–577. doi: 10.1016/j.procbio.2011.12.012
- Choo, W. S. (2018). "Betalains: Application in functional foods" in *Bioactive Molecules in Food. Reference Series in Phytochemistry*, eds J.-M. Mérillon, and K.G. Ramawat (Cham: Springer International Publishing), 1–28.
- Choo, W. S. (2019). "Fruit pigment changes during ripening", in *Encyclopedia of Food Chemistry*, eds P. Varelis, L. Melton, and F. Shahidi (Amsterdam: Elsevier), 117–123.
- Choo, W. S., Dufossé, L., and Morales-Oyervides, L. (2021). Editorial: Sustainable production of bioactive pigments. *Front. Sustain. Food Syst.* 5, 674311. doi: 10.3389/fsufs.2021.674311
- Madsen, H. L., Stapelfeldt, H., Bertelsen, G., and Skibsted, L. H. (1993). Cochineal as a colorant in processed pork meat. Colour matching and oxidative stability. *Food Chem.* 46, 265–271. doi: 10.1016/0308-8146(93)90117-X
- Morales-Oyervides, L., Ruiz-Sánchez, J. P., Oliveira, J. C., Sousa-Gallagher, M. J., Méndez-Zavala, A., Giuffrida, D., et al. (2020). Biotechnological approaches for the production of natural colorants by *Talaromyces/Penicillium*: a review. *Biotechnol. Adv.* 43, 107601. doi: 10.1016/j.biotechadv.2020.107601