

Beyond the genus stereotype. Who were the first toolmarkers in Africa? Crossed views between archaeology and anatomy

Sandrine Prat

► To cite this version:

Sandrine Prat. Beyond the genus stereotype. Who were the first toolmarkers in Africa? Crossed views between archaeology and anatomy. L'anthropologie, 2023, 127 (4), pp.103187. 10.1016/j.anthro.2023.103187. hal-04321679

HAL Id: hal-04321679 https://hal.science/hal-04321679v1

Submitted on 4 Dec 2023 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

- 1 Beyond the genus stereotype. Who were the first tool-markers in Africa ? Crossed views
- 2 between archaeology and anatomy.
- 3

4 Au-delà du stéréotype de genre. Qui étaient les premiers artisans en Afrique ? Regards 5 croisés entre archéologie et anatomie.

6

7 Sandrine Prat

8 UMR 7194 MNHN-CNRS-UPVD, Musée de l'Homme, 17 Place du Trocadéro, 75016 Paris

- 9 <u>sandrine.prat@mnhn.fr</u>
- 10

11 Abstract

The evolutionary history of hominins and archaeological assemblages has become 12 13 considerably more complex in the last twenty-five years due to the contribution of innovative 14 analytical methods and new archaeological and paleoanthropological discoveries. The 15 discoveries unearthed in eastern and southern Africa show a high biological and cultural diversity. But what is the nature of the relationships between hominins and the different 16 17 material cultures? Singularity or plurality of the stone and bone-tool makers and 18 archaeological assemblages? We propose a crossed view on these questions based on 19 investigations of hominin/culture associations in the Late Pliocene and Early Pleistocene 20 fieldwork records, morphology underlying hominins with manual skills as well on 21 interdisciplinary research conducted recently. Our work highlights the plurality of bone and 22 stone tool-makers, in particular Paranthropus, H. habilis and H. erectus/ergaster for Oldowan 23 assemblages. Similarly, in terms of anatomy, the type of dexterity required to make Oldowan 24 assemblages is likely not limited to Homo genus. The frequency of Paranthropus remains 25 associated with Oldowan assemblages is not negligible and insignificant. The occurrences 26 between Oldowan assemblages and Paranthropus seem not to be coincidental and 27 Paranthropus should be therefore considered as a stone and bone-tool maker in the same 28 way as it is for early Homo.

29

30 Keywords :

31 Hominins, material culture, Africa, Plio-Pleistocene

32 33 **Résumé**

L'histoire évolutive des hominines et des assemblages archéologiques est devenue 34 considérablement plus complexe au cours des vingt-cinq dernières années grâce à l'apport de 35 36 méthodes analytiques novatrices et de nouvelles découvertes tant archéologiques que 37 paléoanthropologiques. Les découvertes mises au jour en Afrique orientale et australe 38 montrent une grande diversité tant sur le plan biologique que culturel. Mais quelle est la 39 relation entre les hominines et les différentes cultures matérielles ? Singularité ou pluralité 40 des artisans des artefacts lithiques et osseux et des assemblages archéologiques ? Nous proposons un regard croisé sur ces questionnements à partir d'une part d'une analyse des 41 associations hominines/assemblages archéologiques au Pliocène supérieur et au Pléistocène 42 43 inférieur, et d'autre part de l'anatomie de ces hominines et des recherches interdisciplinaires menées ces dernières années. Notre travail met en évidence une pluralité des fabricants 44 d'outils en os et en pierre, en particulier Paranthropus, H. habilis et H. erectus/ergaster pour 45 les assemblages Oldowayens. De même, d'un point de vue anatomique, le type de dextérité 46 47 requis pour réaliser les assemblages Oldowayens n'est probablement pas limité au genre 48 Homo. La fréquence des restes de Paranthropus associés aux assemblages Oldowayens est

- 49 loin d'être négligeable et insignifiante. Les occurrences entre les assemblages Oldowayens et
- 50 le genre *Paranthropus* ne semblent pas être une coïncidence et le genre *Paranthropus* devrait
- 51 donc être considéré comme un artisan des artefacts lithiques et osseux au même titre que les
- 52 premiers représentants du genre *Homo*.
- 53
- 54 Mots clés :
- 55 hominines ; culture matérielle ; Afrique ; Plio-Pleistocene
- 56

57 Introduction

58 The evolutionary history of hominins and archaeological assemblages has become 59 considerably more complex in the last twenty-five years due to the contribution of innovative 60 analytical methods and new archaeological and paleoanthropological discoveries.

Indeed, discoveries unearthed in eastern and southern Africa indicate high biological and cultural diversity during the Late Pliocene and Early Pleistocene. In this context, can we consider that the manufacture of lithic artefacts is specific to the *Homo* genus and a pertinent defining criterion? The definition of our genus (*Homo* genus) and the criteria used to define it are still widely-debated topics, on anatomical/morphological, behavioural or philosophical levels.

- Among the criteria proposed to define the *Homo* genus by Leakey and his collaborators (Leakey et al., 1964), (i.e., endocranial volume, habitual erect posture, bipedal gait and power and precision grip), manual abilities appeared to be consubstantial in their definition (Leakey et al., 1964). Finally, tools are accorded a special place in the literature in the definition of the *Homo* genus, even if there are not mentioned in the diagnosis of Leakey et al. (1964) but in the following paragraph entitled "Cultural Association" (see paragraph conclusions and discussion).
- 74

75 The place and role of the archaeological remains in this definition (e.g., Prat, 2022) give rise to 76 the question of the possible relationship between the Oldowan industry and the H. habilis 77 specimens. Furthermore, the recent discovery of stone tools dated to 3.3 Ma in Kenya (LOM3 site) (evidence of stone-tools production, Harmand et al., 2015) and cutmarked bones at 78 79 Dikika site dated to 3.4 Ma in Ethiopia (indicating stone-tool use, McPherron et al., 2010), is 80 300,000-500,000 years older than the earliest Oldowan technology (Braun et al., 2019, Plummer et al., 2023) and 500,000 years before the oldest known fossils attributed by some 81 researchers to the Homo genus [Homo sp. remains at Ledi Geraru (Villmoare et al., 2015)]. The 82 stone artefacts from Lomekwi 3 are larger in width, length and thickness than those from 83 Oldowan sites. Two percussion modes have been identified (passive hammer and bipolar 84 percussion). Knappers combined battering activities and core reduction, and may have used 85 mineral blocks in different ways: as anvils, cores in order to produce flakes and/or as pounding 86 tools. The lithic artefacts were found in the same chronological and geographical contexts as 87 88 specimens allocated to Kenyanthropus platyops and are also contemporaneous with individuals attributed to Australopithecus afarensis and Australopithecus deviremeda. These 89 puzzling findings have reopened the debate on the association between lithic artefacts and 90 91 the Homo genus (Harmand et al., 2015; Lewis and Harmand, 2016), and highlighted that 92 species living long before the emergence of the Homo genus also present manipulative and 93 cognitive abilities for tool-making.

94

95 Indeed, far from the once linear perspective of hominin evolution, we currently have a bushy 96 vision of this evolutionary history, with several contemporaneous species or genera in the 97 Pliocene or Early Pleistocene (Figure 1). This points to morphological diversity during these 98 periods. The archaeological data also emphasize diversity in terms of assemblages and a non-99 linear evolution, with the presence of Oldowan and Acheulean artefacts in a same 100 archaeological site (Dana Aoule Gona, DAN 5 and Busidima North Gona, BSN 12 (Semaw et al., 101 2020)). In addition, some Oldowan sites are more recent than Acheulean sites. Therefore, it 102 would probably be a good idea to consider that cultural evolution also branches out in 103 different directions.

- 104
- 105 INSERT FIGURE 1

107 The Oldowan in the broad sense of the term (which can include the early Oldowan, developed Oldowan etc.) overlaps temporally and geographically with different hominin taxa (early 108 109 Homo, Paranthropus, and Australopithecus in eastern Africa and southern Africa (Figure 1). The oldest known Oldowan tools found in situ have been unearthed in Kenya at the site of 110 Nyayanga, older than 2.6 Ma (Plummer et al., 2023, see paragraph 2.1.3.1). In Ethiopia, the 111 Busidima Formation at Gona has yielded the largest number of early Oldowan sites (e.g., 112 113 Roche et Tiercelin, 1977, Semaw et al., 1997, Semaw et al., 2003), dating to 2.6-2.5 Ma, with the localities of EG-10, EG-12, EG-13, EG-24, OGS-6a, OGS-7 yielding over one thousand 114 excavated artefacts (e.g. Semaw et al. 1997; Semaw, 2000; Semaw, 2006; Stout et al. 2005, 115 116 Stout et al. 2010). We can also mention the site of Bokol Dora 1 dated to 2.61-2.58 Ma, Ledi Geraru (Braun et al., 2019), the site of AL 666 at 2.34 Ma (Kimbel et al., 1996; Kimbel et al., 117 1997, see paragraph 2.1.1.1), the Lokalalei complex in Kenya at 2.34 Ma (Kibunjia et al., 1992; 118 Kibunjia, 1994; Roche et al. 1999, Delagnes and Roche, 2005), and the sites from the Lower 119 Omo Valley in the member F (2.32-2.27 Ma (e.g. Omo 57/5 (see paragraph 2.1.3.2.), OmO 120 57/7, Omo 84, Omo 123j, Omo 123k, OMO 123m, Omo 1/E-3, Omo 1/E-4, FtJi1, FtJi2, and 121 122 FtJi5, Merrick et al., 1973; Chavaillon, 1976; Delagnes et al., 2011) and the site of Kanjera South in Kenya, dated to 2 Ma (e.g., Plummer et al., 1999). 123

124

Furthermore, several Oldowan sites have been also discovered in Eurasia in association with 125 126 paleoanthropological remains (Prat, 2018). We can cite Dmanisi in Georgia, where at least five hominin individuals were discovered (Lordkipanidze et al., 2007; Lordkipanidze et al., 2013), 127 in association with Oldowan artefacts (Mgeladze et al., 2011), Sima del Elefante in Spain in the 128 129 Sierra of Atapuerca (1.22 Ma), where the oldest mandible of the Homo genus in Europe was discovered (Parés et al., 2006, Carbonell et al., 2008). We can also mention the Jianshi hominin 130 site in China, also called Longguddong site (with a debatable age, minimum age of 2.14 Ma (Li 131 132 et al., 2007) or 1.52 Ma +/- 0.092-1.05 +/0.04 (ESR dating, mean age of lower layer 8 and layer 4 respectively (Han et al. 2022), with early Homo remains and nine stone artefacts in the 133 134 hominin layer (Li et al., 2017); Yanmou (1.7 Ma), early Homo sp./early H. erectus associated with four Oldowan lithic pieces (Hou et al., 2010; Zhu et al., 2008). At 1.4 Ma, in the Jordan 135 136 Valley (Israel), the site of 'Ubeidyia yielded some early *Homo erectus* specimens (one tooth) 137 associated with Acheulean artefacts (Belmaker et al, 2002; Tobias, 1966; Bar-Yosef and Goren-Inbar, 1993). 138

139

140 The oldest occurrences of Acheulean assemblages are Kokiselei 4 (1.74 Ma, Lepre et al. 2011) 141 and Kokiselei 6, where bifacial shaping emerges (Duke et al., 2021), in the western part of Lake 142 Turkana, Kenya and the Konso Gardula sites in Ethiopia. The latter comprise KGA6-A1, which 143 is the earliest site, bracketed between two tuffs, 1.74 and 1.66 Ma (Beyene et al., 2013) and more typical early Acheulean assemblages, such as for example KGA-4 and KGA-10, dated to 144 ~1.45 Ma, KGA-5, KGA-7 to ~1.4 Ma (e.g., Asfaw et al., 1992, Beyene et al., 2013, see paragraph 145 146 2.1.6.1), and FLK West in Olduvai Gorge in Tanzania dated to circa 1.7 Ma. This latter site 147 contrasts with Kokiselei 4 and Konso as it points to relatively advanced handaxe shaping skills with very symmetric bifacial flakes (Diez-Martin et al., 2015) 148

149

Regarding the *Homo* genus, the first occurrence of early *Homo* is the mandible LD 350-1, Ledi
Geraru, in Ethiopia, which could not be attributed to a specific species (Villmoare et al., 2015).
This is followed by discoveries in Kenya at Lokakalei 1 alpha KNM-WT 42718, 2.33 Ma (see
paragraph 2.1.1.2) (Prat et al., 2005), the temporal bone KNM-BC1 from Chemeron, Kenya,
dated to 2.4 Ma, Chemeron, Kenya, (Hill et al., 1992; Sherwood et al., 2002), the maxilla AL

666-1 dated to 2.34, Hadar, Ethiopia (Kimbel et al., 1996; Kimbel et al., 1997, see paragraph
2.1.1.1), the mandible UR 501 dated to ca. 2.4 Ma from Uraha, Malawi (Schrenk et al., 1993;
Bromage et al., 1995), and teeth from the Omo basin (Ethiopia) dated to 2.3-2.4 Ma (e.g.,
Howell et al., 1987; Suwa et al., 1996; Ramirez- Rozzi, 1997).

- 159
- 160

But what is the nature of relationships between hominins and the different material cultures?Were these early material cultures made by a single or several hominin species?

Here, we address two main aspects. The first is to investigate the hominin/culture associations
in the fieldwork record in the Late Pliocene and Early Pleistocene and the second is related to
the morphology underlying the manual skills of hominins during these periods.

We reflect upon the morphological and archaeological data in the literature and on 166 167 interdisciplinary research conducted in recent years. Data from the literature on hominins, archaeological sites, spatial information and context are taken at face value. However, it is 168 sometimes difficult to find information on fossil descriptions, in particular information 169 170 concerning deposition modes, contexts (in situ, on the surface) and distances from the 171 archaeological remains. In view of this difficulty, despite exhaustive research, it is possible that 172 certain associations are overlooked. Concerning the taxonomical names of hominin remains, 173 the presence of specimens named Homo ergaster or Homo erectus according to different 174 authors, are mentioned here as *Homo ergaster/erectus*. For the archaeological terminology, 175 we have used the terms Oldowan and Acheulean in a broad sense, as used in the cited paper, and we do not make any distinction between early Oldowan, developed Oldowan etc, even if 176 important differences between hominin behaviour at early sites (ca. 2.6-2.3 Ma) in 177 comparison to later ones (ca. 2.0–1.7 Ma) have been attributed to this industrial complex (e.g., 178 179 Harmand 2009a, Roche et al., 2009; Goldman-Neuman and Hovers 2012, Potts 2012). Indeed Oldowan displays considerable differences in terms of technological complexity, groups 180 181 together assemblages with disparate characteristics and has been documented in multiple raw material contexts and physical environments in Africa and Eurasia. A huge variability can 182 be seen in production methods depending on raw material choice (see for example Gallotti, 183 2018 and de Weyver, 2022 for a review). The raw material constraints (linked to the 184 185 landscape) lead to multiple technical systems of stone-tools production (e.g. free hand and bipolar techniques) which did not develop chronologically and are not continuous. Raw 186 187 material seems to be a key factor in the diversity of the Oldowan assemblages (e.g. Schick et al., 2006, Harmand et al., 2009b, Proffitt et al., 2022, Delagnes et al., 2023), 2. Overview of 188 189 archaeological data

190

191 In order to discuss the relationship between hominins and material culture, here we review 192 the eastern and southern African hominin remains from between 3.3 and 1 Ma, and place 193 their relationships into categories of reliability. This catalogue may then serve as a guide for 194 evaluating the association between hominins and archaeology (material culture), in the same 195 way as the taxonomic allocation of postcranial remains for example, by Grine et al in 2022 196 (Grine et al., 2022).

197

198 The relationships between hominins and material culture can be approached on different 199 geographical scales (regional or local).

200

201 Concerning the categories of reliability, we define a low-level confidence (LC) association 202 between archaeological and paleoanthropological remains when the hominin remains are in 203 the same chronological and regional frameworks as the archaeological sites.

- We define high confidence (HC) associations at a more local scale (Table 1). We divide this category into two subcategories depending on the distance between hominin remains and the archaeological findings.
- HC 1: corresponds to an association between lithic and paleoanthropological remains restricted to cases where hominin remains were found in the archaeological site (*in situ* during
- 209 excavations, or on the surface of the site due to erosion processes).
- 210 We divided this category into two subcategories:
- HC 1 A, when the taxonomy is clear;
- HC 1 B, when the taxonomy of the remains is debated (when it concerns for example some postcranial remains, see Grine et al, 2022).
- 214
- HC 2: when hominin remains were found outside the excavation (less than 400 metres away) but in the same chrono-stratigraphical layers. We also divide this category according to the taxonomical confidence of the paleoanthropological remains. HC2 A is when the taxonomy is clear and HC2 B when it is debated.
- 219

220 Concerning the low confidence association between archaeological and hominin remains, we 221 can cite as the 2.5 Ma Hata Member of the Bouri Formation which yielded the specimen defining the species Australopithecus garhi (Asfaw et al., 1999) and evidence of hominin 222 223 processing of large mammal remains (de Heinzelin et al. 1999), but no in situ artefacts, but are 224 sometimes associated with Oldowan site of Gona, 96 kilometres away. Concerning the 225 Lomekwian, some hominin remains were found in the paleontological collecting area LO5 (Harris et al., 1988) in the vicinity of the archaeological site LOM3 (3.3 Ma) (Harmand et al., 226 2015; Skinner et al., 2020), in the same spatiotemporal range as the hominin fossils. These 227 228 include the paratype of Kenyanthropus platyops (KNM-WT 38350) (Leakey et al., 2001) and for example, KNM-WT 64060 (Supplementary Informations in Harmand et al., 2015), a 229 230 hominin molar discovered in 2012 found 600 metres from the LOM3 site. They are deposited 231 in the same chronostratigraphic interval and are pene-contemporaneous with the LOM 3 site. 232

233

234 2.1. High confidence in the association between lithic and paleoanthropological remains in 235 East Africa

236

A high-confidence association (Table 1) between lithic and paleoanthropological remains between 3 and 1.2 Ma in East Africa is noted in 29 cases (22 cases of High Confidence 1 (hominin remains found in the archaeological site) and seven cases of High Confidence 2 (hominin remains found less than 400 metres from the excavation but in the same chronostratigraphical layers) (Table 2).

- 242
- 243 INSERT Table 1
- 244
- 245 INSERT Table 2
- 246

247 **2.1.1.** Association between Oldowan and *Homo habilis* sensu lato

247 248

The oldest evidence of lithic-hominin association between Oldowan and early *Homo* specimens comes from the region of Hadar in Ethiopia. A *Homo* aff. *habilis* maxilla (AL 666-1) was found at the site AL 666, dated to 2.34 Ma, alongside an Oldowan assemblage.

253 2.1.1.1. AL 666 (Oldowan), Ethiopia, Afar Region, AL 666-1 (*Homo* aff. *H. habilis*), 2.34 +/-0.07
254 Ma, High confidence: HC 1 A

255

In Ethiopia, in the Makaamitalu Basin (Kada Hadar, Afar Region), archaeological and hominin 256 257 remains were unearthed at the AL 666 site. Stratigraphically, they lie in the lowermost part of the Busidima Formation, underlying the 2.36 +/- 0.07 Ma BKT-3 tephra (Campisano, 2012) and 258 259 are dated to 2.34 +0.07 Ma. The density of artefacts is low over an area of 20 m2 and mostly represented by flakes. None of the artefacts were retouched (Kimbel et al., 1996, Plummer 260 2004). The raw material was carefully selected, with a high frequency of homogenous and 261 fine-grained raw materials (Goldman-Neuman and Hovers, 2012). According to the 262 discoverers (Kimbel et al, 1996), the lithic assemblage is typical of an Oldowan assemblage. 263 On the surface, at the base of the AL 666 hill, a maxilla (AL 666-1) was discovered. An 264 265 excavation was undertaken to identify the *in situ* horizon, which led to the discovery of *in situ* lithic elements but no additional hominin remains. Thirty-four stone tools including 14 266 excavated in situ were recovered from the AL 666 locality. The hominin maxilla (AL 666-1) 267 268 presents unambiguous affinities with early *Homo* (Kimbel et al., 1996) and has been assigned 269 to Homo aff. H. habilis, based on morphological and metrical comparisons (Kimbel et al. 1997). 270

- 271 2.1.1.2. Lokalalei 1, GaJh5 (Oldowan), Kenya, West Turkana, KNM-WT 42718 (early *Homo*),
 272 2.34 +/-0.04 Ma, High confidence : HC 2 A
- 273

In Kenya, in the western part of the Lake Turkana, the Lokalalei complex of sites (Kibunjia, 274 275 1994) has unearthed one hominin tooth dated to 2.34 +/-0.04 Ma. This tooth KNM-WT 42718 (Prat et al., 2005) is attributed to early *Homo* and was found at Lokalalei alpha, in the vicinity 276 of, but not strictly associated with the Oldowan Lokalalei 1 site. Indeed, this early Homo 277 specimen from Lokalalei 1alpha, is a surface finding discovered 100 m south of the 278 279 archaeological site Lokalalei 1. The duration of accumulation of the 6.2-m- thick fine sandstone 280 and claystone deposits that separate the sites Lokalalei 1alpha and Lokalalei 1 can be estimated at a time scale of 10,000 to 100,000 years. Considering the classical measurement 281 error (0.04 to 0.5 Myr) for fission track and/or K/Ar, 40Ar/39Ar ages obtained on various tuffs 282 283 in Eastern Africa, the two sites have been considered to be chronologically close (Prat et al., 2005). 284

- 285
- 286 2.1.1.3. Fj-1a (Oldowan), Fejej Region, Ethiopia, Fj-1-Hd1, -Hd3, Hd4 (*Homo* aff. *H. habilis*),
 287 1.95-1.9 Ma, High confidence: HC 2 A.
- 288

300

At the northern limit of the Koobi Fora formation, in the Omo-Turkana Basin, the region of 289 290 Fejej yielded the FJ-la archaeological site with faunal and lithic assemblages in a primary context, referred as Oldowan (e.g., Asfaw et al., 1991, de Lumley and Beyene 2004, Barsky et 291 al. 2011). Geochronological data from the sequence indicate an age of nearly 1.9 Ma for the 292 293 FJ-la archaeological level (Chapon et al., 2005). Raw materials (mainly quartz and basalt) were 294 locally available and carefully selected. The direct knapping method was longer and more complex when knapping basalt (Barsky et al. 2011). Located 180 metres from FJ-la, the locality 295 FJ-le (Unit 3) yielded three hominin dental remains belonging to the same individual (Fj-1-Hd1 296 297 lower right first molar; Fj-1-Hd3, lower right second premolar; Fj-1-Hd4, lower left third 298 molar). These remains are attributed to Homo aff. H. habilis based on their morphology and 299 dimensions (de Lumley & Marchal 2004).

2.1.1.4. DK site, Bed I (Oldowan), Olduvai Gorge, Tanzania, OH 24 (*Homo habilis*), circa 1.871.84 Ma, High confidence HC 2 A.

- DK is the oldest site at Olduvai. Artefacts and bones at DK occur in the interval between the Tuff IB and Bed I basalt lava, dated between 1.877 +/- 0.013 Ma and 1.848 +/- 0.003 Ma (Deino, 2012). It unearthed Oldowan assemblage (Leakey, 1971, de la Torre and Mora, 2005). The cranium OH 24 which corresponds to one of the most complete remain allocated to *Homo habilis* has been found 300 metres to the east to the original site (Leakey et al., 1971), but no artefacts.
- 310
- 2.1.1.5 Ewass Oldupa site, Bed I (Oldowan), Olduvai Gorge, Tanzania, OH 65 (*Homo habilis*),
 circa 1.82 Ma, High confidence HC 2 A.
- 313

In Olduvai Gorge Bed I, an almost intact maxilla (OH 65) from Geological locality 64 was 314 315 discovered 350 metres northeast of the Ewass Oldupa site (Geological Locality 63), which contains cut-marked bones and an Oldowan assemblage mainly made on quartzite. Flakes are 316 317 the dominant product (Mercader et al. 2021). All these remains are from strata dated to ~1.82 318 Ma (Blumenschine et al., 2003, Deino, 2012, McHenry, 2012). Labial striations on the anterior 319 teeth suggest that OH 65 was right-handed in manipulating objects during oral processing 320 (Frayer et al., 2016). OH 65 is assigned to the Homo genus, and has been allocated to Homo habilis (Blumenschine et al., 2012, Clarke, 2012). However, Spoor and co-authors argue that 321 its morphology is not compatible with the holotype of Homo habilis (OH 7) and lacks the facial 322 323 morphology observed in Homo rudolfensis (Spoor et al., 2015) contra Blumenschine et al 324 (2012).

325

2.1.1.6 MNK Skull, Middle Bed II (Oldowan), Olduvai Gorge, Tanzania, OH 13, OH 15 (*Homo habilis*), circa 1.67 Ma, High confidence HC 1 A

328

329 MNK Skull (Geological Locality 88, Mary Nicol Korongo, MNK, Middle Bed II) is an archaeological site where two hominins allocated to Homo habilis (OH 13, and OH 15) were 330 discovered in situ and one (OH 14) on the surface. It is dated to circa 1.67 Ma (de la Torre et 331 al., 2021). OH 13 consists of cranial vault bones, parts of the maxilla, mandible with dentition. 332 333 It is one of the paratypes of Homo habilis (Leakey et al., 1964, Tobias 1991). This individual represents the most recent specimen of *Homo habilis* in east Africa (Spoor et al., 2015). OH 15 334 335 consists of dental remains (one upper canine and two third upper molars) and is attributed to H. habilis (Tobias, 1991). Alongside these hominin discoveries, important faunal and 336 337 archaeological assemblages were unearthed by Mary Leakey (Leakey 1971). The stone tool 338 assemblage is typical of the Oldowan, with no technological elements heralding the appearance of the Acheulean, and a debitage dominated by quartizte and an overabundance 339 340 of lava cores (de la Torre et al., 2021).

341 342

343 **2.1.2.** Association between the Oldowan, *Homo habilis* and *Paranthropus boisei*

344

2.1.2.1. FLK level 22 (Oldowan), Olduvai Gorge, Tanzania, OH 5 (*P. boisei*), High confidence
HC1 A; OH 35 (*P. boisei*) or (*H. habilis*), High confidence HC 1 B), OH 6 (*H. habilis*), High
confidence HC 1 A, 1.84 Ma

348

FLK level 22 yielded a dense Oldowan assemblage (more than 255 artefacts) and mammal fossils (more than 3500) (Leakey, 1971) comprising four hominin individuals, including the cranium OH 5, the holotype of the species *Paranthropus boisei* (formerly *Zinjanthropus boisei*, Leakey, 1959). These hominin remains are allocated to *H. habilis* and *P. boisei* (Tobias, 1991). 353 One of these four hominin remains was a surface finding (OH 44, right upper first molar), one (OH 35) represents some postcranial bones (tibia and fibula (Leakey et al., 1964) with an 354 uncertain taxonomical allocation, as they are associated with cranio-dental remains, one is 355 attributed to H. habilis (OH 6, lower third premolar, right upper molar, right upper second 356 357 incisor and six calvaria fragments, Tobias, 1991), and another is OH5, the *P. boisei* holotype. The age of the Zinjanthropus surface is constrained by tuffs IB and IC, dated respectively to 358 359 1.845 +/- 0.002 Ma and 1.839+/- 0.005 Ma (Blumenschine et al, 2003; Blumenschine et al, 2012). 360

2.1.2.2. FLK NN level 3 (Oldowan), Olduvai, Tanzania, OH 7 cranial remains (*H. habilis*): High
confidence HC 1 A; OH 7 hand remains (*H. habilis* or *P. boisei*): High confidence HC 1 B; OH 8
(*H. habilis*): High confidence HC 1 B, 1.84 Ma

365

361

The paleoanthropological remains from site FLK NN level 3, unearthed in Bed I at Olduvai 366 consist of abundant Oldowan stone artefacts and six sets of paleoanthropological remains 367 368 assigned to three catalogue numbers: a) a juvenile mandible, parietals and other isolated cranial fragments (OH 7, the holotype of *H. habilis*); b) a set of 12 hand bones, also catalogued 369 as OH 7; c) 12 foot bones (OH 8); d) an isolated clavicle (OH 48); e) OH 43 two metatarsals; 370 and f) radial shaft (OH 49). The association between the cranial remains and OH 7 hand bones 371 372 seems to have been based on a parsimonious hypothesis (Grine et al., 2012), as for OH 8. 373 Moreover, some researchers have allocated these hand bones to Paranthropus (Moyà-Solà et al., 2008) (see paragraph 3). The date proposed for OH 7 is 1.84 Ma (Deino, 2012). OH 8 was 374 first reported by Day and Napier in 1964 (Day and Napier, 1964), and designated by Leakey et 375 al. in 1964 (Leakey et al., 1694) as one of the paratypes of Homo habilis. 376

377

2.1.2.3. FxJj 38 (Oldowan), Koobi Fora, Kenya, KNM-ER 1805 (*Homo* sp.?), High confidence HC
1 B, and KNM-ER 1806 (*P. boisei*), High confidence HC 1 A, 1.78 and 1.52 Ma or 1.85 Ma

380

381 Two hominin remains were discovered in the Koobi Fora Region, in area 130. The first one (KNM-ER 1805) was found in situ in deposits stratigraphically located below the Okote Tuff 382 383 complex and consists of a cranium and associated mandible. It has been allocated to early Homo (Homo sp., Leakey 1974), but also to other species and genera (see Prat, 2002). It 384 displays a number of unique features differentiating it from other hominins, and should not 385 be considered as a standard Homo habilis male (Prat, 2002). Fifteen metres from this 386 specimen, on the surface, a large mandible (KNM-ER 1806) allocated to P. boisei was 387 discovered (Leakey 1974). Excavations at these two localities unearthed faunal remains, as 388 well as a light scatter of stone artefacts, but no additional hominin remains. Forty metres east 389 390 of these paleoanthropological localities, on the same stratigraphical horizon, lies the archaeological site FxJj 38 North West. The composition of the lithic assemblage, with a large 391 chopper component, is consistent with that of Olduvai Bed I and therefore allocated to the 392 393 Oldowan (Harris & Herbich, 1978) and represents one of the type localities for the Developed 394 Oldowan of Koobi Fora (Mana et al., 2019). These hominin remains and archaeological site belong to the KBS Member, below the base of the Okote Member and above the KBS Tuff. 395 These sites are, according to Mana and co-authors, broadly constrained between 1.78 and 396 397 1.52 Ma (Mana et al., 2019). This is in contradiction with the interpretation of Feibel and 398 collaborators who estimated its age at 1.85 Ma (Feibel et al., 1989). 399

400 **2.1.3.** Association between Oldowan artefacts and *Paranthropus (P. aethiopicus, P. boisei)* 401

402 2.1.3.1. Nyayanga, excavation 3 (Oldowan), Lake Victoria, Kenya, KNM-NG 77316 (*P. boisei*):
403 High confidence: HC 1 A, 3.03 -2.58 Ma

404

In Kenya, the site of Nyayanga, on the southern margin of the Winam Gulf of Lake Victoria has 405 406 revealed one of the oldest known Oldowan assemblages, dated between 3.032 Ma and 2.581 Ma (Plummer et al., 2023). This site yielded 135 artefacts in situ (excavations 3 and 5) and 195 407 on the surface, as well as bones, including cut-marked bones. The flakes were removed from 408 cores using unifacial, bifacial and multifacial reduction, comparable to other Oldowan 409 localities. The use wear on the lithic material, as well macro and micro traces, indicate the 410 processing of plant tissue and faunal remains. In excavation 3, one fragment of a left lower 411 molar (KNM-NG 77316) was found in situ, spatially associated with Oldowan artefacts, and 412 413 one left upper molar (KNM-NG 77315) came from the surface (see paragraph 5). These two 414 teeth, based on the morphology and metrics are assigned to the *Paranthropus* genus.

415

416 2.1.3.2. Omo 57-5 (Oldowan), Lower Omo valley, Ethiopia, Omo 57/5-1972-371 (*P.* aff. *P.*417 *aethiopicus*): high confidence: HC 1 A. 2.32 Ma to 2.23 Ma

418

In Ethiopia, in the Lower Omo Valley, at the archaeological locality Omo 57-5 (Member F of the Shungura Formation), the excavation of 40 m2 unearthed four fragments of hominin teeth, associated with a quartz flake found *in situ* and 16 artefacts (cores and flakes) on the surface (Coppens et al., 1973, Chavaillon, 1976). One of these paleoanthropological remains (Omo 57/5-1972-371, a fragment of a left lower second premolar) has been assigned to *P*. aff. *aethiopicus* based on its morphology (Coppens et al. 1973, Coppens, 1977, Suwa 1990, Wood and Leakey 2011).

Another hominin dental remain (Omo 57/5-1972-319), a left upper first premolar, is considered to be indeterminate (White, 1988) or an indeterminate non-robust hominin (Suwa 1990), or allocated to the *Homo* genus (aff. *Homo* sp., Wood and Leakey 2011). According to de la Torre in 2004, the Omo 57 locality provides clear evidence of flaking activities showing ordered and systematic knapping processes, and does not derive from the fluvial deposition of natural fragments. The lithic assemblage is assigned to the Oldowan (de la Torre, 2004). The Member F deposits are dated to 2.32 Ma to 2.23 Ma (McDougall and Brown, 2008).

433

434 2.1.3.3. Kokiselei 1, FxJh 6 (Oldowan), Kenya, KNM-WT 37744, KNM-WT 37747, KNM-WT
435 37748, KNM-WT 37100 (*P. boisei*): high confidence: HC 1 A. 1.79 Ma

436 The Kokiselei 1 site (FxJh 6) is part of the Kokiselei complex, located in the upper part of the Kaitio member, between the Koobi Fora and KBS tuffs. This site is dated to 1.79 Ma. The 38 437 438 m2 excavation yielded more than 248 faunal remains (Brugal et al., 2003) and 1100 artefacts allocated to the Oldowan (Roche et al., 2003, Arroyo et al., 2020). On the surface of the 439 440 excavation, on the same square, four complete teeth [left upper canine (KNM-WT 37744), left 441 upper first molar (KNM-WT 37747), right upper third molar (KNM-WT 37748), left lower third molar (KNM-WT 37100)] and two molar fragments were discovered in 1997 and 1998. A left 442 443 upper first premolar and a left lower third molar from the same area were collected in 1986 444 and 1987. Based on their morphology and dimensions, all the teeth are attributed to P. boisei 445 (Prat et al., 2003).

446

447 2.1.3.4. FxJj 20 (Oldowan), Koobi Fora, Kenya and KNM-ER 3230 (*P. boisei*): High confidence:
448 HC 1 A. 1.65 +/- 0.05 Ma

At FxJj 20 East, in Koobi Fora Area 131, a hominin mandible with complete dentition (KNM-ER 450 3230) was recovered in situ from the site excavated by J.W.K. Harris in 1975 in tuffaceous 451 floodplain silts of the Okote Tuff complex (Leakey 1976; Harris and Herbich, 1978). Based on 452 the morphologies and metrics of the mandible and teeth, KNM-ER 3230 has been assigned to 453 Paranthropus boisei (Leakey 1976; Leakey and Walker, 1985; Wood, 1991) and dated to 1.65 454 455 +/- 0.05 Ma (Feibel et al., 1989). According to Harris and Herbich, 1978, a low-density scatter 456 of bones and lithic artefacts were found on the same horizon as the hominin, and the vertical distribution of the lithic assemblage shows that the hominin mandible was stratigraphically 457 458 located approximately 10-15 centimetres above the main artefact-bearing horizon (Harris and 459 Herbich, 1978). It should be noted that despite this association, Richard Leakey did not want 460 to associate the two in his 1976 paper. We can wonder whether Richard Leakey would have 461 interpreted the finds in this way if the remains had belonged to the Homo genus (see 462 paragraph 5).

463

2.1.3.5. Chesowanja GnJ1/6E (Oldowan), Chesowanja, Kenya, KNM-CH 304 (*P. boisei*): High
confidence: HC 2 A; Chesowanja GnJ2 (Oldowan), Chesowanja, Kenya, KNM-CH 1, KNM-CH
302, KNM-CH 304 (*P. boisei*): High confidence: HC 2 A 1.42+/-0.07 Ma

467

468 At Chesowanja, near Lake Baringo in Kenya, in the Chemoigut Formation layers, remains 469 allocated to P. boisei were found close to an Oldowan assemblage (Carney et al., 1971; Bishop 470 et al. 1975; Gowlett et al. 1981). The hominin remains consist on a right hemi-face and anterior 471 cranial base (KNM-CH 1); dental fragments (KNM-CH 2) and cranium fragments (KNM-CH 304). 472 The cranium KNM-CH 304 consists of five cranial vault fragments labelled A-E and is allocated 473 to *P. boisei* (Gowlett et al. 1981). It has been found 15 metres from the site GNJ1/6E (40 m² 474 excavated) where 1000 Oldowan artefacts were found in situ with limited vertical distribution. 475 The other specimens also allocated to P. boisei were found in GnJ2, where artefacts scatters occur on the surface, but only one rewarding in situ occurrence (Carney et al., 1971; Bishop et 476 477 al. 1975; Gowlett et al. 1981). According to these authors, this association must be considered to be more than coincidental (see paragraph 5). The basalt member at the base of the 478 479 Chesowanja Formation is dated at 1.42+/-0.07 Ma.

480

2.1.3.6. BK site level 4 (Oldowan), Olduvai Gorge, Tanzania, OH 80 (*P. boisei*): High confidence:
HC 1 A. 1.34 Ma

483

484 At Olduvai Gorge, in Tanzania, in Bed II, the BK Site yielded a partial Paranthropus boisei skeleton (Domínguez-Rodrigo et al., 2013). OH 80 was found in situ in level 4, which is 485 characterized by a large Oldowan assemblage and a huge number of vertebrate fossils, some 486 of which bear butchery marks (Domínguez-Rodrigo et al., 2009). The specimen comprises four 487 488 postcranial bones (distal part of a left humeral diaphysis, proximal half of a right radius, much of the right femoral shaft and part of a tibial midshaft) in direct association with nine upper 489 teeth (left and right first and second incisors, right first premolar, left second premolar, left 490 491 first molar, right second incisor). Based on their morphology, dimensions and section properties, these remains are allocated to P. boisei (Dominguez-Rodrigo et al., 2013). It is the 492 first partial skeleton attributed to this taxon and is dated to 1.34 Ma. 493 494

495 **2.1.4. Association between Oldowan and Homo erectus/ergaster**

497 2.1.4.1. PTK site (Oldowan), Olduvai Gorge, Tanzania, OH 86, (*H. erectus*-like): High
498 confidence: HC 1 B. 1.84 Ma

499

At the PTK (Philip Tobias Korongo) site at Olduvai Gorge in the third level, a manual proximal 500 501 phalanx (OH 86) was unearthed (Domínguez-Rodrigo et al., 2015) alongside a rich Oldowan assemblage (Domínguez-Rodrigo et al., 2015). OH 86, dated to 1.84 Ma, represents the 502 earliest known modern-human like hand anatomy. This specimen contrasts in size and shape 503 not only with Australopithecus or Paranthropus but also with the OH7 hand. Taxonomically 504 505 this specimen is assigned to the Homo genus and differs from OH 7 (H. habilis holotype). 506 However, it is difficult to confidently assign OH 86 to this latter taxon due to the lack of 507 associated craniodental remains and comparative material securely attributed to H. 508 erectus/ergaster. However, Díez-Martín and co-authors considered OH 86 as an H. erectus-509 like hominin in their paper (Diez-Martín et al, 2015).

- 510
- 2.1.4.2. Naiyena Engol 1, FxJh 5, (Oldowan), West Turkana, Kenya, KNM-WT 37745 (aff. *H. ergaster/H. erectus*): High confidence: HC 1 A, 1.75 +/- 0.05 Ma
- 513

The site of Naiyena Engol 1 (NY1 1.75+/-0.05 Ma) is a large site (artefacts collected over a surface of 585 m2 and from the 55 m2 excavation) which has yielded a considerable collection of artefacts (more than 2400 plotted pieces), corresponding to an Oldowan assemblage (Roche et al., 2003). On the surface of the excavated area, a right lower first premolar (KNM-WT 37745) was discovered. Based on its morphology and metrics, this specimen is assigned to *Homo* aff. *H. ergaster* (Prat et al., 2003).

520

2.1.4.3. Melka Kunture Garba IV (level E) (Oldowan), Melka Kunture, Ethiopia, MK 81 GAR IVE
0043 (early *Homo erectus/ H. ergaster*). High confidence: HC 1 A. Slightly older than 1.7 Ma

The Garba IV mandible (MK 81 GAR IVE 0043) from Melka Kubture was found in situ in 1981 524 during the excavation of level E, while opening an area of 2 square metres. Based on its 525 stratigraphical position immediately below the "Grazia tuff" and its normal polarity, this level 526 527 should be around or slightly older than 1.7 Ma (e.g., Morgan et al., 2012, Tamrat et al., 2014, Gallotti and Mussi, 2015). Layer E yielded 504 unworked lithic objects and 718 artefacts. The 528 529 lithic assemblage from Garba IV level E was deposited during a relatively short lapse of time. It is allocated to the Oldowan and shows that hominins had good knowledge of raw material 530 531 suitability (obsidian being the most frequently exploited raw material) (Gallotti and Mussi, 2015). The mandible belongs to a child whose age at death was estimated between 3 and 4 532 years (Condemi, 2004) or between 2.5-3.5 years old based on the tooth development pattern 533 534 (Zanolli et al., 2016). It is allocated to early Homo erectus/Homo ergaster (Condemi 2004). The inner dental morphology also suggests that the Garba IV mandible is a robust early Homo 535 536 specimen (Zanolli et al., 2016).

537

2.1.4.4. Melka Kunture Gombore I (Oldowan), Melka Kunture, Ethiopia, Gombore IB-7594
(*Homo* sp. or *H. erectus*): High confidence: HC 1 B. 1.5-1.6 Ma

540

541 In the Melka Kunture site complex in Ethiopia, a well-preserved distal portion of a left humerus 542 (Gombore IB-7594, formally Melka Kunture 3, or MK3) was discovered in 1976 during the 543 excavation of the site of Gombore I, directed by J. Chavaillon (Chavaillon et al., 1977). This 544 postcranial hominin remain was found in the southern sector of the excavation *in situ* in unit 545 2 of level B, which includes rich faunal and Oldowan assemblages with a predominance of 546 pebble tools, mainly side choppers (Chavaillon and Berthelet, 2004). They are dated to 1.6-1.5

Ma (Morgan et al., 2012; Tamrat et al., 2014, Cazenave et al., 2017). Gombore IB-7594 was 547 548 referred to Homo sp. (Chavaillon et al., 1977), but based on morphometric affinities, Lague and Jungers considered that this specimen could plausibly belong to Paranthropus boisei 549 550 (Lague and Jungers, 1996). However, according to the majority of authors, this specimen is part of the Homo genus (Senut, 1979). Coppens considered that it was too robust to be Homo 551 habilis and referred it to Homo erectus (Coppens, 2004). It is also allocated to Homo sp. based 552 on geometric morphometrics (Di Vincenzo et al. 2015). The study of endostructural 553 554 organization shows that this robust Homo aff. erectus humerus is different to Paranthropus 555 and close to the modern human humerus (Cazenave et al., 2017). 556

557
558 **2.1.5.** Association between the Oldowan, Acheulean and early Homo erectus/Homo ergaster

(early *H. erectus/H. ergaster*), High confidence: HC 1 A. 1.6-1.5 Ma

- 559 560
- 2.1.5.1. DAN 5 (Oldowan and Acheulean), Dana Aoule Gona, Afar Region, Ethiopia, DAN5/P1
- 561 562

In Ethiopia, the Dana Aoule Gona localities in the Afar regions have yielded some hominin 563 remains in association with both Oldowan and Acheulean assemblages, and also some cut-564 marked bones (Semaw et al. 2020). This is the first and unique occurrence of such an 565 association (hominins plus Oldowan and Acheulean) in East Africa. The age of the DAN 5 566 hominin and lithic assemblage is close to ~1.6 to 1.5 Ma. The well-preserved hominin cranium 567 DAN5/P1 was discovered in the DAN 5 locality, with the occipital and left maxilla found in situ 568 569 along with six manuports and one Oldowan core. Other cranial parts and also freshly eroded 570 Oldowan and Acheulean artefacts were collected on the surface of the DAN5 hominin site. 571 Oldowan and Acheulean assemblages were also discovered in the same general stratigraphical 572 context, ~140 m south-southwest (DAN5-South) and ~50 meters west-northwest (DAN5-573 West). The assemblages are composed of unshaped cores, Oldowan and Acheulean (large 574 cutting tools) productions. Based on morphological and metrical data, this cranium is allocated to Homo erectus (Semaw et al., 2020, Baab et al., 2022, Bruner et al., 2023), and shows 575 576 similarities to some Homo erectus crania [Dmanisi (1.77 Ma, Georgia), KNM-ER 42700 (1.6-1.5 577 Ma, Kenya) and KNM-OL 45500 (0.95 Ma, Kenya)] and some differences to Asian Homo 578 erectus.

579

2.1.5.2. BSN12 (Oldowan and Acheulean), Busidima North Gona, Afar Region, Ethiopia,
BSN12/P1 (early *H. erectus/H. ergaster*), High confidence: HC 1 A. 1.26 +/-0.03 Ma

582

583 At BSN12, almost 6 km southwest of DAN5, another hominin (BSN12/P1) was unearthed in direct association with both Oldowan and Acheulean assemblages (Semaw et al. 2020). It is 584 also associated with the Boolihinan Tuff, which is dated, based on geochemical affinities, to 585 1.262 +/-0.034 Ma. The tephra was found cemented on the BSN12/P1 parietal and both 586 587 Oldowan and Acheulean artefacts. The BSN12/P1 hominin also includes some parts of the right orbital margin and frontal squama. Based on its anatomy and dimensions, this robust 588 589 hominin shares Homo erectus features (Semaw et al., 2020, Baab et al., 2022) and is similar to Indonesian and Eastern Asia hominins, as well as to OH 9 and BOU-VP-2/66 (Middle Middle 590 591 Awash, Ethiopia) and UA-31 (Buia, Eritrea).

592

593 **2.1.6.** Association between the Acheulean, *Homo erectus* and *P. boisei*

594

2.1.6.1. KGA 10-1 (Acheulean), Konso-Gardula, Ethiopia, KGA10-506, KGA10-525, KGA10-565,
KGA10-570, KGA10-900, KGA10-1455, KGA10-1720, KGA10-2705 (*H. erectus*): High confidence
HC 1 A; KGA10-2741 (*P. boisei*): High confidence: HC 1 A: 1.4 Ma and KGA 4E (Acheulean), *Homo* sp., High confidence: HC 1 A. 1.44 Ma

599

The first evidence of Acheulean stone tools occurs at 1.76 Ma in the Kokiselei 4 site (Lepre et al. 2011) with no associated hominin remains. The oldest association between *Homo erectus* (KGA 10-1) and an Acheulean assemblage is from Konso-Gardula, dated to 1.4 Ma (Asfaw et al. 1992, Beyene et al., 2013; Beyene et al., 2014). This association have been shown for example in the locality KGA 10, with the evidence of *H. erectus* remains (Suwa et al., 2007) (KGA 10-1 (left mandible with P4-M3, KGA10-620 (right parietal fragment), KGA 10-656 (right parietal and frontal fragment); 1.42 Ma) found less than 50 meters from the excavation of 607 KGA10-A10 which yielded Acheulean picks and cores, and 300 meters west from KGA10-A6 (surface collection, Acheulean cores, handaxes and picks) (Beyene et al., 2014) and at the 608 locality KGA4-A1 with a third upper molar (surface data, KGA4-14, Homo sp, 1.44 ± 0.05 Ma). 609 In the locality KGA10, nine P. boisei were also discovered alongside dry grassland fauna (Suwa 610 611 et al., 1997) (KGA10-506, left maxilla with teeth; KGA10-525, partial skull; KGA10-565, right 612 upper molar; KGA10-570, mandible, KGA10-900, molar fragments; KGA10-1455, left parietal; KGA10-1720, left lower M3; KGA10-2705, right lower M2; KGA10-2741, molar fragments) less 613 than 500 meters from the in situ Acheulean site KGA10-10 and 300 meters from KGA10-A6 614 (surface collection, Acheulean cores, handaxes and picks) (Beyene et al., 2014). Some cut-615 marked and modified bones were also unearthed in this locality (Echassoux, 2012). At KGA13-616 A1, five kilometres from these sites, a bone handaxe was also discovered (Sano et al. 2020). 617 KGA 10 lies stratigraphically between tuffs dated to 1.41 +/0.02 Ma and 1.43 +/- 0.02 Ma. The 618 619 abundance of Acheulean artefacts contrasts with layers bearing H. erectus and/or P. boisei in 620 the Omo-Turkana basin with no Acheulean artefacts.

621

622 **2.2. The case of southern Africa**

623

It should be noted that the depositional and conservation modes of fossil and lithic remains differ between East Africa (open-air sites) and South Africa (karsts). The confidence index (High Confidence) used for remains from East Africa cannot be applied to karstic deposits and the dating of karstic members is controversial. As a result of this depositional mode, even if remains are found in the same stratigraphical member, it is difficult to infer strict contemporaneity between lithic and paleoanthropological remains.

However, some lithic assemblages, and bone artefacts have been discovered in association
with individuals allocated to *P. robustus* and early *Homo* specimens. Sometimes both taxa
were found in the same levels, but sometimes only *P. robustus* remains were discovered.

633 Here, we present some examples. At Sterkfontein, the Member 5 East lower (Member 5 Oldowan infill), dated to around 2.18 ±0.21 Ma (Granger et al., 2015), yielded remains 634 allocated to P. robustus, as well as 3513 lithic artefacts all attributed to the Oldowan (Kuman 635 and Clarke, 2000; Kuman, 2009). On the other hand, in the member 5 West (Sterkfontein 636 637 Acheulean infill), dated between 1.3 and 1.1 Ma (Herries and Shaw, 2011), an Acheulean assemblage was found in association with a bone tool (n=1) and Homo remains (Kuman and 638 639 Clarke, 2000). At Drimolen Main Quarry (MNQ), recently dated to c. 2.04-1.95 Ma (Herries et 640 al., 2020), some early Homo and P. robustus remains, as well as Oldowan stone artefacts and bone tools (n=14) were discovered (Keyser et al., 2000; Moggi-Cecchi et al., 2010; Stammers 641 642 et al., 2018; Backwell and d'Errico, 2008). A similar discovery of six early Homo remains, 113 643 P. robustus, an Oldowan assemblage and bone tools (n=32) (Brain, 1993; Brain and Shipman, 644 1993; d'Errico and Backwell, 2003; Backwell and d'Errico, 2003; Caruana, 2017) was made in the Member 1 of Swartkrans, dated between 2.4 and 1.8 Ma (Pickering et al., 2019). In the 645 Member 2 of Swartkrans, dated to c. 1.4 Ma (Balter et al., 2008), an Acheulean assemblage 646 647 with bone tools (n=11) and both Homo and P. robustus remains (Brain, 1993; d'Errico and 648 Backwell, 2003; Kuman, 2007) were unearthed. In the Member 3 of Swartkrans, dated to circa 0.96 Ma (Gibbon et al., 2014), only one hominin species (P. robustus) was found in association 649 with an Acheulean assemblage and bone tools (n= 41) (Brain, 1993; d'Errico and Backwell, 650 2003; Kuman, 2007). A similar discovery was made at Cooper's D, dated between 1.4 and 1.0 651 652 Ma, in the form of seven P. robustus remains, 49 stone artefacts [not allocated to a specific industry for the moment due to the small number of artefacts (Sutton et al. 2017)] and a bone 653 654 tool (n=1) (Steininger et al., 2008; de Ruiter et al., 2009, Hanon et al., 2021). 655

These examples support the hypothesis that *P. robustus* probably made and/or used stone
and bone tools (e.g., Sterkfontein, Member 5 East lower (Member 5 Oldowan infill), 2.18 ±0.21
Ma, Oldowan; Coopers'D, 1.4-1 Ma, indeterminate for the moment; Swartkrans Member 3, c.
0.96 Ma, Acheulean). The attribution of artefacts to *Homo* rather than *Paranthropus* or both
taxa appears once again to be more ideological than scientific.

661

662 **3. Overview of anatomical/morphological data**

663

The manipulative abilities linked to the evolution of increasingly complex tool technologies in the hominin lineage (Marzke, 1997) can be defined as pad-to-pad precision grasping *sensu* Napier (1956) and Almécija and Alba (2014), and in a broader sense, forceful precision grips and precision handling *sensu* Marzke (Marzke, 1997). Morphological evidence (although limited) suggests that this anatomical ability to use or even make stone tools is not exclusive to the *Homo* genus and is also observed in two different hominin genera: *Australopithecus* (*afarensis, africanus, sediba*) and *Paranthropus* (*robustus* and *boisei*).

671

672 Regarding the Pliocene hominins, no post-cranial element has been allocated to K. platyops, 673 a species in the same temporal and geographical framework as the site of Lomekwi 3 (Harmand et al., 2015). Conversely, a lot of hand bones, including phalanges, carpal and 674 metacarpal bones, attributed to Australopithecus afarensis are also attributed to the same 675 chronological framework as the site of LOM3. They mainly come from the Hadar localities in 676 Ethiopia (e.g., AL 333/333w, AL 438, AL 444, AL 734) (Bush et al., 1982; Drapeau et al., 2005; 677 Johanson et al., 1982; Ward et al., 2012). Different functional interpretations have been put 678 forward for this taxon, based on wrist and hand morphology, as well as intrinsic hand 679 680 proportions. According to Marzke (Marzke, 1983, 1997, 2013, Tocheri et al. 2003), A. afarensis individuals were able to use pad-to-side as well as three-jaw chuck precision grips and 681 682 precision handling (probably less effective precision handling and power squeeze grips). These manipulative abilities (forceful precision grips and precision handling), are required for stone 683 684 tool-making and activities and are considered to be specific to the human hand. They may 685 have been used to apply pressure to tools in activities such as cutting, digging, pounding, 686 chopping or stone-throwing (Marzke et al., 1997). For some authors, A. afarensis is mainly characterized by human-like hand functions, illustrated for example by modern human-like 687 manual proportions with a relatively long thumb and short medial digits (e.g., Alba et al., 2003; 688 Almécija and Alba, 2014, Drapeau et al., 2005; Ward et al., 2012). However, Rolian and Gordon 689 690 (Rolian and Gordon, 2013) suggest that the manual proportions of A. afarensis are intermediary between extant humans and gorillas, and deduce that A. afarensis did not 691 possess the efficient precision grip typical of modern humans. In 2022, Kivell and co-authors 692 693 stressed the importance of acknowledging that precision-grasping abilities in A. afarensis could have evolved in response to manipulative behaviours not linked to lithic tools (e.g., food 694 processing), even if evidence of tool-using behaviours (McPherron et al., 2010) at Dikika 695 696 (Ethiopia), close to an A. afarensis juvenile specimen Dikika 1, and the discovery of the LOM3 697 (Harmand et al., 2015) suggest that stone tool and stone use behaviours may have been part of the behavioural repertoire of A. afarensis (Kivell et al. 2022). 698

Evidence from other non-*Homo* Pliocene hominins, from southern Africa must also be scrutinized. Several phalangeal, carpal and metacarpal remains from the site of Sterkfontein are mostly assigned to *Australopithecus africanus*. Specimens from the Member 4 deposit (recently dated to 3.4 Ma; Granger et al., 2022) indicate that *A. africanus* was able to use a human-like pad-to-pad precision grasp (Green and Gordon, 2008), pad-to-side and handling grips (Marzke, 1997, Ricklan, 1987), but with not the full range of human-like manipulative behaviours observed in later hominins (Green and Gordon, 2008). Furthermore, according to some authors (Kivell et al. 2020), the proportions within each ray of metacarpal and phalange
length are also like those of modern humans. Moreover, based on the internal structure of
metacarpals, Skinner and co-authors (Skinner et al., 2015) confirm that *A. africanus* would
have had the forceful human-like opposition capacity of the thumb and fingers during power
and precision grips.

Still in South Africa, two partial skeletons belonging to the most recent Australopithecus 711 712 species: A. sebiba (Berger et al., 2010), discovered at Malapa and dated to 1.977 Ma (Pickering et al., 2011), indicate that these individuals possessed pad-to-pad precision gripping and hand 713 714 proportions showing enhanced dexterity (Kivell et al., 2011), but with limited force of the 715 thumb. Furthermore, the phalangeal morphology suggests powerful flexion and strong 716 grasping of the fingers. These functional interpretations are also confirmed by recent studies 717 using metacarpal internal trabecular bone analyses (Dunmore et al., 2020) and 718 musculoskeletal modelling (Bardo et al., 2018).

719

720 After 2 Ma, the morphology of the Olduvai and Swartkrans hominin hands discovered in 721 archaeological levels (see paragraph 3) are compatible with tool using and tool-making. For 722 the Olduvai remain (OH 7), human-like precision grips are associated with pad-to-pad precision grasping. Indeed, in the publication defining the species *Homo habilis*, based on the 723 724 OH 7 late juvenile hand bones uncovered at site FLKNN I in Olduvai Gorge (Tanzania), Leakey 725 and co-authors emphasized a well-developed and fully opposable thumb (pollex) and a hand 726 capable of a power grip, as well as a simple and usually well-developed precision grip (Leakey 727 et al., 1964). Hence the OH7 hand has played a crucial role in our perception and 728 understanding of the evolution of hominin tool use and tool-making and therefore on the 729 definition of the *Homo* genus. However, it should be noted that in the initial description in 730 1962, Napier (Napier, 1962) highlighted a mosaic morphology (e.g., modern human-like morphology of the distal pollical phalanx and ape-like carpus and curvature of the phalanges), 731 732 and observed that the thumb was opposable and powerful. In terms of functional 733 interpretation, he emphasized "On anatomical grounds there is no doubt that the Olduvai 734 hand was sufficiently advanced in terms of the basic power and precision grips to have used 735 naturally occurring objects as tools to good advantage. There is less certainty about 736 toolmaking which involves not only a peripheral but also a central intellectual factor as Oakley 737 has long insisted" (Napier, 1962, p. 411). This interpretation contrasts somewhat with that of Leakey, Tobias and Napier 1964 which unlights that "While it is possible that Zinjanthropus 738 739 and Homo habilis both made stone tools, it is probable that the latter was the more advanced 740 tool maker .. (Leakey et al. 1964, p. 9) (see paragraph Discussion and conclusions).

Furthermore, the taxonomical allocation of the OH7 hand has been questioned, as the 741 phalangeal morphologies show some similarities with *Paranthropus* (Moyà-Solà et al., 2008). 742 743 Based on functional analyses, those authors (Moyà-Solà et al., 2008) suggested that the anatomical features linked to human-like precision grasping (previously considered as tool-744 making features by some authors, may be conversely adapted to specialized manual feeding 745 746 techniques in Paranthropus. Furthermore, according to Susman and Creel (Susman and Creel, 747 1979), the mosaic of features observed in the anatomy of OH7 could reflect precision grip abilities (e.g., the modern human-like features of distal phalanges) and frequent climbing (e.g., 748 749 African ape-like features).

750

At the PTK (level 3) site at Olduvai Gorge level 3, a manual proximal phalanx (OH 86) dated to 1.84 Ma (Domínguez-Rodrigo et al., 2015) represents the earliest modern human-like hand anatomy (see paragraph 2.1.4.1). This specimen is assigned to the *Homo* genus and contrasts in size and shape not only with *Australopithecus* or *Paranthropus*, but also with the OH7 hand. The phalangeal features are indicative of more relaxed postural and locomotive selective pressures on the hand and a marked reduction in manual phalangeal arboreal adaptations(Domínguez-Rodrigo et al., 2015).

Similarly, adaptations for tool-using and tool-making are observed in specimens allocated to 758 the Paranthropus genus from the lleret and the Swartkrans hominins. Indeed, the hand 759 760 anatomy of P. boisei specimens (e.g., based on the best Paranthropus hand evidence of the specimen KNM-ER 47000 from Ileret, East Turkana, Kenya, dated to 1.5 Ma), highlights that 761 762 this species did not possess the robust thumb observed in Homo erectus (linked to the intensification of forceful precision grips and tool use) but exhibits the manual dexterity 763 764 required to use or make stone tools (Richmond et al., 2020). According to Moyà-Solà and co-765 authors (who consider that the OH 7 hand bones belong to Paranthropus boisei), 766 Paranthropus individuals exhibit more robust hand bones than Homo and lack the 767 disproportionately hypertrophied first metacarpal characteristic of the Homo genus (Moyà-768 Solà et al., 2008).

769

770 In southern Africa, according to work by Susman (Susman, 1988, 1994, 1998), the remains 771 (phalanges and metacarpals) discovered mainly in Member 1 of the Swartkrans site and possibly but not securely allocated to P. robustus, as early Homo specimens were also found 772 773 in this member, show that these hominins possessed muscles allowing for human-like precision grasping (by stabilizing the opposing the thumb and the fifth finger to the other 774 fingers), which are considered as adaptations for tool-using and tool-making (Susman, 1994, 775 776 1998). For Moyà-Solà and co-authors (Moyà-Solà et al., 2008), the Paranthropus pattern 777 probably reflects an adaptation to manual activities other than tool-making more related to 778 feeding behaviour such as specialized food gathering and processing, as observed in extant 779 hominids (Gérard et al., 2022; Marzke et al., 2015; Neufuss et al., 2018).

780

Lastly, the recent description of a third metacarpal (Kaitio area, West Turkana, Natoo Member,
1.42 Ma) from West Turkana, with a styloid process typical of modern humans, seems to
confirm that enhanced hand function for tool-making and tool-using took place very early on
in the evolution of the *Homo* genus (Ward et al., 2014).

785

786 4. New approaches, how can we go beyond anatomical and archaeological field data?

787 **4.1. Experimental biomechanics and musculoskeletal modelling**

788

The first of the two last points of this review concerns experimental simulations and in particular the required biomechanical capacities and manual skills for percussive movements. Indeed, in addition to the archaeological and paleoanthropological framework, experimental biomechanics and musculoskeletal modelling can shed light on the functional anatomy of the upper limb involved in stone tool production (e.g., Rolian et al., 2011; Tocheri et al., 2003; Key et al., 2019, Kivell et al. 2022).

795 Macchi and co-authors show that muscular demands are similar for passive hammer, bipolar, 796 and free-hand percussion techniques, but that kinematic patterns are more variable. It is 797 interesting to note that free-hand percussion is differentiated from the other two techniques, 798 with more complex motor control demands (Macchi et al., 2021). Furthermore, in order to 799 evaluate the prehensile abilities of a contemporary hominin from the Lomekwi site, a musculoskeletal model of the hand of Australopithecus afarensis was developed by Domalain 800 801 and co-authors (Domalain et al., 2017). They proposed two versions of the Australopithecus afarensis anatomical model based on Homo sapiens and Pan troglodytes. Their results suggest 802 803 a limited capacity of the Australopithecus afarensis hand to bring the pulp of the fifth ray into contact with the surface of a large object. This may have been a limiting factor in exerting a
 large enough grip force to make large lithic artefacts such as those found at Lomekwi 3.

The biomechanics of free-hand knapping (techniques used to produce the Oldowan) received 806 more attention, but are quite limited concerning the hand (Kivell et al., 2022). A wide variety 807 808 of grips were used, with significant emphasis on precision grips (Key et al., 2018; Williams-Hatala et al., 2021). Loads tend to be concentrated on the thumb (pollex) in both the dominant 809 and non-dominant hands, which is probably the best candidate for a manual adaptation for 810 this specific behaviour (Rolian et al., 2011; Williams-Hatala et al., 2018; Williams-Hatala et al., 811 2021; Key & Dunmore, 2015). As summarized by Kivell et al. (2022), the osteology and 812 813 muscular anatomy of the thumb are essential for understanding the adaptative response to stone tool production and use, but we must also consider the fact that different combinations 814 of derived characters can occur at different times and the problems of equifinality with respect 815 816 to behaviour. Furthermore, more data are needed on EMG, kinematics, kinetics of the digits 817 and all the upper limb during free-hand, bipolar and passive hammer knapping. Moreover, the 818 application of new techniques (Manafzadeh & Gatesy, 2021), such as X-ray reconstruction of 819 moving morphology (bi-planar X-ray with X-ROMM), can provide new insights into the 820 reconstruction of joint mobility.

821

822 4.2. Cognitive exploratory studies

823

824 The final section concerns the cognitive abilities of these hominins and the resulting 825 organization of the brain. The artefacts from LOM 3 indicate complex cognitive behaviour with 826 extensive planning and motor control, which may involve the reorganization and/or expansion 827 of several regions of the cerebral cortex (visual and motor) and cerebellum. The findings from 828 the Lomekwi 3 site suggest that the reorganization and/or expansion of these brain regions 829 must have occurred before 3.3 million years ago. This was previously suggested by Ralf 830 Holloway as early as 1983 (Holloway, 1983) on the basis of the endocranial anatomy of the specimen found at Hadar AL 162-28, attributed to A. afarensis. 831

Understanding the evolution of higher order cognitive functions (such as planning through tool-making) requires extrapolating from what we know of modern human and non-human primates to fossil and archaeological data. The link between brain evolution and cognition remains a key scientific challenge, because it is impossible to observe the functional brain activity of extinct hominin species.

837

838 In this context, some exploratory studies using for example fluorodeoxyglucose positron emission tomography (FDG-PET), functional magnetic resonance imaging (fMRI) and image-839 840 based functional near-infrared spectroscopy (fNIRS) on Oldowan and Acheulean productions 841 have been undertaken. The purpose was to discuss which brain regions are the most active during these stone tool productions using different techniques (free-hand percussion and 842 bifacial shaping) and whether advanced executive functions such as cognitive control, 843 844 planning and the working memory are involved. During these experimental studies, the 845 subjects could have different levels of expertise (novices, learning, experts). The subjects could watch videos (Stout et al., 2015, using the fMRI method) or produce stone tools (Stout 846 et al. 2000, Stout et al. 2008, Stout and Chaminade, 2007, using FDG-PET method; Putt 2016, 847 848 Putt et al., 2017, using fNIRS method). According to Stout et al (2015), prefrontal responses 849 are greater in Acheulean as opposed to Oldowan tool-making. The main difference between Oldowan and Acheulean tasks is that Oldowan tool-making relies on the coordination of visual 850 attention and motor control (motor and sensorimotor areas) (Stout et al. 2000). In contrast, 851 according to Putt, Acheulean tool-making requires the integration of higher order motor 852 853 planning, the working memory and auditory feedback mechanisms (Putt 2016, Putt et al.,

854 2017). In order to examine the relationship between stone tool-making and brain function, Stout et al., (2000) and Stout and Chaminade (2007) developed an exploratory study using 855 Position Emission Tomography (FDG-PET). They have shown on one human expert subject 856 (Stout et al, 2000) that the superior parietal region (Brodmann Area (BA) 7, spatial cognition), 857 858 the central sulcus (BA 4 and BA 1, primary motor and somatosensory regions), the postcentral gyrus (BA1, basic sensorimotor processing) and the cerebellum (integration and control of 859 860 motor activity) are activated during the replication of Oldowan technology (Stout et al., 2000), but not the dorsolateral prefrontal cortex. This means that there is no high level of planning, 861 no strategic action planning during free-hand percussion, and that Oldowan tool making does 862 not place exceptional demands on executive functions, such as working memory, condition 863 response selection and inhibition, which are involved in strategic planning (Stout and 864 Chaminade, 2007, study based on 6 right-handed novices subjects). Observed superior 865 866 parietal lobe activity suggests the presence of complex spatial cognition, while activation 867 around the central sulcus reflects basic sensorimotor processing and cerebellar activity represents the integration and control of motor activity (Stout et al., 2000, p. 1222). 868 869 Furthermore, during Oldowan production, the rostral and dorsal intraparietal sulcus (IPS) and ventral premotor cortex (PMC) areas are activated. These zones are linked for example to 870 human visual specialization, manipulation, grip selection and tool-using abilities (Stout and 871 Chaminade, 2007, Stout and Chaminade, 2012). Brain activation data indicate that the initial 872 stages of Oldowan tool making skill acquisition are primarily concerned with perceptual-motor 873 adaptation to task constraints and especially the discovery and exploitation of objects 874 875 affordances, rather than executive planning and problem solving (Stout and Chaminade, 2007). 876

877 During the Acheulean (i.e., handaxe) production, the right inferior prefrontal cortex (Broadmann Area 45) is activated, an area which plays a central role in coordinating flexible, 878 goal-directed behaviour, and is linked to decision making. In contrast, the dorsolateral 879 prefrontal cortex, which is linked to the working memory in tasks that separate planning from 880 execution, is not activated (Stout et al., 2008, 3 experts). According to these authors, the 881 activation of the ventrolateral prefrontal cortex, but not dorsolateral prefrontal cortex 882 indicates that Acheulean tool-making is distinguished by cognitive demands for the 883 coordination of hierarchically organized ongoing action sequences rather than the evaluation 884 885 of action plans (Stout et al. 2008 p. 1946).

886

887 5. Discussion and conclusions

As preliminary remarks, Oldowan assemblages remain relatively scarce before 1.8 Ma, few Acheulean occurrences have been reported before 1.4 Ma, and there is a chronological overlap between Oldowan and Acheulean assemblages, both found sometimes on the same region or event in the same site (DAN 5(Dana Aoule Gona 5) and BSN 12 (Budisima North Gona 12), Semaw et al., 2020). In addition, it should be noted that the number of palaeoanthropological remains associated with archaeological assemblages is very scarce.

- However, our work has highlighted the plurality of artisans of the first material cultures,
 including Pliocene hominins for Lomekwian, and *Paranthropus*, *H. habilis* and *H. erectus/ergaster* for Oldowan.
- 897

Although Oldowan artefacts are often attributed to the *Homo* genus, we show in this article that different taxa (*P. boisei*, *P. aethiopicus*, *P. robustus*, *A. gahri*) overlapped temporally and geographically with these assemblages. More specifically, we observe a lot of High Confidence

- 901 Association between *Paranthropus* remains and Oldowan assemblages (n=11) (n=7 (HC 1 A);n= 2 (HC 1 B) and n=2 (HC 2 A), also sometimes associated with H. habilis (n=3, FLK NN level 3, 902 FLK level 22, FxJj 3), (Table2, Figure 2). Likewise, in terms of anatomy, the type of dexterity 903 required for making Oldowan assemblages is probably not restricted to Homo (Susman, 1994; 904 Marzke, 1997), the anatomy of Paranthropus elements (OH 80) is consistent with stone tool 905 production (Dominguez-Rodrigo et al., 2013), and some hand remains previously allocated to 906 907 H. habilis (e.g. OH 7) are now assigned to Paranthropus according to some authors (Moyà-Solà 908 et al., 2008).
- 909 In our opinion, the occurrences between Oldowan assemblages and *Paranthropus* are not
- 910 coincidental, and it is more than plausible that *Paranthropus* made and used stone tools for 911 cutting bones and/or vegetation. *Paranthropus* should be therefore considered as a stone tool
- 912 maker in the same way as early *Homo*.
- Besides, if after 1.4 Ma *H. erectus* is mainly associated with Acheulean assemblages, our paper also unlights that there are many sites (n=6) where *Homo erectus/ergaster* is associated with
- 915 Oldowan (PTK site Olduvai, Naiyena Engol 1, Melka Kunture Garba IV (level E), Melka Kunture
- 916 Gombore I, DAN 5, BSN 12, e.g. Chavaillon et al., 1977; Chavaillon et Berthelet, 2004; Condemi,
- 917 2004; Coppens, 2004; Domínguez-Rodrigo et al., 2015; Gallotti et Mussi, 2015; Prat et al.,
- 2003; Roche et al., 2003; Semaw et al., 2020), and not with Acheulean in Africa (Table2, Figure
- 2), but also outside Africa (e.g. Dmanisi (e.g. Lordkipanidze et al., 2007, 2013, Mgeladze et al.,
- 2011), 'Ubeidyia (e.g., Belmaker et al., 2002, Tobias, 1966, Bar-Yosef and Goren-Inbar, 1993).
- 921
- 922 INSERT Figure 2
- 923

924 Despite some strong associations between Oldowan assemblages and P. boisei remains, it 925 should be noted that in many cases, it is difficult for authors to admit that P. boisei could have 926 been the stone tool-makers. We can mention notably, the case of Olduvai Gorge. FLK level 22 927 yielded a dense Oldowan assemblage and four hominin individuals, including the cranium OH 928 5, holotype of the species Paranthropus boisei (formerly Zinjanthropus boisei, Leakey, 1959), 929 and one specimen allocated to H. habilis (OH 6) (see paragraph 2.1.2.1). When the OH 5 930 cranium was discovered in this site, it was considered as "the oldest yet discovered maker of 931 stone tools" (Leakey, 1959, p. 493). However, five years later, Louis Leakey changed his mind, based on Homo habilis fossils in association with the Oldowan culture at three other localities 932 933 at Olduvai, such as for example the FLK NN level 3 with P. boisei and H. habilis including the holotype OH 7 (see paragraph 2.1.2.2), and concluded that "while it is possible that 934 Zinjanthropus and Homo habilis both made stone tools, it is probable that the latter was the 935 more advanced tool maker and that the Zinjanthropus skull represents an intruder (or a 936 937 victim) on a Homo habilis living site" (Leakey et al., 1964, p. 9). In five years, with no real argument, the status of *P. boisei* has changed from stone tool-maker to victim. 938

939

940 It is also important to mention that despite the discovery of the *Paranthropus boisei* mandible 941 KNM-ER 3230 *in situ* at the FxJj 20 East site (Koobi Fora Area 131, Kenya) (see paragraph 942 2.1.3.4), Leakey in his 1976 paper did not want to associate the two (Leakey, 1976). We can 943 wonder whether Leakey's interpretation would have been the same if the remains had 944 belonged to the *Homo* genus.

More recently, at the site of Nyayanga (Kenya), a left lower molar (KNM-NG 77316) allocated to *P. boisei* was found *in situ*, spatially associated with Oldowan artefacts (see paragraph 2.1.3.1). The authors (Plummer et al., 2023) highlight a clear association between the lower 948 molar fragment and the artefacts "raising the possibility that Paranthropus made and/or co-949 opted stone tools" (Plummer et al., 2023 p. 5). However, in the same paragraph, the authors mention that even though no early *Homo* remains have been found at Nyayanga, as this taxon 950 occurred in eastern Africa, with the mandible from Ledi Geraru, 1300 kilometres away, dated 951 952 to 2.8 Ma in Ethiopia (Villmoare et al., 2015), the Nyayanga Oldowan artefacts "cannot be definitively attributed to a specific hominin genus "(Plummer et al., 2023 p. 5). This sentence 953 954 illustrates that it is still difficult for some authors to consider that a genus other than Homo could be the Oldowan stone tool makers and they give more importance to remains found 955 956 1300 kilometres away and corresponding to our non-confident (LC) category than to in situ 957 material (confident category HC1 A), which would be the most parsimonious hypothesis from 958 an Occamian perspective. Even if of course, it is always difficult to be sure that the hominins 959 found in some archaeological sites are the actual tool makers, but this hypothesis is commonly 960 used for example for Upper or Middle Palaeolithic sites.

The case of Chesowanja (see paragraph 2.1.3.5), also demonstrates this point. Indeed, the *P. boisei* are not strictly associated with the Oldowan assemblage found nearby, but Gowlett et al. 1981, consider that this association must be considered more than coincidental (Gowlett et al., 1981).

It is interesting here to use the words of Gowlett in 1999. "Although the only hominid remains 965 found at Chesowanja are A. boisei, Gowlett et al., 1981 were inclined to suggest that the maker 966 of the stone industry was H. erectus. If anything, the case for this link has strengthened, 967 through additional finds of *H. erectus* at early dates, the frequency with which *Homo* is found 968 on artefact sites where A. boisei is found, and the continued association of Homo erectus and 969 bifaces following the demise of A. boisei". (Gowlett 1999, p.128)...and "a view seeking simple 970 parsimony would proceed along these lines: that any hominid might in principle make tools; 971 972 that robust australopithecines have well developed thumbs, suggesting tool-making manipulations, and that where there are two occurrences of this hominid close to artefacts, 973 974 this is suggestive a link. Against this, however, one can take a broader view of the arguments: 975 (1) remains of the 'more advanced' Homo are found elsewhere alongside robust 976 australopithecine remains on several sites in several environments; (2) the absence of Homo 977 has no special significance, since hominins do not tend to die in their living area so as to 978 provide makers; (3) Chesowanja represents a developed Oldowan possibly including early 979 picks as found in the Acheulean, and Acheulean industries are commonly associated with Homo ergaster/erectus and Homo sapiens/heidelbergensis." (Gowlett, 1999 p. 130). We can 980 see here that Gowlett changed his mind slightly concerning the perception of P. boisei as a 981 stone tool-maker between his paper published in 1981 and 1999. Whether it is true that the 982 983 place of death is not necessarily the living one, and that the hominin remains found at the site 984 (especially when it is a juvenile) are not necessary those of the hominins who produced the 985 lithic assemblage, the most parsimonious hypothesis merely indicates that members of the same group were present at the archaeological site. Questions of geological scale and 986 sedimentation may also be raised. However, the most parsimonious hypothesis is that the 987 988 stone tool or bone tool makers correspond to the species found at the archaeological site 989 rather than to species not present in the sedimentary levels. This is generally the case when specimens of the *Homo* genus are found, but less systematic for other genus. 990

Additionally, we can emphasize, as shown in this article, that the frequency of *Paranthropus* remains associated with Oldowan assemblages in eastern Africa is not negligible and insignificant (eleven occurrences), that there are quite a lot of sites with *P. boisei* remains not associated with early *Homo* individuals (six occurrences), that some Oldowan sites are associated with *Homo* ergaster/erectus (six evidence).

The plurality of stone tool-makers has been previously highlighted by some researchers (e.g.
Brain et al., 1988; Susman, 1991; Wood, 1997), mainly based on their ability to manufacture

- simple stone tools and on southern African sites, but often the situation is broadly the same
 as it was in 1964. According to Wood and Constantino (2007): "There is no firm evidence that *P. boisei* made and used stone tools, but there is also no firm evidence that they were
 incapable of doing so" (Wood and Constantino, 2007 p.122).
- Several years later, this paper has attempted to review the different sites where *Paranthropus* remains (especially in East Africa) were found *in situ* with Oldowan assemblages alongside anatomical perspectives. We hope that we have shown that there is strong evidence to suggest that *Paranthropus* made and used stone tools and that this taxon should be given more consideration.
- 1007

1008 From a parsimonious perspective, there is more association between *Paranthropus* remains 1009 and lithic artefacts for the oldest sites than between early Homo and lithic artefacts. Based on the evidence presented here, stone tool-making does not seem to be specific to the Homo 1010 genus and cannot define it. Fieldwork data are essential, but the development of 1011 musculoskeletal simulations, kinematic, electro-myographic and electro-encephalographic 1012 1013 experiments (biplanar X-ray, internal morphology) are also promising avenues of research enhance our understanding of the relationship between hominins and material cultures. 1014 Moreover, the unique and specific attribution of a tool-making capacity to the Homo genus 1015 can no longer be deemed valid for these periods. 1016

1017

1018 Acknowledgments

I would like to thanks the reviewers, Stéphanie Bonilauri and Hubert Forestier, as well as the
editors of L'Anthropologie for their invitation to participate to this special issue and Louise
Byrne for the english editing. This work is supported by the French Research Agency (Agence
Nationale de la Recherche) ANR HOMTECH (ANR-17-CE27-0005).

- 1023
- 1024

1025 References

- Alba, D.M., Moyà-Solà, S., Köhler, M., 2003. Morphological affinities of the *Australopithecus afarensis* hand on the basis of manual proportions and relative thumb length. J. Hum. Evol.
 44(2), 225-254. https://doi.org/10.1016/S0047-2484(02)00207-5
- Almécija, S., Alba, D.M., 2014. On manual proportions and pad-to-pad precision grasping in
 Australopithecus afarensis. J. Hum. Evol. 73, 88-92.
 <u>https://doi.org/10.1016/j.jhevol.2014.02.006</u>
- Arroyo, A., Harmand, S., Roche, H., Taylor, N., 2020. Searching for hidden activities: percussive
 tools from the Oldowan and Acheulean of West Turkana, Kenya (2.3–1.76 Ma). J. Archaeol.
 Sci. 123, 105238. <u>https://doi.org/10.1016/j.jas.2020.105238</u>
- 1036 Asfaw, B., Beyene, Y., Semaw, S., Suwa, G., White, T., WoldeGabriel, G., 1991. Fejej: a new 1037 paleoanthropological research area in Ethiopia. J Hum Evol. 21(2), 137-143.
- Asfaw, B., Beyene, Y., Suwa, G., Walter, R. C., White, T. D., WoldeGabriel, G., Yemane, T., 1992.
 The earliest acheulean from Konso-Gardula. Nature 360, 732-735.
- 1040 Asfaw, B., White, T., Lovejoy, O., Latimer, B., Simpson, S., Suwa, G., 1999. *Australopithecus* 1041 *garhi*: a new species of early hominid from Ethiopia. Science 284(5414), 629-635.
- Baab, K. L., Rogers, M., Bruner, E., Semaw, S., 2022. Reconstruction and analysis of the
 DAN5/P1 and BSN12/P1 Gona Early Pleistocene *Homo* fossils. J. Hum. Evol. 162, 103102.
 https://doi.org/10.1016/j.jhevol.2021.103102
- Backwell, L.R., d'Errico, F., 2003. Additional evidence on the early hominid bone tools from
 Swartkrans with reference to spatial distribution of lithic and organic artefacts. S. Afr. J. Sci.
 99, 259-267.
- Backwell, L., d'Errico, F., 2008. Early hominid bone tools from Drimolen, South Africa. J.
 Archaeol. Sci. 35, 2880-2894. <u>https://doi.org/10.1016/j.jas.2008.05.017</u>
- Balter, V., Blichert-Toft, J., Braga, J., Telouk, P., Thackeray, F., Albarède, F., 2008. U-Pb dating
 of fossil enamel from the Swartkrans Pleistocene hominid site, South Africa. Earth Planet.
 Sci. Lett. 267 (1-2), 236–246.
- Bardo, A., Vigouroux, L., Kivell, T.L., Pouydebat, E., 2018. The impact of hand proportions on tool grip abilities in humans, great apes and fossil hominins: A biomechanical analysis using musculoskeletal simulation. J. Hum. Evol. 125, 106-121.
 https://doi.org/10.1016/j.jhevol.2018.10.001
- Barsky, D., Chapon-Sao, C., Bahain, J. J., Beyene, Y., Cauche, D., Celiberti, V., Desclaux, E., de
 Lumley, H., de Lumley, M-A, Marchal, F., Moullé, P-E, Pleurdeau, D., 2011. The early
 Oldowan stone-tool assemblage from Fejej FJ-1a, Ethiopia. J. Afr. Archeaol. 9(2), 207-224.
- Bar-Yosef, O., Goren-Inbar, N., 1993. The Lithic Assemblages of 'Ubeidiya: A Lower Palaeolithic
 Site in the Jordan Valley (Vol. 34). Institute of Archaeology, Hebrew University of Jerusalem.
- Belmaker, M., Tchernov, E., Condemi, S., Bar-Yosef, O., 2002. New evidence for hominid
 presence in the Lower Pleistocene of the Southern Levant. J. Hum. Evol. 43, 43-56.
- Berger, L. R., De Ruiter, D. J., Churchill, S. E., Schmid, P., Carlson, K. J., Dirks, P. H., Kibii, J. M.,
 2010. *Australopithecus sediba*: A new species of *Homo*-like australopith from South Africa.
 Science, 328, 195-204. https://doi.org/10.1126/science.118494
- Beyene, Y., Katoh, S., WoldeGabriel, G., Hart, W. K., Uto, K., Sudo, M., Kondo, M., Hyodo, M.,
 Renne, P.R., Suwa, G., Asfaw, B., 2013. The characteristics and chronology of the earliest
 Acheulean at Konso, Ethiopia. Proc. Natl. Acad, Sci. USA 110(5), 1584-1591.
- 1070

- Beyene, Y., Asfaw, B., Katoh, S., Suwa, G., 2014. Overview of the Archaeological Research at
 Konso. In: Beyne, Y., Asfaw, B., Sano, K., Suwa, G. (Eds.) Konso-Gardula Research Project,
 the University Museum, the University of Tokyo, Tokyo, Japan, p. 9-25.
- 1074 Bishop, W. W., Pickford, M., Hill, A., 1975. New evidence regarding the Quaternary geology, 1075 archaeology and hominids of Chesowanja, Kenya. Nature 258(5532), 204-208.
- Blumenschine, R.J., Peters, C.R., Masa, F.T., Clarke, R.J., Deino, A.L., Hay, R.L., Swiher, C.C.,
 Stanistreet, I.G., Ashley, G.M., McHenry, L.J., Sikes, N.E., van der Merwe, N.J., Tactikos, J.C.,
 Cushing, A.E., Deocampo, D.M., Njau, J.K., Ebert, J.I., 2003. Late Pliocene *Homo* and
 hominid land use from western Olduvai gorge, Tanzania. Science 299, 1217-1221.
- Blumenschine, R. J., Stanistreet, I. G., Njau, J. K., Bamford, M. K., Masao, F. T., Albert, R. M.,
 Stollhofen, H., Andrews, P., Prassack, K.A., McHenry, L.J., Fernández-Jalvo, Y. Camilli, E.L.,
 Ebert, J. I. (2012). Environments and hominin activities across the FLK Peninsula during *Zinjanthropus* times (1.84 Ma), Olduvai Gorge, Tanzania. J. Hum. Evol 63(2), 364-383.
- Brain, C.K., 1993. Swartkrans. A Cave's Chronicle of Early Man. Transvaal Museum, Pretoria.
- Brain, C.K., Shipman, P., 1993. The Swartkrans bone tools. In: Brain, C.K. (Ed.), Swartkrans: A
 Cave's Chronicle of Early Man. Transvaal Museum Monograph, Pretoria, p. 195-215.
- Brain, C.K., Churcher, C.S., Clark, J.D., Grine, F.E., Shipman, P., Susman, R.L., Turner, A.,
 Watson, V. 1988. New evidence of early hominids, their culture and environment from the
 Swartkrans cave, South Africa. S. Afr. J. Sci. 84, 828-835.
- Braun, D.R., Aldeias, V., Archer, W., Arrowsmith, J.R., Baraki, N., Campisano, C.J., Deino, A.L.,
 DiMaggio, E.N., Dupont-Nivet, G., Engda, B., Feary, D.A., Garello, D.I., Kerfelew, Z.,
 McPherron, S.P., Patterson, D.B., Reeves, J.S., Thompson, J.C., Reed, K.E., 2019. Earliest
 known Oldowan artifacts at >2.58 Ma from Ledi-Geraru, Ethiopia, highlight early
 technological diversity. Proc. Natl. Acad. Sci. U S A. 116(24), 11712-11717.
 https://doi.org/10.1073/pnas.1820177116
- Bromage, T. G., Schrenk, F., Zonneveld, F. W., 1995. Paleoanthropology of the Malawi Rift: an
 early hominid mandible from the Chiwondo Beds, northern Malawi. J. Hum. Evol. 28(1), 71 1098
- Brugal, J. P., Roche, H., Kibunjia, M., 2003. Faunes et paléoenvironnements des principaux
 sites archéologiques plio-pléistocènes de la formation de Nachukui (Ouest-Turkana,
 Kenya). C. R. Palevol. 2(8), 675-684. <u>https://doi.org/10.1016/j.crpv.2003.09.028</u>
- Bruner, E., Holloway, R., Baab, K. L., Rogers, M. J., Semaw, S., 2023. The endocast from Dana
 Aoule North (DAN5/P1): A 1.5 million year-old human braincase from Gona, Afar, Ethiopia.
 Am. J. Biol. Anthropol., 1-10. https://doi.org/10.1002/ajpa.24717
- Bush, M. E., Lovejoy, C. O., Johanson, D. C., Coppens, Y., 1982. Hominid carpal, metacarpal,
 and phalangeal bones recovered from the Hadar formation: 1974–1977 collections. Am. J.
 Phys. Anthropol. 57, 651-677. <u>https://doi.org/10.1002/ajpa.1330570410</u>
- Campisano, C. J., 2012. Geological summary of the Busidima formation (Plio-Pleistocene) at
 the Hadar paleoanthropological site, Afar Depression, Ethiopia. J. Hum. Evol. 62(3), 338352.
- 1111 Carbonell, E., Bermúdez de Castro, J. M., Parés, J. M., Pérez-González, A., Cuenca-Bescós, G.,
- 1112 Ollé, A., Mosquera, M., Huguet, R., van der Made, J., Rosas, A., Sala, R., Vallverdú, J., García,
- 1113 N., Granger, D.E., Martinón-Torres, M., Rodríguez, X. P., Stock, G. M., Vergès, J. M., Allué,
- 1114 E., Burjachs, F., Cáceres, I., Canals, A., Benito, A., Díez, C., Lozano, M., Mateos, A., Navazo,

- 1115 M., Rodríguez, J., Rosell, J., Arsuaga, J. L., 2008. The first hominin of Europe. Nature 452, 465-469.
- Carney, J., Hill, A., Miller, J. A., Walker, A., 1971. Late australopithecine from Baringo district,
 Kenya. Nature 230 (5295), 509-514.
- Caruana, M.V., 2017. Lithic Production Strategies in the Oldowan Assemblages from
 Sterkfontein Member 5 and Swartkrans Member 1, Gauteng Province, South Africa. J. Afr.
 Archaeol. 15, 1-19.
- Cazenave, M., Braga, J., Oettlé, A., Thackeray, J. F., De Beer, F., Hoffman, J., Endalamaw, M.,
 Redae, B.E., Macchiarelli, R., 2017. Inner structural organization of the distal humerus in
 Paranthropus and *Homo*. C. R. Palevol. 16(5-6), 521-532.
- Chapon, C., Bahain, J. J., Lumley, H. D., Perrenoud, C., Vandamme, D., 2005. Stratigraphie,
 sédimentologie et magnétostratigraphie du site oldowayen de Fejej FJ-1, Sud-Omo,
 Ethiopie: Premiers résultats. Quaternaire.16(3), 143-152.
- Chavaillon, J., 1976. Evidence for the technical practices of Early Pleistocene hominids,
 Shungura Formation, Lower Omo Valley, Ethiopia. In: Coppens, Y., Howell, F.C., Isaac, G.L.I.,
 Leakey, R. (Eds.), Earliest Man and Environments in the Lake Rudolf Basin: Stratigraphy,
 Paleoecology and Evolution. University of Chicago Press, Chicago and London, p. 565-573.
- Chavaillon, J., Berthelet, A., 2004. The archaeological sites of Melka Kunture. In: Chavaillon, J.,
 Piperno, M. (Eds.), Studies on the Early Paleolithic Site of Melka Kunture, Ethiopia. IIPP,
 Florence, p. 25-80.
- 1135 Chavaillon, J., Chavaillon, N., Coppens, Y., Senut, B., 1977. Présence d'Hominidé dans le site 1136 Oldowayen de Gomboré I à Melka-Kunturé. C.R.Acad. Serie D 285, 961-963.
- 1137 Clarke, R. J., 2012. A *Homo habilis* maxilla and other newly-discovered hominid fossils from 1138 Olduvai Gorge, Tanzania. J. Hum. Evol. 63(2), 418-428.
- 1139 Condemi, S., 2004. The garba IV E mandible. In: Chavaillon, J., Piperno, M. (Eds.), Studies on
 1140 the Early Paleolithic Site of Melka Kunture, Ethiopia. IIPP, Florence, p.687-701.
- 1141 Coppens, Y., 1977. Hominid remains from the Plio/Pleistocene formations of the Omo Basin, 1142 Ethiopia. J. Hum. Evol *6*(2), 169-173.
- 1143 Coppens, Y., 2004. The hominids of Melka Kunture. Some general reflections. In: Chavaillon,
 1144 J., Piperno, M. (Eds.), Studies on the Early Paleolithic site of Melka Kunture, Ethiopia, IIPP,
 1145 Florence, p. 685-686.
- Coppens, Y., Chavaillon J, Beden M., 1973. Résultats de la nouvelle mission de l'Omo
 (campagne 1972). Découverte de restes d'Hominides et d'une industrie sur éclats. C. R.
 Acad. Sci. Série D. 276, 161-164.
- d'Errico, F., Backwell, L.R., 2003. Possible evidence of bone tool shaping by Swartkrans early
 hominids. J. Archaeol. Sci. 30, 1559-1576.
- De Heinzelin, J. D., Clark, J. D., White, T., Hart, W., Renne, P., WoldeGabriel, G., Beyene, Y.
 Vrba, E., 1999. Environment and behavior of 2.5-million-year-old Bouri hominids. Science 284(5414), 625-629.
- de Lumley, H., Beyne Y. (Eds) (2004). Les sites préhistoriques de la région de Fejej, Sud-Omo,
 Éthiopie, dans leur contexte stratigraphique et paléontologique Éditions Recherche sur les
 civilisations.
- de Lumley, M.-A., Marchal, F., 2004. Les restes d'hominidés du site de Fejej FJ-1. In : de
 Lumley, H., Beyne Y. (Eds.), Les sites préhistoriques de la région de Fejej, Sud-Omo,
 Ethiopie, dans leur contexte stratigraphique et paléontologique, Editions Recherche sur les
 Civilisations, Paris, France, p. 340-383.
 - 26

- de Ruiter, D.J., Pickering, R., Steininger, C.M., Kramers, J.D., Hancox, P.J., Churchill, S. E.,
 Berger, L.R., Backwell, L., 2009. New *Australopithecus robustus* fossils and associated U-Pb
 dates from Cooper's Cave (Gauteng, South Africa). J. Hum. Evol. 56, 497-513.
- de la Torre, I., 2004. Omo revisited: evaluating the technological skills of Pliocene hominids.
 Curr. Anthrop. 45(4), 439-465.
- de la Torre, I., Benito-Calvo, A., Martín-Ramos, C., McHenry, L. J., Mora, R., Njau, J. K., Pante,
 M.C., Stanistreet, I.G., Stollhofen, H., 2021. New excavations in the MNK Skull site, and the
 last appearance of the Oldowan and *Homo habilis* at Olduvai Gorge, Tanzania. J. Anthropol.
 Archaeol. 61, 101255, <u>https://doi.org/10.1016/j.jaa.2020.101255</u>
- de la Torre, I., Mora, R., 2005. Technological Strategies in the Lower Pleistocene at Olduvai
 Beds I & II. ERAUL, Liege, Belgium.
- de Weyer, L. 2022. Early Human Lithic Technologies: Multiple Technical Histories from Africa
 to East Asia. In : Holl A.F.C. (Ed.) *Studying Africa and Africans Today*, BP International,
- 1174 Kolkata, India, pp. 167-188. https://doi.org/10.9734/bpi/mono/978-93-5547-847-4/CH11
- Day, M.H., Napier, J.R., 1964. Hominid fossils from Bed I, Olduvai Gorge, Tanganyika: Fossilfoot bones. Nature 201, 967-970.
- Deino, A.L., 2012. 40Ar/39Ar dating of Bed I, Olduvai Gorge, Tanzania, and the chronology of
 early Pleistocene climate change. J. Hum. Evol. 63, 251-273.
- 1179 Delagnes, A., Boisserie, J. R., Beyene, Y., Chuniaud, K., Guillemot, C., Schuster, M., 2011. Archaeological investigations in the Lower Omo Valley (Shungura Formation, Ethiopia): 1180 1181 new data and perspectives. J. Hum. Evol., 61(2), 215-222. https://doi.org/10.1016/j.jhevol.2011.03.008 1182
- Delagnes, A., Brenet, M., Gravina, B., Santos, F., 2023. Exploring the relative influence of raw
 materials, percussion techniques, and hominin skill levels on the diversity of the early
 Oldowan assemblages: Insights from the Shungura Formation, Lower Omo Valley, Ethiopia.
 Plos one, 18(4), e0283250. https://doi.org/10.1371/journal.pone.0283250
- Delagnes, A., Roche, H., 2005. Late Pliocene hominid knapping skills: the case of Lokalalei 2C, 1187 West Turkana, Evol. 435-472. 1188 Kenya. Hum. 48(5), J. 1189 https://doi.org/10.1016/j.jhevol.2004.12.005Diez-Martín, F., Sánchez Yustos, Ρ., 1190 Uribelarrea, D., Baquedano, E., Mark, D. F., Mabulla, A., Fraile, C., Duque, J., Pérez-GonzáA., 1191 Yravedra, J., Egeland, C.P., Organista, E., Domínguez-Rodrigo, M., 2015. The origin of the Acheulean: The 1.7 million-year-old site of FLK West, Olduvai Gorge (Tanzania). Scientific 1192 1193 reports, 5(1), 17839. DOI: 10.1038/srep17839.
- 1194Di Vincenzo, F., Rodriguez, L., Carretero, J. M., Collina, C., Geraads, D., Piperno, M., Manzi, G.,11952015. The massive fossil humerus from the Oldowan horizon of Gombore I, Melka Kunture1196(Ethiopia,>11971.39Ma).Quat.Sci.Rev.122,207-221.1197https://doi.org/10.1016/j.quascirev.2015.05.014
- Domalain, M., Bertin, A., Daver, G., 2017. Was Australopithecus afarensis able to make the
 Lomekwian stone tools? Towards a realistic biomechanical simulation of hand force
 capability in fossil hominins and new insights on the role of the fifth digit. C. R. Palevol. 16,
 572-584.
- Domínguez-Rodrigo, M., Mabulla, A., Bunn, H. T., Barba, R., Diez-Martín, F., Egeland, C. P.,
 Espílez, E., Egeland, A., Yravedra, J., Sánchez, P., 2009. Unraveling hominin behavior at
 another anthropogenic site from Olduvai Gorge (Tanzania): new archaeological and
 taphonomic research at BK, Upper Bed II. J. Hum. Evol. 57(3), 260-283.

- Domínguez-Rodrigo, M., Pickering, T. R., Almécija, S., Heaton, J. L., Baquedano, E., Mabulla,
 A., Uribelarrea, D., 2015. Earliest modern human-like hand bone from a new> 1.84-million year-old site at Olduvai in Tanzania. Nat. Comm., 6(1), 7987. DOI: 10.1038/ncomms8987
- 1209 Domínguez-Rodrigo, M., Pickering, T.R., Baquedano, E., Mabulla, A., Mark, D.F., Musiba, C.,
- Bunn, H.T., Uribelarrea, D., Smith, V., Diez-Martin, F., Pérez-González, A., Sánchez, P., 1210 Santonja, M., Barboni, D., Gidna, A., Ashley, G., Yravedra, J., Heaton, J.L., Arriaza, M.C., 1211 2013. First partial skeleton of a 1.34-million-year-old Paranthropus boisei from Bed II, 1212 PLoS 1213 Olduvai Gorge, Tanzania. One. 8(12), e80347. https://doi.org/10.1371/journal.pone.0080347 1214
- Drapeau, M.S., Ward, C.V., Kimbel, W.H., Johanson, D.C., Rak, Y. 2005. Associated cranial and
 forelimb remains attributed to *Australopithecus afarensis* from Hadar, Ethiopia. J Hum Evol.
 48, 593-642, https://doi.org/10.1016/j.jhevol.2005.02.005
- Duke, H., Feibel, C., Harmand, S., 2021 Before the Acheulean: The emergence of bifacial
 shaping at Kokiselei 6 (1.8 Ma), West Turkana, Kenya. J Hum Evol. 159, 103061.
 <u>https://doi.org/10.1016/j.jhevol.2021.103061</u>
- Dunmore, C.J., Skinner, M.M., Bardo, A., Berger, L.R., Hublin, J.J., Pahr, D.H., Rosas, A.,
 Stephens, N.B., Kivell, T.L. 2020. The position of *Australopithecus sediba* within fossil
 hominin hand use diversity. Nat Ecol Evol. 4(7), 911-918. <u>https://doi.org/10.1038/s41559-</u>
 020-1207-5
- 1225 Echassoux, A., 2012. Comportements de subsistance et modifications osseuses à l'aube de
 1226 l'Acheuléen à Konso, Éthiopie. L'Anthropologie, 116(3), 291-320.
 1227 <u>https://doi.org/10.1016/j.anthro.2012.06.002</u>
- Frayer, D. W., Clarke, R. J., Fiore, I., Blumenschine, R. J., Pérez-Pérez, A., Martinez, L. M., 1228 Estebaranz, F., Holloway, R., Bondioli, L., 2016. OH-65: The earliest evidence for right-1229 1230 handedness in the fossil record. J Hum Evol. 100, 65-72. https://doi.org/10.1016/j.jhevol.2016.07.002 1231
- Feibel, C.S, Brown, F.H., McDougall, I., 1989. Stratigraphic context of fossil hominids from the
 Omo Group deposits: Northern Turkana Basin, Kenya and Ethiopia. Am. J. Phys. Anthrop.
 78, 595-622. <u>https://doi.org/10.1002/ajpa.1330780412</u>
- Gallotti, R., 2018. Before the Acheulean in East Africa: An overview of the Oldowan lithic
 assemblages. In: Gallotti, R. & Mussi M. (Eds.). *The Emergence of the Acheulean in East Africa and Beyond: Contributions in Honor of Jean Chavaillon*, Vertebrate Paleobiology and
- 1238 Paleoanthropology Series, Springer, Cham, Switzerland, p. 13–32.
- Gallotti, R., Mussi, M., 2015. The unknown Oldowan:~ 1.7-million-year-old standardized
 obsidian small tools from Garba IV, Melka Kunture, Ethiopia. *PLoS One*, *10*(12), e0145101.
 <u>https://doi.org/10.1371/journal.pone.0145101</u>
- Gérard, C., Bardo, A., Guéry, J.P., Pouydebat, E., Simmen, B., Narat, V., 2022. Manipulative
 repertoire of bonobos (*Pan paniscus*) in spontaneous feeding situation. Am J Primatol.
 84(7), e23383. <u>https://doi.org/10.1002/ajp.23383</u>
- Gibbon, R.J., Pickering, T.R., Sutton, M.B., Heaton, J.L., Kuman, K., Clarke, R.J., Brain, C.K.,
 Granger, D.E., 2014. Cosmogenic nuclide burial dating of hominin-bearing Pleistocene cave
 deposits at Swartkrans, South Africa. Quat. Geochronol. 24, 10-15
- 1248Goldman-Neuman T, Hovers E., 201). Raw material selectivity in Late Pliocene Oldowan sites1249in the Makaamitalu Basin, Hadar, Ethiopia. J Hum Evol. 62(3):353-66.1250https://doi.org/10.1016/j.jhevol.2011.05.006

- Gowlett, J.A.J., 1999. Lower and Middle Pleistocene archaeology of the Baringo Basin. In:
 Andrews, P. and Banham, P. (Eds) Late Cenozoic environments and hominid evolution: a
 tribute to Bill Bishop. Geological Society, London, p. 123-141.
- 1254 Gowlett, J. A., Harris, J. W., Walton, D., Wood, B. A., 1981. Early archaeological sites, hominid 1255 remains and traces of fire from Chesowanja, Kenya. Nature, 294(5837), 125-129.
- Granger, D. E., Gibbon, R. J., Kuman, K., Clarke, R. J., Bruxelles, L., Caffee, M. W., 2015. New
 cosmogenic burial ages for Sterkfontein Member 2 *Australopithecus* and Member 5
 Oldowan. Nature, 522(7554), 85-88.
- Granger, D.E., Stratford, D., Bruxelles, L., Gibbon, R.J., Clarke, R.J., Kuman, K., 2022.
 Cosmogenic nuclide dating of *Australopithecus* at Sterkfontein, South Africa. Proc Natl Acad
 Sci U S A. 119(27), e2123516119. <u>https://doi.org/10.1073/pnas.2123516119</u>
- 1262 Green, D.J., Gordon, A.D., 2008. Metacarpal proportions in *Australopithecus africanus*. J Hum 1263 Evol. 54(5), 705-719. <u>https://doi.org/10.1016/j.jhevol.2007.10.007</u>
- Grine, F. E., Mongle, C. S., Fleagle, J. G., Hammond, A. S., 2022. The taxonomic attribution of
 African hominin postcrania from the Miocene through the Pleistocene: Associations and
 assumptions. J Hum Evol. 173, 103255. <u>https://doi.org/10.1016/j.jhevol.2022.103255</u>
- Han, F., Bahain, J. J., Shao, Q., Sun, X., Voinchet, P., Xiao, P., Huang, M., Li, M., Yin, G., 2022.
 The Chronology of Early Human Settlement in Three Gorges Region, China—Contribution
 of Coupled Electron Spin Resonance and Uranium-Series Dating Method. Frontiers in Earth
- 1270 Science, 10, 939766. https://doi.org/10.3389/feart.2022.939766
- Hanon, R., Francesco d'Errico, F., Backwell, L., Prat, S., Péan, S., Patou-Mathis, M., 2021. New
 evidence of bone tool use by Early Pleistocene hominins from Cooper's D, Bloubank Valley,
 South Africa. J. Archaeol. Sci. Rep. 39, 103129,
 https://doi.org/10.1016/j.jasrep.2021.103129.
- Harmand, S., 2009a. Raw material selectivity and techno-economic behaviours at Oldowan
 Acheulean sites in the West Turkana region, Kenya. In: Adams, B. & Blades B. (Eds.), Lithic
 Materials and Paleolithic Societies, Blackwell Publishing, Oxford, United Kingdom, p. 3-14.
- Harmand S., 2009 b. Variability in raw material selectivity at the late Pliocene sites of Lokalalei,
 West Turkana, Kenya. In: Hovers, E. & Braun, D.R. (Eds.), Interdisciplinary Approaches to
 the Oldowan. New-York: Springer; 2009. p. 85–97.
- Harmand, S., Lewis, J.E., Feibel, C.S., Lepre, C.J., Prat, S., Lenoble, A., Boës, X., Quinn, R.L.,
 Brenet, M., Arroyo, A., Taylor, N., Clément, S., Daver, G., Brugal, J-P., Leakey, L., Mortlock,
 R.A., Wright, J.D., Lokorodi, S., Kirwa, C., Kent, D.V., Roche, H., 2015. 3.3-million-year-old
 stone tools from Lomekwi 3, West Turkana, Kenya. Nature 521(7552), 310-315.
 https://doi.org/10.1038/nature14464
- Harris, J. M., Brown, F.H., Leakey, M.G. 1988. Stratigraphy and paleontology of Pliocene and
 Pleistocene localities west of Lake Turkana, Kenya. Contr. Sci. Nat. Mus. Los Angeles 399,
 1-128.
- Harris, J. W., Herbich, I., 1978. Aspects of early Pleistocene hominid behaviour east of Lake
 Turkana, Kenya. Geological Society, London, Special Publications, 6(1), 529-547.
- Herries, A. I., Shaw, J., 2011. Palaeomagnetic analysis of the Sterkfontein palaeocave deposits:
 Implications for the age of the hominin fossils and stone tool industries. J. Hum. Evol.
 60(5):523-539. <u>https://doi.org/10.1016/j.jhevol.2010.09.001</u>
- Herries, A.I.R., Martin, J.M., Leece, A.B., Adams, J.W., Boschian, G., Joannes-Boyau, R.,
 Edwards, T.R., Mallett, T., Massey, J., Murszewski, A., Neubauer, S., Pickering, R., Strait,
 D.S., Armstrong, B.J., Baker, S., Caruana, M.V., Denham, T., Hellstrom, J., Moggi-Cecchi, J.,
 Mokobane, S., Penzo-Kajewski, P., Rovinsky, D.S., Schwartz, G.T., Stammers, R.C., Wilson,

- 1298 C., Woodhead, J., Menter, C., 2020. Contemporaneity of *Australopithecus*, *Paranthropus*,
 1299 and early *Homo erectus* in South Africa. Science 368, eaaw7293.
 1300 <u>https://doi.org/10.1126/science:aaw7293</u>
- Hill, A., Ward, S., Deino, A., Curtis, G., Drake, R. 1992. Earliest *Homo*. Nature, 355(6362), 719722. DOI: <u>10.1038/355719a0</u>
- Holloway, R.L., 1983. Cerebral brain endocast pattern of *Australopithecus afarensis* hominid.
 Nature 303(5916), 420-422. <u>https://doi.org/10.1038/303420a0</u>
- Hou, Y. M., Zhao, L. X. 2010. An archeological view for the presence of early humans in China.
 Quat. Int. 223–224, 10-19. <u>https://doi.org/10.1016/j.quaint.2009.09.025</u>
- Howell, F. C., Haesaerts, P., de Heinzelin, J., 1987. Depositional environments, archeological
 occurrences and hominids from Members E and F of the Shungura Formation (Omo basin,
 Ethiopia). J. Hum. Evol. 16(7-8), 665-700. <u>https://doi.org/10.1016/0047-2484(87)90019-4</u>
- Johanson, D. C., Lovejoy, C. O., Kimbel, W. H., White, T. D., Ward, S. C., Bush, M. E., Latimer,
 B. M., Coppens, Y., 1982. Morphology of the Pliocene partial hominid skeleton (A.L. 288-1)
- 1312
 from the Hadar formation, Ethiopia. Am. J. Phys. Anthrop., 54, 403-452.

 1313
 <u>https://doi.org/10.1002/ajpa.1330570403</u>
- Key, A. J., Dunmore, C. J., 2015. The evolution of the hominin thumb and the influence exerted
 by the non-dominant hand during stone tool production. J. Hum. Evol., 78, 60-69.
 <u>https://doi.org/10</u>. 1016/j.jhevol.2014.08.006
- 1317 Key, A. J., Dunmore, C. J., Marzke, M. W., 2019. The unexpected importance of the fifth digit
 1318 during stone tool production. Scientific Reports, 9, 1–8. <u>https://doi.org/10.1038/s41598-</u>
 1319 <u>019-53332-w</u>
- 1320 Key, A., Merritt, S. R., Kivell, T. L., 2018. Hand grip diversity and frequency during the use of
 1321 lower Palaeolithic stone cutting-tools. J. Hum. Evol. 125, 137–158.
 1322 <u>https://doi.org/10.1016/j.jhevol</u>. 2018.08.006
- 1323 Keyser, A.W., Menter, C.G., Moggi-Cecchi, J., Pickering, T.R., Berger, L.R., 2000. Drimolen: a
 1324 new hominid-bearing site in Gauteng, South Africa. S. Afr. J. Sci. 96, 193-197.
- Kibunjia, M., 1994. Pliocene archaeological occurrences in the Lake Turkana basin. J. Hum.
 Evol. 27, 159-171 <u>https://doi.org/10.1006/jhev.1994.1040</u>
- Kibunjia, M., Roche, H., Brown, F.H., Leakey, R.E.F., 1992. Pliocene and Pleistocene
 archaeological sites west of Lake Turkana, Kenya. J. Hum. Evol., 23, 431-438.
 <u>https://doi.org/10.1016/0047-2484(92)90091-M</u>
- Kimbel, W. H., Johanson, D. C., Rak, Y., 1997. Systematic assessment of a maxilla of *Homo* from
 Hadar, Ethiopia. Am. J. Phys. Anthrop. 103(2), 235-262.
- 1332 Kimbel, W. H., Walter, R. C., Johanson, D. C., Reed, K. E., Aronson, J. L., Assefa, Z., Marean, C.
 1333 W., Eck, G. G., Bobe, R., Hovers, E., Rak, Y., Vondra, C., Yemane, T., York, D., Chen, Y.,
- Evensen, N. M., Smith, P. E., 1996. Late Pliocene *Homo* and Oldowan Tools from the Hadar
 Formation (Kada Hadar Member), Ethiopia. J. Hum. Evol. 31, 549-561.
 <u>https://doi.org/10.1006/jhev.1996.0079</u>
- Kivell, T. L., Baraki, N., Lockwood, V., Williams-Hatala, E. M., Wood, B. A., 2022. Form, function
 and evolution of the human hand. Am. J. Biol. Anthrop. DOI: 10.1002/ajpa.24667
- Kivell, T. L., Kibii, J. M., Churchill, S. E., Schmid, P., Berger, L. R., 2011. Australopithecus sediba
 hand demonstrates mosaic evolution of locomotor and manipulative abilities. Science
 333(6048), 1411-1417. https://doi.org/10.1126/science.1202625
- Kivell, T. L., Ostrofsky, K. R., Richmond, B. G., Drapeau, M. S. M., 2020. Metacarpals and manual
 phalanges. In: Zipfel, B., Richmond, B. G., & Ward, C. V. (Eds.), Hominin postcranial remains

- 1344 from Sterkfontein, South Africa, 1936-1995, Oxford University Press, Oxford, United 1345 Kingdom, p. 106-143. <u>https://doi.org/10.1093/oso/9780197507667.003.0009</u>.
- Kuman, K., 2007. The earlier Stone Age in South Africa: site context and the influence of cave
 studies. In: Pickering, T. R., Schick, K., & Toth, N., (Eds.), Breathing Life Into Fossils:
 Taphonomic Studies in Honor of C. K. (Bob) Brain. Stone Age Institute Press, Gosport, USA,
 p. 181-198.
- Kuman, K., 2009. The oldowan industry from sterkfontein caves, south africa. In: Schick, K. &
 Toth, N. (Eds.), The Cutting Edge: new approaches to the archaeology of human origins,
 Stone Age Institute Press, Gosport, USA, p. 151-169.
- Kuman, K. Clarke, R. J. 2000. Stratigraphy, artefact industries and hominid associations for
 Sterkfontein, Member 5. J. Hum. Evol. 38, 827-847.
 <u>https://doi.org/10.1006/jhev.1999.0392</u>
- Lague, M. R., Jungers, W. L., 1996. Morphometric variation in Plio-Pleistocene hominid distal
 humeri. Am. J. Phys. Anthrop. 101(3), 401-427.
- 1358 Leakey, L.S.B., 1959. A new fossil skull from Olduvai. Nature 184, 491-493.
- Leakey, L.S.B., Tobias, P.V., Napier, J.R., 1964. A new species of the genus *Homo* fromOlduvai Gorge. Nature 202, 7-9.
- Leakey, M.D., 1971. Olduvai Gorge. Vol 3. Excavations in Beds I and II, 1960–1963. Cambridge
 University Press, Cambridge, United Kingdom.
- Leakey, M.D., Clarke, R.J., Leakey, L.S.B., 1971. New Hominid skull from Bed I, Olduvai Gorge,
 Tanzania. Nature 232, 308-312.
- Leakey, M.G., Spoor, F., Brown, F.H., Gothogo, P.N., Kiairie, Ch., Leakey, L.N., McDougall, I.,
 2001. New hominin genus from eastern Africa shows diverse middle Pliocene lineages.
 Nature. 410, 433-439. <u>https://doi.org/10.1038/35068500</u>
- Leakey, R. E. F., 1974. Further evidence of lower Pleistocene hominids from east Rudolf, North
 Kenya, 1973. Nature, 248(5450), 653-656.
- Leakey, R.E.F., 1976. New hominids fossils from the Koobi Fora formation in northern Kenya,Nature 261, 574-576.
- Leakey, R. E. F., Walker, A. C., 1985. Further hominids from the Plio-Pleistocene of Koobi Fora,
 Kenya. Am. J. Phys. Anthrop. 67(2), 135-163.
- Lepre, C. J., Roche, H., Kent, D. V., Harmand, S., Quinn, R. L., Brugal, J.-P., Texier, P.-J., Lenoble,
 A., Feibel, C. S. 2011. An earlier origin for the Acheulian. Nature 477, 82-85.
- Lewis, J.E., Harmand, S. 2016. An earlier origin for stone tool making: implications for cognitive
 evolution and the transition to *Homo*. Philos Trans R Soc Lond B Biol Sci. 371(1698),
 20150233. <u>https://doi.org/10.1098/rstb.2015.0233</u>
- Li, H., Li, C., Kuman, K., 2017. Longgudong, an Early Pleistocene site in Jianshi, South China,
 with stratigraphic association of human teeth and lithics. Sci. China Earth Sci. 60, 452-462.
- Lordkipanidze, D., de León, M. S. P., Margvelashvili, A., Rak, Y., Rightmire, G. P., Vekua, A.,
 Zollikofer, C. P. E., 2013. A Complete Skull from Dmanisi, Georgia, and the Evolutionary
 Biology of early *Homo*. Science 342, 326-331.
- Lordkipanidze D., Jashashvili, T., Vekua, A., de León, M. S. P., Zollikofer, C. P. E., Rightmire, G.
 P., Pontzer, H., Ferring, R., Oms, O., Tappen, M., Bukhsianidze, M., Agusti, J., Kahlke, R.,
 Kiladze, G., Martinez-Navarro, B., Mouskhelishvili, A., Nioradze, M., Rook, L., 2007.
- 1387 Postcranial evidence from early *Homo* from Dmanisi, Georgia. Nature 449, 305-310.

- Macchi, R., Daver, G., Brenet, M., Prat, S., Hugheville, L., Harmand, S., Lewis, J., Domalain, M.,
 2021. Biomechanical demands of percussive techniques in the context of early stone
 toolmaking. J. R. Soc. Interface. 18, 20201044. <u>https://doi.org/10.1098/rsif.2020.1044</u>
- Mana, S., Hemming, S., Kent, D. V., Lepre, C. J., 2019. Temporal and stratigraphic framework
 for paleoanthropology sites within East-Central Area 130, Koobi Fora, Kenya. Frontiers in
 Earth Science, 7, 230. <u>https://doi.org/10.3389/feart.2019.00230</u>
- Manafzadeh, A. R., Gatesy, S. M., 2021. Paleobiological reconstructions of articular function
 require all six degrees of freedom. J. Anat. 239, 1516-1524.
 <u>https://doi.org/10.1111/joa.13513</u>
- Marzke, M.W., 1983. Joint functions and grips of the *Australopithecus afarensis* hand, with
 special reference to the region of the capitate. J. Hum. Evol. 12, 197-211.
- Marzke, M.W., 1997. Precision grips, hand morphology, and tools. Am J Phys Anthropol.102(1), 91-110.
- Marzke, M.W., 2013. Tool making, hand morphology and fossil hominins. Phil. Trans. R. Soc.
 Lond. B Biol Sci. 368(1630), 20120414. https://doi.org/doi: 10.1098/rstb.2012.0414
- Marzke, M.W., Marchant, L.F., McGrew, W.C., Reece, S.P., 2015. Grips and hand movements
 of chimpanzees during feeding in Mahale Mountains National Park, Tanzania. Am. J. Phys.
 Anthropol. 156(3), 317-26. <u>https://doi.org/10.1002/ajpa.22651</u>
- 1406 McDougall, I. A. N., Brown, F. H., 2008. Geochronology of the pre-KBS tuff sequence, Omo 1407 Group, Turkana Basin. J. Geol. Soc. London 165(2), 549-562.
- McHenry, L. J., 2012. A revised stratigraphic framework for Olduvai Gorge Bed I based on tuff
 geochemistry. J. Hum. Evol. *63*(2), 284-299. <u>https://doi.org/10.1016/j.jhevol.2011.04.010</u>
- McPherron, S. P., Alemseged, Z., Marean, C. W., Wynn, J. G., Reed, D., Geraads, D., Bobe, R.,
 Béarat, H. A., 2010. Evidence for stone-tool-assisted consumption of animal tissues before
 3.39 million years ago at Dikika, Ethiopia. Nature 466, 857-860.
- Mercader, J., Akuku, P., Boivin, N., Bugumba, R., Bushozi, P., Camacho, A., Carter, T. Clarke, S.,
 Cueva-Temprana, A., Durkin, P. Favreau, J., Fella, K., Haberle, S., Hubbard, S., Inwood, J.,
 Itambu, M., Koromo, S., Lee, P., Mohammed, A., Mwanbwiga, A., Olesilau, L., Patalano, R.,
 Roberts, P., Rule, S., Saladie, P., Siljedal, G., Soto, M., Umbsaar, J., Petraglia, M., 2021.
 Earliest Olduvai hominins exploited unstable environments~ 2 million years ago. Nat.
 Comm. 12(1), 3. doi: 10.1038/s41467-020-20176-2.
- Merrick, H.V., de Heinzelin, J., Haesaerts, P., Howell, F.C., 1973. Archaeological occurrences of
 Early Pleistocene age from the Shungura Formation, Lower Omo Valley, Ethiopia. Nature
 242, 572-575.
- Mgeladze, A., Lordkipanidze, D., Moncel, M.-H., Despriee, J., Chagelishvili, R., Nioradze, M.,
 Nioradze, G., 2011. Hominin occupations at the Dmanisi site, Georgia, Southern Caucasus:
 Raw materials and technical behaviours of Europe's first hominins. J. Hum. Evol. 60, 571596. https://doi.org/10.1016/j.jhevol.2010.10.008
- Moggi-Cecchi, J., Menter, C., Boccone, S., Keyser, A., 2010. Early hominin dental remains from
 the Plio-Pleistocene site of Drimolen, South Africa. J. Hum. Evol. 58 (5), 374-405.
 <u>https://doi.org/10.1016/j.jhevol.2010.01.006</u>
- Moyà-Solà, S., Kohler, M., Alba, D.M., Almécija, S., 2008. Taxonomic attribution of the Olduvai
 hominid 7 manual remains and the functional interpretation of hand morphology in robust
 australopithecines. Folia Primatol 79(4), 215-50. <u>https://doi.org/10.1159/000113458</u>
- 1432 Morgan, L. E., Renne, P. R., Kieffer, G., Piperno, M., Gallotti, R., Raynal, J. P., 2012. A 1433 chronological framework for a long and persistent archaeological record: Melka Kunture,
- 1434 Ethiopia J. Hum. Evol. 62(1), 104-115. <u>https://doi.org/10.1016/j.jhevol.2011.10.007</u>

- 1435 Napier J.R., 1956. The prehensile movements of the human hand. J Bone Joint Surg Br. 38 1436 B(4), 902-13. <u>https://doi.org/10.1302/0301-620X.38B4.902</u>
- 1437 Napier, J. R., 1962. Fossil hand bones from Olduvai Gorge. Nature 196, 409-411.
 1438 https://doi.org/10.1038/196409a0
- Neufuss, J., Robbins, M., Baeumer, J., Hulme, T., Kivell, T., 2018. Manual skills for food
 processing by mountain gorillas in Bwindi Impenetrable National Park, Uganda. Biol. J. Linn.
 Soc.127(3), 543-562.
- Parés, J. M., Pérez-González, A., Rosas, A., Benito, A., Bermúdez de Castro, J. M., Carbonell, E.,
 Huguet, R., 2006. Matuyama-age lithic tools from the Sima del Elefante site, Atapuerca
 (northern Spain). J. Hum. Evol. 50, 163-169. <u>https://doi.org/10.1016/j.jhevol.2005.08.011</u>
- Pickering, R., Dirks, P. H., Jinnah, Z., De Ruiter, D. J., Churchill, S. E., Herries, A. I., Woodhead,
 J. D. H., Berger, L. R., 2011. *Australopithecus sediba* at 1.977 ma and implications for the
 origins of the genus *Homo*. Science 333, 1421-1423. https://doi.org/10.1126/science.
- Pickering, T. R., Heaton, J. L., Clarke, R. J., Stratford, D., 2019. Hominin vertebrae and upper
 limb bone fossils from Sterkfontein Caves, South Africa (1998-2003 excavations). Am J Phys
 Anthropol. 168, 459-480.
- Plummer T., 2004. Flaked stones and old bones: biological and cultural evolution at the dawn
 of technology. Am J Phys Anthropol. 125, Suppl 39, 118-64. doi: 10.1002/ajpa.20157.
- 1453Plummer, T., Bishop, L. C., Ditchfield, P., Hicks, J., 1999. Research on Late Pliocene oldowan1454sites at kanjera south, Kenya. J. Hum. Evol. 36(2), 151-170.1455https://doi.org/10.1006/jhev.1998.0256
- Plummer, T., Oliver, J. S., Finestone, E. M., Ditchfield, P. W., Bishop, L. C., Blumenthal, S. A.,
 Lemorini, C., Caricola, I., Bailey, S., Herries, A.L.R., Parkinson, J.A., Whitfield, E., Hertel, F.,
 Kinyanjui, R.N., Vincent, T., Li., Y., Louys, J., Frost, S.R., Braun, D.R., Reeves, J.S., Early,
 E.D.G., Onyango, B., Lamela-Lopez, R., Forrest, F.L., He, H., Lane, T.P., Frouin, M., Nomade,
 S., Wilson, E.P., Bartilol, S., Rotich N.K., Potts, R., 2023. Expanded geographic distribution
 and dietary strategies of the earliest Oldowan hominins and *Paranthropus*. Science,
- 1462 379(6632), 561-566. DOI: 10.1126/science.abo7452
- Potts, R., 2012. Evolution and environmental change in early human prehistory. Annu. Rev.Anthropol., 41, 151-167.
- Prat, S., 2002. Anatomical study of the skull of the Kenyan specimen KNM-ER 1805: a re-evaluation of its taxonomic allocation? C. R. Palevol, 1(1), 27-33.
- Prat, S., 2018. First hominin settlements out of Africa. Tempo and dispersal mode: Review and
 perspectives. C. R. Palevol. 17(1-2), 6-16. <u>https://doi.org/10.1016/j.crpv.2016.04.009</u>
- Prat, S., 2022. Emergence of the genus *Homo*: From concept to taxonomy. L'Anthropologie,
 126(4), 103068. <u>https://doi.org/10.1016/j.anthro.2022.103068</u>
- Prat, S., Brugal, J.-P., Roche, H., Texier, P.-J. 2003. Nouvelles découvertes de dents d'hominidés
 dans le membre Kaitio de la formation de Nachukui (1,65–1,9 Ma), Ouest du lac Turkana
 (Kenya). C. R. Palevol, 2, 685-693.
- Prat, S., Brugal, J. P., Tiercelin, J. J., Barrat, J. A., Bohn, M., Delagnes, A., Harmand, S., Kimeu,
 K., Kibunjia, M., TExier, P-J., Roche, H., 2005. First occurrence of early *Homo* in the Nachukui
 formation (West Turkana, Kenya) at 2.3-2.4 Myr. J. Hum. Evol. 49(2), 230-240.
 https://doi.org/10.1016/j.jhevol.2005.03.009
- Proffitt, T, Bargalló, A, de la Torre, I. 2022. The effect of raw material on the identification of
 knapping skill: A case study from Olduvai Gorge, Tanzania. J Archaeol Method Theory.
- 1480 29(1): 50-82. https://doi.org/10.1007/s10816-021-09511-z

- 1481
- Putt S.S., 2016. Human brain activity during stone tool production: tracing the evolution of cognition and language, Doctor of Philosophy, University of Iowa.
- Putt, S. S., Wijeakumar, S., Franciscus, R. G., Spencer, J. P., 2017) The functional brain networks
 that underlie Early Stone Age tool manufacture. Nat. Human Behav. 1(6), 0102.
- 1486 Ramirez-Rozzi, F., 1997. Les hominides du Plio-Pleistocene de la vallee de l'Omo. Cahiers de 1487 paleoanthropologie. CNRS editions, Paris, France.
- Richmond, B.G., Green, D.J., Lague, M.R., Chirchir, H., Behrensmeyer, A.K., Bobe, R., Bamford,
 M.K., Griffin, N.L., Gunz, P., Mbua, E., Merritt, S.R., Pobiner, B., Kiura, P., Kibunjia, M.,
 Harris, J.W.K., Braun, D.R., 2020. The upper limb of *Paranthropus boisei* from Ileret, Kenya.
 J Hum Evol. 141, 102727. https://doi.org/10.1016/j.jhevol.2019.102727
- 1492 Ricklan, D. E., 1987. Functional anatomy of the hand of *Australopithecus africanus*. J. Hum 1493 Evol.16, 643-664.
- Roche, H, Blumenschine, R.J, Shea, J.J., 2009. Origins and adaptations of early genus *Homo*:
 what archeology tells us. In: Grine, F.E., Fleagle, J.G., Leakey, R.E. (Eds), The First Humans:
 Origin and Early Evolution of the Genus *Homo*. Springer, Dordrecht, p. 135–147.
- Roche, H., Brugal, J.-P., Delagnes, A., Feibel, C., Harmand, S., Kibunjia, M., Prat, S., Texier, P.-J.
 2003. Les sites archéologiques plio-pléistocènes de la formation de Nachukui, Ouest-
- 1499 Turkana, Kenya: bilan synthétique 1997-2001. C.R. Palevol. 2, 663-673.
- Roche, H., Delagnes, A., Brugal, J-P., Feibel, C.S., Kibunjia, M., Mourre, V., Texier, P-J., 1999.
 Early hominid stone tool production and technical skill 2.34 Myr ago, in West Turkana,
 Kenya. Nature 399, 57-60. <u>https://doi.org/10.1038/19959</u>
- 1503Roche, H., Tiercelin, J.J., 1977. Découverte d'une industrie lithique ancienne *in situ* dans la1504formation d'Hadar, Afar central, Ethiopie. C.R. Acad. Sci. Paris D. 284, 174-187.
- Rolian, C., Gordon, A.D., 2013. Reassessing manual proportions in *Australopithecus afarensis*.
 Am J Phys Anthropol. 152(3), 393-406. <u>https://doi.org/10.1002/ajpa.22365</u>
- Rolian, C., Lieberman, D. E., Zermeno, J. P. 2011. Hand biomechanics during simulated stone
 tool use. J. Hum. Evol. (1), 26-41. <u>https://doi.org/10.1016/j.jhevol.2011.01.008</u>
- Sano, K., Beyene, Y., Katoh, S., Koyabu, D., Endo, H., Sasaki, T., Asfaw, B., Suwa, G., 2020. A
 1.4-million-year-old bone handaxe from Konso, Ethiopia, shows advanced tool technology
 in the early Acheulean. Proc. Nat. Acad. Sci. USA, 117(31), 18393-18400.
- Schick, K, Toth, N., 2006. An Overview of the Oldowan Industrial Complex: The sites and the
 nature of their evidence. In: Toth N. & Schick K. (Eds.): The Oldowan: Case Studies into the
 Earliest Stone Age, Stone Age Institute Press, Gosport, USA, p 3-42.Schrenk, F., Bromage,
 T. G., Betzler, C. G., Ring, U., Juwayeyi, Y. M., 1993. Oldest *Homo* and Pliocene biogeography
- 1516 of the Malawi rift. Nature, 365(6449), 833-836.
- Senut, B., 1979. Comparaison des Hominidés de Gomboré IB et de Kanapoi; deux pièces du genre *Homo*? Bull. Mem. Soc. Anthropol. Paris, XIII Série 6, 111-117.
- Semaw, S., Renne, P., Harris, J.W.K., Feibel, C.S., Bernor, R.L., Fesseha, N., Mowbrav, K., 1997.
 2.5-million-year-old stone tools from Gona, Ethiopia, Nature 385, 333-336.
- 1521 Semaw, S., Rogers, M. J., Simpson, S. W., Levin, N. E., Quade, J., Dunbar, N., W. C. McIntosh, I.
- 1522 Cáceres, G. E. Stinchcomb, R. L. Holloway, F. H. Brown, R. F. Butler, D. Stout, Everett, M.,
- 1523 2020. Co-occurrence of Acheulian and Oldowan artifacts with *Homo erectus* cranial fossils
- 1524 from Gona, Afar, Ethiopia. Sci. Adv. 6(10), eaaw4694. DOI: 10.1126/sciadv.aaw4694
- Semaw, S., Rogers, M.J., Quade, J., Renne, P.R., Bulter, R.F., Domínguez-Rodrigo, M., Stout, D.,
 Hart, W.S., Pickering, T., Simpson, S.W. 2003. 2.6-Million-year-old stone tools and

- 1527
 associated bones from OGS-6 and OGS-7, Gona, Afar, Ethiopia, J. Hum. Evol, 45, 169-177.

 1528
 https://doi.org/10.1016/S0047-2484(03)00093-9
- Semaw, S., 2000. The world's oldest stone artefacts from Gona, Ethiopia: their implications for
 understanding stone technology and patterns of human evolution between 2.6-1.5 million
 years ago. *Journal of Archaeological Science*, 27(12), 1197-1214.
- Semaw, S. 2006. The oldest stone artifacts from Gona (2.6-2.5 Ma), Afar, Ethiopia: implications
 for understanding the earliest stages of stone knapping. In Toth N. & Schick K. (Eds.): The
 Oldowan: Case Studies into the Earliest Stone Age, Stone Age Institute Press, Gosport, USA,
 p. 43-75.
- Sherwood, R. J., Ward, S. C., Hill, A., 2002. The taxonomic status of the Chemeron temporal
 (KNM-BC 1). J. Hum. Evol. 42(1-2), 153-184. <u>https://doi.org/10.1006/jhev.2000.0409</u>
- Skinner, M.M., Stephens, N.B., Tsegai, Z.J., Foote, A.C., Nguyen, N.H., Gross, T., Pahr, D.H.,
 Hublin, J.J., Kivell, T.L., 2015. Human evolution. Human-like hand use in *Australopithecus africanus*. Science. 347(6220), 395-399. <u>https://doi.org/10.1126/science.1261735</u>.
- Skinner, M. M., Leakey, M. G., Leakey, L. N., Manthi, F. K., Spoor, F., 2020. Hominin dental
 remains from the Pliocene localities at Lomekwi, Kenya (1982–2009). J. Hum. Evol. 145,
 102820. <u>https://doi.org/10.1016/j.jhevol.2020.102820</u>
- Spoor, F., Gunz, P., Neubauer, S., Stelzer, S., Scott, N., Kwekason, A., Dean. C., 2015.
 Reconstructed *Homo habilis* type OH 7 suggests deep-rooted species diversity in early
 Homo, Nature, 519: 83-86. <u>https://doi.org/10.1038/nature14224</u>
- Stammers, R.C., Caruana, M.V., Herries, A.I.R., 2018. The first bone tools from Kromdraai and
 stone tools from Drimolen, and the place of bone tools in the South African Earlier Stone
 Age. Quat. Int. 495, 87–101
- Steininger, C., Berger, L.R., Kuhn, B.F., 2008. A partial skull of *Paranthropus robustus* from
 Cooper's Cave, South Africa. S. Afr. J. Sci. 104, 143–146
- 1552Stout, D., Chaminade, T., 2007. The evolutionary neuroscience of tool making.1553Neuropsychologia,45(5),1091-1100.
- 1554 <u>https://doi.org/10.1016/j.neuropsychologia.2006.09.014</u>
- Stout, D., Chaminade, T., 2012. Stone tools, language and the brain in human evolution. Philos.
 Trans. R. Soc. B: Biol. Sci. 367(1585), 75-87. doi: <u>10.1098/rstb.2011.0099.</u>
- 1557Stout, D., Hecht, E., Khreisheh, N., Bradley, B., Chaminade, T., 2015. Cognitive demands of1558LowerPaleolithictoolmaking.PLoSOne,10(4),e0121804.1559https://doi.org/10.1371/journal.pone.0121804
- Stout, D., Quade, J., Semaw, S. and Levin, N.E. 2005. Raw material selectivity of the earliest
 stone toolmakers at Gona, Afar, Ethiopia. J. Hum. Evol, 48, 365-380.
- 1562Stout, D., Semaw, S., Rogers, M.J. and Cauche, D. 2010. Technological variation in the earliest1563OldowanfromGona,Afar,Ethiopia.J.Hum.Evol.58,474-491.1564https://doi.org/10.1016/j.jhevol.2010.02.005
- Stout, D., Toth, N., Schick, K., Chaminade, T., 2008. Neural correlates of Early Stone Age
 toolmaking: technology, language and cognition in human evolution. Philos. Trans. R. Soc.
 B: Biol. Sci., 363(1499), 1939-1949. doi: 10.1098/rstb.2008.0001
- Stout, D., Toth, N., Schick, K., Stout, J., Hutchins, G., 2000. Stone tool-making and brain
 activation: position emission tomography (PET) studies. J. Archaeol. Sci., 27(12), 12151223.
- 1571Susman, R.L., 1988. Hand of Paranthropus robustus from Member 1, Swartkrans: fossil1572evidencefortoolbehavior.Science240(4853),781-784.1573https://doi.org/10.1126/science.3129783

- Susman, R.L., 1991. Who made the Oldowan tools? Fossil evidence for tool behavior in Plio Pleistocene hominids. J. Anthrop. Res. 47, 129–151.
- Susman, R.L., 1994. Fossil evidence for early hominid tool use. Science. 265(5178), 1570-1573.
 <u>https://doi.org/10.1126/science.8079169</u>
- Susman, R.L., 1998. Hand function and tool behavior in early hominids. J Hum Evol. 35(1), 2346. <u>https://doi.org/10.1006/jhev.1998.0220</u>
- Susman, R. L., Creel, N., 1979. Functional and morphological affinities of the subadult hand 1580 from Olduvai 1581 (OH 7) Gorge. Am. J. Phys. Anthrop. 51, 311-331. https://doi.org/10.1002/ajpa. 1330510303 1582
- Sutton, M.B., Kuman, K., Steininger, C., 2017. Early Pleistocene stone artefacts from Cooper's
 cave, South Africa. South African Archaeol. Bull. 72, 156–161.
- Suwa, G., White, T. D., Howell, F. C., 1996. Mandibular postcanine dentition from the Shungura
 Formation, Ethiopia: Crown morphology, taxonomic allocations, and Plio-Pleistocene
 hominid evolution. Am. J. Phys. Anthrop. 101(2), 247-282.
- Suwa, G., 1990. A comparative analysis of hominid dental remains from the Shungura andUsno Formations, Omo Valley, Ethiopia. Ph. D Thesis. University of California, Berkeley.
- Suwa, G., Asfaw, B., Beyene, Y., White, T. D., Katoh, S., Nagaoka, S., Nakaya, H., Uzawa, K.,
 Renne, P., WoldeGabriel, G., 1997. The first skull of *Australopithecus boisei*. Nature
 389(6650), 489-492.
- Suwa, G. E. N., Asfaw, B., Haile-Selassie, Y., White, T. I. M., Katoh, S., WoldeGabriel, G., Hart, W.K.,
 Nakaya, H., Beyene, Y., 2007. Early pleistocene *Homo erectus* fossils from Konso, southern Ethiopia.
 Anthropol. Sci. 115(2), 133-151.
- Tamrat, E., Thouveny, N., Taieb, M., Brugal, J. P., 2014. Magnetostratigraphic study of the
 Melka Kunture archaeological site (Ethiopia) and its chronological implications. Quat. Inter.
 343, 5-16.
- 1599 Tobias, P.V. 1966. Fossil Hominid Remains from Ubeidiya, Israel. Nature 211, 130–133.
- Tobias, P.V., 1991. Olduvai gorge Vol 4a et 4b. Homo habilis: skulls, endocasts and teeth.
 Cambridge University Press, Cambridge, United Kingdom.
- Tocheri, M. W., Marzke, M. W., Liu, D., Bae, M., Jones, G. P., Williams, R. C., Razdan, A., 2003.
 Functional capabilities of modern and fossil hominid hands: Three-dimensional analysis of
 trapezia. Am. J. Phys. Anthrop. 122, 101-112. <u>https://doi</u>. org/10.1002/ajpa.10235
- Villmoare, B. Kimbel, W.H., Seyoum, C., Campisano, C.J., DiMaggio, E.N., Rowan, J., Braun D.R.,
 Arrowsmith, Reed, K.E., 2015. Early *Homo* at 2.8 Ma from Ledi-Geraru, Afar, Ethiopia.
 Science 347, 1352-1354. <u>https://doi.org/10.1126/science.aaa1343</u>
- Ward, C.V., Kimbel, W.H., Harmon, E.H., Johanson, D.C., 2012. New postcranial fossils of
 Australopithecus afarensis from Hadar, Ethiopia (1990-2007). J Hum Evol. 63(1), 1-51.
 <u>https://doi.org/10.1016/j.jhevol.2011.11.012</u>
- Ward, C. V., Tocheri, M. W., Plavcan, J. M., Brown, F. H., Manthi, F. K., 2014. Early Pleistocene
 third metacarpal from Kenya and the evolution of modern human-like hand morphology.
 Proc. Natl Acad. Sci. USA 111, 121-124.
- Williams-Hatala, E. M., Hatala, K. G., Key, A., Dunmore, C. J., Kasper, M., Gordon, M., Kivell, T.
 L., 2021. Kinetics of stone tool production among novice and expert tool makers. Am. J.
 Phys. Anthrop. 174, 714-727. <u>https://doi.org/10.1002/ajpa.24159</u>
- Williams-Hatala, E. M., Hatala, K. G., Gordon, M., Key, A., Kasper, M., Kivell, T. L., 2018. The
 manual pressures of stone tool behaviors and their implications for the evolution of the
 human hand. J. Hum. Evol. 119, 14-26. https://doi.org/10.1016/j.jhevol.2018.02.008

- Wood, B.A., 1991. Koobi Fora Project Vol 4: Hominid Remains From Koobi Fora, ClarendonPress, Oxford, United Kingdom.
- 1622 Wood, B.A. 1997. The oldest whodunnit in the world. Nature 385:292–293.
- Wood, B., Constantino, P., 2007. *Paranthropus boisei*: fifty years of evidence and analysis. Am.
 J. Phys. Anthrop. 134(S45), 106-132.
- Wood, B., Leakey, M., 2011. The Omo-Turkana Basin fossil hominins and their contribution to
 our understanding of human evolution in Africa. Evol. Anthrop. 20(6), 264-292.
- Zanolli, C., Dean, M.C., Assefa, Y., Bayle, P., Braga, J., Condemi, S., Endalamaw, M., Engda 1627 Redae, B., Macchiarelli, R., 2016. Structural organization and tooth development in a Homo 1628 aff. erectus juvenile mandible from the Early Pleistocene site of Garba IV at Melka Kunture, 1629 1630 Ethiopian highlands. Am. J. Phys. Anthropol. 162(3), 533-549, http://dx.doi.org/10.1002/ajpa.23135 1631
- 1632 Zhu, R. X., Potts, R., Pan, Y. X., Yao, H. T., Lü, L. Q., Zhao, X., Gao, X., Chen, L. W., Gao, F., Deng,
 1633 C. L., 2008. Early evidence of the genus *Homo* in East Asia. J. Hum. Evol. 55, 1075-1085.
- 1634 <u>https://doi.org/10.1016/j.jhevol.2008.08.005</u>
- 1635

1636	Figure and table lists
1637	
1638 1639	Figure 1: Chronological repartition of hominin species and cultural assemblage during the Pliocne and early Pleistocene in Africa.
1640	
1641 1642	Figure 1. Répartition chronologique des espèces d'hominines et des assemblages Itihiques en Afrique au cours du Pliocene et du Pléistocène inférieur.
1643	
1644	Figure 2. Localisation of the archaeological sites in association (HC1 and HC 2) with hominin remains.
1645	
1646 1647	Figure 2 : Localisation des sites archéologiques en association (HC1 and HC 2) avec des restes d'hominines.
1648	
1649 1650	Table 1 : Categories of attribution and association between hominin remains and archaeological assemblages.
1651	
1652 1653	Tableau 1: Catégories d'attribution et d'association entre les restes d'hominines et les assemblages archéologiques.
1654	
1655 1656	Table 2: Association between hominins remains and archaeological assemblage with a High Confidence level: HC (local scale).
1657	
1658 1659	Tableau 2: Association entre les restes d'hominines et les assemblages archéologiques ayant un niveau de confidence important HC (échelle locale).

Category	Definition
HC1	The hominin remains have been found in the archaeological site (<i>in situ</i> during
	the excavation, or at the surface of the site due to erosion processes)
HC 1 A	when the taxonomy of the hominin remain is clear
HC 1 B	when the taxonomy of the hominin remain is debated
HC 2	The hominin remains have been found few meters (less than 400 meters) from
	the excavation but in the same chrono-stratigraphical layers
HC 2 A	when the taxonomy of the hominin remain is clear
HC 2 B	when the taxonomy of the hominin remain is debated

1662 Table 1 : Categories of attribution and association between hominin remains and

1663 archeological assemblages.

1664

- 1665 Tableau 1: Catégories d'attribution et d'association entre les restes d'hominines et les
- 1666 assemblages archéologiques.

1668	
1669	


Age (Ma)	Region	Locality	Industry	References	Hominins	References	Category of reliability
3.03 -2.58	Lake Victoria (Kenya)	Nyayanga, excavation 3	Oldowan	Plummer et al., 2023	KNM-NG 77316 Paranthropus	Plummer et al., 2023	HC 1 A
2.34 +/- 0.07	Hadar (Ethiopia)	AL 666	Oldowan	Kimbel et al., 1996	AL 666-1 Homo aff. H. habilis	Kimbel et al., 1997	HC 1 A
2.34 +/- 0.04	West Turkana (Kenya)	Lokalalei 1, Gajh5	Oldowan	Prat et al., 2005	KNM-WT 42718 Early <i>Homo</i>	Prat et al., 2005	HC 2 A
2.32 Ma to 2.23	Omo Valley (Ethiopia)	Omo 57	Oldowan	Coppens et al., 1973, Chavaillon, 1976	Omo 57/5-1972-371, P. aff. aethiopicus	Coppens, 1977, Suwa, 1990; Wood et Leakey 2011	HC 1 A
1.95-1.90	Fejej (Ethiopia)	FJ-1a	Oldowan	e.g. de Lumley and Beyene 2004	Fj-1-Hd1, Hd3, Hd4, <i>Homo</i> aff. <i>H. habilis</i>	de Lumley & Marchal 2004	HC 2 A
1.87-1.84	Olduvai Gorge (Tanzania)	DK	Oldowan	Leakey, 1971, de la Torre and Mora, 2005	OH 24 Homo habilis	Leakey et al., 1971	HC 2 A
1.84	Olduvai (Tanzania)	FLK NN level 3	Oldowan	Leakey et al., 1964 Leakey, 1971	OH 7 cranial remains <i>H. habilis</i> (Holotype)	Leakey et al. 1964	HC 1 A
1.84	Olduvai (Tanzania)	FLK NN level 3	Oldowan	Leakey et al., 1964, Leakey, 1971	OH 7 hand bones H. habilis or Paranthropus	Leakey et al. 1964 Moyà-Solà et al., 2008	HC 1 B
1.84	Olduvai (Tanzania)	FLK NN level 3	Oldowan	Leakey et al., 1964, Leakey , 1971	OH 8 foot bones <i>H. habilis</i> (paratype)	Leakey et al. 1964 Day and Napier, 1964	HC 1 B
1.84	Olduvai Gorge (Tanzania)	FLK level 22	Oldowan	Leakey, 1971	OH 5 Holotype of <i>P. boisei</i>	Leakey, 1959	HC 1 A
1.84	Olduvai Gorge (Tanzania)	FLK level 22	Oldowan	Leakey, 1971	OH 6 H. habilis	Tobias, 1991	HC 1 A
1.84	Olduvai Gorge (Tanzania)	FLK level 22	Oldowan	Leakey, 1971	OH 35 P. boisei or H. habilis	Leakey, 1964	HC 1 B

1.84	Olduvai Gorge (Tanzania)	PTK site	Oldowan	Domínguez-Rodrigo, 2015	OH 86 <i>Homo erectus</i> -like	Domínguez-Rodrigo, 2015 Díez-Martín et al, 2015	HC 1 B
Circa 1.82	Olduvai Gorge (Tanzania)	Ewass Oldupa	Oldowan	Mercader et al. 2021	OH 65 Homo habilis	Blumenshine et al., 2012. Clarke, 2012	HC 2 A
1.79	West Turkana (Kenya)	Kokiselei 1 FxJh6	Oldowan	Roche et al., 2003	KNM-WT 37744, 37747, 37748, 37100 <i>P. boisei</i>	Prat et al., 2003	HC 1 A
1.78-1.52 (Mana et al., 2019) 1.85 (Feibel et al., 1989)	Koobi Fora (Kenya)	FxJj 38	Oldowan	Harris et Herbich, 1978	KNM-ER 1805 <i>Homo</i> sp. but taxonomical allocation under discussion	Leakey, 1974 Prat, 2002	HC 1 B
1.78-1.52 (Mana et al., 2019) 1.85 (Feibel et al., 1989)	Koobi Fora (Kenya)	FxJj 38	Oldowan	Harris et Herbich, 1978	KNM-ER 1806 P. boisei	Leakey, 1974 Wood, 1991	HC 1 A
1.75 +/- 0.05	West Turkana (Kenya)	Naiyena Engol 1 FxJh5	Oldowan	Roche et al., 2003	KNM-WT 37745 <i>Homo</i> aff. <i>H. ergaster</i>	Prat et al., 2003	HC 1 A
Slightly older than 1.7	Melka Kunture (Ethiopia)	Garba IV, level E	Oldowan	Gallotti and Mussi, 2015	MK 81 GAR IVE 0043 early Homo erectus/ H. ergaster	Condemi, 2004	HC 1 A
Circa 1.67	Olduvai Gorge (Tanzania)	MNK Skull	Oldowan	Leakey, 1971 de la Torre et al., 2021	OH 13 Paratype Homo habilis OH 15, Homo habilis	Leakey et al., 1964, Tobias 1991	HC 1 A
1.65 +/- 0.05	Koobi Fora (Kenya)	FxJj 20	Oldowan	Harris et Herbich, 1978	KNM-ER 3230 P. boisei	Wood, 1991	HC 1 A
1.6-1.5	Melka Kunture (Ethiopia)	Gombore I	Oldowan	Chavaillon et al., 1977 Chavaillon and Berthelet 2004	Gombore IB-7594 early Homo erectus Homo sp.	Coppens, 2004 Chavaillon et al., 1977	HC 1 B

						Di Vincenzo et al., 2015	
1.6-1.5	Afar Region (Ethiopia)	DAN 5	Oldowan and Acheulean	Semaw et al. 2020	DAN5/P1 early Homo erectus	Semaw et al., 2020, Baab et al., 2022, Bruner et al., 2023	HC 1 A
1.44	Konso-Gardula (Ethiopia)	KGA 4E	Acheulean	Asfaw et al., 1992 Beyene et al., 2013	KGA 4 E <i>Homo</i> sp.	Asfaw et al., 1992	HC 1 A
1.42 +/0.07	Chesowanja (Kenya)	GNJ1/6E Gn J2	Oldowan Oldowan	Gowlett et al. 1981	KNM-CH 1, 302, 304 P. boisei	Carney et al., 1971, Bishop et al., 1975, Gowlett et al. 1981,	HC 2A
1.4	Konso-Gardula (Ethiopia)	KGA 10-A10	Acheulean	Asfaw et al. 1992 Beyene et al., 2013 Beyene et al., 2014	KGA 10-1; KGA KGA10- 620 (right parietal fragment), KGA 10-656 <i>Homo erectus</i>	Asfaw et al. 1992 Suwa et al., 2007	HC 2 A
1.4	Konso-Gardula (Ethiopia)	KGA 10-A10	Acheulean	Asfaw et al., 1992 Beyene et al., 2013 Beyene et al., 2014	KGA10-506, KGA10-525 ; KGA10-565, KGA10-570, KGA10-900, KGA10-1455, KGA10-1720, KGA10- 2705, KGA10-2741 <i>P. boisei</i>	Suwa et al., 1997	HC 2 A
1.34	Olduvai Gorge (Tanzania)	BK site level 4, Bed II	Oldowan	Dominguez-Rodrigo et al., 2009	OH 80 P. boisei	Dominguez-Rodrigo et al., 2013	HC 1 A
1.26 +/-0.03	Afar Region (Ethiopia)	BSN12	Oldowan and Acheulean	Semaw et al. 2020	BSN12/P1 Homo erectus	Semaw et al., 2020, Baab et al., 2022	HC 1 A

1671 Table 2 : Association between hominins remains and archeological assemblage with a High Confidence level: HC (local scale)

1673 Tableau 2: Association entre les restes d'hominines et les assemblages archéologiques ayant un niveau de confidence important HC (échelle locale)



Paranthropus

early Homo

Homo sp.

early Homo erectus

Oldowan (