



HAL
open science

Correction to: An accurate set of H₃O⁺ – H₂ collisional rate coefficients for non-LTE modelling of warm interstellar clouds

Sándor Demes, François Lique, Alexandre Faure, Floris F S van der Tak

► To cite this version:

Sándor Demes, François Lique, Alexandre Faure, Floris F S van der Tak. Correction to: An accurate set of H₃O⁺ – H₂ collisional rate coefficients for non-LTE modelling of warm interstellar clouds. *Monthly Notices of the Royal Astronomical Society*, 2024, 527 (2), pp.2573-2574. 10.1093/mnras/stad3281 . hal-04323928

HAL Id: hal-04323928

<https://hal.science/hal-04323928v1>

Submitted on 5 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Correction to: An accurate set of $\text{H}_3\text{O}^+ - \text{H}_2$ collisional rate coefficients for non-LTE modelling of warm interstellar clouds

by Sándor Demes¹,¹★ François Lique,¹★ Alexandre Faure²,²★ and Floris F. S. van der Tak³

¹Univ Rennes, CNRS, IPR (Institut de Physique de Rennes) - UMR 6251, F-35000 Rennes, France

²IPAG, Université Grenoble Alpes & CNRS, CS 40700, F-38058 Grenoble, France

³SRON Netherlands Institute for Space Research & Kapteyn Astronomical Institute, University of Groningen, 9747 AD Groningen, the Netherlands

Key words: errata, addenda – astrochemistry – molecular data – molecular processes – radiative transfer – methods: laboratory: molecular – ISM: molecules.

We have discovered errors in our recent paper Demes et al. (2022). Indeed, the supplementary material contains some incorrect theoretical data due to a software bug in the calculations of high energy cross sections. The cross sections for *para*- H_3O^+ collision with *ortho*- H_2 are affected at total energies above 800 cm^{-1} . Consequently, most of the high temperature rate coefficients for the *para*- $\text{H}_3\text{O}^+ + \text{ortho}$ - H_2 nuclear spin configurations are affected too. At kinetic temperatures lower than $T_{\text{kin}} = 100 \text{ K}$ the impact due to the correction of the cross sections is usually very minor: the average relative change in the magnitude of the rate coefficients is less than 0.5 per cent including all transitions in the analysis and the largest state-to-state change is below 4 per cent. These corrections are much smaller than the global accuracy of the collisional data (25–30 per cent) estimated by Demes et al. (2022). While the average relative difference for the rate coefficients at temperatures 150, 200, and 300 K are also low (0.5, 1.5, and 5.1 per cent, respectively), some specific transitions are significantly impacted. In particular, the largest state-specific correction at 150 K is about 15 per cent, while at 200 and 300 they reach ~ 41 and ~ 71 per cent, respectively. Except for some particular cases, the magnitude of the rate coefficients decreased after the corrections have been made in the cross section calculations. Large corrections (i.e. those above the global accuracy of the data) are almost entirely related to high-lying states (with $j_k^e \geq 5_3^+$, see table A1 in the Appendix of the original paper).

Regarding the radiative transfer modellings, the corrected data have a moderate impact, except for some specific transitions. Thus, most of the radiation temperatures (T_{R}) reported by Demes et al.

(2022) are well-reproduced with the corrected rate coefficients. At lower kinetic temperatures (50 and 100 K), the largest correction in the ratios $T_{\text{R}}^{\text{curr.work}}/T_{\text{R}}^{\text{LAMDA}}$ (see fig. 6 in the original paper) is less than 0.5 per cent and on average less than 0.1 per cent. At 200 and 300 K we found slightly larger impact due to the corrections, up to ~ 8 per cent for specific transitions with a mean of 0.5–4.0 per cent, depending on the temperature, H_2 and column densities. It is worth to mention however that at 10^6 cm^{-3} molecular and $N = 10^{16} \text{ cm}^{-2}$ column densities we found two strongly impacted transitions with larger changes in the T_{R} -ratios. In particular, after the corrections they decrease by ~ 37 per cent for the 364 GHz $3_2^+ \rightarrow 2_2^-$ transition, while for the 4241 GHz $4_1^- \rightarrow 3_1^+$ transition at 300 K the T_{R} -ratio decreases by a factor of ~ 40 . This strong impact is however due to the large opacity of the line (> 100) which makes the convergence of RADEX difficult.

We would like to emphasise that the changes found due to the subsequent corrections do not affect the general discussion or the conclusion of the original paper. The corrected magnitude of the cross sections, rate coefficients, and radiation temperatures fully supports the original findings of the paper and only gives quantitative corrections. The updated set of collisional data is made available within the current supplementary material online as well as through the EMAA, LAMDA, and BASECOL databases.

Since the corrections of the data affected some values plotted in fig. 6 of the original paper, we provide the updated figures in the present notice as Fig. 1.

The authors apologize for any inconvenience due to the possible use of the incorrect supplementary data.

* E-mail: sandor.demes@univ-rennes.fr (SD); francois.lique@univ-rennes.fr (FL); alexandre.faure@univ-grenoble-alpes.fr (AF)




Figure 1. [Correction to fig. 6 originally published by Demes et al. (2022)] The relative ratio of radiation temperatures at 50, 100, 200, and 300 K kinetic temperatures for the majority of the observed radiative H_3O^+ transitions calculated for diffuse [panel (a), $n(\text{H}_2) = 10^3 \text{ cm}^{-3}$] and dense [panel (b), $n(\text{H}_2) = 10^6 \text{ cm}^{-3}$] molecular cloud conditions. The $T_R^{\text{curr.work}}$ data is computed from our new set of rate coefficients, while the $T_R^{\text{LAMDA}^*}$ set is calculated based on the corresponding collisional data from the LAMDA data base. The colour bars correspond for radiative transfer calculations in the optically thin environments ($N = 10^{13} \text{ cm}^{-3}$), while the hatch-filled bars show the same quantities, but for optically thick regions ($N = 10^{16} \text{ cm}^{-3}$).

REFERENCE

Demes S., Lique F., Faure A., van der Tak F. F. S., 2022, *MNRAS*, 518, 3593

SUPPORTING INFORMATION

Supplementary data are available at *MNRAS* online.

H3O+ - H2 RADEX input with rate coefficients.zip

H3O+ - H2 collisional cross sections.zip

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.