



# **Yellow-legged gull populations (*Larus michahellis*) link the history of landfills to soil eutrophication and time-related vegetation changes on small Mediterranean islands**

Clémentine Mutillod, Teddy Baumberger, Pascale Prudent, Arne Saatkamp, Eric Vidal, Lidwine Le-Mire-Pecheux, Laurence Affre

## **► To cite this version:**

Clémentine Mutillod, Teddy Baumberger, Pascale Prudent, Arne Saatkamp, Eric Vidal, et al.. Yellow-legged gull populations (*Larus michahellis*) link the history of landfills to soil eutrophication and time-related vegetation changes on small Mediterranean islands. *Science of the Total Environment*, 2023, 878, pp.162948. <10.1016/j.scitotenv.2023.162948>. <hal-04326871>

**HAL Id: hal-04326871**

**<https://hal.science/hal-04326871v1>**

Submitted on 26 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# Science of the Total Environment

## Yellow-legged gull populations (*Larus michahellis*) link the history of landfills to soil eutrophication and time-related vegetation changes on small Mediterranean islands --Manuscript Draft--

|                               |   |
|-------------------------------|---|
| <b>Manuscript Number:</b>     | STOTEN-D-23-00547R1   |
| <b>Article Type:</b>          | Research Paper  |
| <b>Keywords:</b>              | Long-term diachronic survey; seabirds; Trace elements; Plant community; land-use history  |
| <b>Corresponding Author:</b>  | Clémentine Mutillod, Ph.D. student<br>Mediterranean Institute of Biodiversity and Marine and Continental Ecology<br>Marseille, FRANCE   |
| <b>First Author:</b>          | Clémentine Mutillod, Ph.D. student  |
| <b>Order of Authors:</b>      | Clémentine Mutillod, Ph.D. student  |
|                               | Teddy Baumberger  |
|                               | Pascale Prudent   |
|                               | Arne Saatkamp   |
|                               | Eric Vidal  |
|                               | Lidwine Le-Mire-Pecheux   |
|                               | Laurence Affre  |
| <b>Abstract:</b>              | <p>Seabird colonies have a strong influence on both the physical and chemical soil parameters and plant communities of the islands where they settle to nest. Scientists have studied the effects of the demographic explosion of seabird populations, but few have explored the long-term effects when the colonies were in decline. The aim of this study was to investigate diachronic changes over a 24 year period of soil parameters, floristic composition and plant functional types (Raunkiaer growth forms and Grime life strategies) up to the decrease of the number of nesting yellow-legged gulls (<i>Larus michahellis</i> Naumann, 1840) on Mediterranean islands. We used 78 permanent plots to survey the vegetation and the soil parameters on 9 islands and one mainland area within the Calanques National Park (south east of France), for three periods (i.e., 1997, 2008, 2021).</p> <p>Since 1997, the increase of nesting gulls has caused a nitrogen and pH increase and organic carbon and C/N ratio decrease, although the values were still higher than mainland plots without nesting gulls. This has led to changes in plant species composition e.g., higher values of N favouring the development of ruderal plant species, still present in high frequency in 2021. Furthermore, plant species highly tolerant to disturbances (i.e., R Grime strategy) in harsh environments were still favoured even after the decline of gull abundance. However, both the frequency of the chamaephytes and the vegetation cover has increased with the decline of gull colony. In 2021, measures of trace elements' concentrations and calculation of pollution load index (Cu, Pb and Zn) reveals relatively low multi-contamination levels on the mainland and the archipelagos.</p> <p>On naturally oligotrophic and semi-arid Mediterranean islands, gull colonies induce a persistent alteration in soil characteristics that still influences plant communities (composition and functional types), 11 years after the decline in bird abundance.</p> |
| <b>Response to Reviewers:</b> |   |

**Title: Yellow-legged gull populations (*Larus michahellis*) link the history of landfills to soil eutrophication and time-related vegetation changes on small Mediterranean islands**

Clémentine Mutillod<sup>1\*</sup>, Teddy Baumberger<sup>2</sup>, Pascale Prudent<sup>3</sup>, Arne Saatkamp<sup>1</sup>, Eric Vidal<sup>1,4</sup>, Lidwine Le-Mire-Pecheux<sup>5</sup>, Laurence Affre<sup>1</sup>

<sup>(1)</sup> Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE Institut Méditerranéen de Biodiversité et d'Écologie marine et continentale, Marseille, France

<sup>(2)</sup> Société EcoStat, Baudinard-sur-Verdon, France


<sup>(3)</sup> Aix Marseille Univ, CNRS, LCE Laboratoire de Chimie de l'Environnement, Marseille, France

<sup>(4)</sup> UMR ENTROPIE (IRD, Université de la Réunion, IFREMER, Université de la Nouvelle-Calédonie, CNRS), BP A5, Nouméa CEDEX 98848, New Caledonia

<sup>(5)</sup> Calanques National Park, Marseille, France

\* Contact details: [clementine.mutillod@gmail.com](mailto:clementine.mutillod@gmail.com) IMBE, IUT Avignon, 337 chemin des Meinajariés, Site Agroparc BP 61207, 84911 Avignon cedex 09, France

**Graphical abstract**



**Title: Yellow-legged gull populations (*Larus michahellis*) link the history of landfills to soil eutrophication and time-related vegetation changes on small Mediterranean islands**

Clémentine Mutillod<sup>1\*</sup>, Teddy Baumberger<sup>2</sup>, Pascale Prudent<sup>3</sup>, Arne Saatkamp<sup>1</sup>, Eric Vidal<sup>1,4</sup>, Lidwine Le-Mire-Pecheux<sup>5</sup>, Laurence Affre<sup>1</sup>

<sup>(1)</sup> Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE Institut Méditerranéen de Biodiversité et d'Écologie marine et continentale, Marseille, France

<sup>(2)</sup> Société EcoStat, Baudinard-sur-Verdon, France

<sup>(3)</sup> Aix Marseille Univ, CNRS, LCE Laboratoire de Chimie de l'Environnement, Marseille, France

<sup>(4)</sup> UMR ENTROPIE (IRD, Université de la Réunion, IFREMER, Université de la Nouvelle-Calédonie, CNRS), BP A5, Nouméa CEDEX 98848, New Caledonia

<sup>(5)</sup> Calanques National Park, Marseille, France

\* Contact details: [clementine.mutillod@gmail.com](mailto:clementine.mutillod@gmail.com) IMBE, IUT Avignon, 337 chemin des Meinajariés, Site Agroparc BP 61207, 84911 Avignon cedex 09, France

**Highlights**

- Soil parameters alterations (notably nutrients increase) induced by seagulls persist after the colony declines.
- Ruderal plant species and more chamaephytes still occurred after the gull colony decline.
- Vegetation cover increased with the decline of gull colony.

**Title: Yellow-legged gull populations (*Larus michahellis*) link the history of landfills to soil  
eutrophication and time-related vegetation changes on small Mediterranean islands**

Clémentine Mutillod<sup>1\*</sup>, Teddy Baumberger<sup>2</sup>, Pascale Prudent<sup>3</sup>, Arne Saatkamp<sup>1</sup>, Eric Vidal<sup>1,4</sup>, Lidwine  
Le-Mire-Pecheux<sup>5</sup>, Laurence Affre<sup>1</sup>

<sup>(1)</sup> Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE Institut Méditerranéen de Biodiversité  
et d'Écologie marine et continentale, Marseille, France

<sup>(2)</sup> Société EcoStat, Baudinard-sur-Verdon, France

<sup>(3)</sup> Aix Marseille Univ, CNRS, LCE Laboratoire de Chimie de l'Environnement, Marseille, France

<sup>(4)</sup> UMR ENTROPIE (IRD, Université de la Réunion, IFREMER, Université de la Nouvelle-Calédonie,  
CNRS), BP A5, Nouméa CEDEX 98848, New Caledonia

<sup>(5)</sup> Calanques National Park, Marseille, France

\* Contact details: [clementine.mutillod@gmail.com](mailto:clementine.mutillod@gmail.com) IMBE, IUT Avignon, 337 chemin des Meinajariés,  
Site Agroparc BP 61207, 84911 Avignon cedex 09, France

**Abstract**

Seabird colonies have a strong influence on both the physical and chemical soil parameters and plant  
communities of the islands where they settle to nest. Scientists have studied the effects of the  
demographic explosion of seabird populations, but few have explored the long-term effects when the  
colonies were in decline. The aim of this study was to investigate diachronic changes over a 24 year  
period of soil parameters, floristic composition and plant functional types (Raunkiaer growth forms  
and Grime life strategies) up to the decrease of the number of nesting yellow-legged gulls (*Larus  
michahellis* Naumann, 1840) on Mediterranean islands. We used 78 permanent plots to survey the  
vegetation and the soil parameters on 9 islands and one mainland area within the Calanques National  
Park (south east of France), for three periods (i.e., 1997, 2008, 2021).

Since 1997, the increase of nesting gulls has caused a nitrogen and pH increase and organic carbon and  
C/N ratio decrease, although the values were still higher than mainland plots without nesting gulls. This  
has led to changes in plant species composition e.g., higher values of N favouring the development of

ruderal plant species, still present in high frequency in 2021. Furthermore, plant species highly tolerant to disturbances (i.e., R Grime strategy) in harsh environments were still favoured even after the decline of gull abundance. However, both the frequency of the chamaephytes and the vegetation cover has increased with the decline of gull colony. In 2021, measures of trace elements' concentrations and calculation of pollution load index (Cu, Pb and Zn) reveals relatively low multi-contamination levels on the mainland and the archipelagos.

On naturally oligotrophic and semi-arid Mediterranean islands, gull colonies induce a persistent alteration in soil characteristics that still influences plant communities (composition and functional types), 11 years after the decline in bird abundance.

**Key words:** Long-term diachronic survey, seabirds, trace elements, plant community, land-use history.

## 1 Introduction

Manmade global change has had a profound impact on biodiversity (Vié, Hilton-Taylor and Stuart, 2009), especially in the Mediterranean basin which is considered one of the biodiversity hotspots (Cunningham and Beazley, 2018). An important part of Mediterranean plant diversity is linked to the occurrence of 10,000 small islands, islands being refuges for rare and threatened mainland plant species (Médail, 2017). However, the Mediterranean basin has been subjected to an increase in human activity, particularly in coastal and island habitats (UNEP/MAP-Plan Bleu, 2009). Human activities alter ecosystem properties and services (Hooper *et al.*, 2005), especially when they affect biotic factors (Chapin *et al.*, 1997).

In this insular context, populations of the yellow-legged gull (*Larus michahellis*, hereinafter referred to as gulls) have greatly increased over the past few decades (Thibault *et al.*, 1996; Bellout *et al.*, 2021). Indeed, the high availability and predictability of food resources (Bosch *et al.*, 1994), provided by open compartments of landfills during their operation on the close mainland, is known to have driven and shaped gull colony distribution and demographic explosion on the studied islands (Duhem *et al.*, 2002, 2003, 2007, 2008). Gulls have therefore been considered as a superabundant bird species in the Mediterranean basin (Vidal, Médail & Tatoni, 1998) that conflict with island biodiversity conservation

programmes (e.g. Médail, 2017; De La Peña-Lastra *et al.*, 2021; Mouga *et al.*, 2021). Gulls, like other seabirds, are chemical and physical ecosystem engineers (Ellis, 2005; Grant *et al.*, 2022). Due to the high spatial heterogeneity of their feeding, nesting and defecation behaviour, seabirds can have contrasting impacts on ecosystems. On the one hand, their presence can lead to an increase of plant biomass and height (Ellis, 2005) and seed dispersal (Nogales *et al.*, 2001; Calvino-Cancela, 2011). On the other hand, high seabird densities can prevent plant growth owing to the uprooting of plants for nest building, trampling and soil compaction (Sobey and Kenworthy, 1979). Many studies have shown that large seabird colonies can alter soil characteristics, increasing the quantity of nitrogen, phosphorus, potassium, magnesium, total salts (Sigurdsson and Magnusson, 2010; Bouyahmed and Moulai, 2018; Aerts *et al.*, 2020), heavy metals (García *et al.*, 2002; Ziółek, Bartmiński and Stach, 2017; Otero *et al.*, 2018), organic matter (García *et al.*, 2002; Ghermaoui, Hassaine and Moulai, 2016) and soil depth (Ishida, 1997; Mulder and Keall, 2001). They also usually lead to a decrease of soil pH (Mulder and Keall, 2001; Sigurdsson and Magnusson, 2010). Other soil characteristics can also be altered, but the trend can be both upward or downward such as for calcium, carbon content, water retaining capacity and soil moisture (Ellis, 2005). These different alterations may cause changes in plant communities – with high pressure and duration, exotic, ruderal or nitrophilous plant species may develop that may lead to a reduction of the abundance of native plant species (Vidal *et al.*, 1998, 2000; Baumberger *et al.*, 2012; Bou, Vilar and Carreras, 2020). The reduction of food availability following the closure of open compartments of landfills is indicated as the main driver for a long term reduction on gull population size (Duhem, 2007, 2008) and the negative density-dependent breeding investment contributed to population decline (Real *et al.*, 2017).

Few studies (Baumberger *et al.*, 2012; Otero *et al.*, 2015; De La Peña-Lastra *et al.*, 2021) have investigated the long-term patterns of change in plant communities and soil factors after the decrease of gull populations, showing that the environmental effects of the colonies may persist for a long time. For example, high concentrations of phosphorus (P) have been observed 5 (De La Peña-Lastra *et al.*, 2021) and 14 years (Otero *et al.*, 2015) after the decline of gull populations. The amount of P generated by the colony did not decrease and can be considered as an irreversible impact for the plant community (Otero *et al.*, 2015; De La Peña-Lastra *et al.*, 2021). The Mediterranean basin has a particular climate with low

precipitation. In these semi-arid ecosystems, alterations linked to trophic shift can lead to a persisting imprint of eutrophication in the soil (Saatkamp, Henry and Dutoit, 2020) and the impact on plant communities is linked to higher P, C:N and organic matter. The threats to the Mediterranean islands are exacerbated by climate change because of the limited habitat availability and climatic range limitations which may hamper the expansion of the species' range (Vogiatzakis, Mannion and Sarris, 2016). Our study focused on nine islands in the two Marseille archipelagos, Frioul and Riou (Calanques National Park, SE France), which are under the influence of breeding gulls (*Larus michahellis*) in comparison with the nearby mainland coastal area without gull colonies. Previous studies have shown that the high pressure of gull colonies leads to a ruderalisation of the floristic community (Vidal *et al.*, 1998, 2000). This pattern has persisted (Baumberger *et al.*, 2012) despite four open landfills having been closed in the Marseille area during the last decade, which has caused a decline in the nesting gull population (Calanques National Park monitoring). We have extended monitoring of the vegetation changes up to 24 years and added information with additional analysis of nutrients and trace elements in the soil according to the fluctuations in the nesting gull density. Because of the strong relationships within the gull/soil/plant system, we expected to demonstrate a resilience in plant communities despite the persistent effect of gull colonies on soil chemistry.

## 2 Material and Methods

### 2.1 Study area

The study was conducted on two archipelagos and on the mainland area near Marseille (SE France) (Fig.1). The sampling took place on two archipelagos of Marseille: the Riou archipelago (43°11'N, 5°21'E; five islands and two islets), and the Frioul archipelago (43°16'N, 103 5°18'E; four islands) (Fig. 1). The Riou archipelago islands studied are from 50 to 3525m distant from the mainland (Table 1). Concerning the Frioul archipelago, the two islands studied are 1800 and 2800m away from the mainland. The mainland area was considered as a habitat not subjected to gull pressure (no gulls nesting on the studied mainland area). These islands and the mainland area are included in natural protected areas



managed by the Calanques National Park (CNP). Situated within the thermo-Mediterranean bioclimatic belt (Quézel & Médail, 2003), these islands and the mainland area have semi-arid climatic conditions, with lower precipitations on the islands (e.g., Planier, an island located 9km from Marseille: 342mm) compared to the mainland (Marseille 524mm), however little reliable data exists (Météo France). The studied sites are composed of compact limestone on the surface of which soils are limited to thin proto-rendzina patches, and are calcareous, skeletal and oligotrophic (Vidal, 1998). The plants on the islands and the mainland coastal area are subjected to stresses both hydric (low precipitation and frequent strong wind episodes) and haline (with salt input from the sea and the toxicity of the salt within the tissues). The islands are mainly covered by low vegetation from halophyte plants to matorral shrub species (Baumberger *et al.*, 2012). Vegetation on the mainland shows a similar composition, with however higher species richness compared to islands of similar size (Martín-Queller *et al.*, 2017).

Yellow-legged gull status is considered as Least Concern in the IUCN European Red List, it is a protected specie in France and a determining specie of natural areas of ecological, faunistic and floristic interest (INPN MNHN, 2023). Gull colonies are nesting in the Mediterranean basin and the Atlantic coast from Morocco to the southern Brittany (Duhem, 2004). Gull colonies can be found on all the islands close to Marseille. However, nesting gull abundance and densities are not homogenous between the archipelagos, and depend on local habitats. The gull population increased in the Riou and Frioul archipelagoes between 1997 and 2005, with a growth rate ranging from 6% for the Riou archipelago to 177% for the Frioul archipelago (Duhem *et al.*, 2008). The fluctuations of gull population size in the two archipelagos are presented in figure 2, between 1923 and 2021. Data were compiled by the CNP. Gulls' census on the islands of Marseille is part of a CNP program taking place every five years as well as the national census every ten years. Gull estimation was carried out by counting the number of nests (empty and with eggs) exhaustively on each island until 2021.

In 2021, exhaustive counting was done on only Grand Congloué and Tiboulén de Maire. For the rest of the islands, estimation was done by distance sampling, an effective method for many gull species (Barbraud *et al.*, 2014).

The main available anthropogenic food resources, four landfills (with open compartments during their operation) located within a radius of 60 kms, were closed in 2008, 2010 and 2013 and led to a decline in gull population size (Calanques National Park monitoring). The decrease of easily accessible food resources led to the decline of the nesting population by 63% in the Riou archipelago and 37% in the Frioul archipelago between 2005 and 2015 (Fig. 2). However, an increase was recorded in 2021 with the difference being about 1100 nesting couples.

## 2.2 Sampling design

The vegetation was surveyed on the 56 island plots and on the 22 mainland plots in 1997, 2008 and in 2021 (Tab. 1). The number of plots was proportional to the size of the island in order to take into account the heterogeneity of the zones studied and the different vegetation types. A plot consisted of two concentric circles, a first one of 500m<sup>2</sup> where environmental variables and gull density were measured, and, a second of 100m<sup>2</sup> where floristic inventories and soil sampling took place (Vidal, 1998).

## 2.3 Mesological parameters

Environmental variables were measured in the 500 m<sup>2</sup> circles for each plot to better qualify and quantify the environmental effects (see table 2 for further details). Measures and estimations were made during the floristic inventories from April to May in 1997, 2008 and 2021. The environmental variables measured were the elevation above sea level, the distance from the coastline, the slope, the exposure, and the cover rate of each substrate type: rock, block, stone, bare ground, the vegetation cover of the different layers.

Gull nest density (number of nests) was estimated and we also considered the cover of the resting places per plot being an indication of gull pressure.

## 2.4 Floristic inventories

Three plant species inventories (1997, 2008 and 2021) were carried out within the same 78 100m<sup>2</sup> concentric plots from April to May using Tison *et al.* flora (2014). The plant cover in the different

layers was estimated (2008, 2021) for each species present on each plot using the 6 cover abundance categories (0: cover = 0%, 1: 0% < cover < 10%, 2: 10% ≤ cover < 25%, 3: 25% ≤ cover < 50%, 4: 50% ≤ cover < 75%, 5: cover ≥ 75 %). The following easily accessible plant functional types, i.e., Raunkiaer growth form (Raunkiaer, 1934) and Grime life strategy (Grime, 1977), for each plant species, were extracted from the BASECO data base (Gachet, Véla and Taton, 2005).

## 2.5 Soil, droppings sampling and chemical analysis

The soil was sampled in 1997 (56 plots only on the islands) and in 2021 (56 plots on the islands and 22 on the mainland). In 1997, organic carbon, total nitrogen, and pH were measured within the plots. In 2021, soil parameters included organic carbon, total nitrogen, pH, conductivity, total phosphorus, potassium, sodium, copper, lead, zinc and stable isotopes ( $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$ ).

The soil samplings were performed once between March and April, when most of the gulls were nesting and after the winter rainfall. Within each plot, five soil subsamples of 200mL each were collected (one in the center and four on each side, approximately 50cm from the 100m<sup>2</sup> radius boundary) and pooled in a composite soil sample, in order to make a composite sample representative of the plot. Soil subsamples were collected from the top 15cm layer after litter removal and sieved at 2mm directly on site. The soil samples were dried before analysis (at 40°C up to constant weight). Dry gull droppings were sampled on nine different plots in the Frioul and Riou archipelagos in 2021. They were collected on rocks to avoid contact with soil.

To assess the soil organic matter percentage (SOM), the method of Mathieu & Pieltain (2003) was used: the samples were first weighed and then heated several times at high temperature. The samples were heated first at 220°C for 16 hours (bound water elimination), then heated again at least three times at 450°C for four hours (loss of organic matter by combustion, and SOM was estimated by mass loss between 220 and 450°C). The organic carbon was estimated using the values of the organic matter multiplied by 0.50, based on the assumption that the soil organic matter contains 50% carbon (Pribyl, 2010).

To determine the pH and the conductivity, the samples were extracted with osmosed water (ratio 1:5 V/V), then agitated for at least one hour, decanted for one more hour and finally the conductimeter (WTW Multi 3420) and the pH-meter (Orion 2 Star Thermo Scientific SM30B) could be used (according to norm ISO 11265 (1995) and ISO 10390 (2005) respectively; Baize and Girard, 1992).

The total nitrogen, carbon,  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  content in ground soil samples (ground to less than 0.2mm with RETSH 22, agate mortar) and droppings samples (n = 9, not composites) were performed on an elemental analyzer (Flash HT Plus, Thermo Fisher Scientific) coupled with a mass spectrometer (Delta V Advantage, Thermo Fisher Scientific). Elemental analysis of carbon and nitrogen were performed in a combustion reactor (1,000°C) under a carrier gas (Helium: 100mL/min) with samples weighing approximately 800µg sealed in tin capsules, introduced via an automatic injector. The gases were separated on a chromatographic column in an oven at 45°C before being quantified via the thermal conductivity detector (TCD) of the elemental analyzer. The gases were injected into the Delta V spectrometer for isotopic analysis. The calibration of the catharometric detector and the ratio-isotopic mass spectrometer was performed with a secondary standard, EDTA (tetra-acetic ethylene diamine acid). The analyses (samples or standards) of the ratio  $^{13}\text{C}/^{12}\text{C}$  were calculated in delta value which represents the deviation from an international standard and is calculated as follows :

$$\delta^{13}\text{C} = \left( \frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right) \times 1000 (\text{‰})$$

The international standard for  $\delta^{13}\text{C}$  is Vienna PeeDee Belemnite (VPDB),  $R_{\text{standard}} = 0.011237$ . The  $\delta^{15}\text{N}$  is expressed in the same way in relation to atmospheric nitrogen with a conventional value of 0‰. The results of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  analyses are produced with an uncertainty of +/- 0.5 ‰. The  $\delta^{15}\text{N}$  and the  $\delta^{13}\text{C}$  were analysed because they can give an indication of the trophic transfer, being robust and effective in tracing the path of the organic matter and the trophic relationships within the ecosystems (Post, 2002).

The analysis of the other elements (Cu, K, Na, P, Pb and Zn) in the soil and the droppings samples was carried out after acid mineralization. Samples were digested in a microwave mineralizer (Milestone Start D) using aqua regia (1/3  $\text{HNO}_3$  (Fisher chemical, Trace metal grade 67-69%) and 2/3 HCl (Fisher chemical, Trace metal grade 35-38%)). After mineralization, the samples were filtered through a 0.45µm

cellulose ester membrane filter and stored at 4°C until analysis (mineralization protocol for pseudo-total concentration determination in soil, adapted from norm ISO 11466, 1995). Then element concentrations in samples were carried out by ICP-OES (Inductive Coupled Plasma, Optical Emission Spectrometry, Jobin Yvon Horiba Spectra 2000). Quality controls and accuracy were checked using standard soil reference materials (CRM 049-050 from RTC-USA), with accuracies within 100±10%).

Concerning Cu, Pb and Zn, considered as potentially toxic elements (PTE), contamination factors in soil (CF) were calculated as followed:  $CF_{PTE} = [PTE]_{\text{pseudo-total in soil}} / [PTE]_{\text{local background value}}$

where  $[PTE]_{\text{local background value}}$  are 43, 66 and 7.5 mg.kg<sup>-1</sup> for Pb, Zn and Cu respectively (Affholder *et al.*, 2014). The potential multi-contamination level of soils was estimated by calculating an integrative index i.e. pollution load index (PLI) according to the formula (Affholder *et al.*, 2014; Rashed, 2010):

$$PLI = \sqrt[3]{CF_{Pb} \times CF_{Zn} \times CF_{Cu}}$$

## 2.6 Statistical analysis

Since plant names have been revised between different study periods, we followed the taxonomy of Gargominy and collaborators (Gargominy *et al.*, 2020). In order to understand patterns of change of soil parameters, we used boxplots in pairs for contrasting years, and separated to compare Riou, Frioul and the mainland. We used Student tests for variables with normal distribution and homoscedasticity or pairwise Wilcoxon in the other cases. To measure the potential effects of the gulls on the soil parameters, linear models were used (*stats* package). We pooled all the plots from the two archipelagos and considered the number of counted nests or the cover of resting places as explaining variables for the soil parameters measured in 2021. We excluded mainland plots from this analysis to avoid confounding effects. When conditions were not met for model residuals (normality, homoscedasticity and independency), outliers were removed and/or the variables were log transformed. To compare Na and conductivity mean values properly without the confounding effect of the sea shore distance, we selected a subsample of plots from the mainland area. We selected only plots within 188m distance from the sea shore, which is the maximum distance from the sea shore of island plots.

With the aim of describing the flora trajectory and to know whether there was a flora niche alteration due to climate change or only to gulls pressure, different analyses were carried out. Non-metrical multidimensional scaling (NMDS) was performed to view the alterations in the flora on the islands between the three dates. Using the coordinates of groups in the NMDS, we used Wilcoxon paired tests to verify whether the changes are significantly different. To quantify the changes, mean comparisons were made between years with paired Wilcoxon tests, using the number of species, Grime life strategies and Raunkiaer growth forms.

In order to compare the vegetation cover between 2008 and 2021, we performed an Ascending Hierarchical Classification (AHC) (*stats* package) with the data of 2021 (mesological and soil parameters) in order to have different groups of plots and to carry out paired Wilcoxon tests between them.

The observations made on the islands could be explained by more global alterations such as climate change (Heilmeier, 2019; Ahrens *et al.*, 2020). To comprehend the flora ecological requirement changes with the environmental alterations due to gulls pressure or/and climate change between 1997, 2008 and 2021, we compared Ellenberg-Pignatti factors (Organic Matter, Nutrient, Soil Acidity, Humidity, Light, Salinity, Temperature) with paired pairwise Wilcoxon tests and Generalised Additive Models (GAM) (*mgcv* package). The variables included were the different Pignatti factors: organic matter, nutrient, soil acidity, humidity, light, salinity, temperature; and the explained variable was the number of nests. As variables responses are expected to be non-linear, we used Generalized Additive Models with gaussian family to evaluate the statistical link between Pignatti factors and the number of nests. The indicators were adapted for the Mediterranean species by Saatkamp *et al.* (2022), so that the bioindications given by plants fit better than the initial values.

All statistical analyses were performed with R software (R version 4.2.2: R Core Team, 2022). Null hypotheses were rejected when the p-value was below 0.05.

## 3 Results

### 3.1 Nesting gulls and dynamic of the soil parameters

The number of nests has decreased since 1997 (Fig. 3. A). The cover of resting places decreased less on the Frioul archipelago compared to the Riou archipelago between 2008 and 2021 (Fig. 3. B).

Soil acidity decreased during the study period (see Supplementary Materials Table S1 for a summary of all means values  $\pm$  SD), with mean pH in 2021 significantly higher than in 1997 (Fig. 4. A.,  $p < 0.001$ ). Soil pH was lower on Riou islands (mean  $\pm$  standard deviation,  $7.89 \pm 0.47$ ,  $n = 30$ ) compared to the mainland ( $8.18 \pm 0.28$ ,  $n = 22$ ) in 2021 ( $p < 0.05$ ). There was no significant difference between soil pH values on Riou and Frioul islands whatever the year considered.

Organic carbon in soils of the archipelagos decreased during the study period (Fig. 4. B,  $p < 0.01$ ), but was not significantly different between islands in 1997 ( $p=0.332$ ) and 2021 ( $p=0.091$ ), respectively, and between islands and the mainland in 2021 ( $64.07 \pm 27.01$ ,  $53.06 \pm 20.70$  and  $56.06 \pm 31.12$  mg/g for Riou, Frioul ( $n = 26$ ) and the mainland respectively; mainland vs Frioul  $p=0.70$ , mainland vs Riou  $p=0.34$ , Frioul vs Riou  $p=0.09$ ).

Soil total nitrogen content of Frioul and Riou samples was significantly higher in 2021 than in 1997 (Fig. 4. C,  $p < 0.01$ ). Soil total nitrogen on Riou islands ( $9.14 \pm 3.63$  mg/g) was significantly higher than on the mainland ( $5.86 \pm 2.45$  mg/g) and also than on Frioul islands ( $7.18 \pm 2.84$  mg/g) in 2021 ( $p < 0.001$ ). Total nitrogen values for Riou and Frioul in 1997 were not significantly different from the mainland in 2021 ( $p=0.161$  and  $p=0.088$ , respectively). Total nitrogen content of droppings ( $44.50 \pm 8.34$  mg/g) was significantly higher than in any soils ( $p < 0.001$ ).

The C/N ratio was significantly lower for Riou ( $13.57 \pm 6.86$ ) and Frioul ( $17.95 \pm 6.88$ ) in 2021 than the values for Riou ( $21.97 \pm 13.22$ ) and Frioul ( $18.24 \pm 2.88$ ) in 1997, and than the mean value for the mainland ( $19.03 \pm 6.15$ ) in 2021 (Fig. 4. D,  $p < 0.001$ ).

## 3.2 Additional chemical characterisation of soils and impacts of gull presence on edaphic parameters

No difference was shown for the soil conductivity between the archipelagos ( $p=0.96$ ), nor for islands *versus* mainland (mainland *vs* Frioul  $p=0.99$ , mainland *vs* Riou  $p=0.23$ ), mean values from 0.64 to 1.07 mS/cm, not presented here, Table 3).

Regarding nutrient concentrations in soils, the mean amount of P in soils of the mainland (Table 3) was significantly lower than in the soils of the archipelagos (Fig. 5. A, ( $p < 0.001$ ), and total P concentrations were significantly higher in the soils of Riou compared to Frioul (Fig. 5. A,  $p < 0.01$ ). The concentration of P appeared to be higher in droppings than in the soils. The K values were significantly higher in the soils of the Riou archipelago than on the mainland (Fig. 5. B,  $p < 0.05$ ), but there was no significant difference between the soils of the archipelagos ( $p=0.58$ ). The droppings appeared to be less concentrated in K than the soils. No significant difference was observed for Na concentration whatever the soils on the islands or on the mainland (Fig. 5. C, mainland *vs* Frioul  $p=0.33$ , mainland *vs* Riou  $p=0.08$ , Frioul *vs* Riou  $p=0.87$ ), and Na contents were higher in droppings.

Regarding potentially toxic elements (PTE), Pb soil concentrations were higher on the mainland than on the archipelagos (Fig. 5. D,  $p < 0.01$  between the mainland and Riou and  $p < 0.001$  between the mainland and Frioul). Pb concentrations were lower in droppings (mean of 4.34 mg/kg). With regards to Cu and Zn soil concentrations, no significant difference was observed whatever the soils on the islands or on the mainland (Cu: mainland *vs* Frioul  $p=0.99$ ; mainland *vs* Riou  $p=0.44$ ; Frioul *vs* Riou  $p=0.77$ ; and Zn: mainland *vs* Frioul  $p=0.15$ ; mainland *vs* Riou  $p=0.56$ ; Frioul *vs* Riou  $p=0.61$ ) (Fig. 5. E, F). Cu and Zn contents were higher in droppings than in soils. In order to estimate the potential soil contamination by PTE, contamination factors (CF) were calculated for Cu, Pb and Zn, in relation to the local contamination background proposed for the Calanques hills (Affholder et al, 2014). The highest CF values (Table 4) were measured for Pb in mainland soils ( $8.23 \pm 6.90$ ) and for Cu in soils of the Frioul archipelago ( $6.87 \pm 12.94$ ), and these values reflected a high contamination ( $CF > 5$ ). CF values for Pb in soils of the archipelagos reflected a low contamination rate (low contamination estimated when  $CF < 3$ ,  $3.22 \pm 1.83$  at Riou and  $2.24 \pm 2.28$  at Frioul), and those for Cu in soils of Riou and the mainland



reflected a medium contamination rate (medium when  $3 < CF < 5$ ,  $4.26 \pm 1.48$  at Riou and  $3.68 \pm 1.57$  on the mainland). Concerning Zn, CF values reflected a low contamination rate whatever the sites (values between 2.39 and 3.22). The pollution load index (PLI) for Cu, Pb and Zn estimates the potential multi-contamination level of soils. This index, revealed a relatively low multi-contamination level whatever the sites studied, with PLI values around 2 ( $2.10 \pm 0.30$  at Riou,  $2.10 \pm 0.54$  at Frioul, and  $2.33 \pm 0.57$  on the mainland).

It was noticeable that the mean amounts of Cu, Zn, Na and P showed a tendency to be higher in the droppings than in soils (Fig. 5. E, D, F). The contrary was observed for K and Pb (Fig. 5. B, E).

The analysis of the stable isotopes showed an enriched signature of the  $\delta^{15}N$  for soils of the two archipelagos compared to the reference plots of the mainland (Fig. 6.), with higher values for Riou ( $> 5\%$ ). Concerning the  $\delta^{13}C$ , values did not show a clear pattern.

Analyses of potential impacts of a colony of gulls on physical and chemical parameters of the islands' soil in 2021 showed that the concentrations of K were significantly and positively correlated with the number of nests ( $p < 0.01$ , Adjusted  $R^2 = 0.1123$ ), P concentrations ( $p < 0.05$ , Adjusted  $R^2 = 0.08533$ ) and Na concentrations ( $p < 0.05$ , Adjusted  $R^2 = 0.08361$ ). These results also showed that the pH values were significantly and negatively correlated with the number of nests ( $p < 0.05$ , Adjusted  $R^2 = 0.09594$ ). Moreover, the resting places were positively correlated with conductivity ( $p < 0.05$ , Adjusted  $R^2 = 0.1728$ ).

Concerning the other parameters, no significant correlations were observed neither with the number of nests (Cu:  $p=0.51$ , Pb:  $p=0.99$ , Zn:  $p=0.82$ , Conductivity:  $p=0.08$ , N:  $p=0.27$ , Organic Matter:  $p=0.65$ ) nor the resting places (Cu:  $p=0.64$ , P:  $p=0.22$ , Pb:  $p=0.39$ , pH:  $p=0.41$ , K:  $p=0.27$ , Zn:  $p=0.44$ , Na:  $p=0.09$ , N:  $p=0.95$ , Organic Matter:  $p=0.73$ ).

### 3.3 Changes in plant communities

NMDS drew a pattern of the flora community based on the three successive surveys conducted on the islands (Fig. 7.). The first axis separated the saline environment with salt-adapted plant species (*Camphorosma monspeliaca*, *Crithmum maritimum*, *Limonium echioides*, *Pancratium maritimum*) on the left, to the less saline environment represented by other plant species (*Avena barbata*, *Lonicera implexa*, *Rosmarinus officinalis*, *Urospermum picroides*) on the right. The second axis showed a gradient from the ruderal plant community (top: *Hordeum murinum*, *Malva arborea*, *Senecio leucanthemifolius*, *Sonchus asper subsp. glaucescens*) to the more stress-tolerant plant community (bottom: *Asteriscus aquaticus*, *Brachypodium retusum*, *Sedum littoreum*, *Teucrium flavum*).

Ruderal plants increased in the vegetation over time as highlighted by increased scores of recent compared to ancient vegetation data on the second axis. The flora in 1997 was less ruderal than in 2008 and in 2021 ( $p < 0.01$ ). The flora in 2008 and 2021 was also different ( $p < 0.01$ ) with more ruderal plant species in 2021.

No pattern of change was detected on the salinity axis (axis 1) when using the coordinates of the NMDS, (1997 vs 2008  $p=0.11$ , 1997 vs 2021  $p=0.44$ , 2008 vs 2021  $p=0.19$ ).

The Pignatti nutrient ecological indicator indicated by the flora community on the islands was significantly lower in 2021 and 2008 compared to 1997 ( $p < 0.05$ ). The humidity factor decreased in 2021 compared to 1997 on the islands ( $p < 0.05$ ). The light factor decreased in 2021 compared to 1997 and 2008 on the islands ( $p < 0.05$ ). The soil acidity indicated by the flora community was not significantly different between years for islands plots (1997 vs 2008  $p=0.48$ , 1997 vs 2021  $p=0.16$ , 2008 vs 2021  $p=0.057$ ). Generalized Additive Models (GAM) showed no relation with the number of nests counted in 2021 (nutrient  $p=0.72$ , humidity  $p=0.64$ , light  $p=0.06$ , soil acidity  $p=0.83$ ).

The modification of the temperature indicated by the flora community on the islands showed a diminution in 2021 compared to 1997 and 2008 ( $p < 0.01$ ). In addition, there was a positive correlation with the number of nests ( $p < 0.05$ ). However, a diminution of the temperature in 2008 and 2021 ( $p < 0.05$ ) was registered with the mainland plots, where there were no gull nests.

The fluctuation of the salinity factor indicated by the flora community was significantly lower in 2008 than in 1997 on the islands ( $p < 0.05$ ). The GAM showed a positive correlation with the number of nests ( $p < 0.01$ ). Concerning the mainland, no other difference has been observed ( $p > 0.15$ ).

Results of mean comparisons showed that the total number of species was not different between years on the islands (1997 vs 2008:  $p=0.669$ , 1997 vs 2021:  $p=0.2446$ , 2008 vs 2021:  $p=0.5376$ ). However, for the mainland, differences were observed between 1997 and 2008 ( $p < 0.05$ ) and between 2008 and 2021 ( $p < 0.001$ ) with the lowest value in 2008.

When analysing Grime plant strategies, ruderals (the most strongly represented Ruderals (R): *Sonchus tenerrimus*, Ruderal-Competitives (RC): *Parietaria judaica*, Ruderal-Stress tolerants (RS): *Lobularia maritima*, Stress tolerant- Ruderals (SR): *Euphorbia linifolia*) increased between 1997 and 2008 ( $p < 0.001$ ) and this pattern remained in 2021 ( $p < 0.001$ ). Moreover, intermediate plant strategies (Competitive-Stress tolerant-Ruderals (CSR), such as *Cistus albidus*, *Euphorbia characias*, *Lotus cytisoides*) decreased significantly on the islands from 1997 to 2008 ( $p < 0.05$ ) and slightly increased again to 2021, with no significant difference between 1997/2021 ( $p=0.06277$ ) and 2008/2021 ( $p=0.8016$ ).

Island vegetation showed a significant increase of geophytes, hemicryptophytes and therophytes ( $p < 0.001$ ) in 2008, remaining constant to 2021. Chamaephytes on the islands decreased from 1997 to 2008 ( $p < 0.001$ ) and afterwards remained stable ( $p > 0.005$ ). Chamaephytes on the mainland continuously decreased from 1997 through 2008 ( $p < 0.01$ ) to 2021 ( $p < 0.001$ ), and annual plants increased continuously. No significant pattern of change was detected for geophytes (1997 vs 2008  $p=0.38$ , 1997 vs 2021  $p=0.59$ , 2008 vs 2021  $p=0.85$ ) and hemicryptophytes (1997 vs 2008  $p=0.77$ , 1997 vs 2021  $p=0.037$ , 2008 vs 2021  $p=0.99$ ) on the mainland. No pattern of change was detected for the nanophanerophytes and the phanerophytes at any site.

The vegetation cover was significantly higher on two groups of plots in 2021 than in 2008 ( $p < 0.01$  for the first group and  $p < 0.05$  for the second group) (for more details and photographs, see Supplementary Materials Fig. S1 and S2).

## 4 Discussion

### *Dynamic of the soil parameters and characterisation*

Comparing the fluctuation of soil parameters between 1997 and 2021, we recorded an increase of pH values. This means that the soil was less acidic in 2021 than 24 years ago. Considering the values of pH, it was still approaching neutrality so little impact on the flora was expected. Conversely, the amount of N had increased because of droppings accumulation and limited leaching. When decomposing, the organic matter increases total N concentration in soil. Moreover, organic matter decomposition explains its decrease and the decrease of C/N values. These quantities of N could help plant development as seen on a recently emerged volcanic island (Sigurdsson and Magnusson, 2010; Aerts *et al.*, 2020). However, in the case of the Marseille archipelagos, where a flora community was already established, this could trigger a lasting alteration of plant composition. The C/N ratio plays a role in the balance of the bacterial and fungal communities of soil, with an increase of the bacterial community when C/N decreases (Grosso, Bååth and De Nicola, 2016). Moreover, soil organic matter is considered as a good indicator of soil quality and high levels of organic matter are associated with nutrient availability and increased water retention capacity (Nyiraneza *et al.*, 2017).

The values of P measured in 2021 showed that the soils in the archipelagos are still enriched with, and this observation was corroborated by other studies which showed that P enrichment by seabirds is an irreversible process (Myrcha & Tatur, 1991; Holdaway *et al.*, 1999; Sun *et al.*, 2000; Otero *et al.*, 2015; De la Peña-Lastra *et al.*, 2021). Moreover, the values in both reference and colonies plots were comparable to the study of De la Peña-Lastra *et al.* (2020). The lower coverage of plants when the gull colony was at its demographic peak, implying a decline in nutrient retention (Hanrahan *et al.*, 2021), might also explain the high values of P and its accumulation in the soils. The soils of the archipelago of Riou showed the highest levels of P. This observation could be explained by the fact that the colony was first established on Riou before expanding to Frioul (Baumberger *et al.*, 2012). Hence, its impact on the soil and the flora lasted longer and today the colony on Riou is still larger.

The values of conductivity and Na concentrations measured in soils in 2021 showed no differences between the mainland and the islands, which means gull presence today has no impact with regards to

these characteristics. Usually, conductivity and Na concentrations have been shown to increase with the presence of seabird colonies (Ellis, 2005; Zwolicki *et al.*, 2015). However, another study also did not register salinization (De la Peña-Lastra *et al.*, 2021), probably due in their case to high precipitations and leaching (Otero *et al.*, 2015).

In 2021, we also added to the analysis trace elements (or PTE) in soils, and the determination of pollution load index (PLI), integrating Cu, Pb and Zn, reveals relatively low multi-contamination levels on the mainland and the archipelagos, with values around 2.2. These PLI values correspond to those calculated for soils in the Calanques hills (mean value of 2.3) some distance away from the urbanized area of Marseille and from current and former industrial activities (Affholder *et al.*, 2014). We observed higher values for the amount of Pb in soils of the mainland than in those of the archipelagos. We can argue that the origin is mostly anthropic, notably with the use of leaded gasoline, coal burning and waste incineration (Erel, Veron and Halicz, 1997; Komárek *et al.*, 2008), but mainly with the former lead smelter sites located on the south-east coast of Marseille (Daumalin *et al.*, 2016; Heckenroth *et al.*, 2016, 2022; Affholder *et al.*, 2014; ). The values of Cu and Pb are comparable to the results of a previous study on the impact of waterbird colonies in the Mediterranean semiarid floodplain (Laguna *et al.*, 2021), but we cannot argue for a direct effect of the gulls on these PLI. However, the literature shows the importance of pollutants transport, such as Cu, Pb, Zn, by seabirds (Shoji *et al.*, 2019; Castro *et al.*, 2021; Grant *et al.*, 2022).

Moreover, the values of K, Na, P and conductivity on the islands showed greater amplitudes, which led to a greater spatial variation for the chemical composition of soils, plant richness and assemblages (Wait, Aubrey and Anderson, 2005).

The results of stable isotopes  $\delta^{15}\text{N}$  reflected higher values for the two archipelagos and the droppings, with values around 10‰ for the droppings and even higher than 15‰ for the Riou archipelago. The values of previous studies were on the same scale (Ellis, 2005). High  $\delta^{15}\text{N}$  derived from the droppings, and hence explained the values of the archipelagos, the seabirds being positioned high in the trophic chain (García *et al.*, 2002; Wait, Aubrey and Anderson, 2005; Laguna *et al.*, 2021). The values of the mainland higher than 0‰, derived from the matter of the vegetation (Post, 2002). There was no

significant difference concerning the  $\delta^{13}\text{C}$ , which resulted mostly from the autochthonous organic matter even on the island (Torres *et al.*, 2012).

An increase of pH was observed but there was still a negative link with gull density, as in other studies (Vidal, 1998; Ellis, 2005). Moreover, cover of the resting places had a positive link with conductivity. Except for pH, K, P, Na, we did not find the same links as in 1997 with positive correlations between nest density and the organic matter nor the nitrogen content (Vidal, 1998). Moreover, the results also indicated that the amounts of Na, P and Zn were higher in the droppings than in soils. According to these observations, we could expect existing links between gull densities and today's values in the islands' soils. However, with the absence of links as shown before, we believe that the levels of nutrients and other elements found today could still be linked with the density of nests of the last decades. Actual observations have been shown to retrace land-use history and be linked to it even millennia after in dry ecosystem (Mizutani *et al.*, 1991; Saatkamp, Henry and Dutoit, 2020). As previously seen, changes in soil P driven by seabird colonies have persistent effects (Myrcha & Tatur, 1991; Holdaway *et al.*, 1999; Sun *et al.*, 2000; Otero *et al.*, 2015; De la Peña-Lastra *et al.*, 2021). The remanence in soil is also the case for other nutrients and contaminants such as N or Selenium (Hawke & Wu, 2012; Jaunatre *et al.*, 2014; De la Peña-Lastra *et al.*, 2021). The retention and so the remanence time for nutrients or contaminants can vary with several factors, such as successional state and ecosystem type, demand or retention capacity, land-use history, type of soils, topography, climate and the rate, timing and type of deposition as was shown for N inputs (Matson *et al.*, 2002).

## *Vegetation dynamics*

### *Lag dynamics between soil and vegetation*

The previous hypothesis, according to which the current observations would be explained by previous gull pressure, is reinforced by the study of the flora community because we highlighted that the plant community on the island is still different from 1997, being as ruderal as in 2008. This observation could find an echo in a previous study (Purschke *et al.*, 2012) which found that current factors alone could not explain today's flora pattern. This finding is similar to those of a recent study (Saatkamp, Henry and Dutoit, 2020) where it was demonstrated that the use of land two thousand years

ago by the Romans is still modelling the vegetation and the soil today. Furthermore, we can also surmise that the plant community can be firstly influenced by the mesological parameters such as the distance from the mainland, the surface area, the topography, the perimeter and the maximum altitude of the island (MacArthur and Wilson, 1963; Whittaker *et al.*, 2017), the direct effects of the seabird colonies (Bancroft, Roberts and Garkaklis, 2005; Wait, Aubrey and Anderson, 2005), and secondly by the induced soil alterations (Duda *et al.*, 2020). It appears that at the level of the species and the functional traits, there have been no significant changes since 2008. Indeed, there are still more ruderal species strategies (with the most represented R: *Sonchus tenerrimus*, RC: *Parietaria judaica*, RS: *Lobularia maritima*, SR: *Euphorbia linifolia*) and more geophytes, hemicryptophytes, and therophytes in 2021 than in 1997 on the islands. The differences observed on the islands means that the plant species which are highly tolerant to disturbances in harsh environment are still favoured even with the decline of the pressure induced by the presence of the gull (Baumberger *et al.*, 2012).

The decrease of the chamaephytes recorded in 2008 compared to 1997 is no longer occurring in 2021 on the island. This is noticeable because certain of the heritage/endangered plant species of the archipelagos are chamaephytes and specialists of this environment (e.g., *Astragalus tragacantha*, *Limonium pseudominutum*, *Frankenia hirsuta*, *Euphorbia pithyusa*, *Plantago subulata*). However, further analyses and studies are needed to conclude more precisely on the dynamics of each of the species mentioned above. Nevertheless, this observation still permit to underline the vulnerability of these species (Bou, Vilar and Carreras, 2020) being not ruderal and jeopardized by the eutrophication of the soil. These observations also raise questions regarding their protection and the support of the viable populations until the habitat returns, or not, to a previous state (Knoerr, 1960 ; Molinier 1936, 1939 in Vidal, Médail, Tatoni *et al.*, 1998). On this time scale, we may inquire about the expression of the seed banks, which are considered essential for plant communities because of their contribution to ecological processes, in terms of resilience, the persistence of the community and the restoration of the ecosystems (del Cacho and Lloret, 2012; Kiss *et al.*, 2018).

An increase in the vegetation cover was registered between 2008 and 2021 especially on island plots. As supported by the previous results on functional types, it was not a sign of the resilience of the functional flora community such as there was in 1960 (Knoerr 1960 in Vidal, Médail, Tatoni *et al.*,

1998) – with more stress-tolerant species. However, it was a sign of the resilience of the flora developed since then, being more ruderal. Concerning the mainland plots where vegetation cover increased, we can assume that in 24 years and without major disturbances (fire, rockslide), the plants followed a natural dynamic and so the aboveground vegetation cover increased.

#### *Interactive effect of gulls and climate change*

In the context of the alteration of the biota (e.g., the yellow-legged gull colony and so the flora) linked to the land use (e.g., landfills), it was interesting to explore whether the alteration of the disturbance is modulated by climate change (Perring *et al.*, 2016), or whether they may act in synergy. This could also offer us a basis for deciding on future management actions (Perring *et al.*, 2015). Using the bioindicators of Pignatti (Schwabe, Kratochwil and Pignatti, 2007; Saatkamp *et al.*, 2022) helped us to characterise the requirements of plant species between the three dates and underline the results obtained from the soil analyses. In fact, we could highlight that since 1997 there has been a decline of the nutrient factor on the islands, not correlated with the gull density in 2021. This observation is corroborated by the chemical results. Schaffers and Sýkora (2000) found that the nutrient factor is not directly linked to the nitrogen but is correlated with the productivity. Previous studies have shown that gull colonies can lead to an increase in soil moisture, which can be explained by greater water retention due to higher amounts of organic matter (Ishida, 1997; García *et al.*, 2002). The decrease of the productivity, soil moisture and light all imply a diminution of the vegetation cover before 2021. It will be interesting to see in the future whether the increase of the cover observed today will trigger this factor.

We could also observe differences on the mainland concerning the temperature, with plants indicating a decrease in 2021 compared to 2008. The fact that there were no significant differences between 2021 and 1997 on the mainland, regarding the temperature, could perhaps point to a fluctuation being due to the meteorology, with dryer or wetter years. However, we could also note that there were fewer CSR strategies in 2008 than in 1997 on the islands and fewer in 2021 compared to 1997 on the mainland. This could be explained by a natural fluctuation, with the renewal rate of these plants, which here in the Calanques National Park are *Cistus albidus*, *Cistus monspeliensis*, *Cistus salvifolius*, *Euphorbia characias*, *Halimione portulacoides*, and *Lotus cytisoides* (mostly nanophanerophytes). On the



mainland, we also observed the increase of the therophytes. These observations could also be linked with the increase of more generalist and fewer thermophilous species, the closure of the vegetation or the pattern with less rainfall (Kimball *et al.*, 2010).

Climate change in the region and its impacts on flora have been synthesized in details in a recently accepted work (Saatkamp *et al.*, 2023 *in press*), showing that there is an overall warming and drying in lowland areas, with a stronger effect inland compared to coastal areas. It seems however that climate change effects are overridden by trophic changes and long-term succession in the present data-set, since there is no clear effect of thermophilization and xerophytisation. Moreover, higher temperatures during the cold season could increase the presence of plants from higher altitudes or latitudes but not thermophilous species because on the islands this type of plant is already present. Further investigations should be undertaken linked with the precipitation and the temperature as measured and not only bio-indicated.

## 5 Conclusions

We highlight that seabird populations provide a link between landfills history and soil eutrophication and time-related vegetation changes on small Mediterranean islands. We observed a remanence of the effects on soil parameters. However, we were expecting more changes in plant community composition but apart from the cover, the community remained similar to 11 years ago, with more ruderal plants.

It is important to follow the patterns of change in gull populations, the flora and the physical and chemical parameters of the soil to better understand the resistance and resilience of the ecosystem post-disturbance. Furthermore, gull populations showed a slight increase in 2021. It is necessary to determine whether this a natural fluctuation between years, a stochastic event, or whether the gulls found another food resource (such as the rubbish bins in coastal cities) leading to a new demographic explosion in the coming years.

## Acknowledgements

We are grateful to the Calanques National Park for the operational and financial support which enabled us to carry out this study, with the team of the *Secteur Littoral Ouest et Archipels*: Lorraine Anselme, Camille Brunet, Timothee Cuchet, Jean-Patrick Durand, Lionel Laso, Alain Mante, Olivier Montigny, Célia Pastorelli, Aurélien Robin; the team of *Secteur Monts et Vallons de Marseille* with Patrice d’Onofrio and the *Pôle Connaissances Scientifiques* with Olivier Ferreira. We also thank Daniel Pavon, Frédéric Médail, Thierry Taton, Véronique Bonnet and Christiane Rolando from the *Institut Méditerranéen de Biodiversité et d’Ecologie* for their help in identifying some of the plant species. Many thanks to Laurent Vassalo, Didier Gori and Patrick Hohener from the *Laboratoire de Chimie de l’Environnement* for their technical and scientific assistance in soil chemical analyses. Finally, we wish to thank Michael Paul (N°4500621889) and Cate Evans (N°4500634762) for reviewing the English language, and the two anonymous reviewers who helped us improve this article.

## Bibliography

- Aerts, R. *et al.* (2020) ‘Effects of sea birds and soil development on plant and soil nutritional parameters after 50 years of succession on Surtsey’, *Surtsey research*, 14(June), pp. 85–90. doi: 10.33112/surtsey.14.7.
- Affholder, M.-C. *et al.* (2014) "As, Pb, Sb and Zn transfer from soil to root of wild rosemary: do native symbionts matter?", *Plant and Soil*, 382, pp. 219-236. doi: 10.1007/s11104-014-2135-4
- Ahrens, C. W. *et al.* (2020) ‘Plant functional traits differ in adaptability and are predicted to be differentially affected by climate change’, *Ecology and Evolution*, 10(1), pp. 232–248. doi: 10.1002/ece3.5890.
- Baize, D. and Girard, M.-C. (1992) *Référentiel pédologique. Principaux sols d’Europe.*, Versailles, INRA Editions.
- Bancroft, W. J., Roberts, J. D. and Garkaklis, M. J. (2005) ‘Burrowing seabirds drive decreased diversity and structural complexity, and increased productivity in insular-vegetation communities’, *Australian Journal of Botany*, 53, pp. 231–241. Available at: papers2://publication/uuid/E56F92BF-

590 EC8C-4E56-B629-6C5BEA669C14.

591 Barbour, M. G. (1978) 'Salt spray as a microenvironmental factor in the distribution of beach plants at  
592 point reyes, California', *Oecologia*, 32(2), pp. 213–224. doi: 10.1007/BF00366073.

593 Baumberger, T. *et al.* (2012) 'Plant community changes as ecological indicator of seabird colonies'  
594 impacts on Mediterranean Islands', *Ecological Indicators*, 15(1), pp. 76–84. doi:  
595 10.1016/j.ecolind.2011.09.009.

596 Barbraud, C., Fortin, M., Charbonnier, Y., Delord, K., Gadennne, H., Thiebot, J. B., & Gélinaud, G.  
597 (2014). A comparison of direct and distance sampling methods to estimate abundance of nesting  
598 gulls. *Ardeola*, 61(2), 367-377.

599 Bellout, S. *et al.* (2021) 'Population growth for yellow-legged gulls in morocco changes in the  
600 population size of yellow-legged gull *larus michahellis* at Essaouira and Mogador Island, West-  
601 Central Morocco', *Marine Ornithology*, 49(1), pp. 101–107.

602 Bosch, M., Oro, D. and Ruiz, X. (1994) 'Dependence of yellow-legged gulls (*Larus cachinnans*) on  
603 food from human activity in two Western Mediterranean colonies', *Avocetta*, 18(January), pp. 135–  
604 139.

605 Bou, J., Vilar, L. and Carreras, J. (2020) 'Habitat changes over 45 years on the small Mediterranean  
606 archipelago of the Medes Islands (Catalonia, NE Spain)', *Plant Biosystems*, 0(0), pp. 1–11. doi:  
607 10.1080/11263504.2020.1810806.

608 Bouyahmed, H. and Moulaï, R. (2018) 'Influence of the yellow-legged gull on soil, composition and  
609 organisation of plant communities on some small Islands of Algeria', *Zoology and Ecology*, 28(4), pp.  
610 300–312. doi: 10.1080/21658005.2018.1527604.

611 del Cacho, M. and Lloret, F. (2012) 'Resilience of Mediterranean shrubland to a severe drought  
612 episode: The role of seed bank and seedling emergence', *Plant Biology*, 14(3), pp. 458–466. doi:  
613 10.1111/j.1438-8677.2011.00523.x.

614 Calvino-Cancela, M. (2011) 'Gulls (*Laridae*) as frugivores and seed dispersers', *Plant Ecology*,

615 212(7), pp. 1149–1157. doi: 10.1007/s11258-011-9894-2.

616 Castro, M. F., Neves, J. C., Francelino, M. R., Schaefer, C. E. G., & Oliveira, T. S. (2021). Seabirds  
617 enrich Antarctic soil with trace metals in organic fractions. *Science of the Total Environment*, 785,  
618 147271.

619 Chapin, F. S. *et al.* (1997) ‘Biotic control over the functioning of ecosystems’, *Science*, 277(5325), pp.  
620 500–504. doi: 10.1126/science.277.5325.500.

621 Cunningham, C. and Beazley, K. F. (2018) ‘Changes in human population density and protected areas  
622 in terrestrial global biodiversity hotspots, 1995-2015’, *Land*, 7(4), pp. 1–20. doi:  
623 10.3390/land7040136.

624 Daumalin, X. *et al.* (2016) *Les calanques industrielles de Marseille et leurs pollutions - Une histoire*  
625 *au présent*. Ref<sup>2</sup>C.

626 De La Peña-Lastra, S. *et al.* (2021) ‘Effects of a yellow legged gull (*Larus michahellis*) colony on soils  
627 and cliff vegetation in the Atlantic Islands of Galicia National Park (NW Spain)’, *Catena*, 199  
628 (December 2020). doi: 10.1016/j.catena.2020.105115.

629 De la Peña-Lastra, S., Affre, L. and Otero, X. L. (2020) ‘Soil nutrient dynamics in colonies of the  
630 yellow-legged seagull (*Larus michahellis*) in different biogeographical zones’, *Geoderma*, 361 (July  
631 2019). doi: 10.1016/j.geoderma.2019.114109.

632 Duda, M. P., Glew, J. R., Michelutti, N., Robertson, G. J., Montevecchi, W. A., Kissinger, J. A., ... &  
633 Smol, J. P. (2020). Long-term changes in terrestrial vegetation linked to shifts in a colonial seabird  
634 population. *Ecosystems*, 23, 1643-1656.

635 Duhem, C. *et al.* (2002) ‘Influence de l’accessibilité des ressources anthropiques sur les paramètres  
636 reproducteurs de deux colonies de goélands leucophées *Larus michahellis*’, *Revue d’Ecologie (La*  
637 *Terre et la Vie)*, 57(3–4), pp. 343–353.

638 Duhem, C. *et al.* (2003) ‘Opportunistic feeding responses of the yellow-legged gull *larus michahellis*  
639 to accessibility of refuse dumps’, *Bird Study*, 50(1), pp. 61–67. doi: 10.1080/00063650309461291.

640 Duhem, C. (2004) *Goélands surabondants et ressources alimentaires anthropiques: cas des colonies*  
641 *insulaires de Goélands leucophées du littoral provençal. Ecologie, Environnement. Université de*  
642 *droit, d'économie et des sciences - Aix-Marseille III. Français.*

643 Duhem, C. *et al.* (2007) 'Distribution of breeding sites and food constrains size and density of yellow-  
644 legged gull colonies', *Ecoscience*, 14(4), pp. 535–543. doi: 10.2980/1195-  
645 6860(2007)14[535:DOBSAF]2.0.CO;2.

646 Duhem, C. *et al.* (2008) 'Effects of anthropogenic food resources on yellow-legged gull colony size on  
647 Mediterranean islands', *Population Ecology*, 50(1), pp. 91–100. doi: 10.1007/s10144-007-0059-z.

648 Ellis, J. C. (2005) 'Marine birds on land: A review of plant biomass, species richness, and community  
649 composition in seabird colonies', *Plant Ecology*, 181(2), pp. 227–241. doi: 10.1007/s11258-005-7147-  
650 y.

651 Erel, Y., Veron, A. and Halicz, L. (1997) 'Tracing the transport of anthropogenic lead in the  
652 atmosphere and in soils using isotopic ratios', *Geochimica et Cosmochimica Acta*, 61(21), pp. 4495–  
653 4505. doi: 10.1016/S0016-7037(97)00353-0.

654 Gachet, S., Véla, E. and Tatoni, T. (2005) 'BASECO: A floristic and ecological database of  
655 Mediterranean French flora', *Biodiversity and Conservation*, 14(4), pp. 1023–1034. doi:  
656 10.1007/s10531-004-8411-5.

657 García, L. V. *et al.* (2002) 'Seagull influence on soil properties, chenopod shrub distribution, and leaf  
658 nutrient status in semi-arid Mediterranean islands', *Oikos*, 98(1), pp. 75–86. doi: 10.1034/j.1600-  
659 0706.2002.980108.x.

660 Gargominy, O. *et al.* (2020) 'TAXREF v14.0, référentiel taxonomique pour la France : méthodologie,  
661 mise en oeuvre et diffusion', p. 126. Available at:  
662 <https://inpn.mnhn.fr/telechargement/referentielEspece/taxref/14.0/menu>.

663 Ghermaoui, M., Hassaine, K. and Moulaï, R. (2016) 'Influence du Goéland leucophée *Larus*  
664 *michahellis* sur les formations végétales ouvertes du littoral de Rachgoun (Ouest Oranie, Algérie)',

665 *Revue d'écologie*, 71(3), pp. 250–265.

666 Grant, M. L., Bond, A. L., & Lavers, J. L. (2022). The influence of seabirds on their breeding, roosting  
667 and nesting grounds: A systematic review and meta- analysis. *Journal of Animal Ecology*, 91(6),  
668 1266-1289. doi: 10.1111/1365-2656.13699

669 Grime, J. (1977) 'Evidence for the existence of three primary strategies in plants and its relevance to  
670 ecological and.pdf', *The American Naturalist*, 111(982), pp. 1169–1194.

671 Grosso, F., Bååth, E. and De Nicola, F. (2016) 'Bacterial and fungal growth on different plant litter in  
672 Mediterranean soils: Effects of C/N ratio and soil pH', *Applied Soil Ecology*, 108, pp. 1–7. doi:  
673 10.1016/j.apsoil.2016.07.020.

674 Hanrahan, B. R., Tank, J. L., Speir, S. L., Trentman, M. T., Christopher, S. F., Mahl, U. H., & Royer,  
675 T. V. (2021). Extending vegetative cover with cover crops influenced phosphorus loss from an  
676 agricultural watershed. *Science of The Total Environment*, 801, 149501.

677 Hawke, D. J., & Wu, J. R. (2012). Soil selenium in a forested seabird colony: distribution, sources,  
678 uptake by plants, and comparison with non-seabird sites. *Soil Research*, 50(7), 588-595.

679 Heckenroth, A. *et al.* (2022) "Coronilla juncea, a native candidate for phytostabilization of potentially  
680 toxic elements and restoration of Mediterranean soils", *Scientific reports*, 12, 10033. doi:  
681 10.1038/s41598-022-141339-4.

682 Heckenroth, A. *et al.* (2016) 'Selection of native plants with phytoremediation potential for highly  
683 contaminated Mediterranean soil restoration: Tools for a non-destructive and integrative approach',  
684 *Journal of Environmental Management*, 183, pp. 850–863. doi: 10.1016/j.jenvman.2016.09.029.

685 Heilmeier, H. (2019) 'Functional traits explaining plant responses to past and future climate changes',  
686 *Flora: Morphology, Distribution, Functional Ecology of Plants*, 254(April), pp. 1–11. doi:  
687 10.1016/j.flora.2019.04.004.

688 Holdaway, R. N., Hawke, D. J., Causer, J. E., & Ogden, S. (1999). Soil indicators of pre-European  
689 seabird breeding in New Zealand at sites identified by predator deposits. *Soil Research*, 37(1), 103-

690 114.

691 Hooper, D. U. *et al.* (2005) ‘Effect of biodiversity on ecosystem functioning: a consensus of current  
692 knowledge’, *Science*, 75(1), pp. 3–35. doi: 10.1126/science.305.5687.1101.

693 INPN MNHN, 2023 March 1. Statuts d'Evaluation, de Protection et de Menace  
694 [https://inpn.mnhn.fr/espece/cd\\_nom/199374/tab/statut#ancStatutEspece](https://inpn.mnhn.fr/espece/cd_nom/199374/tab/statut#ancStatutEspece)

695 Ishida, A. (1997) ‘Seed germination and seedling survival in a colony of the common cormorant,  
696 *Phalacrocorax carbo*’, *Ecological Research*, 12(3), pp. 249–256. doi: 10.1007/BF02529454.

697 Jaunatre, R., Buisson, E., & Dutoit, T. (2014). Can ecological engineering restore Mediterranean  
698 rangeland after intensive cultivation? A large-scale experiment in southern France. *Ecological*  
699 *Engineering*, 64, 202-212.

700 Kimball, S. *et al.* (2010) ‘Contemporary climate change in the Sonoran Desert favors cold-adapted  
701 species’, *Global Change Biology*, 16(5), pp. 1555–1565. doi: 10.1111/j.1365-2486.2009.02106.x.

702 Kiss, R. *et al.* (2018) ‘Grassland seed bank and community resilience in a changing climate’,  
703 *Restoration Ecology*, 26(June), pp. S141–S150. doi: 10.1111/rec.12694.

704 Komárek, M. *et al.* (2008) ‘Lead isotopes in environmental sciences: A review’, *Environment*  
705 *International*, 34(4), pp. 562–577. doi: 10.1016/j.envint.2007.10.005.

706 Laguna, C. *et al.* (2021) ‘Nutrient enrichment and trace element accumulation in sediments caused by  
707 waterbird colonies at a Mediterranean semiarid floodplain’, *Science of the Total Environment*, 777, p.  
708 145748. doi: 10.1016/j.scitotenv.2021.145748.

709 MacArthur, R. H. and Wilson, E. O. (1963) ‘An Equilibrium Theory of Insular Zoogeography’,  
710 *Evolution*, 17(4), pp. 373–387.

711 Mathieu, C. and Pieltain, F. (2003) *Analyse chimique des sols Méthodes choisies*. Editions T. Edited  
712 by Lavoisier.

713 Matson, P., Lohse, K. A., & Hall, S. J. (2002). The globalization of nitrogen deposition: consequences

714 for terrestrial ecosystems. *Ambio*, 113-119.

715 Martín- Queller, E., Albert, C. H., Dumas, P. J., & Saatkamp, A. (2017). Islands, mainland, and  
716 terrestrial fragments: How isolation shapes plant diversity. *Ecology and evolution*, 7(17), 6904-6917.

717 Médail, F. (2017) ‘The specific vulnerability of plant biodiversity and vegetation on Mediterranean  
718 islands in the face of global change’, *Regional Environmental Change*, 17(6), pp. 1775–1790. doi:  
719 10.1007/s10113-017-1123-7.

720 Médail, F., & Vidal, É. (1998). Organisation de la richesse et de la composition floristiques d’îles de la  
721 Méditerranée occidentale (sud-est de la France). *Canadian Journal of Botany*, 76(2), 321-331.

722 Météo France, 2023 March 1, Données publiques.  
723 [https://donneespubliques.meteofrance.fr/?fond=contenu&id\\_contenu=37](https://donneespubliques.meteofrance.fr/?fond=contenu&id_contenu=37)

724 Mouga, T. *et al.* (2021) ‘Recent efforts to recover armeria berlengensis, an endemic species from  
725 berlingas archipelago, Portugal’, *Plants*, 10(3), pp. 1–19. doi: 10.3390/plants10030498.

726 Mulder, C. P. H. and Keall, S. N. (2001) ‘Burrowing seabirds and reptiles: Impacts on seeds, seedlings  
727 and soils in an island forest in New Zealand’, *Oecologia*, 127(3), pp. 350–360. doi:  
728 10.1007/s004420000600.

729 Myrcha, A., & Tatur, A. (1991). Ecological role of the current and abandoned penguin rookeries in the  
730 land environment of the maritime Antarctic. *Polish Polar Research*, 3-24.

731 Nogales, M. *et al.* (2001) ‘Ecological and biogeographical implications of Yellow-Legged Gulls  
732 (*Larus cachinnans* Pallas) as seed dispersers of *Rubia fruticosa* Ait. (Rubiaceae) in the Canary  
733 Islands’, *Journal of Biogeography*, 28(9), pp. 1137–1145. doi: 10.1046/j.1365-2699.2001.00622.x.

734 Nyiraneza, J. *et al.* (2017) ‘Changes in soil organic matter over 18 yr in Prince Edward Island,  
735 Canada’, *Canadian Journal of Soil Science*, 97(4), pp. 745–756. doi: 10.1139/cjss-2017-0033.

736 Otero, X. L. *et al.* (2015) ‘Phosphorus in seagull colonies and the effect on the habitats. The case of  
737 yellow-legged gulls (*Larus michahellis*) in the Atlantic Islands National Park (Galicia-NW Spain)’,  
738 *Science of the Total Environment*, 532, pp. 383–397. doi: 10.1016/j.scitotenv.2015.06.013.



739 Otero, X. L. *et al.* (2018) 'Trace elements in biomaterials and soils from a Yellow-legged gull (*Larus*  
740 *michahellis*) colony in the Atlantic Islands of Galicia National Park (NW Spain)', *Marine Pollution*  
741 *Bulletin*, 133(May), pp. 144–149. doi: 10.1016/j.marpolbul.2018.05.027.

742 Perring, M. P. *et al.* (2015) 'Advances in restoration ecology: Rising to the challenges of the coming  
743 decades', *Ecosphere*, 6(8). doi: 10.1890/ES15-00121.1.

744 Perring, M. P. *et al.* (2016) 'Global environmental change effects on ecosystems: The importance of  
745 land-use legacies', *Global Change Biology*, 22(4), pp. 1361–1371. doi: 10.1111/gcb.13146.

746 Post, D. M. (2002) 'Using stable isotopes to estimate trophic position: Models, methods, and  
747 assumptions', *Ecology*, 83(3), pp. 703–718. doi: 10.1890/0012-  
748 9658(2002)083[0703:USITET]2.0.CO;2.

749 Pribyl, D. W. (2010) 'A critical review of the conventional SOC to SOM conversion factor',  
750 *Geoderma*, 156(3–4), pp. 75–83. doi: 10.1016/j.geoderma.2010.02.003.

751 Purschke, O. *et al.* (2012) 'Linking landscape history and dispersal traits in grassland plant  
752 communities', *Oecologia*, 168(3), pp. 773–783. doi: 10.1007/s00442-011-2142-6.

753 Quézel, P., & Médail, F. (2003). *Ecologie et biogéographie des forêts du bassin méditerranéen*.  
754 Elsevier.

755 Rashed, M.N. (2010) "Monitoring of contaminated toxic and heavy metals from mine tailings through  
756 age accumulation in soil and some wild plants at Southeast Egypt", *J. Hazard. Mater.*, 178, pp. 739-  
757 746. doi: 10.1016/j.hazmat.2010.01.147.

758 Raunkiaer, C. (1934). The life forms of plants and statistical plant geography; being the collected  
759 papers of C. Raunkiaer. *The life forms of plants and statistical plant geography; being the collected*  
760 *papers of C. Raunkiaer*.

761 R Core Team (2022). R: *A language and environment for statistical computing*. R Foundation for  
762 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

763 Real, E., Oro, D., Martínez- Abraín, A., Igual, J. M., Bertolero, A., Bosch, M., & Tavecchia, G.

764 (2017). Predictable anthropogenic food subsidies, density- dependence and socio- economic factors  
 765 influence breeding investment in a generalist seabird. *Journal of Avian Biology*, 48(11), 1462-1470.  
 766 doi: 10.1111/jav.01454.

767 Saatkamp, A., Argagnon, O., Noble, V., Finocchiaro, M., Meineri, E. (2023, *in press*). Climate change  
 768 impacts on Mediterranean vegetation are amplified at low altitudes. *Global Ecology and*  
 769 *Biogeography*.

770 Saatkamp, A., Falzon, N., Argagnon, O., Noble, V., Dutoit, T., & Meineri, E. (2022). Calibrating  
 771 ecological indicator values and niche width for a Mediterranean flora. *Plant Biosystems-An*  
 772 *International Journal Dealing with all Aspects of Plant Biology*, 1-11. doi:  
 773 10.1080/11263504.2022.2104399.

774 Saatkamp, A., Henry, F. and Dutoit, T. (2020) ‘Romans Shape Today’s Vegetation and Soils: Two  
 775 Millennia of Land-Use Legacy Dynamics in Mediterranean Grasslands’, *Ecosystems*. doi:  
 776 10.1007/s10021-020-00581-w.

777 Schaffers, A. P. and Sýkora, K. V. (2000) ‘Reliability of Ellenberg indicator values for moisture,  
 778 nitrogen and soil reaction: a comparison with field measurements’, *Journal of Vegetation Science*, 11,  
 779 pp. 225–244.

780 Schwabe, A., Kratochwil, A. and Pignatti, S. (2007) ‘Plant indicator values of a high-phytodiversity  
 781 country (Italy) and their evidence, exemplified for model areas with climatic gradients in the southern  
 782 inner Alps’, *Flora: Morphology, Distribution, Functional Ecology of Plants*, 202(5), pp. 339–349. doi:  
 783 10.1016/j.flora.2006.09.002.

784 Shoji, A., Elliott, K. H., Aris- Brosou, S., Mizukawa, H., Nakayama, S. M. M., Ikenaka, Y., ... &  
 785 Watanuki, Y. (2019). Biotransport of metallic trace elements from marine to terrestrial ecosystems by  
 786 seabirds. *Environmental toxicology and chemistry*, 38(1), 106-114.

787 Sigurdsson, B. D. and Magnusson, B. (2010) ‘Effects of seagulls on ecosystem respiration, soil  
 788 nitrogen and vegetation cover on a pristine volcanic island, Surtsey, Iceland’, *Biogeosciences*, 7(3),

789 pp. 883–891. doi: 10.5194/bg-7-883-2010.

790 Sobey, D. G. and Kenworthy (1979) ‘The Relationship between Herring Gulls and the Vegetation of  
791 their Breeding Colonies’, *Journal of Ecology*, 67(2), pp. 469–496.

792 Sun, L., Xie, Z. & Zhao, J. A 3,000-year record of penguin populations. *Nature* 407, 858 (2000).  
793 <https://doi.org/10.1038/35038163>

794 Thibault, J. C. *et al.* (1996) ‘Recent trends in breeding marine birds of the Mediterranean region with  
795 special reference to Corsica’, *Waterbirds*, 19(SUPPL.1), pp. 31–40. doi: 10.2307/1521943.

796 Thompson, K. R. and Rothery, P. (1991) ‘A census of the black-browed albatross *Diomedea*  
797 *melanophrys* population on Steeple Jason Island, Falkland Islands’, *Biological Conservation*, 56(1),  
798 pp. 39–48. doi: 10.1016/0006-3207(91)90087-P.

799 Tison, J. M., Jauzein, P., Michaud, H., & Michaud, H. (2014). *Flore de la France méditerranéenne*  
800 *continentale* (Vol. 1). Turriers: Naturalia publications.

801 Torres, I. C. *et al.* (2012) ‘Stable isotope ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) values of sediment organic matter in  
802 subtropical lakes of different trophic status’, *Journal of Paleolimnology*, 47(4), pp. 693–706. doi:  
803 10.1007/s10933-012-9593-6.

804 UNEP/MAP-Plan Bleu (2009) *State of the Environment and Development in the Mediterranean -*  
805 *2009, UNEP/MAP-Plan Bleu, Athens.*

806 Vidal, E. *et al.* (1998) ‘Functional analysis of the newly established plants induced by nesting gulls on  
807 Riou archipelago (Marseille, France)’, *Acta Oecologica*, 19(3), pp. 241–250. doi: 10.1016/S1146-  
808 609X(98)80028-3.

809 Vidal, E. (1998) *Organisation des phytocénoses en milieu insulaire méditerranéen perturbé. Analyse*  
810 *des inter-relations entre les colonies de Goélants leucophées et la végétation des îles de Marseille.*  
811 UNIVERSITÉ DE DROIT, D’ÉCONOMIE ET DES SCIENCES D’AIX-MARSEILLE III (AIX-  
812 MARSEILLE.

813 Vidal, E. *et al.* (2000) ‘Seabirds drive plant species turnover on small Mediterranean islands at the

814 expense of native taxa', *Oecologia*, 122(3), pp. 427–434. doi: 10.1007/s004420050049.

815 Vidal, E., Médail, F. and Taton, T. (1998) 'Is the yellow-legged gull a superabundant bird species in  
816 the Mediterranean? Impact on fauna and flora, conservation measures and research priorities',  
817 *Biodiversity and Conservation*, 7(8), pp. 131–141. doi: 10.1023/A.

818 Vié, J. C., Hilton-Taylor, C. and Stuart, S. N. (Eds. . (2009) *Wildlife in a changing world: an analysis*  
819 *of the 2008 IUCN Red List of threatened species*, IUCN.

820 Vogiatzakis, I. N., Mannion, A. M. and Sarris, D. (2016) 'Mediterranean island biodiversity and  
821 climate change: the last 10,000 years and the future', *Biodiversity and Conservation*, 25(13), pp.  
822 2597–2627. doi: 10.1007/s10531-016-1204-9.

823 Wait, D. A., Aubrey, D. P. and Anderson, W. B. (2005) 'Seabird guano influences on desert islands:  
824 Soil chemistry and herbaceous species richness and productivity', *Journal of Arid Environments*,  
825 60(4), pp. 681–695. doi: 10.1016/j.jaridenv.2004.07.001.

826 Whittaker, R. J. *et al.* (2017) 'Island biogeography: Taking the long view of nature's laboratories',  
827 *Science*, 357(6354). doi: 10.1126/science.aam8326.

828 Ziółek, M., Bartmiński, P. and Stach, A. (2017) 'The influence of seabirds on the concentration of  
829 selected heavy metals in organic soil on the Bellsund Coast, Western Spitsbergen', *Arctic, Antarctic,*  
830 *and Alpine Research*, 49(4), pp. 507–520. doi: 10.1657/AAAR0016-024.

831 Zwolicki, A., Barcikowski, M., Barcikowski, A. *et al.* Seabird colony effects on soil properties and  
832 vegetation zonation patterns on King George Island, Maritime Antarctic. *Polar Biol* **38**, 1645–1655  
833 (2015). <https://doi.org/10.1007/s00300-015-1730-z>

834

835

**Title: Yellow-legged gull populations (*Larus michahellis*) link the history of landfills to soil eutrophication and time-related vegetation changes on small Mediterranean islands**

Clémentine Mutillod<sup>1\*</sup>, Teddy Baumberger<sup>2</sup>, Pascale Prudent<sup>3</sup>, Arne Saatkamp<sup>1</sup>, Eric Vidal<sup>1,4</sup>, Lidwine Le-Mire-Pecheux<sup>5</sup>, Laurence Affre<sup>1</sup>

<sup>(1)</sup> Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE Institut Méditerranéen de Biodiversité et d'Écologie marine et continentale, Marseille, France

<sup>(2)</sup> Société EcoStat, Baudinard-sur-Verdon, France

<sup>(3)</sup> Aix Marseille Univ, CNRS, LCE Laboratoire de Chimie de l'Environnement, Marseille, France

<sup>(4)</sup> UMR ENTROPIE (IRD, Université de la Réunion, IFREMER, Université de la Nouvelle-Calédonie, CNRS), BP A5, Nouméa CEDEX 98848, New Caledonia

<sup>(5)</sup> Calanques National Park, Marseille, France

\* Contact details: [clementine.mutillod@gmail.com](mailto:clementine.mutillod@gmail.com) IMBE, IUT Avignon, 337 chemin des Meinajariés, Site Agroparc BP 61207, 84911 Avignon cedex 09, France

## TABLES

Table 1: Data of studied islands and the coastal part of the Calanques National Park (CNP), and number of plots surveyed in 1997, 2008 and 2021.

|                      | Islands           | Area (ha) | Distance to mainland sea shore (m) | Maximum altitude (m) | Number of plots |
|----------------------|-------------------|-----------|------------------------------------|----------------------|-----------------|
| Frioul Archipelago   | Pomègues          | 89        | 2800                               | 86                   | 13              |
|                      | Ratonneau         | 95        | 1800                               | 74                   | 13              |
| Riou Archipelago     | Riou              | 90.3      | 3100                               | 190                  | 16              |
|                      | Maïre             | 27.6      | 50                                 | 141                  | 4               |
|                      | Plane             | 15        | 2100                               | 22                   | 3               |
|                      | Jarre             | 18        | 800                                | 57                   | 3               |
|                      | Jarron            | 3         | 800                                | 33                   | 1               |
|                      | Grand Congloué    | 2         | 3525                               | 50                   | 1               |
|                      | Tiboulén de Maïre | 2.3       | 525                                | 47                   | 2               |
| Mainland part of CNP |                   | 8157.8    |                                    | 645                  | 22              |

Table 2: Mesological parameters measured in the 500 m<sup>2</sup> circles for each plot.

| Environmental variables measured in the 500 m <sup>2</sup> circles  | Measures and estimations made   |
|---|---|
| Altitude  | GPS measures.   |
| Exposure  | 1: North, 2: North-East, 3: West, 4: South-East, 5: South   |
| Distance from the coastline   | Reflects the amount of salt deposited on the ground (Barbour, 1978).  |
| Slope   | Slope was estimated visually using 6 categories: 0: flat, 1: 0% < slope < 10%, 2: 0% ≤ slope < 25%, 3: 25% ≤ slope < 50%, 4: 50% ≤ slope < 75%, 5: slope ≥ 75 %.  |
| Cover rate of each substrate type:<br>rock,<br>block i.e. Ø > 20 cm,<br>stone i.e. Ø < 20 cm,<br>bare ground  | Cover was estimated visually using respectively 6 categories: 0: cover = 0%, 1: 0% < cover < 10%, 2: 10% ≤ cover < 25%, 3: 25% ≤ cover < 50%, 4: 50% ≤ cover < 75%, 5: cover ≥ 75 %.  |
| Vegetation cover and the different layers (High arborescent h (height) > 10 m, Arborescent 10 > h > 4 m, High shrub 4 > h > 2 m, Shrub 2 > h > 0.5 m, Herbaceous h < 0.5 m) |   |
| Gull nest density (number of nests)   | Is considered as an indicator of gull pressure intensity. Following Thompson & Rothery (1991), gull nest density was measured by counting the number of both occupied – with eggs – and empty nests during the egg-laying period in the 78 plots for the three years 1997, 2008 and 2021. |
| Cover of the resting places per plot  | Measured on a semi-quantitative basis using the 6 cover abundance categories previously described, used as a measure of pressure by gulls.  |

Table 3: Complementary soil parameters measured on archipelagos and mainland plots, and on gulls' droppings (mean ± SD) in 2021.

| Area      | Conductivity (mS/cm) | P (mg/g dry matter (DM)) | K (mg/gDM)        | Na (mg/gDM)       | Cu (mg/gDM)   | Pb (mg/gDM)     | Zn (mg/gDM)     | Number of plots |
|-----------|----------------------|--------------------------|-------------------|-------------------|---------------|-----------------|-----------------|-----------------|
| Frioul    | 1.07 ± 1.78          | 1671.42 ± 939.93         | 8897.83 ± 2663.84 | 1467.50 ± 1531.58 | 48.12 ± 90.62 | 96.50 ± 98.26   | 150.18 ± 82.03  | 26              |
| Riou      | 0.75 ± 0.35          | 3300.35 ± 2020.55        | 9705.64 ± 2597.42 | 1339.63 ± 714.84  | 29.85 ± 10.37 | 133.48 ± 81.75  | 157.64 ± 58.31  | 30              |
| Mainland  | 0.64 ± 1.19          | 472.26 ± 205.22          | 8165.27 ± 2692.98 | 1125.71 ± 2246.57 | 25.74 ± 10.99 | 337.90 ± 299.36 | 212.70 ± 138.27 | 22              |
| Droppings | NA                   | 6767.72 ± 5106.96        | 4217.13 ± 3671.31 | 3665.08 ± 1401.69 | 71.53 ± 72.36 | 4.34 ± 14.03    | 396.13 ± 146.16 | 9               |

Table 4: Contamination factors calculated for Cu, Pb, Zn and the Pollution Load Index for Frioul and Riou archipelagos and the mainland area (mean ± SD).

| Area     | CF Cu        | CF Pb       | CF Zn       | PLI         |
|----------|--------------|-------------|-------------|-------------|
| Frioul   | 6.87 ± 12.94 | 2.24 ± 2.28 | 2.28 ± 1.24 | 2.10 ± 0.54 |
| Riou     | 4.26 ± 1.48  | 3.22 ± 1.83 | 2.39 ± 0.88 | 2.10 ± 0.30 |
| Mainland | 3.68 ± 1.57  | 8.23 ± 6.90 | 3.22 ± 2.09 | 2.33 ± 0.56 |

Title: **Yellow-legged gull populations (*Larus michahellis*) link the history of landfills to soil eutrophication and time-related vegetation changes on small Mediterranean islands**

Clémentine Mutillod<sup>1\*</sup>, Teddy Baumberger<sup>2</sup>, Pascale Prudent<sup>3</sup>, Arne Saatkamp<sup>1</sup>, Eric Vidal<sup>1,4</sup>, Lidwine Le-Mire-Pecheux<sup>5</sup>, Laurence Affre<sup>1</sup>

<sup>(1)</sup> Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE Institut Méditerranéen de Biodiversité et d'Écologie marine et continentale, Marseille, France

<sup>(2)</sup> Société EcoStat, Baudinard-sur-Verdon, France

<sup>(3)</sup> Aix Marseille Univ, CNRS, LCE Laboratoire de Chimie de l'Environnement, Marseille, France

<sup>(4)</sup> UMR ENTROPIE (IRD, Université de la Réunion, IFREMER, Université de la Nouvelle-Calédonie, CNRS), BP A5, Nouméa CEDEX 98848, New Caledonia

<sup>(5)</sup> Calanques National Park, Marseille, France

\* Contact details: [clementine.mutillod@gmail.com](mailto:clementine.mutillod@gmail.com) IMBE, IUT Avignon, 337 chemin des Meinajariés, Site Agroparc BP 61207, 84911 Avignon cedex 09, France

**FIGURES**




Fig. 1: Location of the islands in the Frioul archipelago and the Riou Archipelago (Marseille, south-eastern France). The dark squares show the localisation of the plots on the islands studied (56) and the nearby mainland areas (22).






Fig. 2: Gull (*Larus michahellis*) population dynamics in the Riou and Frioul archipelagos from 1923 to 2021, and indications of the opening and closing of the landfills in the vicinity of Marseille.




Fig. 3. A. Mean number of nests (mean + SD) at plots in 1997, 2008 and 2021 on Frioul (n = 26) and Riou (n = 30) islands. B. Mean cover of resting places in 2008 and 2021 at the same sites.




Fig. 4. Changes in soil parameters during the monitoring period (1997 to 2021) on Frioul (dark grey) and Riou (light grey) islands and the mainland (yellow) and data from droppings (light blue) (for total nitrogen in 2021), non-significant (ns): p-value > 0.05, \*: p-value < 0.05, \*\*: p-value < 0.01, \*\*\*: p-value < 0.001. A. The pH of the soil. B. Soil organic carbon (mg/g dry matter (DM)). C. The total nitrogen (mg/g DM). D. The soil C/N ratio.









Fig. 5. Boxplots of the soil measured physical and chemical parameters in 2021 in the Frioul (n = 26) and Riou (n = 30) archipelagos, on the nearby mainland (n = 22) (south-eastern France) and in the droppings (n = 9), non-significant (ns): ns: p-value > 0.05, \*: p-value < 0.05, \*\*: p-value < 0.01, \*\*\*: p-value < 0.001, \*\*\*\* p-value < 0.0001. A. The total phosphorus (mg/kg DM). B. The soil potassium (mg/kg DM). C. The soil sodium (mg/kg DM). D. The soil copper (mg/kg DM). E. The soil lead (mg/kg DM). F. The soil zinc (mg/kg DM).




Fig. 6. The Delta ( $\delta$ ) of  $^{13}\text{C}$  versus  $\delta^{15}\text{N}$  (means  $\pm$  SD) of stable isotopic composition in soil samples from the Frioul (n = 26) and Riou (n = 30) archipelagos, the nearby mainland (n = 22) (south-eastern France) and droppings samples (n = 9).





Fig. 7. Non-metrical multidimensional scaling (NMDS) of plant community composition of the three years of the survey (1997, 2008 and 2021) in the Frioul and Riou archipelagos.



Click here to access/download

**Supplementary Material**

MutillodEtal\_SupplementaryMaterial\_STOTEN\_Revised.  
docx



Title: **Yellow-legged gull populations (*Larus michahellis*) link the history of landfills to soil eutrophication and time-related vegetation changes on small Mediterranean islands**

Clémentine Mutillod<sup>1\*</sup>, Teddy Baumberger<sup>2</sup>, Pascale Prudent<sup>3</sup>, Arne Saatkamp<sup>1</sup>, Eric Vidal<sup>1,4</sup>, Lidwine Le-Mire-Pecheux<sup>5</sup>, Laurence Affre<sup>1</sup>

<sup>(1)</sup> Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE Institut Méditerranéen de Biodiversité et d'Écologie marine et continentale, Marseille, France

<sup>(2)</sup> Société EcoStat, Baudinard-sur-Verdon, France

<sup>(3)</sup> Aix Marseille Univ, CNRS, LCE Laboratoire de Chimie de l'Environnement, Marseille, France

<sup>(4)</sup> UMR ENTROPIE (IRD, Université de la Réunion, IFREMER, Université de la Nouvelle-Calédonie, CNRS), BP A5, Nouméa CEDEX 98848, New Caledonia

<sup>(5)</sup> Calanques National Park, Marseille, France

\* Contact details: [clementine.mutillod@gmail.com](mailto:clementine.mutillod@gmail.com) IMBE, IUT Avignon, 337 chemin des Meinajariés, Site Agroparc BP 61207, 84911 Avignon cedex 09, France

### **Declaration of interests**

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

**Title: Yellow-legged gull populations (*Larus michahellis*) link the history of landfills to soil eutrophication and time-related vegetation changes on small Mediterranean islands**

Clémentine Mutillod<sup>1\*</sup>, Teddy Baumberger<sup>2</sup>, Pascale Prudent<sup>3</sup>, Arne Saatkamp<sup>1</sup>, Eric Vidal<sup>1,4</sup>, Lidwine Le-Mire-Pecheux<sup>5</sup>, Laurence Affre<sup>1</sup>

<sup>(1)</sup> Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE Institut Méditerranéen de Biodiversité et d'Écologie marine et continentale, Marseille, France

<sup>(2)</sup> Société EcoStat, Baudinard-sur-Verdon, France

<sup>(3)</sup> Aix Marseille Univ, CNRS, LCE Laboratoire de Chimie de l'Environnement, Marseille, France

<sup>(4)</sup> UMR ENTROPIE (IRD, Université de la Réunion, IFREMER, Université de la Nouvelle-Calédonie, CNRS), BP A5, Nouméa CEDEX 98848, New Caledonia

<sup>(5)</sup> Calanques National Park, Marseille, France

\* Contact details: [clementine.mutillod@gmail.com](mailto:clementine.mutillod@gmail.com) IMBE, IUT Avignon, 337 chemin des Meinajariés, Site Agroparc BP 61207, 84911 Avignon cedex 09, France

**Authors Contributions Statement**

**Clémentine Mutillod:** Conceptualization, Methodology, Formal analysis, Investigation, Resources, Data Curation, Writing - Original Draft, Visualization

**Teddy Baumberger:** Conceptualization, Methodology, Formal analysis, Investigation, Resources, Data Curation, Writing - Review & Editing

**Pascale Prudent:** Conceptualization, Methodology, Investigation, Resources, Data Curation, Writing - Review & Editing, Funding acquisition

**Arne Saatkamp:** Conceptualization, Methodology, Formal analysis, Resources, Writing - Review & Editing

**Eric Vidal:** Conceptualization, Methodology, Investigation, Resources, Data Curation, Writing - Review & Editing

**Lidwine Le-Mire-Pecheux:** Conceptualization, Methodology, Resources, Supervision, Project administration, Funding acquisition

**Laurence Affre:** Conceptualization, Methodology, Resources, Writing - Review & Editing, Supervision, Project administration, Funding acquisition