



**HAL**  
open science

## Using priority effects for grassland restoration: Sequential sowing can promote subordinate species

Aure Durbecq, Armin Bischoff, Elise Buisson, Emmanuel Corcket, Renaud Jaunatre

### ► To cite this version:

Aure Durbecq, Armin Bischoff, Elise Buisson, Emmanuel Corcket, Renaud Jaunatre. Using priority effects for grassland restoration: Sequential sowing can promote subordinate species. *Applied Vegetation Science*, 2023, 26 (4), pp.1-12. 10.1111/avsc.12748 . hal-04327974

**HAL Id: hal-04327974**

**<https://hal.science/hal-04327974>**

Submitted on 6 Dec 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# Using priority effects for grassland restoration: Sequential sowing can promote subordinate species

Aure Durbecq<sup>1,2,3</sup>  | Armin Bischoff<sup>2</sup>  | Elise Buisson<sup>2</sup>  | Emmanuel Corcket<sup>2</sup>  |  
Renaud Jaunatre<sup>1</sup> 

<sup>1</sup>Université Grenoble Alpes, INRAE, LESSEM, St-Martin-d'Hères, France

<sup>2</sup>Aix Marseille Université, Avignon Université, CNRS, IRD, IMBE, Marseille, Avignon, France

<sup>3</sup>Environmental consultancy ECO-MED, Marseille, France

## Correspondence

Aure Durbecq, I.U.T Avignon, Campus Jean Henri Fabre, 337 Chemin des Meinajaries, BP 61207, 84911 Avignon Cedex 9, France.  
Email: [aure.durbecq@gmail.com](mailto:aure.durbecq@gmail.com)

## Funding information

Association Nationale de la Recherche et de la Technologie; CIFRE; RTE

## Abstract

**Aims:** The influence of priority effects on plant community succession is increasingly discussed in community ecology. However, most grassland restoration approaches involving seed addition transfer all target species at the same time. Our objective was to tackle the question: does the order of arrival influence the establishment of restored communities?

**Location:** La-Bâtie-Neuve, Southern Alps, France.

**Methods:** We applied sequential sowing using two groups, one set of dominant species and one set of subordinate species, each comprising three different perennial plant species. We tested four sowing treatments: control (without any sowing), two sequential sowing treatments (dominants first or subordinates first) and synchronous sowing. We analysed plant cover each year for three years after sowing and calculated priority and earliness indices (the third year) for each group and each sown species.

**Results:** Manipulating the order of arrival shaped community composition and trajectories. Some species of both groups were positively affected by being sown first compared to being sown synchronously. However, dominant and subordinate groups differed in their earliness index, showing a significant benefit for subordinates to be sown first. The subordinate species *Onobrychis viciifolia* and *Plantago lanceolata* established only when they were sown first, while *Festuca cinerea* showed greater establishment when sown first and simultaneously, compared to late sowing. The dominant *Anthyllis vulneraria* was not affected by date or type of sowing. However, the cover of the most dominant *Bromopsis erecta* was lower when being sown second, allowing a control of its dominance by delayed sowing.

**Conclusion:** The strength of priority effects differs between species, which may depend on niche characteristics or microenvironment, influencing (negatively or positively) the establishment of late-arriving species and affecting their competitive abilities. Our study provides evidence that plant community assembly was influenced by the order of arrival, but demonstrated a strong species-specific response to priority effects.

Co-ordinating Editor: Péter Török

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Applied Vegetation Science* published by John Wiley & Sons Ltd on behalf of International Association for Vegetation Science.

## KEYWORDS

community assembly, early-arriving species, ecological restoration, late-arriving species, mesophilous grassland, niche modification, stepwise sowing, time-advanced

## 1 | INTRODUCTION

Priority effects occur when the order of species arrival in a habitat changes biotic and abiotic properties of the microenvironment and thus affects the recruitment, establishment, growth and/or reproduction of late-arriving species (von Gillhaussen et al., 2014; Temperton et al., 2016; Weidlich et al., 2018). Priority effects can be either positive (i.e., facilitative; Bertness & Shumway, 1993) or negative (i.e., inhibitory, competitive; Cole, 1983; Fukami, 2015). Most of the studied priority effects, however, show that a competitive advantage for the first-arriving species prevails because of their greater size or density compared with late-arriving species (i.e., size-asymmetric competition; Grman & Suding, 2010; Wainwright et al., 2012). Many other mechanisms may drive priority effects, such as niche pre-emption or niche modification (Fukami, 2015; Helsen et al., 2016), plant–soil feedbacks and soil legacies (Bever, 2003; Grman & Suding, 2010; Fukami & Nakajima, 2013; van der Putten et al., 2013) or allelopathic effects (Levine et al., 2003). Furthermore, the strength of priority effects can be mediated by both biotic and abiotic factors, such as species identity (von Gillhaussen et al., 2014; Cleland et al., 2015; Wilsey et al., 2015; Stuble & Souza, 2016; Werner et al., 2016) or overlapping niches (Vannette & Fukami, 2014), predation pressure (Chase et al., 2009) and habitat size or productivity (Fukami, 2004; Orrock & Fletcher Jr., 2005; Chase, 2010; Kardol et al., 2013). In productive environments, priority effects may be very strong, resulting in alternative stable states (Chase, 2003; Hobbs & Norton, 2004; Kardol et al., 2013; Weidlich et al., 2017). Priority effects are, however, poorly studied in less productive environments (Wilsey, 2020). Finally, priority effects do not only influence the early community assembly but also shape the potential community trajectories, including their species composition and associated functions (Grman & Suding, 2010; von Gillhaussen et al., 2014; Werner et al., 2016; Stuble & Young, 2020).

Translating the concepts of community assembly and trajectories into relevant methods to drive ecosystem dynamics is a core approach in restoration ecology (Temperton et al., 2016). If an ecosystem has been degraded, damaged or destroyed, restoration managers try to assist its recovery; this can include the recovery of species composition and ecosystem functions (Gann et al., 2019). Changing initial biotic or abiotic conditions and species pools may help to accelerate restoration toward the reference plant community (Garrouj et al., 2019). Despite growing evidence of the importance of priority effects in early plant succession and ecological restoration (van der Putten et al., 2013; Vaughn & Young, 2015), their long-term consequences (Fukami, 2004; Švamberková et al., 2019; García-Girón et al., 2022) and their potential to improve target species establishment (Young et al., 2001, 2017;

Fukami & Nakajima, 2011; Werner et al., 2016), most restoration approaches use a simultaneous transfer of target species (Török et al., 2018).

Consequently, there is an increasing number of field and greenhouse studies testing priority effects as a restoration tool. Testing priority effects involves sequential sowing of different species or species groups in order to analyse their influence on plant community dynamics. Species groups may be based on their origin (native vs exotic, Goodale & Wilsey, 2018; Grman & Suding, 2010; Hess et al., 2022; Ploughe et al., 2020; Yu et al., 2020), on their life cycle type (annual vs perennial, Schantz et al., 2015; Vaughn & Young, 2015), or on their functional group (grasses vs legumes vs non-legume forbs; Delory et al., 2019; Stuble & Souza, 2016; Stuble et al., 2017; von Gillhaussen et al., 2014; Weidlich et al., 2017, 2018). Other studies examined priority effects by mixing these groups (Cleland et al., 2015; Lang et al., 2017; Mason et al., 2013; Schantz et al., 2018; Stuble & Young, 2020; Werner et al., 2016; Wilsey, 2020; Young et al., 2017). Alternatively, species may be assigned to groups according to their importance in reference plant communities, for example in distinguishing dominant (species frequently occurring in high abundance) and subordinate species (frequently occurring in low abundance; Grime, 1998). Dominant species are expected to play more important functional roles in communities due to their relative abundance, but subordinate species may also contribute significantly to community diversity and represent impactful ecosystem functions, like N fixation (Mariotte et al., 2013; Mariotte, 2014), particularly in semi-natural grasslands (Gibson, 2009; Werner et al., 2016). For example, subordinate species can increase plant community resistance against drought (Mariotte et al., 2013), probably via mycorrhizal fungal associations that improve plant productivity and positive plant–soil feedback (de Vries et al., 2012, 2018; Mariotte et al., 2012). Contrary to dominant species, subordinate species are expected to show, on average, a low competitive response (tolerance to competition by other species) and lower competitive effect (competition pressure exerted on other species; Goldberg & Landa, 1991; Keddy et al., 2002; Wang et al., 2010). Such competitive properties may explain why subordinate species may be more sensitive to priority effects than dominant species (Sarneel et al., 2016). Delaying the arrival of dominant species may thus improve the establishment of less abundant subordinate species (Mariotte et al., 2012; Young et al., 2017). Historically contingent coexistence has recently been described as possible assembly dynamics that allows coexistence: the coexistence of species A and B is only possible if B is introduced after A in the community whereas only B persists if A and B are introduced together or if A is introduced after B (Song et al., 2021). These assembly dynamics have not only been described theoretically but also experimentally in spider mite communities: the coexistence of the two species was

only possible when the weakest species was introduced first, otherwise only the strongest survived (Fragata et al., 2022). Our objective was to test sequential sowing of dominant and subordinate grassland species and its effect on the restored community assembly.

Plant dominance (relative abundance) in communities may be determined by plant cover or frequency in vegetation surveys. We first identified sets of dominant or subordinate species in our study area using abundances in vegetation surveys of undisturbed semi-natural grasslands. Then we tested priority effects by sowing dominants and subordinates both at the same time and sequentially. Temporal changes in dominance are supposed to occur through autogenic processes involving competitive interactions between plants, depending on life history and adaptive strategies (Grime, 1973; Huston & Smith, 1987). We therefore hypothesised that the order of arrival of the dominants and subordinates changes the species composition of the plant community. More specifically, as dominants have a strong competitive ability (both competitive effect and response), we expected that sowing dominants first or together with subordinates would lead to reduction in cover, or competitive exclusion of subordinates. In contrast, sowing subordinates first should favour their establishment without preventing the establishment of dominants (Figure 1). We further expect that the sowing treatment affects community assembly and spontaneous species establishment.

## 2 | MATERIALS AND METHODS

### 2.1 | Site description

The experiment was carried out in a degraded montane grassland at La-Bâtie-Neuve, in the upper Durance valley ('Haute-Durance') of the Southern French Alps (44°57'93" N, 6°20'77" E). The site is located at 1270 metres above sea level (a.s.l.), on an approximately 20° west-facing slope. The average annual precipitation is 947 mm. July is the warmest month with an estimated average temperature of 15.5°C, and January is the coldest with an average temperature of -3°C (estimated from Chorges meteorological station, 863 m a.s.l., 10 km from La-Bâtie-Neuve). Frost can occur from September to May. The soil is stony on calcareous bedrock. The grasslands in this area are extensively grazed by cattle. The recent construction of a high-voltage transmission line in the valley involved a degradation of the soil structure (and vegetation) affected by stripping, compacting and decompacting the soil to create transitory access tracks and construction platforms. At the end of the work, the stockpiled soil was moved back to recreate the original slope.

### 2.2 | Experimental setup

To test the effect of the order of species arrival on the community assembly, we applied sequential sowing using two groups of perennial plant species according to their frequency of occurrence in 54 plots of typical undisturbed grasslands of the 'Haute-Durance' valley

(Durbecq et al., 2020). Both groups included contrasting functional types represented by three plant families: Poaceae, Plantaginaceae and Fabaceae. The first group includes the 'dominant species' of each family, that is the most abundant species family in the studied grasslands. The second group represents the 'subordinate species', that is, the less abundant species in each of the studied families (Table 1). Dominant species were thus defined within plant families and may thus be less abundant than subordinates of another family. All these species are common in grasslands belonging to the habitat type "Semi-natural dry grasslands of *Festuco-Brometalia* and scrubland facies on calcareous substrates" (N6210, EU directive habitat 92/43/EEC, Calaciura & Spinelli, 2008).

### 2.3 | Experimental design

The experiment included four treatments on bare soil:


- (i) Control (without any sowing: it allowed us to estimate the importance of spontaneous colonisation)
- (ii) S+D: synchronous sowing (the six species were sown together in 2018 = year 0)
- (iii) D1<sup>st</sup>: dominant species (D) were sown in year 0 and subordinate species (S) in 2019 = first year
- (iv) S1<sup>st</sup>: subordinate species were sown in year 0 and dominant species in first year.

Each treatment was replicated in ten spatially randomised plots (total  $n=40$ ; Appendix S1). The 2018 sowing date was 19<sup>th</sup> October, and the 2019 sowing date was 9<sup>th</sup> October. Autumn sowing was chosen to allow dormancy breaking by cold stratification. Seeds of local origin according to the French label 'Végétal local' (Malaval et al., 2015) were obtained from a local seed company (Phytosem, Gap, France). The number of sown seeds was adjusted to a potential density of 100 individuals/m<sup>2</sup> for each sown species, based on seed producer expertise (Table 1). Plot size was 2 m × 2 m, separated by unsown margins of 50 cm. To prepare the seed bed, the plots were manually raked before and after the first sowing in year 0 to favour seedling recruitment. The experimental site was fenced to prevent seedling damage by cattle grazing. The plots were neither watered nor fertilised.

The dominant Plantaginaceae *Plantago media* did not establish in any treatment of the first year and was also absent the next three years. We therefore removed this species from the analyses.

### 2.4 | Data collection

Plant community surveys were carried out on all plots. The percentage cover of all vascular species was visually estimated in subplots of 1 m × 1 m placed in the centre of each experimental plot ( $n=40$ ; Appendix S1) in June 2019, 2020 and 2021 (first, second and third years; Appendix S2).



**FIGURE 1** Hypothesised patterns of priority effects. Plant interactions depending on which species arrives first (1<sup>st</sup>), dominants (D), subordinates (S) or when they are sown synchronously (S+D), resulting in (a) competitive response and (b) competitive effect (Goldberg & Landa, 1991). (c) Hypothesised response of species interactions when subordinates are sown first (S1<sup>st</sup>), when subordinates and dominants are sown synchronously (S+D) and when dominants are sown first (D1<sup>st</sup>). (d) Assembly dynamics showing three types of potential assembly dynamics in our study using the graph-based approach suggested by Song et al. (2021). Dominants are shown in red and subordinates in purple. Competitive response (proportion of focal species cover when grown with other species compared to monoculture) and effect (i.e., proportion of other species cover when grown with the focal species compared to monoculture) are displayed according to the hypotheses drawn for each species group (dominant or subordinate). Hypothesis 1: subordinates show a weak competitive effect and therefore do not or hardly affect other species regardless of arrival date (a, purple line). Hypothesis 2: subordinates show a weak competitive response and therefore establish less well in pre-established communities (assuming that pre-established species are already well developed and therefore exert higher resource pre-emption), but resist when sown first (b, purple line; assuming that they are already well developed and able to withstand resource pre-emption). Hypothesis 3: dominant species show a strong competitive effect and therefore strongly affect other species when arriving first, but to a lesser degree when arriving later (a, red line; because at least in a first phase they are less developed than the early-arriving species). Hypothesis 4: dominant species show a highly competitive response and are therefore not affected by other species, even if sown later (b, red line).

## 2.5 | Statistical analyses

In order to compare the plant community composition of the four treatments, Non-Metric Multidimensional Scaling (NMDS) was performed based on Bray–Curtis distances (Borcard et al., 2011) using R package *vegan* (Oksanen et al., 2020). A permutation multivariate analysis of variance (PERMANOVA, Anderson, 2017) with 9,999 permutations was used to analyse whether the community composition was significantly different between treatments (R package *vegan*).

To explore the effect of sequential sowing on community assembly, linear models (LM) were fitted with sowing treatment (D+S, D1<sup>st</sup>, S1<sup>st</sup>) as fixed effect and total plant species cover (i.e., community) as response variable. We tested the significance of the LM

by an *F*-test using the ‘Anova’ function of the R package *car* (Fox & Weisberg, 2019). As unsown species cover did not comply with the assumptions of LM (normal distribution, homoscedasticity), we ran generalised linear models (GLM) using a quasibinomial distribution to reduce overdispersion. In case of a significant treatment effect, multiple least-square mean comparisons were run using a Tukey adjustment in order to test differences between treatment levels (*emmeans* package, Lenth, 2022).

We further tested for each sown species the difference between being sown one year before other species and being sown synchronously. The proportion of individual species cover to total plant cover was used as response variable. Relative cover was used to correct for great differences in total cover that may bias dominance evaluation. Since data did not comply with the assumptions

**TABLE 1** Characteristics of the dominant and subordinate species sown

| TAXA                         | Composition group | Frequency of occurrence | Average cover (%) | CSR strategy | No. of seeds/m <sup>2</sup> |
|------------------------------|-------------------|-------------------------|-------------------|--------------|-----------------------------|
| POACEAE                      |                   |                         |                   |              |                             |
| <i>Bromopsis erecta</i>      | Dominant (D)      | 0.93                    | 32.13             | S/SC         | 555                         |
| <i>Festuca cinerea</i>       | Subordinate (S)   | 0.26                    | 21.51             | S/SR         | 2000                        |
| PLANTAGINACEAE               |                   |                         |                   |              |                             |
| <i>Plantago media</i>        | Dominant (D)      | 0.68                    | 5.42              | C/CR         | 1250                        |
| <i>Plantago lanceolata</i>   | Subordinate (S)   | 0.40                    | 1.22              | C/SC         | 533                         |
| FABACEAE                     |                   |                         |                   |              |                             |
| <i>Anthyllis vulneraria</i>  | Dominant (D)      | 0.28                    | 3.07              | CS/CSR       | 400                         |
| <i>Onobrychis viciifolia</i> | Subordinate (S)   | 0.14                    | 1.04              | CS/CSR       | 710                         |

Note: Competitive (C), stress-tolerant (S) and ruderal (R) strategies according to Erschbamer (2007), Grime (1977), and Pierce et al. (2013). Number of sown seeds per m<sup>2</sup> adjusted according to seed producer expertise on field germination (Phytosem SAS, Gap, France). Frequency of occurrence and average cover according to Durbecq et al. (2020) (n = 18).

of LM, differences between the three sequential sowing treatments were analysed using GLM with quasibinomial distribution to reduce overdispersion.

We used two indices to assess historical contingency processes. The priority index (PI – equation 1) measures priority effects sensu stricto (effect of early species on late-arriving species) and the earliness index (EI – equation 2), which indicates the benefit of arriving early compared to synchronous arrival with other species. Similar to the additive neighbour effect intensity index (Díaz-Sierra et al., 2017), these indices are standardised, symmetric (additive symmetry), and bounded between -1 (competitive exclusion of late-arriving species [PI] or by species sown synchronously [EI]) and +2 (obligate facilitation by early-arriving species [PI] or by species sown synchronously [EI]; Díaz-Sierra et al., 2017; Delory et al., 2019). The combination of both indices provides information on the symmetry of biotic interactions according to sowing time as suggested by Cleland et al. (2015), where PI is very close to their *secondary effect* and EI to their *priority effect*.

$$PI_{ni} = 2 \frac{Y_{ni}^{Late} - \overline{Y_n^{Sync}}}{Y_n^{Sync} + |Y_{ni}^{Late} - \overline{Y_n^{Sync}}|} \quad (1)$$

$$EI_{ni} = 2 \frac{Y_{ni}^{Early} - \overline{Y_n^{Sync}}}{Y_n^{Sync} + |Y_{ni}^{Early} - \overline{Y_n^{Sync}}|} \quad (2)$$

$Y_{ni}^{Early}$  and  $Y_{ni}^{Late}$  are the cover of a species (Figure 5c,d), or a group of species (Figure 5a,b), sown early and late in plot *i*, respectively, *n* years after being sown.  $\overline{Y_n^{Sync}}$  is the average cover of that same species, or group of species when sown synchronously. As the time between sowing events is one year, we used plant cover of the same age and not necessarily that of the same year. This may result in a bias if environmental factors such as meteorological conditions differ between years. This bias is, however, lower than using plant cover of the same year but different age.

We tested whether the estimated means were significantly different from zero by graphically analysing 95% confidence intervals. We then analysed for each index the differences between the species and between the two groups of species (dominants vs subordinates) using GLM for PI with Gamma distribution (log-link function) and LM for EI. All analyses were performed using R 3.5.3 (R Core Team, 2021).


### 3 | RESULTS

#### 3.1 | Community assemblages

The NMDS ordination showed a clear separation of the community composition between the four treatments (df = 3;  $F = 9.321$ ;  $p < 0.001$ , Figure 2), even after three years. Along the first axis of the NMDS, the three sequential sowing treatments were characterised by the sown species on the far right, while ruderal species, such as *Lactuca serriola* L., *Taraxacum* sp., *Medicago lupulina* L. and some mesophilous grassland species, such as *Coronilla minima* L., *Lotus corniculatus* L., *Poterium sanguisorba* L. and *Hippocrepis comosa* L. dominated the control on the left. The second axis separated the three sowing treatments, placing the synchronous sowing (D+S) in the middle of the two sequential sowing treatments with S1<sup>st</sup> on the upper and D1<sup>st</sup> on the lower part of this axis.

#### 3.2 | Effect of sequential sowing on vegetation cover

In 2021, three years after the first sowing, the three seed addition treatments increased total plant cover and limited the development of species emerging from the soil seed bank or seed rain (Appendix S2, Appendix S3). Total cover contribution of unsown




**FIGURE 2** Plant species composition according to different sowing sequences using NMDS (NMDS stress: 0.24). Green, orange, blue and grey colour indicates D1<sup>st</sup>, S1<sup>st</sup>, Synchronous and Control sequences respectively. Arrows indicate changes in the position of the plot barycentre according to year (2019, 2020 and 2021). The third-year plots are shown as full dots and polygon group for each treatment. Only the 14 species of which cover is most correlated with plot position are shown and sown species are in bold.

species was 30% in the control and was much lower in D+S, D1<sup>st</sup> and S1<sup>st</sup> treatments (8%; **Figure 3**). The two Poaceae (both the relatively dominant and the relatively subordinate) and the subordinate *Onobrychis viciifolia* dominated the synchronous sowing (D+S). In the D1<sup>st</sup> treatment, the dominant *Bromopsis erecta* represented more than half of the total community cover whereas the cover of both the dominant *Anthyllis vulneraria* and the three subordinates was much smaller (<5%). As expected, subordinates best developed when sown first (S1<sup>st</sup>). In particular, the covers of the subordinate species *Plantago lanceolata* and *Onobrychis viciifolia* were considerably higher in the S1<sup>st</sup> than in other treatments.

In the third year, the total cover of the dominant Poaceae *Bromopsis erecta* was significantly higher in both D+S and D1<sup>st</sup> treatments with an average cover of 34%, compared to its cover in the S1<sup>st</sup> treatment being approximately 5% (**Figure 4**, Appendix S4). There was no significant difference in *Bromopsis erecta* cover between D+S and D1<sup>st</sup> treatments. No significant treatment effect was found for the dominant Fabaceae *Anthyllis vulneraria*. The three subordinate species (*Festuca cinerea*, *Onobrychis viciifolia* and *Plantago lanceolata*) performed significantly better in S1<sup>st</sup> than in D1<sup>st</sup>. However, the cover of the subordinate Poaceae *Festuca cinerea* was not significantly different between S1<sup>st</sup> and D+S. Additionally, the cover of *Onobrychis viciifolia* and *Plantago lanceolata* was significantly lower in D+S than in S1<sup>st</sup> (when first sown).


### 3.3 | Priority and earliness indices

The priority index (PI) and the earliness index (EI) showed that the establishment of both dominants and subordinates was affected by



**FIGURE 3** Species cover contribution to the total cover in the four treatments, in the third year. D, dominant species; S, subordinate species. D+S, dominant and subordinate species sown synchronously; D1<sup>st</sup>, dominant species sown first (subordinates second); S1<sup>st</sup>, subordinate species sown first (dominants second) and Control (no sowing). *Plantago media* did not germinate.

sequential sowing (**Figure 5**). Dominant and subordinate groups significantly differed in their EI ( $df = 1$ ,  $F = 16.16$ ,  $p < 0.001$ ) but not in their PI ( $df = 1$ ,  $F = 0.18$ ,  $p = 0.677$ ). PI was negative for both groups of species (**Figure 5a**) indicating that the early sown species induced negative effects on later sown species. A PI significantly different from zero would show that both the dominant *Bromopsis erecta* and the subordinates *Onobrychis viciifolia* and *Plantago lanceolata* were negatively affected when arriving late, but PI did not significantly differ between species (**Figure 5c**). Accordingly, over all dominants



**FIGURE 4** Sown species cover in the third year in the three sequential sowing treatments with: D + S, dominant and subordinate species sown synchronously; D1<sup>st</sup>, dominant species sown first (subordinates second); and S1<sup>st</sup>, subordinate species sown first (dominants second). Error bars are  $\pm$ SE and a common letter indicates the absence of significant differences between treatments. Details in Appendix S1.

and subordinates, the PI was significantly negative but there was no significant difference between the two groups (Figure 5a). The EI of the subordinates, in particular of *Festuca cinerea* and *Plantago lanceolata*, was significantly different from zero whereas the EI of the dominants was not (Figure 5b,d,  $df=4$ ,  $F=5.10$ ,  $p<0.01$ ). The EI was significantly higher in subordinates than in dominants indicating that they benefited more from being sown first.

## 4 | DISCUSSION


We tested priority effects applied in grassland restoration using synchronous and sequential sowing with a one-year time difference. Manipulating the order of arrival influenced the community composition and trajectories three years after the first sowing. Both dominant and subordinate species showed a reduced cover when sown secondly compared to being sown synchronously. Being sown first, however, increased cover compared to being sown synchronously only for subordinates. The two Poaceae showed the same treatment response: the dominant *Bromopsis erecta* and its relative subordinate *Festuca cinerea* better established when sown first compared with both other treatments (sown later or synchronously).

We hypothesised that sowing subordinate species first not decrease the establishment of the dominant species, but we showed the opposite. The most abundant dominant species *Bromopsis erecta* is known to be a stress-tolerant competitor (Grime, 1979) and showed a significant disadvantage when sown second. This result suggests that *Bromopsis erecta* was affected by direct competition or indirect effects of already established vegetation. The competitive response of *Bromopsis erecta* depended on the time of arrival in a community (Cleland et al., 2015). The relative susceptibility of *Bromopsis erecta* to competition has already been highlighted under different environmental constraints (Corcket et al., 2003; Liancourt et al., 2005).

The abundance of this dominant species in dry grasslands is also related to stress tolerance (Corcket et al., 2003; Liancourt et al., 2005). The second hypothesis, that subordinate species are favoured by being sown first, was partly supported by our results. *Onobrychis viciifolia* and *Plantago lanceolata* established well only when sown first. However, the subordinate *Festuca cinerea* was not favoured by sowing first compared to synchronous sowing. These priority effects may have been due to the stronger competitive effect of the dominants, in particular of *Bromopsis erecta* (hypothesis 3). Similar patterns were found in another study on *Onobrychis viciifolia*, showing a particularly weak competitiveness when sown after exotic species (Hess et al., 2020). Additionally, a priority effect experiment in riparian communities showed that subordinate species were more affected by the strength of priority effects (Sarneck et al., 2016). Furthermore, priority effects both favoured subordinate species when sown first, as shown by their highest EI, and hampered the growth of the most dominant species, *Bromopsis erecta*, allowing a higher level of species coexistence in the community. In this paper we differentiated dominant and subordinate species by their relative abundances within each functional group, not overall. For instance, *Festuca cinerea* has a lower average cover than *Bromopsis erecta* but also has a higher average cover than the other dominants (*Plantago media* and *Anthyllis vulneraria*). This may explain why its cover is not different when sown synchronously or when it is sown first. In the same way, we expected the communities resulting from sowing simultaneously and dominants first to be more similar if the 'dominants' were competitively dominant across all species.

The different response of species to the order of arrival may be explained by the niche components hypothesis (Vannette & Fukami, 2014), which states that the strength of priority effects depends on niche components (overlap, impact and requirements). Niche requirements differ between species and may influence (negatively or positively) their recruitment and/or establishment when





**FIGURE 5** (a) Priority Index (PI) and (b) Earliness Index (EI) calculated for the cover of dominants (*Bromopsis erecta*+*Anthyllis vulneraria*) and subordinates (*Festuca cinerea*+*Onobrychis viciifolia*+*Plantago lanceolata*) after two years. (c) PI and (d) EI calculated for the sown species cover in 2021. For each species, values are estimated marginal means  $\pm$ 95% confidence intervals ( $n=10$ ). White dots indicate that confidence intervals included zero, black dots indicate that there was no overlap between confidence intervals and zero. A letter in common between two bars indicates the absence of significant differences between dominant and subordinate groups, or between two species ( $p < 0.05$ ).

arriving late. Our results showed a lower performance of late-arriving species that were probably hampered by niche modification and size-asymmetric competition of the pre-established vegetation (Vannette & Fukami, 2014; Wilsey, 2020), or by niche pre-emption (Fukami, 2015). Hypotheses three and four were partly validated since the dominant *Bromopsis erecta* had a negative effect on the development of other sown species. The cover of the dominant Fabaceae *Anthyllis vulneraria* was not affected by the sowing treatments or was potentially negatively affected by being sown together with *Bromopsis erecta*, and was thus not higher when the species

was sown first. *Anthyllis vulneraria* is a dispersal-limited, late-successional but fast-growing species that may benefit from being sown in a pre-established plant community (Erschbamer, 2007; Marcante et al., 2009). Furthermore, its roots are longer and deeper than those of the sown subordinate Fabaceae *Onobrychis viciifolia* and of the other sown species (Jungk, 1993), thus avoiding competition for soil resources. In our study, *Anthyllis vulneraria* was not negatively affected by the pre-established community when sown after subordinate species (PI close to zero), which confirms a high recruitment capacity in established vegetation (Vannette & Fukami, 2014),

unless it was affected by *Bromopsis erecta*. Positive effects of already established vegetation on the recruitment niche (e.g., living cover protecting young seedlings and improving water retention; Donath et al., 2006; Durbecq et al., 2022; Wright et al., 2014) may have favoured the dominant *Anthyllis vulneraria* when sown second or synchronously. Alternatively, plant–soil feedback such as interactions with rhizobia may have contributed to the priority effects (de Vries et al., 2012, 2018; Fry et al., 2017; Fukami, 2015; Fukami & Nakajima, 2013; Goldstein & Suding, 2014). The mechanisms of underlying priority effects need to be better understood, with a particular focus on plant–soil feedback and below-ground interactions (Bever, 2003; Fry et al., 2017; Fukami & Nakajima, 2013; Hess et al., 2020; Weidlich et al., 2018). Plant–soil interactions are species-specific (Bezemer et al., 2006; van der Voorde et al., 2011) and the strength of priority effects depends on both the order of arrival and species identity (Cleland et al., 2015; Stuble & Souza, 2016; von Gillhausen et al., 2014; Wilsey et al., 2015; Werner et al., 2016). Therefore, an advance of a particular species of one year may change soil biota and nutrient status, as well as productivity by niche modification (Vannette & Fukami, 2014).

In conclusion, regardless of underlying mechanisms, our study provides evidence that plant community assembly was influenced by the order of arrival, potentially changing community composition and trajectories. In several cases, coexistence may need sequential arrival in a community (Fragata et al., 2022; Song et al., 2021). Thus, the order of species arrival may be a useful tool to optimise seed-sowing approaches in ecological restoration. Our results were in agreement with those of other studies indicating strong variations between species and thus highly species-specific priority effects (Cleland et al., 2015; Stuble & Souza, 2016). The next step in the use of priority effects in restoration ecology would be to obtain a better understanding of the predictability of priority effects (Song et al., 2021) and to identify the need for sequential sowing which may differ depending on species and environmental characteristics. Our distinction between relatively (within each family) dominant and subordinate species was a first attempt to categorise species that would need to be sown sequentially and those that are competitive in simultaneous sowing. A more specific categorisation based on functional traits of species, especially traits related to germination and early-stage survival and growth may provide more results (Torrez et al., 2017). To be applied in ecological restoration, further studies are needed to assess priority effects of target species in different habitat types with different plant communities. Furthermore, a higher number of species of different functional groups needs to be tested to evaluate a potential application in ecological restoration. Such knowledge would allow practitioners to establish lists of species that need to be sown first, second or synchronously in order to develop the plant community structure toward identified references.

#### AUTHOR CONTRIBUTIONS

Aure Durbecq, Armin Bischoff, Elise Buisson and Renaud Jaunatre conceived the project and conducted the field surveys; Aure

Durbecq and Renaud Jaunatre analysed the data and drafted the manuscript. All authors (Aure Durbecq, Renaud Jaunatre, Armin Bischoff, Elise Buisson and Emmanuel Corcket) improved the text and gave final approval for publication.

#### ACKNOWLEDGEMENTS

We thank Alice Dupré-La-Tour, Daniel Pavon, Julien Planche, Margaux d'Ambly, Léo Rocher, Chloé Malik, Gaëlle Carrière, and François-Marie Martin for their assistance. We also thank Benjamin Delory for the discussion on priority effects and the three reviewers for their suggestions.

#### FUNDING INFORMATION

Financial support was provided by the environmental consultancy ECO-MED (Ecologie & Médiation), RTE (Electricity Transmission Network) and ANRT (National Agency of Research and Technology via the grant CIFRE N°2017/0478).

#### CONFLICT OF INTEREST STATEMENT

The authors declare that they have no conflict of interests.

#### DATA AVAILABILITY STATEMENT

Code and data are publicly available at [https://forgemia.inra.fr/renaud.jaunatre/durance\\_pe](https://forgemia.inra.fr/renaud.jaunatre/durance_pe).

#### ORCID

Aure Durbecq  <https://orcid.org/0000-0002-9201-1916>

Armin Bischoff  <https://orcid.org/0000-0003-2865-8720>

Elise Buisson  <https://orcid.org/0000-0002-3640-8134>

Emmanuel Corcket  <https://orcid.org/0000-0002-8586-2202>

Renaud Jaunatre  <https://orcid.org/0000-0001-6970-8304>

#### REFERENCES

- Anderson, M.J. (2017) Permutational multivariate analysis of variance (PERMANOVA). In: Balakrishnan, N., Colton, T., Everitt, B., Piegorsch, W., Ruggeri, F. & Teugels, J.L. (Eds.) *Wiley StatsRef: statistics reference online*. Auckland, New Zealand: Massey University. Available from: <https://doi.org/10.1002/9781118445112.stat07841>
- Bertness, M.D. & Shumway, S.W. (1993) Competition and facilitation in marsh plants. *The American Naturalist*, 142, 718–724. Available from: <https://doi.org/10.1086/285567>
- Bever, J.D. (2003) Soil community feedback and the coexistence of competitors: conceptual frameworks and empirical tests. *New Phytologist*, 157, 465–473. Available from: <https://doi.org/10.1046/j.1469-8137.2003.00714.x>
- Bezemer, T.M., Lawson, C.S., Hedlund, K., Edwards, A.R., Brook, A.J., Igual, J.M., et al. (2006) Plant species and functional group effects on abiotic and microbial soil properties and plant–soil feedback responses in two grasslands. *Journal of Ecology*, 94, 893–904. Available from: <https://doi.org/10.1111/j.1365-2745.2006.01158.x>
- Borcard, D., Gillet, F. & Legendre, P. (2011) *Numerical ecology with R*. New York: Springer Science & Business Media.
- Calaciura, B. & Spinelli, O. (2008) Management of Natura 2000 habitats. 6210 semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (\*important orchid sites). Technical report 2008 12/24, European Commission.

- Chase, J.M. (2003) Community assembly: when should history matter? *Oecologia*, 136, 489–498. Available from: <https://doi.org/10.1007/s00442-003-1311-7>
- Chase, J.M. (2010) Stochastic community assembly causes higher biodiversity in more productive environments. *Science*, 328, 1388–1391. Available from: <https://doi.org/10.1126/science.1187820>
- Chase, J.M., Biro, E.G., Ryberg, W.A. & Smith, K.G. (2009) Predators temper the relative importance of stochastic processes in the assembly of prey metacommunities. *Ecology Letters*, 12, 1210–1218. Available from: <https://doi.org/10.1111/j.1461-0248.2009.01362.x>
- Cleland, E.E., Esch, E. & McKinney, J. (2015) Priority effects vary with species identity and origin in an experiment varying the timing of seed arrival. *Oikos*, 124, 33–40. Available from: <https://doi.org/10.1111/oik.01433>
- Cole, B.J. (1983) Assembly of mangrove ant communities: patterns of geographical distribution. *Journal of Animal Ecology*, 52(2), 339–347. Available from: <https://doi.org/10.2307/4557>
- Corcket, E., Liancourt, P., Callaway, R. & Michalet, R. (2003) The relative importance of competition for two dominant grass species as affected by environmental manipulations in the field. *Écoscience*, 10(2), 186–194. Available from: <https://doi.org/10.1080/11956860.2003.11682766>
- de Vries, F.T., Griffiths, R.I., Bailey, M., Craig, H., Girlanda, M., Gweon, H.S. et al. (2018) Soil bacterial networks are less stable under drought than fungal networks. *Nature Communications*, 9, 3033. Available from: <https://doi.org/10.1038/s41467-018-05516-7>
- de Vries, F.T., Liiri, M.E., Bjørnlund, L., Bowker, M.A., Christensen, S., Setälä, H.M. et al. (2012) Land use alters the resistance and resilience of soil food webs to drought. *Nature Climate Change*, 2, 276–280. Available from: <https://doi.org/10.1038/nclimate1368>
- Delory, B.M., Weidlich, E.W.A., von Gillhausen, P. & Temperton, V.M. (2019) When history matters: the overlooked role of priority effects in grassland overyielding. *Functional Ecology*, 33, 2369–2380. Available from: <https://doi.org/10.1111/1365-2435.13455>
- Díaz-Sierra, R., Verwijmeren, M., Rietkerk, M., de Dios, V.R. & Baudena, M. (2017) A new family of standardized and symmetric indices for measuring the intensity and importance of plant neighbour effects. *Methods in Ecology and Evolution*, 8, 580–591. Available from: <https://doi.org/10.1111/2041-210X.12706>
- Donath, T.W., Hölzel, N. & Otte, A. (2006) Influence of competition by sown grass, disturbance and litter on recruitment of rare flood-meadow species. *Biological Conservation*, 130, 315–323. Available from: <https://doi.org/10.1016/j.biocon.2005.12.022>
- Durbecq, A., Jaunatre, R., Buisson, E., Cluchier, A. & Bischoff, A. (2020) Identifying reference communities in ecological restoration: the use of environmental conditions driving vegetation composition. *Restoration Ecology*, 28, 1445–1453. Available from: <https://doi.org/10.1111/rec.13232>
- Durbecq, A., Rocher, L., Jaunatre, R., la Tour, A.D., Buisson, E. & Bischoff, A. (2022) Mountain grassland restoration using hay and brush material transfer combined with temporary wheat cover. *Ecological Engineering*, 174, 106447. Available from: <https://doi.org/10.1016/j.ecoleng.2021.106447>
- Erschbamer, B. (2007) Winners and losers of climate change in a central alpine glacier. *Foreland. Arctic, Antarctic, and Alpine Research*, 39(2), 237–244. Available from: [https://doi.org/10.1657/1523-0430\(2007\)39\[237:WALOC\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2007)39[237:WALOC]2.0.CO;2)
- Fox, J. & Weisberg, S. (2019) *An R companion to applied regression*, 3rd edition. Thousand Oaks, CA: Sage.
- Fragata, I., Costa-Pereira, R., Kozak, M., Majer, A., Godoy, O. & Magalhães, S. (2022) Specific sequence of arrival promotes coexistence via spatial niche pre-emption by the weak competitor. *Ecology Letters*, 25, 1629–1639. Available from: <https://doi.org/10.1111/ele.14021>
- Fry, E.L., Pilgrim, E.S., Tallowin, J.R.B., Smith, R.S., Mortimer, S.R., Beaumont, D.A. et al. (2017) Plant, soil and microbial controls on grassland diversity restoration: a long-term, multi-site mesocosm experiment. *Journal of Applied Ecology*, 54, 1320–1330. Available from: <https://doi.org/10.1111/1365-2664.12869>
- Fukami, T. (2004) Assembly history interacts with ecosystem size to influence species diversity. *Ecology*, 85, 3234–3242. Available from: <https://doi.org/10.1890/04-0340>
- Fukami, T. (2015) Historical contingency in community assembly: integrating niches, species pools, and priority effects. *Annual Review of Ecology, Evolution, and Systematics*, 46, 1–23. Available from: <https://doi.org/10.1146/annurev-ecolsys-110411-160340>
- Fukami, T. & Nakajima, M. (2011) Community assembly: alternative stable states or alternative transient states? *Ecology Letters*, 14, 973–984. Available from: <https://doi.org/10.1111/j.1461-0248.2011.01663.x>
- Fukami, T. & Nakajima, M. (2013) Complex plant–soil interactions enhance plant species diversity by delaying community convergence. *Journal of Ecology*, 101, 316–324. Available from: <https://doi.org/10.1111/1365-2745.12048>
- Gann, G.D., McDonald, T., Walder, B., Aronson, J., Nelson, C.R., Jonson, J. et al. (2019) International principles and standards for the practice of ecological restoration. Second edition. *Restoration Ecology*, 27, S1–S46. Available from: <https://doi.org/10.1111/rec.13035>
- García-Girón, J., Lindholm, M., Heino, J., Toivonen, H. & Alahuhta, J. (2022) Historical contingency via priority effects counteracts environmental change on metacommunity dynamics across decades. *Limnology and Oceanography*, 67, S38–S53. Available from: <https://doi.org/10.1002/lno.11699>
- Garrouj, M., Alard, D., Corcket, E., Marchand, L. & Benot, M.-L. (2019) The effects of management on vegetation trajectories during the early-stage restoration of previously arable land after hay transfer. *Ecology and Evolution*, 9, 13776–13786. Available from: <https://doi.org/10.1002/ece3.5798>
- Gibson, D.J. (2009) *Grasses and grassland ecology*. New York: Oxford University Press.
- Goldberg, D.E. & Landa, K. (1991) Competitive effect and response: hierarchies and correlated traits in the early stages of competition. *Journal of Ecology*, 79, 1013–1030. Available from: <https://doi.org/10.2307/2261095>
- Goldstein, L.J. & Suding, K.N. (2014) Applying competition theory to invasion: resource impacts indicate invasion mechanisms in California shrublands. *Biological Invasions*, 16, 191–203. Available from: <https://doi.org/10.1007/s10530-013-0513-0>
- Goodale, K.M. & Wilsey, B.J. (2018) Priority effects are affected by precipitation variability and are stronger in exotic than native grassland species. *Plant Ecology*, 219, 429–439. Available from: <https://doi.org/10.1007/s11258-018-0806-6>
- Grime, J.P. (1973) Competitive exclusion in herbaceous vegetation. *Nature*, 242, 344–347. Available from: <https://doi.org/10.1038/242344a0>
- Grime, J.P. (1977) Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *The American Naturalist*, 111(982), 1169–1194. Available from: <https://doi.org/10.1086/283244>
- Grime, J.P. (1979) *Plant strategies and vegetation processes*. Chichester: John Wiley & Sons. Hammer EC. Available from: <https://agris.fao.org/agris-search/search.do?recordID=XF2016008771>
- Grime, J.P. (1998) Benefits of plant diversity to ecosystems: immediate, filter and founder effects. *Journal of Ecology*, 86, 902–910. Available from: <https://doi.org/10.1046/j.1365-2745.1998.00306.x>
- Grman, E. & Suding, K.N. (2010) Within-year soil legacies contribute to strong priority effects of exotics on native California grassland communities. *Restoration Ecology*, 18, 664–670. Available from: <https://doi.org/10.1111/j.1526-100X.2008.00497.x>
- Helsen, K., Hermy, M. & Honnay, O. (2016) A test of priority effect persistence in semi-natural grasslands through the removal of plant functional groups during community assembly. *BMC Ecology*, 16, 22. Available from: <https://doi.org/10.1186/s12898-016-0077-9>

- Hess, M.C.M., Buisson, E., Fontes, H., Bacon, L., Sabatier, F. & Mesléard, F. (2020) Giving recipient communities a greater head start and including productive species boosts early resistance to invasion. *Applied Vegetation Science*, 23(3), 340–352. Available from: <https://doi.org/10.1111/avsc.12502>
- Hess, M.C.M., Mesléard, F., Young, T., de Freitas, B., Haveneers, N. & Buisson, E. (2022) Ltering native community assembly history influences the performance of an annual invader. *Basic and Applied Ecology*, 59, 70–81. Available from: <https://doi.org/10.1016/j.baae.2022.01.004>
- Hobbs, R.J. & Norton, D.A. (2004) Ecological filters, thresholds, and gradients in resistance to ecosystem reassembly. In: *Assembly rules and restoration ecology: bridging the gap between theory and practice*. Washington, DC: Island Press, pp. 72–95.
- Huston, M. & Smith, T. (1987) Plant succession: life history and competition. *The American Naturalist*, 130(2), 168–198. Available from: <https://doi.org/10.1086/284704>
- Jungk, A. (1993) Kutschera, L., E. Lichtenegger, M. Sobotik: Wurzelatlas mitteleuropäischer Grünlandpflanzen. Band 2: Pteridophyta und Dicotyledoneae (Magnoliopsida). Teil 1: Von L. Kutschera und E. Lichtenegger: Morphologie, Anatomie, Ökologie, Verbreitung, Soziologie, Wirtschaft; mit 851 S., 364 Abb., 137 z. T. farb. Taf. Verlag Gustav Fischer 1992; Preis Ln. DM 390, 00. *Zeitschrift für Pflanzenernährung und Bodenkunde*, 156, 97. Available from: <https://doi.org/10.1002/jpln.19931560118>
- Kardol, P., Souza, L. & Classen, A.T. (2013) Resource availability mediates the importance of priority effects in plant community assembly and ecosystem function. *Oikos*, 122, 84–94. Available from: <https://doi.org/10.1111/j.1600-0706.2012.20546.x>
- Keddy, P., Nielsen, K., Weiher, E. & Lawson, R. (2002) Relative competitive performance of 63 species of terrestrial herbaceous plants. *Journal of Vegetation Science*, 13, 5–16. Available from: <https://doi.org/10.1111/j.1654-1103.2002.tb02018.x>
- Lang, M., Hanslin, H.M., Kollmann, J. & Wagner, T. (2017) Suppression of an invasive legume by a native grass – high impact of priority effects. *Basic and Applied Ecology*, 22, 20–27. Available from: <https://doi.org/10.1016/j.baae.2017.06.005>
- Lenth, R. (2022) Emmeans: estimated marginal means, aka least-squares means. R package version 1.7.2.
- Levine, J.M., Vilà, M., Antonio, C.M.D., Dukes, J.S., Grigulis, K. & Lavorel, S. (2003) Mechanisms underlying the impacts of exotic plant invasions. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270, 775–781. Available from: <https://doi.org/10.1098/rspb.2003.2327>
- Liancourt, P., Corcket, E. & Michalet, R. (2005) Stress tolerance abilities and competitive responses in a watering and fertilization field experiment. *Journal of Vegetation Science*, 16, 713–722. Available from: <https://doi.org/10.1111/j.1654-1103.2005.tb02414.x>
- Malaval, S., Bischoff, A., Hédont, M., Provendier, D., Boutaud, M., Dao, J. et al. (2015) Végétal local: une marque française pour la conservation de la flore indigène. In: Eurogard VII. European Botanic Gardens Congress, Paris, France. Available from: <https://hal-univ-avignon.archives-ouvertes.fr/hal-01778344/file/Malaval%20et%20al.%202018.pdf>
- Marcante, S., Winkler, E. & Erschbamer, B. (2009) Population dynamics along a primary succession gradient: do alpine species fit into demographic succession theory? *Annals of Botany*, 103(7), 1129–1143. Available from: <http://www.jstor.org/stable/43576197>
- Mariotte, P. (2014) Do subordinate species punch above their weight? Evidence from above- and below-ground. *New Phytologist*, 203, 16–21. Available from: <https://doi.org/10.1111/nph.12789>
- Mariotte, P., Buttler, A., Johnson, D., Thébault, A. & Vandenberghe, C. (2012) Exclusion of root competition increases competitive abilities of subordinate plant species through root–shoot interactions. *Journal of Vegetation Science*, 23, 1148–1158. Available from: <https://doi.org/10.1111/j.1654-1103.2012.01432.x>
- Mariotte, P., Vandenberghe, C., Kardol, P., Hagedorn, F. & Buttler, A. (2013) Subordinate plant species enhance community resistance against drought in semi-natural grasslands. *Journal of Ecology*, 101, 763–773. Available from: <https://doi.org/10.1111/1365-2745.12064>
- Mason, T.J., French, K. & Jolley, D. (2013) Arrival order among native plant functional groups does not affect invasibility of constructed dune communities. *Oecologia*, 173, 557–568. Available from: <https://doi.org/10.1007/s00442-013-2628-5>
- Oksanen, J., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., Minchin, P.R. et al. (2020) Vegan: community ecology package. R Package Version 2.5-7. Available from: <https://CRAN.R-project.org/package=vegan>
- Orrock, J.L. & Fletcher, R.J., Jr. (2005) Changes in community size affect the outcome of competition. *The American Naturalist*, 166, 107–111. Available from: <https://doi.org/10.1086/430641>
- Pierce, S., Brusa, G., Vagge, I. & Cerabolini, B.E.L. (2013) Allocating CSR plant functional types: the use of leaf economics and size traits to classify woody and herbaceous vascular plants. *Functional Ecology*, 27, 1002–1010. Available from: <https://doi.org/10.1111/1365-2435.12095>
- Ploughe, L.W., Carlyle, C.N. & Fraser, L.H. (2020) Priority effects: how the order of arrival of an invasive grass, *Bromus tectorum*, alters productivity and plant community structure when grown with native grass species. *Ecology and Evolution*, 10, 13173–13181. Available from: <https://doi.org/10.1002/ece3.6908>
- R Core Team. (2021) *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Available from: <https://www.R-project.org/>
- Sarneel, J.M., Kardol, P. & Nilsson, C. (2016) The importance of priority effects for riparian plant community dynamics. *Journal of Vegetation Science*, 27, 658–667. Available from: <https://doi.org/10.1111/jvs.12412>
- Schantz, M.C., Sheley, R.L. & James, J.J. (2015) Role of propagule pressure and priority effects on seedlings during invasion and restoration of shrub-steppe. *Biological Invasions*, 17, 73–85. Available from: <https://doi.org/10.1007/s10530-014-0705-2>
- Schantz, M.C., Sheley, R.L. & James, J.J. (2018) Effects of propagule pressure and priority effects on seedling recruitment during restoration of invaded grassland. *Journal of Arid Environments*, 150, 62–70. Available from: <https://doi.org/10.1016/j.jaridenv.2017.12.001>
- Song, C., Fukami, T. & Saavedra, S. (2021) Untangling the complexity of priority effects in multispecies communities. *Ecology Letters*, 24, 2301–2313. Available from: <https://doi.org/10.1111/ele.13870>
- Stuble, K.L., Fick, S.E. & Young, T.P. (2017) Every restoration is unique: testing year effects and site effects as drivers of initial restoration trajectories. *Journal of Applied Ecology*, 54, 1051–1057. Available from: <https://doi.org/10.1111/1365-2664.12861>
- Stuble, K.L. & Souza, L. (2016) Priority effects: natives, but not exotics, pay to arrive late. *Journal of Ecology*, 104, 987–993. Available from: <https://doi.org/10.1111/1365-2745.12583>
- Stuble, K.L. & Young, T.P. (2020) Priority treatment leaves grassland restoration vulnerable to invasion. *Diversity*, 12, 71. Available from: <https://doi.org/10.3390/d12020071>
- Švamberková, E., Doležal, J. & Lepš, J. (2019) The legacy of initial sowing after 20 years of ex-arable land colonisation. *Oecologia*, 190, 459–469. Available from: <https://doi.org/10.1007/s00442-019-04415-y>
- Temperton, V.M., Baasch, A., von Gillhausen, P. & Kirmer, A. (2016) Assembly theory for restoring ecosystem structure and functioning: timing is everything? In: Palmer, M.A., Zedler, J.B. & Falk, D.A. (Eds.) *Foundations of restoration ecology*. Washington, DC: Island Press/Center for Resource Economics, pp. 245–270. Available from: [https://doi.org/10.5822/978-1-61091-698-1\\_9](https://doi.org/10.5822/978-1-61091-698-1_9)
- Török, P., Helm, A., Kiehl, K., Buisson, E. & Valkó, O. (2018) Beyond the species pool: modification of species dispersal, establishment, and

- assembly by habitat restoration. *Restoration Ecology*, 26, S65–S72. Available from: <https://doi.org/10.1111/rec.12825>
- Torrez, V., Mergeay, J., Meester, L.D., Honnay, O. & Helsen, K. (2017) Differential effects of dominant and subordinate plant species on the establishment success of target species in a grassland restoration experiment. *Applied Vegetation Science*, 20, 363–375. Available from: <https://doi.org/10.1111/avsc.12302>
- van der Putten, W.H., Bardgett, R.D., Bever, J.D., Bezemer, T.M., Casper, B.B., Fukami, T. et al. (2013) Plant–soil feedbacks: the past, the present and future challenges. *Journal of Ecology*, 101, 265–276. Available from: <https://doi.org/10.1111/1365-2745.12054>
- Vannette, R.L. & Fukami, T. (2014) Historical contingency in species interactions: towards niche-based predictions. *Ecology Letters*, 17, 115–124. Available from: <https://doi.org/10.1111/ele.12204>
- Vaughn, K.J. & Young, T.P. (2015) Short-term priority over exotic annuals increases the initial density and longer-term cover of native perennial grasses. *Ecological Applications*, 25, 791–799. Available from: <https://doi.org/10.1890/14-0922.1>
- van de Voorde, T.F.J., van der Putten, W.H. & Martijn Bezemer, T. (2011) Intra- and interspecific plant–soil interactions, soil legacies and priority effects during old-field succession. *Journal of Ecology*, 99, 945–953. Available from: <https://doi.org/10.1111/j.1365-2745.2011.01815.x>
- von Gillhausen, P., Rascher, U., Jablonowski, N.D., Plückers, C., Beierkuhnlein, C. & Temperton, V.M. (2014) Priority effects of time of arrival of plant functional groups override sowing interval or density effects: a grassland experiment. *PLoS ONE*, 9, e86906. Available from: <https://doi.org/10.1371/journal.pone.0086906>
- Wainwright, C.E., Wolkovich, E.M. & Cleland, E.E. (2012) Seasonal priority effects: implications for invasion and restoration in a semi-arid system. *Journal of Applied Ecology*, 49, 234–241. Available from: <https://doi.org/10.1111/j.1365-2664.2011.02088.x>
- Wang, C., Long, R., Wang, Q., Liu, W., Jing, Z. & Zhang, L. (2010) Fertilization and litter effects on the functional group biomass, species diversity of plants, microbial biomass, and enzyme activity of two alpine meadow communities. *Plant and Soil*, 331, 377–389. Available from: <https://doi.org/10.1007/s11104-009-0259-8>
- Weidlich, E.W.A., von Gillhausen, P., Delory, B.M., Blossfeld, S., Poorter, H. & Temperton, V.M. (2017) The importance of being first: exploring priority and diversity effects in a grassland field experiment. *Frontiers in Plant Science*, 7, 2008. Available from: <https://doi.org/10.3389/fpls.2016.02008>
- Weidlich, E.W.A., von Gillhausen, P., Max, J.F.J., Delory, B.M., Jablonowski, N.D., Rascher, U. et al. (2018) Priority effects caused by plant order of arrival affect below-ground productivity. *Journal of Ecology*, 106, 774–780. Available from: <https://doi.org/10.1111/1365-2745.12829>
- Werner, C.M., Vaughn, K.J., Stuble, K.L., Wolf, K. & Young, T.P. (2016) Persistent asymmetrical priority effects in a California grassland restoration experiment. *Ecological Applications*, 26, 1624–1632. Available from: <https://doi.org/10.1890/15-1918.1>
- Wilsey, B. (2020) Restoration in the face of changing climate: importance of persistence, priority effects, and species diversity. *Restoration Ecology*, 29(S1), e13132. Available from: <https://doi.org/10.1111/rec.13132>
- Wilsey, B.J., Barber, K. & Martin, L.M. (2015) Exotic grassland species have stronger priority effects than natives regardless of whether they are cultivated or wild genotypes. *New Phytologist*, 205, 928–937. Available from: <https://doi.org/10.1111/nph.13028>
- Wright, A., Schnitzer, S.A. & Reich, P.B. (2014) Living close to your neighbors: the importance of both competition and facilitation in plant communities. *Ecology*, 95, 2213–2223. Available from: <https://doi.org/10.1890/13-1855.1>
- Young, T.P., Chase, J.M. & Huddleston, R.T. (2001) Community succession and assembly comparing, contrasting and combining paradigms in the context of ecological restoration. *Ecological Restoration*, 19, 5–18. Available from: <https://doi.org/10.3368/er.19.1.5>
- Young, T.P., Stuble, K.L., Balachowski, J.A. & Werner, C.M. (2017) Using priority effects to manipulate competitive relationships in restoration. *Restoration Ecology*, 25, S114–S123. Available from: <https://doi.org/10.1111/rec.12384>
- Yu, H., Yue, M., Wang, C., Roux, J.J.L., Peng, C. & Li, W. (2020) Priority effects and competition by a native species inhibit an invasive species and may assist restoration. *Ecology and Evolution*, 10, 13355–13369. Available from: <https://doi.org/10.1002/ece3.6938>

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**Appendix S1.** Experimental design with the 40 plots including the ten replicates of the four randomly distributed treatments: D+S: dominant and subordinate species were sown synchronously; D1st: dominant species were sown first (subordinates second); S1st: subordinate species were sown first (dominants second); and Control (no sowing).

**Appendix S2.** Total species cover and unsown species cover (species from the seed bank/seed rain) in the four treatments in 2019, 2020 and 2021.

**Appendix S3.** Percentage mean cover of spontaneously emerging plant species.

**Appendix S4.** Effect of the sowing treatments on sown species cover in 2021 (Figure 4).

**How to cite this article:** Durbecq, A., Bischoff, A., Buisson, E., Corcket, E. & Jaunatre, R. (2023) Using priority effects for grassland restoration: Sequential sowing can promote subordinate species. *Applied Vegetation Science*, 26, e12748. Available from: <https://doi.org/10.1111/avsc.12748>