

Contribution of Fenix high-resolution hyperspectral airborne images to the socio-morphological characterisation of Toulouse

Sébastien Gadal, Thomas Gloaguen

▶ To cite this version:

Sébastien Gadal, Thomas Gloaguen. Contribution of Fenix high-resolution hyperspectral airborne images to the socio-morphological characterisation of Toulouse. 8e colloque scientifique du Groupe Hyperspectral de la Société Française de Photogrammétrie et de Télédétection, Dec 2023, Paris, France. hal-04329105

HAL Id: hal-04329105

https://hal.science/hal-04329105

Submitted on 7 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Contribution of Fenix high-resolution hyperspectral airborne images to the socio-morphological characterisation of Toulouse

ONERA



Sébastien Gadal, Thomas Gloaguen

Aix-Marseille Université, CNRS ESPACE UMR 7300

Vegetation

population cat.)






INTRODUCTION

In this presentation, the contributions of high metric resolution hyperspectral images from the FENIX sensor (2 m) from the CAMCATT airborne campaign (ONERA) covered in June 2021 in Toulouse will be presented to characterise urban generate morpho-social pattern indicators. These indicators are essential in urban planning to understand the relationships between urban forms, functions, populations, and uses. The construction of this type of morpho-social (and morpho-economic) indicators is usually made by integrating socioeconomic and demographic databases with high resolution remote sensing images. Hyperspectral images make a significant contribution compared with multispectral images, thanks to their ability to discriminate between urban geographical objects: soils, vegetation, types of materials, etc.

METHODOLOGY

408 spectral Hyperspectral bands (2 m) Linear signature Unmixing Alg. extraction routine



The methodological approach, which was exploratory and involved testing several approaches, focused on the individual identification of urban components (spectral analysis and machine learning extraction) and urban structures/forms: morphologies and deep learning object recognition, data classification and urban-object data fusion.

database




Fig. 1 – Overall accuracy of urban land cover mapping with vegetation (left) and built-ups (right) objects. The level of mapping accuracy is better than that of reference data such as Street Tree Layer (yellow linear) for vegetation or Open Street Map (blue linear) for built-ups.

RESULTS




Fig. 2 – Urban Land Cover Morphology of Toulouse Results of processing with the predictive deep-learning model using the Multi-Layer Perceptron algorithm (left) and principal component analyses in T-mode (centre) and S-mode eigenvector (right).





Fig. 3 – Extraction of geographic objects in dense urban environments Identification of the built-up structure (left), water surfaces and humidity (right) using a Support Vector Machine classifier, and vegetation (centre) using a Linear Assemblist Transformed into Principal Component Analyses of the green and infrared bands.



Areas with a lower ratio of flats to houses are those with more vegetation and larger, newer buildings, inhabited mainly by pensioners or managers and intellectual workers.

CONCLUSIONS

- Metric hyperspectral images can be used to better identify the geographical objects that make up the urban fabric in dense built-up areas. They make it possible to identify urban objects and environmental indices (trees, buildings, materials, water features, humidity, etc.) that are absent from geographic databases and more precise spatially.
- The urban morphologies, socio-demographic and economic structures identified are perfectly spatially correlated.
- The exploratory methodologies show a strong correlation between the results and the spectral sensitivity of the images. Certain algorithms, whatever the spectral domain and spatial resolution (2 and 4 meters), such as ACP Tmode and Fuzzy Artmap NNA, are remarkably consistent in their ability to recognise urban objects and morphologies.