



HAL
open science

Additions to the late Eocene Süngülü mammal fauna in Easternmost Anatolia and the Eocene-Oligocene transition at the periphery of Balkanatolia

Grégoire Métais, Pauline Coster, Alexis Licht, Faruk Ocakoğlu, K. Christopher Beard

► To cite this version:

Grégoire Métais, Pauline Coster, Alexis Licht, Faruk Ocakoğlu, K. Christopher Beard. Additions to the late Eocene Süngülü mammal fauna in Easternmost Anatolia and the Eocene-Oligocene transition at the periphery of Balkanatolia. *Comptes Rendus. Palevol*, 2023, 22 (35), pp.711-727. 10.5852/cr-palevol2023v22a35 . hal-04335099

HAL Id: hal-04335099

<https://hal.science/hal-04335099>


Submitted on 11 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Additions to the late Eocene Süngülü mammal fauna
in Easternmost Anatolia and the Eocene-Oligocene
transition at the periphery of Balkanatolia**

Grégoire MÉTAIS, Pauline COSTER, Alexis LICHT,
Faruk OCAKOĞLU & K. Christopher BEARD



DIRECTEURS DE LA PUBLICATION / PUBLICATION DIRECTORS :
Gilles Bloch, Président du Muséum national d'Histoire naturelle
Étienne Ghys, Secrétaire perpétuel de l'Académie des sciences

RÉDACTEURS EN CHEF / EDITORS-IN-CHIEF : Michel Laurin (CNRS), Philippe Taquet (Académie des sciences)

ASSISTANTE DE RÉDACTION / ASSISTANT EDITOR : Adenise Lopes (Académie des sciences ; cr-palevol@academie-sciences.fr)

MISE EN PAGE / PAGE LAYOUT : Audrina Neveu (Muséum national d'Histoire naturelle ; audrina.neveu@mnhn.fr)

RÉVISIONS LINGUISTIQUES DES TEXTES ANGLAIS / ENGLISH LANGUAGE REVISIONS : Kevin Padian (University of California at Berkeley)

RÉDACTEURS ASSOCIÉS / ASSOCIATE EDITORS (*, *took charge of the editorial process of the article/a pris en charge le suivi éditorial de l'article*):

Micropaléontologie/*Micropalaeontology*

Lorenzo Consorti (Institute of Marine Sciences, Italian National Research Council, Trieste)

Paléobotanique/*Palaeobotany*

Cyrille Prestianni (Royal Belgian Institute of Natural Sciences, Brussels)

Métazoaires/*Metazoa*

Annalisa Ferretti (Università di Modena e Reggio Emilia, Modena)

Paléochthyologie/*Palaeoichthyology*

Philippe Janvier (Muséum national d'Histoire naturelle, Académie des sciences, Paris)

Amniotes du Mésozoïque/*Mesozoic amniotes*

Hans-Dieter Sues (Smithsonian National Museum of Natural History, Washington)

Tortues/*Turtles*

Walter Joyce (Universität Freiburg, Switzerland)

Lépidosauromorphes/*Lepidosauromorphs*

Hussam Zaher (Universidade de São Paulo)

Oiseaux/*Birds*

Eric Buffetaut (CNRS, École Normale Supérieure, Paris)

Paléomammalogie (mammifères de moyenne et grande taille)/*Palaeomammalogy (large and mid-sized mammals)*

Lorenzo Rook* (Università degli Studi di Firenze, Firenze)

Paléomammalogie (petits mammifères sauf Euarchontoglires)/*Palaeomammalogy (small mammals except for Euarchontoglires)*

Robert Asher (Cambridge University, Cambridge)

Paléomammalogie (Euarchontoglires)/*Palaeomammalogy (Euarchontoglires)*

K. Christopher Beard (University of Kansas, Lawrence)

Paléoanthropologie/*Palaeoanthropology*

Aurélien Mounier (CNRS/Muséum national d'Histoire naturelle, Paris)

Archéologie préhistorique (Paléolithique et Mésolithique)/*Prehistoric archaeology (Palaeolithic and Mesolithic)*

Nicolas Teyssandier (CNRS/Université de Toulouse, Toulouse)

Archéologie préhistorique (Néolithique et âge du bronze)/*Prehistoric archaeology (Neolithic and Bronze Age)*

Marc Vander Linden (Bournemouth University, Bournemouth)

RÉFÉRÉS / REVIEWERS : <https://sciencepress.mnhn.fr/fr/periodiques/comptes-rendus-palevol/referes-du-journal>

COUVERTURE / COVER :

Made from the Figures of the article.

Comptes Rendus Palevol est indexé dans / *Comptes Rendus Palevol is indexed by:*

- Cambridge Scientific Abstracts
- Current Contents® Physical
- Chemical, and Earth Sciences®
- ISI Alerting Services®
- Geoabstracts, Geobase, Georef, Inspec, Pascal
- Science Citation Index®, Science Citation Index Expanded®
- Scopus®.

Les articles ainsi que les nouveautés nomenclaturales publiés dans *Comptes Rendus Palevol* sont référencés par / *Articles and nomenclatural novelties published in Comptes Rendus Palevol are registered on:*

- ZooBank® (<http://zoobank.org>)

Comptes Rendus Palevol est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris et l'Académie des sciences, Paris
Comptes Rendus Palevol is a fast track journal published by the Museum Science Press, Paris and the Académie des sciences, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish:*

Adansonia, Geodiversitas, Zoosystema, Anthropolozologica, European Journal of Taxonomy, Naturae, Cryptogamie sous-sections *Algologie, Bryologie, Mycologie*.

L'Académie des sciences publie aussi / *The Académie des sciences also publishes:*

Comptes Rendus Mathématique, Comptes Rendus Physique, Comptes Rendus Mécanique, Comptes Rendus Chimie, Comptes Rendus Géoscience, Comptes Rendus Biologies.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél. : 33 (0)1 40 79 48 05 / Fax : 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <https://sciencepress.mnhn.fr>

Académie des sciences, Institut de France, 23 quai de Conti, 75006 Paris.

© This article is licensed under the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>)
ISSN (imprimé / print) : 1631-0683/ ISSN (électronique / electronic) : 1777-571X

Additions to the late Eocene Süngülü mammal fauna in Easternmost Anatolia and the Eocene-Oligocene transition at the periphery of Balkanatolia

Grégoire MÉTAIS

Centre de Recherche en Paléontologie – Paris (CR2P),
CNRS, Sorbonne Université, Muséum national d'Histoire naturelle,
case postale 38, 57 rue Cuvier, F-75231 Paris, cedex 05 (France)
gregoire.metais@mnhn.fr (corresponding author)

Pauline COSTER

Réserve Naturelle Nationale Géologique du Luberon, Parc naturel régional du Luberon,
Luberon UNESCO Global Geopark, 60 Place Jean Jaurès, F-84400 Apt (France)
pauline.coster@parcduluberon.fr

Alexis LICHT

CEREGE, Technopole Environnement Arbois-Méditerranée, Aix Marseille University,
CNRS, IRD, INRAE, CEREGE, BP 80, 13545 Aix-en-Provence, cedex 04 (France)
licht@cerge.fr

Faruk OCAKOĞLU

Department of Geological Engineering, Eskişehir Osmangazi University,
Meşelik Yerleşkesi, 26040, Odunpazarı, Eskişehir (Turkey)
focakoglu@gmail.com

K. Christopher BEARD

Biodiversity Institute and Department of Ecology and Evolutionary Biology,
University of Kansas, 1345 Jayhawk Blvd, Lawrence, KS 66045 (United States)
chris.beard@ku.edu

Submitted on 4 April 2023 | Accepted on 11 July 2023 | Published on 11 December 2023

[urn:lsid:zoobank.org:pub:E41471F9-7425-482F-9A32-118346B7FC66](https://zoobank.org/pub:E41471F9-7425-482F-9A32-118346B7FC66)

Métais G., Coster P., Licht A., Ocakoğlu F. & Beard K. C. 2023. — Additions to the late Eocene Süngülü mammal fauna in Easternmost Anatolia and the Eocene-Oligocene transition at the periphery of Balkanatolia. *Comptes Rendus Palevol* 22 (35): 711-727. <https://doi.org/10.5852/cr-palevol2023v22a35>

ABSTRACT

The Eocene-Oligocene transition marks a period of dramatic global climatic change correlated with pronounced mammalian faunal change. Fossil evidence is indispensable for studying the distribution of taxa through time, and determining how abiotic parameters shaped ancient biodiversity. Here we report ruminant artiodactyls and a new anthropoid primate from Süngülü, a locality in Eastern Anatolia that has yielded a diversified and largely endemic assemblage of rodents. Three taxa of

KEY WORDS
Balkanatolia,
Ruminantia,
anthropoid primate,
Eocene-Oligocene
transition,
new genus,
new species.

ruminants are recognized, the tragulid *Iberomeryx parvus* Gabunia, 1964, a larger species of *Iberomeryx* Gabunia, 1964, and a bachitheriid referred to cf. *Bachitherium* sp. A lower molar is identified as the new eosimiid primate *Sungulusimias unayae* n. gen., n. sp., which is the first occurrence of Paleogene anthropoids in western Asia. The lower molar of *Sungulusimias unayae* n. gen., n. sp. is characterized by protoconid and metaconid closely spaced and of similar height and volume, paraconid cuspidate and nearly connate with metaconid, strong mesiobuccal cingulid, and entoconid without strong connection to hypoconulid via the postcristid. The composition of this assemblage together with rodents indicates a probable Latest Eocene age for Süngülü, although an early Oligocene age cannot be completely ruled out. The rodent fauna from Süngülü suggests that endemism persisted at the periphery of Balkanatolia until the latest Eocene while Eastern Anatolia was situated in a strategic corridor for faunal exchanges between eastern Asia, Indo-Pakistan and Europe. During the Eocene-Oligocene transition, Balkanatolia probably functioned as a “holding pen” where various taxa were confined for significant intervals of time before proceeding to colonize Western Europe at the Grande Coupure.

RÉSUMÉ

Nouvelles données sur la faune de mammifères de l'Éocène supérieur de Süngülü en Anatolie orientale, et sur la transition Éocène-Oligocène à la périphérie de la Balkanatolie.

La transition Éocène-Oligocène est marquée par une période de changements climatiques globaux spectaculaires corrélés à des changements prononcés de la faune mammalienne. Les fossiles sont indispensables pour étudier la distribution des taxons dans le temps et comprendre comment les paramètres abiotiques ont façonné la biodiversité passée. Nous présentons ici les artiodactyles ruminants et un nouveau primate anthropoïde de Süngülü, une localité de l'Anatolie orientale qui a livré un assemblage diversifié et largement endémique de rongeurs. Trois taxons de ruminants sont identifiés, le tragulidé *Iberomeryx parvus* Gabunia, 1964, une espèce plus grande d'*Iberomeryx* Gabunia, 1964, et un bachitherridé laissé en cf. *Bachitherium* sp. Une molaire inférieure (probablement m2) est identifiée comme un nouveau primate eosimiidé *Sungulusimias unayae* n. gen., n. sp. qui est la première occurrence d'anthropoïde paléogène en Asie occidentale. La m2 de *Sungulusimias unayae* n. gen., n. sp. est caractérisée par un protoconide et métaconide rapprochés ainsi que de même hauteur et volume, un fort cingulide méso-labial, et un entoconide sans forte postcristide reliée à l'hypoconide. La composition de cet assemblage de mammifères indique un âge probable de l'Éocène supérieur pour Süngülü, bien qu'un Oligocène basal ne puisse pas être complètement exclu. La faune de rongeurs de Süngülü suggère que l'endémisme a persisté à la périphérie de la Balkanatolie jusqu'à l'Éocène supérieur, tandis que l'Anatolie orientale était située dans un corridor stratégique pour les échanges de faune entre l'Asie, l'Indo-Pakistan et l'Europe. Au cours de la transition Éocène-Oligocène, la Balkanatolie a probablement fonctionné comme un sas où certains taxons ont été confinés durant l'Éocène supérieur avant de coloniser l'Europe de l'Ouest à la Grande Coupure.

MOTS CLÉS
Balkanatolie,
Ruminantia,
primate anthropoïde,
transition Éocène-
Oligocène,
genre nouveau,
espèce nouvelle.

INTRODUCTION

From the Late Cretaceous to the late Paleogene, the area spanning from the Balkans to Pakistan has experienced multiple collision and deformation events associated with the closure of the Neotethys Ocean and convergence between the Eurasian and African/Arabian Plates. These events have resulted in the formation of long, semi-continuous strips of land, bound by the Paratethys to the north and the Neotethys to the south, displaying a complex pattern of evolving seaways and land-bridges (e.g. Popov *et al.* 2004; Barrier *et al.* 2018). The complex and dynamically changing paleogeographic configuration of this area shaped its biogeographic history as well as faunal exchanges between Africa, Asia, Europe, and India. Paleontological data suggest that a low-topography land mass covering the Balkans and Anatolia—recently named Balkanatolia—existed during at least the early and middle Eocene, which was populated by endemic and anachronis-

tic mammals reflecting its isolation from mainland Eurasia (Kappelman *et al.* 1996; Métais *et al.* 2018; Licht *et al.* 2022; Beard *et al.* 2023). This geographic isolation seems to have ceased through various stages of biogeographic connection with Eurasia by the late Eocene.

Eastern Anatolia consists of three terranes that are all attributed to the Balkanatolian landmass: the eastern Pontides, the eastern Taurides and the Armenia Block (van Hinsbergen *et al.* 2020). During the Eocene, the area was separated from other emerged lands further east (the Cimberian blocks of Iran) by remnant seaways connecting the Neotethys to the Paratethys on both sides of the Lesser Caucasus. Eastern Anatolia hosts multiple intermontane pull-apart basins and grabens, the sedimentary fill of which is often attributed to “undifferentiated Neogene” in geological reports (Şengör *et al.* 2008). Older, Oligocene deposits commonly display an overall regressive trend, with shallow marine deposits passing upward to detrital and gypsiferous facies (Adamia *et al.* 2010);




FIG. 1. — **A**, Location map of the Süngülü, Ardahan province, Eastern Anatolia; **B**, simplified geological map of the area redrawn after the 1/500 000 scale geological map of Turkey.

this trend is attributed to the onset of regional deformation following the final closure of the Neotethys and the drying-out of the remnant seaways.

The Posof-Ahalsikhe Basin (Fig. 1) lies at the edge of the eastern Pontides in the Lesser Caucasus and was filled with a detrital sequence dated as Oligocene based on marine molluscs (Yılmaz *et al.* 2001); the top of the sedimentary sequence consists of continental coal deposits and red clastics that have yielded the Benara mammal fauna generally considered as “middle” (Lucas & Emry 1999) or late Oligocene (Gabunia 1964). Only two other localities have yielded late Paleogene mammals in Easternmost Anatolia: 1) the late Oligocene Güngörmez Formation in the Kağızman-Tuzluca Basin has yielded sparse mammal remains, including large rhinos (Sen *et al.* 2011) and ruminants (Métais *et al.* 2015); and 2) the Süngülü locality, of debated late Eocene-early Oligocene age, in the Posof-Ahalsikhe Basin near the Turkish-Georgian border, has produced a diverse assemblage of small mammals, of which only the rodents have been described previously (de Bruijn *et al.* 2003). Paleogene mammals from Eastern Anatolia remain barely known, although they are of utmost significance for understanding the early evolution and distribution of modern faunas during an interval marked by strong climatic pulses. This dearth of paleontological data hampers attempts to reconstruct the complex biogeographic history of this region.

The aim of this paper is twofold: 1) describing the artiodactyl and primate fossils known from the Süngülü locality; and 2) analyzing their biochronological and paleobiogeographical implications in the context of the dismantling of Balkanatolia.

MATERIAL AND METHODS

The fossil material described herein was collected by screen washing about 3.5 tons of sediment (de Bruijn *et al.* 2003). The mammal remains have been collected from two successive beds: a ± 40 cm thick tuffite and the overlying ± 30 cm of white silty limestone with silicified nodules, but no significant difference in the faunal assemblages of these two beds has been detected (de Bruijn *et al.* 2003). The Süngülü mammal assemblage is dominated by rodents, and apart from artiodactyls and a single primate, it also includes insectivorous mammals and reptiles (de Bruijn *et al.* 2003). The specimens are permanently deposited in the collections of the MTA museum of Natural History, Ankara; casts of these fossils are available at the University of Utrecht (Netherlands). The measurements of teeth or postcrania are taken to measure the maximum length and width (maximum crown height was measured from the base of the crown to the tip of enamel for teeth). Measurements have been realized thanks to a caliper (precision 0.1 mm). The dental terminology follows Barmann & Rössner (2011) and Ni *et al.* (2016), the postcranial terminology follows Martinez & Sudre (1995).

ABBREVIATION

GABI Great American Biotic Interchange.

SYSTEMATIC PALAEOLOGY

Class MAMMALIA Linnaeus, 1758
 Order ARTIODACTYLA Owen, 1848
 Suborder RUMINANTIA Scopoli, 1777
 Family TRAGULIDAE Milne-Edwards, 1864

Genus *Iberomeryx* Gabunia, 1964

Iberomeryx Gabunia, 1964: 179.

TYPE SPECIES. — *Iberomeryx parvus* Gabunia, 1964.

DIAGNOSIS (from Mennecart *et al.* 2021). — Small-sized ruminant with upper molars possessing the following combination of characters: well-marked parastyle and mesostyle in small-column shape; strong paracone rib; metacone rib absent; metastyle absent; unaligned external walls of metacone and paracone; strong postprotocrista stopping against the anterior side of the premetaconulecrista; continuous lingual cingulum, stronger under the protocone. Lower dental formula is primitive (3-1-4-3) with non-molarized premolars. Tooth c is adjacent to i3. Tooth p1 is single-rooted, reduced and separated from c and p2 by short diastemata. The premolars have a well-developed anterior conid. Teeth p2-p3 display a distally bifurcated mesolabial conid. Tooth p3 is the largest premolar. Tooth p4 displays no mesolingual conid and a large posterior valley. Regarding the lower molars, the trigonid and talonid are lingually open with a trigonid more tapered than the talonid. The anterior fossa is open, due to a forward orientation of the preprotocristid and the presence of a paraconid. The internal postprotocristid is oblique and the external postprotocristid reaches the prehypocristid. The internal postprotocristid, postmetacristid and preentocristid are fused and Y-shaped. Protoconid and metaconid display a weak *Tragulus* Brisson, 1762 fold and a well-developed *Dorcatherium* fold, respectively. The mandible displays a regularly concave ventral profile in lateral view, a marked incisura vasorum, a strong mandibular angular process, a vertical ramus, and a stout condylar process.

TYPE LOCALITY. — Benara, Georgia, late Oligocene.

Iberomeryx parvus Gabunia, 1964

Iberomeryx parvus Gabunia, 1964: 182.

TYPE MATERIAL. — **Holotype**. PIT-7-08, fragmentary maxilla preserving left M1-M3, Georgian National Museum, Tbilisi, Georgia.

Iberomeryx sp.
 (Fig. 2P-Q)

REFERRED MATERIAL. — Sü-2016, left m3 (third lobe, hypoconulid lacking)

LOCALITY. — Outcrop of unnamed rock unit consisting of c. 40 cm thick tuffite bed containing gastropod operculae superposed by c. 30 cm thick white silty limestone with silicified nodules exposed in a streambed roughly two km northwest of Süngülü, Ardahan Province, Turkey (de Bruijn *et al.* 2003: fig. 1). Latest Eocene according to de Bruijn *et al.* (2018, 2019).

DESCRIPTION

The m3 is incomplete since the third lobe bearing the hypoconulid is damaged, and the mesial margin and labial flank of the protoconid are broken (Fig. 2). The molar is brachyo-




FIG. 2. — Dental remains and post-cranial material of ruminants from Süngülü: **A-O**, *Iberomeryx* sp. cf. *I. parvus*: **A, B**, Sü-2002, left p3 in occlusal (**A**) and lingual (**B**) views; **C**, Sü-2020, left p4 in occlusal view; **D**, Sü-2005, talonid of left p4 in occlusal view; **E, F**, Sü-2003, left p4 in occlusal (**E**) and lingual (**F**) views; **G**, Sü-2010, lingual half of a right upper molar in occlusal view; **H, I**, Sü-2009, right m1 in occlusal (**H**) and labial (**I**) views; **J-L**, Sü-2014, right astragalus in anterior (**J**), medial (**K**) and lateral (**L**) views; **M-O**, Sü-2013, right astragalus in anterior (**M**), medial (**N**) and lateral (**O**) views; **P, Q**, *Iberomeryx* sp., Sü-2016, left m3 (third lobe lacking) in occlusal (**P1**, **P2** is the line drawing, **blue areas** how missing parts) and labial (**Q**) views; **R, S**, cf. *Bachitherium* sp.: **R**, Sü-2017, lingual half of a right upper molar in occlusal view; **S**, Sü-2019, talonid of a left p4 in occlusal view (**S1**, **S2** is the line drawing, **blue areas** how missing parts). The **grey areas** show the parts of the teeth that are broken, and missing. Abbreviations: **acf**, astragalo-calcaneal facet; **daf**, distal astragalar facet; **imf**, internal malleolus facet. Scale bars: A-I, P-S, 1 mm; J-O, 4 mm.

dont, and the protoconid is slightly taller than the metaconid, which is more mesially positioned than the protoconid. The premetacristid is absent. There is a strong wear facet on the preprotocristid and the mesial part of the molar is broken, hampering evaluation of the extension of this crest mesially and lingually; it is worth noting that the metaconid is not conical but instead shows an incipient labial crest, unlike the condition typically observed on m3 of *Iberomeryx*. The short internal postprotocristid is lingually oriented and joins the short and posterolabially oriented internal postmetacristid, but they do not reach the mesially oriented and elongated

preentocristid that stops at the base of the posterior wall of the trigonid. The external postprotocristid is barely visible (but present) and fused with the distal portion of the prehypocristid. The metaconid displays a short and marked external postmetacristid. Although attenuated (on its labial side), the M-structure is present (Fig. 2). The entoconid is more mesially positioned than the hypoconid, it displays a strong wear facet on its posterior face, and it is not transversely compressed as in *Iberomeryx parvus*. The posthypocristid extends lingually and its distal extremity is connected to a mesio-distally extended pre-hypoconulid cristid. There is a crest extending mesio-

TABLE 1. — Measurements (in mm) of the ruminant dental material from Süngülü discussed in the text. The asterisks designates estimated values.

Taxon	Rank	Specimen	Length	Width
<i>Iberomeryx</i> sp. cf. <i>I. parvus</i>	p3	Sü-2002	3.8	1.4
	p4	Sü-2003	4	1.7
	p4 (talonid)	Sü-2005	–	1.6
	m1	Sü-2009	3.9	1.8-2.0
	lingual M1 or M2	Sü-2010	3.7*	–
<i>Iberomeryx</i> sp.	m3	Sü-2016	7.1*	3.6
cf. <i>Bachitherium</i> sp.	M1 or M2	Sü-2017	5.4	–
	p4	Sü-2019	–	2.5
<i>Sungulusimias unayae</i> n. gen., n. sp.	m2	Sü-2020	3	2.4

distally at the base of the posterior wall of the entoconid which is most probably a remnant of the internal pre-hypoconulid cristid, suggesting that the hypoconulid was connected to the rest of the talonid by two cristids.

TAXONOMIC ATTRIBUTION

Even considering intraspecific size variability, this m3 is roughly twice as large as would be expected for an m3 that would match the size of Sü-2009 (m1). The current sample from Süngülü does not allow us to estimate the size variability of *Iberomeryx* sp. cf. *I. parvus*, but the size of this m3 surpasses the size variability observable in the extant tragulid *Tragulus kanchil* (Raffles, 1821), for example (Meijaard & Groves 2004). Moreover, the morphology of this m3 is more bunodont and bulkier than the m1 of *Iberomeryx* sp. cf. *I. parvus* described below. The combination of an external postprotocristid, an internal postprotocristid, an internal postmetacristid and an external postmetacristid corresponds to the tragulid M-structure or Σ- structure, justifying identification of this molar as that of a tragulid (Mottl 1961; Rössner 2007). However, given the paucity of the present material and pending additional data, we prefer to assign this tooth to *Iberomeryx* sp.

cf. *Iberomeryx parvus* Gabunia, 1964
(Fig. 2A-I, J-O)

REFERRED MATERIAL. — Sü-2002, left p3; Sü-2003, left p4; Sü-2005, talonid of a left p4; Sü-2009, right m1; Sü-2010, lingual part of a right M1 or M2; Sü-2020, left p4; Sü-2013, right astragalus; Sü-2014, right astragalus.

LOCALITY. — Outcrop of unnamed rock unit consisting of c. 40 cm thick tuffite bed containing gastropod operculae superposed by c. 30 cm thick white silty limestone with silicified nodules exposed in a streambed roughly two km northwest of Süngülü, Ardahan Province, Turkey (de Bruijn *et al.* 2003: fig. 1). Latest Eocene according to de Bruijn *et al.* (2018, 2019).

DESCRIPTION

Dentition

The p3 is elongated with a small paraconid, and a hypoconid lower than the protoconid. A sharp crest joins the apex of the protoconid to the hypoconid, and is bifurcated distally, so that the two crests form a circular, postero-lingually opened

depression. The paraconid of p4 is more pronounced than on p3, and the crest joining the paraconid to the protoconid corresponds to a wear facet. Two crests extend backward from the apex of the protoconid; the better expressed labial crest joins the distal border of the tooth, whereas the weaker lingual crest stops at the mid-point of the posterior half of the tooth, leaving the talonid distolingually opened.

The lower molar is brachyodont, and only the hypoconid is completely crescent-shaped. The trigonid is narrower than the talonid, and the protoconid is taller than the metaconid. The anterior fossa is widely antero-lingually open due to the absence of a premetacristid. The preprotocristid extends mesially and is notched; it forms the anterior-most part of the lower molars. The internal postprotocristid is lingually oriented and short, and it joins the very short and posterolabially oriented internal postmetacristid to form a cristid at the rear of the trigonid which is oriented toward the mesially oriented preentocristid. The external postprotocristid is very faint and extends toward the prehypocristid without connecting it because of transverse breakage of Sü-2009. The metaconid is rounded mesially and displays a short external postmetacristid. The combination of an external postprotocristid, an internal postprotocristid, an internal postmetacristid and an external postmetacristid forms a M-structure or Σ- structure which is characteristic of tragulids. The posthypocristid is short and lingually oriented. The entoconid is labio-lingually compressed and its distal side is rounded (devoid of any crest), thus leaving the posterior fossa open on its postero-lingual corner. There are both mesial and distal cingulids and an ectostylid on Sü-2009.

The fragmentary upper molar (Sü-2010) only preserves the lingual part. The relatively large size of the metaconule suggests it is either a M1 or a M2. The postprotocrista is short, straight, and postero-labially oriented whereas the preprotocrista is labially oriented. The premetaconulecrista extends labially. There is a strong cingulum surrounding the protocone and extending along the mesial border of the tooth; however, there is no protoconule as in *Nalameryx* (Métais *et al.* 2009).

Postcranials

Two astragali of similar size are referred to this small taxon on the basis of both their size and morphology. The astragali display non-aligned proximal and distal trochlea, the distal one being slightly rotated medially. The distal trochlea does not show any ridge separating the articular surfaces of the

TABLE 2. — Measurements (in mm) of the ruminant postcranial material from Süngülü discussed in the text, and all referred to *Iberomeryx* sp. cf. *I. parvus*. The asterisks designates estimated values when material is slightly broken or heavily worn/digested.

Taxon	Specimen	Length	Mesio-lateral width	Dorso-plantar height
<i>Iberomeryx</i> sp. cf. <i>I. parvus</i>	Sü-2014	10	5.46	5.2*
<i>Iberomeryx</i> sp. cf. <i>I. parvus</i>	Sü-2013	9.2	5	5*

cuboid and the navicular. The astragalo-calcaneal facet is well-developed, and there is a deep fossa for the fibular condyle of the calcaneum. The distal astragalar facet is generally well-developed as it is in primitive Pecora (Métais *et al.* 2016). The distal part of the internal malleolus facet is wide. Sü-2013 is slightly smaller than Sü-2014, but they are similar in morphology, and we attribute these slight metric differences to intraspecific variation.

Measurements

Provided in Table 1 for the dental material, and in Table 2 for postcranial material.

TAXONOMIC ATTRIBUTION

The dental material displays the characteristics of the genus *Iberomeryx*, and the astragali strongly resemble those reported from the early late Oligocene Kızılırmak Formation, central Anatolia, Turkey (Métais *et al.* 2016). The genus *Iberomeryx* is known by three species: *I. miaoi* Mennecart, Aiglstorfer, Li, Li & Wang, 2021 recently described from material first identified as *Lophiomeryx gracilis* Miao, 1982 (Mennecart *et al.* 2021) and restricted to the late Eocene beds of the Shinao Basin, southern China; *I. minor* (Filhol, 1882) is known from early Oligocene (MP 23-24) localities of Western Europe (Mennecart *et al.* 2011); *I. parvus* is known from the late Oligocene of Georgia (Gabunia 1964) and Central Anatolia (Métais *et al.* 2016). The material from Süngülü is identified as pertaining to *I. parvus* on the basis of the combination of these diagnostic features: lower molars lack a metastylid, and show a thin mesial cingulum, p4 with a distinct disto-lingual notch. However, the teeth of type material of *I. parvus* are higher crowned than that of the material from Süngülü which is also slightly smaller. Therefore, owing to these morphological and size differences, and pending additional fossil data, we prefer to refer our fossil material to *Iberomeryx* sp. cf. *I. parvus*.

Family BACHITHERIIDAE Janis, 1987

Genus *Bachitherium* Filhol, 1882

cf. *Bachitherium* sp.

(Fig. 2R-S)

REFERRED MATERIAL. — Sü-2017, lingual half of a right molar (M1 or M2); Sü-2019, talonid of a left p4.

DIAGNOSIS (from Métais & Vislobokova 2007). — Elongated muzzle with small orbits placed relatively posteriorly (the anterior border of the orbit is situated above M2), postorbital bar partly formed by the jugal apophysis, sagittal crest developed and extending posteriorly

into temporal crests, small auditory bullae, lacrimal fossa absent, strong paraoccipital apophysis, limited postero-lateral exposure of the mastoid, ethmoidal fissure moderately developed, post-glenoid apophysis well-developed; the angular region of the mandible is extended posteriorly and upward, and the coronoid apophysis is much higher than the articular joint suggesting strong abductor muscles. Dental Formula: 0/3 1/1 3/4 3/3, upper canine tusk-like, slightly curved posteriorly, and occludes against the anterior side of the caniniform p1, P1 lost, very long diastema between C and P2, lower incisors as the incisiform canine are small, small diastema between c and p1, and long diastema between p1 and p2. Radius and ulna separate, tibia and fibula partially fused, distal extremity reduced to a malleolar bone, Mc III and IV unfused, Mt III and IV fused proximally, lateral metatarsals absent, astragalus with unparallel trochlea, crural index close to that of the extant genus *Moschus*.

LOCALITY. — Outcrop of unnamed rock unit consisting of c. 40 cm thick tuffite bed containing gastropod operculae superposed by c. 30 cm thick white silty limestone with silicified nodules exposed in a streambed roughly two km northwest of Süngülü, Ardahan Province, Turkey (de Bruijn *et al.* 2003: fig. 1). Latest Eocene according to de Bruijn *et al.* (2018, 2019).

DESCRIPTION

The lingual half of the molar preserves highly worn protocone and metaconule; there is no lingual cingulum or entostyle, and there is both a mesial and a distal cingulum; the enamel is relatively thick and smooth. The talonid of p4 which is tentatively referred to the same taxon as the fragmentary upper molar (mostly on the basis of large size) displays in occlusal view a pattern of crests that sets it apart from tragulids and lophiomerycids. The top of the protoconid is preserved and the transversely wide talonid is delimited by two crests extending posteriorly; the labial postprotocristid reaching the posterior margin, where it forms the posterior border of the tooth; the lingual postprotocristid seems to be interrupted at the level of a marked transverse crest which is connected to the labial postprotocristid but without totally reaching the lingual border of the tooth. The disto-lingual border of the tooth is badly damaged but it does not appear that a lingual opening existed. Posterior to the protoconid, there are enamel foldings.

TAXONOMIC ATTRIBUTION

The lack of a lingual cingulum on the upper molar Sü-2017 is a character seen in *Bachitherium* (Geraads *et al.* 1987). If the premolar Sü-2019 belongs to the same taxon as we hypothesize here, its morphology is also consistent with a p4 of *Bachitherium*. This “gutter-like” p4 showing transverse cristids subdividing the talonid into two basins is also present in the type species *Bachitherium curtum* Filhol, 1877 (MNHN-Qu3917-holotype), and in *Bachitherium insigne* Filhol, 1882 (MNHN-Qu3902; MNHN-Qu3918-holotype). However, the p4 of *B. curtum* and *B. insigne* display not

one but two neocristids extending transversely from the lingual and labial postprotocristids respectively, but they do not fuse and leave a notch, medially situated, between the extremities of these two crests. This morphology is not what is observed on Sü-2019, where a single transverse neocristid extends transversely from the labial postprotocristid to reach the lingual wall of the posterior valley, which is devoid of a neocristid. We consider that Sü-2017 and Sü-2019 are closer to *Bachitherium* than to any other ruminant taxon, but the morphology of Sü-2019 does not match any species of *Bachitherium* known so far. Given the paucity of the current fossil material, we refer Sü-2019 to cf. *Bachitherium* sp. The morphology of Sü-2019 is also unlike the p4 of *Bachitherium thracensis* Mennecart, Geraads, Spassov & Zagorchev, 2018a from the late Eocene of Bulgaria, in which the gutter-like talonid of p4 lacks a transverse cristid (Mennecart *et al.* 2018a). Thus, if our assumption is correct, it worth noting that there are two forms of bachitheriids in the late Eocene of Balkanatolia, suggesting a diversification of the family in this biogeographic province before its dispersal to Western Europe at MP 23 (“*Bachitherium* dispersal event” of Mennecart *et al.* 2018a, b).

Order PRIMATES Linnaeus, 1758

Suborder ANTHROPOIDEA Mivart, 1864

Family EOSIMIIDAE Beard, Qi, Dawson, Wang & Li, 1994

Genus *Sungulusimias* n. gen.

[urn:lsid:zoobank.org:act:B7E863C3-4D08-4FFB-BF9D-C74C72927E23](https://doi.org/10.31233/osf.io/zt9q2)

TYPE SPECIES. — *Sungulusimias unayae* n. sp.

Sungulusimias unayae n. gen., n. sp.
(Fig. 3A-C)

[urn:lsid:zoobank.org:act:F21B9D32-30F3-4A64-B0C7-F89DDB9ED92C](https://doi.org/10.31233/osf.io/zt9q2)

TYPE MATERIAL. — **Holotype.** Sü-2021, isolated right m2, only known specimen.

DIAGNOSIS. — Differs from *Eosimias* Beard, Qi, Dawson, Wang & Li, 1994, *Phenacopithecus* Beard & Wang, 2004 and *Bahinia* Jaeger, Thein, Benammi, Chaimanee, Soe, Lwin, Tun, Wai & Ducrocq, 1999 in having m2 with protoconid and metaconid more closely spaced and of similar height and volume, paraconid more cuspidate and more nearly connate with metaconid, stronger mesio Buccal cingulid, and entoconid without strong connection to hypoconulid via the postcristid. Differs from *Phileosimias* Marivaux, Antoine, Hassan Baqri, Benammi, Chaimanee, Crochet, de Franceschi, Iqbal, Jaeger, Métais, Roohi & Welcomme, 2005 in having M₂ with paraconid fully lingual in position and hypoconulid distobuccal in position, being located closer to the hypoconid than the entoconid.

ETYMOLOGY. — Generic name from the village of Süngülü and the Latin *simias* (ape). Specific name in recognition of Prof. Engin Ünay, who discovered the type locality and helped collect the holotype and associated fauna from Süngülü.

TYPE LOCALITY. — Outcrop of unnamed rock unit consisting of c. 40 cm thick tuffite bed containing gastropod operculae superposed by c. 30 cm thick white silty limestone with silicified nodules exposed in a streambed roughly two km northwest of Süngülü, Ardahan Province, Turkey (de Bruijn *et al.* 2003: fig. 1). Latest Eocene according to de Bruijn *et al.* (2018, 2019).

DESCRIPTION

The holotype is an isolated right m2 (L = 3.0 mm; W = 2.4 mm). The crown is subrectangular in occlusal outline, although the trigonid is slightly narrower than the talonid. A relatively well-developed mesio Buccal cingulid extends from the base of the paraconid to the level of the hypoflexid. All three trigonid cusps are present, roughly similar in size, and distinctly cuspidate. The protoconid is situated internally rather than peripherally on the trigonid. As a result, the protoconid and metaconid are closely approximated. The metaconid is located slightly posterior to the level of the protoconid, and these two cusps are connected by a protocristid that runs somewhat obliquely with respect to the long axis of the tooth. A short but relatively trenchant paracristid arcs mesiolingually to connect the protoconid with the paraconid. The paraconid is fully lingual in position and almost connate with the metaconid. A short premetacristid fills the gap between the paraconid and metaconid. The postvallid is essentially vertical, separating the trigonid from the broader talonid. The hypoconid is the dominant talonid cusp, being situated near the distobuccal corner of the tooth. It gives rise to a straight, moderately trenchant cristid obliqua, which joins the postvallid near the lingual base of the protoconid. The hypoflexid is moderately deep but partly filled by the mesio Buccal cingulid. The lingual side of the talonid is marked by a relatively tall, isolated entoconid. A notch separates the lingual base of the postvallid from the entoconid, although a weak preentocristid is present. Distobuccally, the entoconid is not strongly connected to the hypoconulid by the postcristid. Instead, the hypoconulid projects slightly distally beyond the rest of the talonid and is more closely associated with the hypoconid than the entoconid.

COMPARISONS

Sü-2021 differs from primitive adapiform (e.g. *Donrussellia* Szalay, 1976) and omomyid primates (e.g. *Teilhardina* Simpson, 1940, *Steinius* Bown & Rose, 1984) in having the paraconid and metaconid of m2 closely approximated but not fully connate and in having an enlarged, distally expansive hypoconulid (Rose & Bown 1991). Most late early Eocene and younger adapiforms (adapids, notharctids, and sivaladapids) and stem lemuriforms (e.g. *Djebelemur* Hartenberger & Marandat, 1992) have reduced or completely lost the paraconid on m2 (Godinot 2014). Sivaladapid adapiforms (e.g. *Yunnanadapis* Ni, Li, Li & Beard, 2016, *Laomaki* Ni, Li, Li & Beard, 2016) retain a large hypoconulid on their lower molars, but this structure is invariably closely approximated with the entoconid in this clade, and it does not project distally beyond the rest of the talonid as it does in *Sungulusimias* (Ni *et al.* 2016). Anaptomorphine omomyids, best documented from the early and middle Eocene of North America, are much




FIG. 3. — The eosimiid primate from Süngülü *Sungulusimias unayae* n. gen., n. sp.: **A-C**, Sü-2021, holotype, right m2, in occlusal (**A**), labial (**B**), and lingual (**C**) views. Scale bar: 1 mm.

more bunodont than *Sungulusimias* and they never possess the enlarged, distally expansive molar hypoconulid found in the latter taxon (Godinot 2014). North American and Asian omomyines, including the middle Eocene *Nesomomys* Beard, Métais, Ocakoglu & Licht, 2020 from Balkanatolia (Beard *et al.* 2021), are typically less bunodont than their anaptomorphine relatives, but they differ from *Sungulusimias* in having lower molars with trigonids much narrower than talonids (e.g. *Nesomomys*, *Hemiacodon* Marsh, 1872) and much smaller, less bulbous hypoconulids (e.g. *Shoshonius* Granger, 1910, *Mytonius* Robinson, 1968). The European microchoerid radiation, which persisted on the Iberian Peninsula as recently as the early Oligocene (Köhler & Moyà-Solà 1999), differs from *Sungulusimias* in having reduced both paraconid and hypoconulid on m2, with certain taxa (e.g. *Pseudoloris* Stehlin, 1916) showing enhanced molar shearing related to insectivory and others (e.g. *Microchoerus* Wood, 1846) developing heavily crenulated enamel in relation to frugivorous adaptations (Godinot 2014).

The distinctive trigonid and hypoconulid morphology of Sü-2021 allows it to be identified as an eosimiid rather than any other anthropoid clade. Like eosimiids, and in contrast to other stem anthropoids, Sü-2021 retains a lingual, distinctly cuspidate paraconid on m2. Certain stem anthropoids from Africa (e.g. *Biretia* de Bonis, Jaeger, Coiffait & Coiffait, 1988, *Proteopithecus* Simons, 1989) sometimes retain small paraconids on m1, but these taxa rarely retain even vestigial

paraconids on m2 and their overall molar morphology is more bunodont than that of *Sungulusimias* (Seiffert 2012). Among stem anthropoid taxa that are generally recognized as eosimiids, the trigonid morphology of Sü-2021 most closely resembles that of *Phileosimias* from the Oligocene of Pakistan (Marivaux *et al.* 2005), from which it differs in having a fully lingual paraconid. In other eosimiids (*Eosimias*, *Phenacopithecus* and *Bahinia*), the paraconid of m2 is typically less cuspidate and farther removed from the metaconid. The hypoconulid of Sü-2021 flares distobuccally, as it does in *Eosimias* and *Phenacopithecus*. In *Bahinia*, the talonids of m1-2 lack distinct hypoconulids. In *Phileosimias* and most early African anthropoids, the lower molar hypoconulids can be substantial and cuspidate, but they are located closer to the entoconid than the hypoconid, in contrast to the more central placement of the hypoconulid in *Sungulusimias*.

REMARKS

Sungulusimias is the first record of stem anthropoids in Anatolia and only the second example of Paleogene primates from there (Beard *et al.* 2021). Other eosimiids are known from China (Beard *et al.* 1994, 1996; Beard & Wang 2004; Ni *et al.* 2016), Myanmar (Jaeger *et al.* 1999) and Pakistan (Marivaux *et al.* 2005), so *Sungulusimias* is also the westernmost record of this group. Discovery of Anatolian eosimiids is not unexpected, given that eosimiids are well-documented from farther east along the Eurasian Neotethyan margin and

that eosimiiforms colonized the African/Arabian Plate prior to its tectonic collision with Eurasia near the Oligocene-Miocene boundary (Chaimanee *et al.* 2012). However, *Sungulusimias* is apparently younger than the oldest known African anthropoids, which date to the late middle or early late Eocene (Seiffert *et al.* 2005; Jaeger *et al.* 2010; Marivaux *et al.* 2014). What role, if any, *Sungulusimias* and its collateral relatives may have played in the colonization of Africa by early anthropoids must await better understanding of its anatomy and phylogenetic relationships.

DISCUSSION

AGE OF THE SÜNGÜLÜ FAUNA

The ruminant fauna from Süngülü does not bring definitive arguments about the age of this important fossil assemblage. The earliest occurrences of tragulids are known from the late Eocene of Southeast Asia with *Archaeotragulus krabiensis* Métais, Chaimanee, Jaeger & Ducrocq, 2001 from Krabi (Métais *et al.* 2001), and *Iberomeryx miaoi* is now reported from the late Eocene of China (Mennecart *et al.* 2021). The two other species of *Iberomeryx*, *I. minor* and *I. parvus*, are known from the early and late Oligocene, respectively. Lucas & Emry (1999) suggested that the age of the Benara locality, which has yielded the type of *I. parvus*, may equally be correlated with MP 23 (early Oligocene), which is consistent with the occurrence in the Benara fauna of taxa such as the ruminant *Lophiomeryx* Pomel, 1854 and the anthracotheriid *Anthracotherium* Cuvier, 1822, which are known in western Europe from MP 22 and MP 21, respectively (BiochroM'97 1997). Gabunia (1964) assigned a late Oligocene age to the Benara mammals largely because they immediately overlie and are interbedded with marine to brackish water sandstones that contain a *Corbula*-dominated bivalve assemblage which is supposed to record a well-documented early Chattian marine transgression in the eastern Paratethys. The primate *Sungulusimias* likewise provides an ambiguous biostratigraphic signal. Most eosimiids (including *Eosimias*, *Phenacopithecus* and *Bahinia*) are middle Eocene in age, but *Phileosimias* from Pakistan is Oligocene (Marivaux *et al.* 2005). *Sungulusimias* combines molar features that occur separately in Eocene eosimiids (such as its distinctive hypoconulid morphology) and Oligocene *Phileosimias* (particularly its trigonid morphology). It is possible that *Sungulusimias* is a primitive relative of *Phileosimias*, but this hypothetical relationship cannot be tested adequately until additional material of *Sungulusimias* is recovered.

The rodent assemblage from Süngülü consists of a mixture of endemic and mostly Asian taxa showing a dominance of muroids and ctenodactyloids (de Bruijn *et al.* 2003). It is worth noting that the typically European theridomyids, pseudosciurids and glirids are missing in Süngülü, suggesting an absence of biogeographic relationships with western Europe at that time. The endemic character of rodents from Süngülü makes biochronological correlations with faunas from Central Asia very difficult. According to de Bruijn *et al.* (2003), the cricetid *Eucricetodon* (*Atavocricetodon*) *kurthi*

from Süngülü matches in size and morphology *A. minusculus* from Caijiachong (southern China), which is now considered to be late Eocene in age (Mennecart *et al.* 2021). In their chronostratigraphic scheme of Paleogene faunas from Anatolia, the Balkans and Southern Asia, van de Weerd *et al.* (2021) and Marković *et al.* (2018: fig. 8) considered Süngülü as latest Eocene.

THE LEAKY BALKANATOLIAN HOLDING PEN AND ITERATIVE FAUNAL CHANGE AROUND THE GRANDE COUPURE

The newly recognized Balkanatolian biogeographic province is defined on the basis of multiple Eocene sites in the Balkans and central Anatolia (Licht *et al.* 2022). During the middle Eocene, central Anatolia hosted strongly endemic faunas characterized by a mélange of anachronistic taxa such as the pleuraspidotheriid *Hilalia* and invasive clades including embriothopods and anatoliadelphid metatherians (Métais *et al.* 2017, 2018; Beard *et al.* 2023). The insular nature of these middle Eocene Balkanatolian faunas is underscored by the absence of artiodactyls, perissodactyls, carnivorans, hyaenodontids and rodents, which are otherwise ubiquitous across Laurasia at this time (Métais *et al.* 2018). The only insular Balkanatolian mammal showing Asian affinities is the omomyid primate *Nesomomys*, which is thought to have reached Balkanatolia by rafting (Beard *et al.* 2021). We currently lack middle Eocene paleontological sites between central Anatolia and South Asia (Pakistan), and therefore the eastward extent of Balkanatolian endemism remains unknown. The eastern boundary of Balkanatolia is hypothetically placed in the Lesser Caucasus, where remnant seaways separated Anatolian terranes from the Cimmerian blocks of Iran during most of the Eocene (Popov *et al.* 2004; Barrier *et al.* 2018; Palcu & Krijgsman 2023). While no body fossils of mammals have been so far reported from the Eocene of Iran, several late Eocene localities have yielded mammal footprints organized in trackways that have been interpreted as being made by perissodactyls and proboscideans (Abbassi *et al.* 2016). The Süngülü locality, on the eastern Pontides, would have been west of these seaways and thus part of Balkanatolia, although its younger age and relatively cosmopolitan fauna shows that it postdates the height of Balkanatolian endemism. Licht *et al.* (2022) have proposed that the progressive retreat of adjacent seaways in the late middle Eocene, *c.* 40–38 Ma, opened a terrestrial pathway for faunal exchanges between Balkanatolia and the Cimmerian blocks, favoring the dispersal of East Asian mammals as far as western Balkanatolia (i.e., Romania, Bulgaria). Moreover, Licht *et al.* (2022) hypothesized that subsequent westward prolongation of this dispersal pathway could have instigated the Grande Coupure roughly 34 Ma, when mammals of East Asian affinity invaded western Europe. Although there is currently no overlap in faunal composition between the central Anatolian Büyükteflekköy fauna reported by Licht *et al.* (2022), which is dominated by large ungulates, and the eastern Anatolian Süngülü fauna dominated by micromammals and small ruminants that we focus on here, this overall scenario is equally applicable to both of these late Eocene Anatolian faunas.

It is instructive to compare the pattern of faunal turnover observed in western Europe on either side of the Grande Coupure with that which has been documented for the Great American Biotic Interchange (GABI) during the late Neogene (e.g. Woodburne 2010). In both cases mammalian faunal turnover was severe, and in both cases it was caused by paleogeographic changes linked to tectonics, eustasy, and/or climate change. However, the nature and complexity of potential pathways for dispersal were very different. In the case of GABI, dispersal ultimately entailed traversing the Panamanian Isthmus connecting North and South America. In the case of the Grande Coupure and closely related dispersal episodes in Eurasia, potential pathways for dispersal are more numerous, reflecting the dynamic paleogeography of Eurasia during the latter part of the Eocene and Oligocene. For example, Mennecart *et al.* (2018b: 9) have stressed that two dispersal episodes are discernible across Eurasia during the late Eocene-early Oligocene, while emphasizing that associated dispersal pathways may have been different in each case. Classically, it is postulated that the earlier dispersal episode – the Grande Coupure *sensu stricto* – utilized a northern dispersal pathway linking Central Asia with western Europe across northern latitudes (mainly comprising the modern Russian Federation), while a more recent dispersal episode – designated the *Bachitherium* event by Mennecart *et al.* (2018b) – traversed a more southern pathway that included Balkanatolia. Likewise, the mechanisms underlying these two events are postulated to be different – including the drying out of the Turgäi Strait with respect to the Grande Coupure and tectonic uplift associated with the Alpine orogeny connecting Balkanatolia with western Europe in the case of the *Bachitherium* event. However, several authors (Heissig 1979; Böhme *et al.* 2013) suggested that the alpine tectonics created intermittent land bridge that probably connected the Island Europe and Southwestern Asia, and thus another potential dispersal pathway to explain the sudden appearance of Asian mammals at MP 21 in Western Europe.

Closer examination of faunal changes in western Europe during the late Eocene and early Oligocene suggests a more complex pattern of mammalian dispersal that probably included at least four separate “events” (Table 3). The first of these, which occurred prior to the Grande Coupure during MP 18, records the first appearance in western Europe (La Débruge, France) of the anthracothere *Elomeryx*, which is also known from Balkanatolia terranes to the east (de Bonis 1964; Nikilov & Heissig 1985). A separate anthracothere, *Prominatherium*, is relatively widely distributed across western Balkanatolia during the late Eocene. The Grande Coupure itself comprises the second event, characterized by the first appearances in western Europe at MP 21 of a large number of small- to large-bodied mammals (Table 3), which share only 7% of common genera with the late Eocene Balkanatolian mammal fauna, possibly suggesting a different geographic source for the MP 21 fauna of Western Europe. The third and fourth dispersal events are relatively minor inputs of Asian taxa of much lower magnitude than the major MP 21 turnover that was initially documented in the Paris Basin by Stehlin (1909).

The fourth dispersal event corresponds to the *Bachitherium* event of Mennecart *et al.* during MP 23, when *Bachitherium* and *Iberomeryx* first appear in Western Europe.

Finer resolution of the GABI dispersals during the late Neogene of North and South America has also resulted in recognition of four separate dispersal episodes within GABI (Woodburne 2010). These iterative dispersal events are thought to have been facilitated by fluctuations in eustatic sea level and oscillating climatic conditions, which would have rendered Central America more or less porous to invasive taxa hailing from higher latitudes in either North or South America. At times, Central America and adjacent southern Mexico functioned as a “holding pen” where various taxa were confined for significant intervals of time before proceeding to colonize higher latitudes (Woodburne 2010). Here, we propose that the concept of a holding pen also applies to Balkanatolia during the late Eocene and early Oligocene, thereby posing a biogeographically simpler hypothesis than that of Mennecart *et al.* (2018b) to explain the iterative pattern of dispersal observed in western Europe during this interval.

The new data presented here and by Licht *et al.* (2022) show that Balkanatolia was colonized by multiple clades of Asian mammals by the late Eocene. Although these late Eocene Asian mammals were virtually on Europe’s doorstep, with the notable exception of *Elomeryx* and *Phaneromeryx* (Table 3), very few of them were immediately able to overcome the biogeographic barrier posed by remnant peri-Alpine seaways blocking unfettered access to western Europe. Some taxa, including brontotheres and eosimiid anthropoids, apparently never colonized western Europe, either succumbing to extinction or being regionally extirpated before leaving the Balkanatolian holding pen. Indeed, Balkanatolia seems to have functioned as a biogeographic cul-de-sac for many of its late Eocene mammalian taxa that never reached Western Europe (Table 3). Most of the taxa that first appeared in Western Europe at MP 21 have no direct ancestry among the late Eocene taxa known so far from Balkanatolia, thus suggesting that another pathway was used for these MP 21 newcomers to reach Western Europe. However, we cannot rule out that this hypothesis may be biased by the relatively poor fossil data currently available for the late Eocene faunas from Balkanatolia. The impact of the Eocene-Oligocene boundary on the Balkanatolian mammal fauna is difficult to evaluate because earliest Oligocene faunas are poorly represented there, with the exception of an early Oligocene rodent assemblage from Serbia (de Bruijn *et al.* 2018, 2019, 2023; Marković *et al.* 2018; Wessels *et al.* 2020). However, as in the case of GABI, the Balkanatolian holding pen must have been porous to some degree, and its proximity to Western Europe may explain most, if not all, of the iterative dispersal events outlined in Table 3, including the Grande Coupure. The Balkanatolian holding pen continued to function into the early Oligocene, as suggested by the recent discovery of diatomyid rodents from the early Oligocene of Southeastern Serbia (Marković *et al.* 2018), a family that is common in Oligocene faunas of Pakistan (Marivaux & Welcomme 2003), India (Nanda &

TABLE 3. — First appearance of mammal genera in Balkanatolia and Western Europe around the Grande Coupure. Four phases can be differentiated in the faunal turnover that occurred during the Eocene-Oligocene transition. Taxa in **bold** are those reported from Sungülü. References refer to recent publications regarding systematic revision or updated occurrences of each taxon.

Phase	First appearance		Family	Reference	Locality
	Western Europe	Balkanatolia			
Phase 0					
Bartonian (MP16-17?)	Hyaenodontinae [Solé <i>et al.</i> (2022)]	<i>Sivatitanops</i>	Brontotheriidae	Nikolov & Heissig (1985)	Kameno, Tchernomore
		<i>Cadurcodon</i>	Amyndodontidae	Nikolov & Heissig (1985), Tissier <i>et al.</i> (2018)	Kameno
		<i>Elomeryx</i> <i>Bakalovia</i>	Anthracotheriidae Anthracotheriidae	Nikolov (1967) Nikolov & Heissig (1985)	Tchernomore Tchernomore
Phase 0-1					
Localities that could be either Bartonian or Priabonian		<i>Prohyracodon</i>	Hyracodontidae	Heissig (1990)	Mottnig
		<i>Anthracohyus</i>	Anthracotheriidae	Heissig (1990)	Mottnig
		<i>Amyndodontopsis</i>	Amyndodontidae	Tissier <i>et al.</i> (2018)	Dorog
		<i>Amyndodon</i>	Amyndodontidae	Tissier <i>et al.</i> (2018)	Tapioszele
		<i>Prominatherium</i>	Anthracotheriidae	Grandi & Bona (2017)	Mt Promina
		<i>Forstercooperia</i> <i>Bachitherium</i>	Hyracodontidae Bachitheriidae	Mennecart <i>et al.</i> (2018a, b) Mennecart <i>et al.</i> (2018a, b)	Bobov Dol Thrace Basin
Phase 1					
MP18-20 (early late Eocene)	<i>Elomeryx</i> [de Bonis (1964)] <i>Phaneromeryx</i> [Métais & Vislobokova (2007)] <i>Cynodictis</i> [Le Verger <i>et al.</i> 2020]	cf. <i>Bachitherium</i> sp.	Bachitheriidae	this study	Sungulu
		<i>Iberomeryx</i>	Tragulidae	this study	Sungulu
		<i>Sellamynodon</i>	Amyndodontidae	Tissier <i>et al.</i> (2018)	Dobarca
		<i>Embolotherium</i>	Brontotheriidae	Licht <i>et al.</i> (2022)	Büyükteflekk
		<i>Brachydiastematherium</i>	Brontotheriidae	Mihlbachler <i>et al.</i> (2004)	Mera
		<i>Debruijnia</i>	Spalacidae	de Bruijn <i>et al.</i> (2023)	Zvonce
		<i>Bustrania</i>	Muroidea	de Bruin <i>et al.</i> (2019)	Buštranje
		<i>Mogilia</i>	Muroidea	Wessels <i>et al.</i> (2018)	Buštranje, Zvonce
		<i>Pseudocricetodon</i>	Muroidea	de Bruijn <i>et al.</i> (2003, 2018); Baciú & Hartenberger (2001)	Sungülü, Buštranje
		<i>Paracricetodon</i>	Muroidea	de Bruijn <i>et al.</i> (2003, 2018)	Sungülü, Buštranje
		<i>Witenia</i>	Muroidea	de Bruijn <i>et al.</i> (2003, 2018, 2019)	Sungülü, Buštranje
		<i>Edirnella</i>	Muroidea	de Bruijn <i>et al.</i> (2003, 2018)	Sungülü, Zvonce
		<i>Eucricetodon</i>	Muroidea	de Bruijn <i>et al.</i> (2003); Baciú & Hartenberger (2001)	Sungülü
		cf. <i>Lignitella</i>	Muroidea	de Bruijn <i>et al.</i> (2003)	Sungülü
		<i>Heosminthus</i>	Dipodidae	de Bruijn <i>et al.</i> (2003)	Sungülü
		<i>Ottomania</i>	Baluchimyine	de Bruijn <i>et al.</i> (2003)	Sungülü
<i>Confiniummys</i>	Baluchimyine	de Bruijn <i>et al.</i> (2003)	Sungülü		
<i>Sungulusimias</i>	Eosimiidae	this study	Sungülü		
Phase 2					
MP21 (earliest Oligocene)	<i>Gelocus</i>	–	Geolocidae	Blondel (2001)	–
	<i>Paragelocus</i>	–	Geolocidae	Blondel (2001)	–
	<i>Pseudogelocus</i>	–	Geolocidae	Blondel (2001)	–
	<i>Anthracotherium</i>	–	Anthracotheriidae	Pandolfi <i>et al.</i> (2016)	–
	<i>Anthracochœrus</i>	–	Anthracotheriidae	Pandolfi <i>et al.</i> (2016)	–
	<i>Bothriodon</i>	–	Anthracotheriidae	Brunet & Vianey-Liaud (1987)	–
	<i>Propalaeochoerus</i>	–	Suidae	Pandolfi <i>et al.</i> (2016)	–
	<i>Cadurcotherium</i>	–	Amyndodontidae	Tissier <i>et al.</i> (2018)	–
	<i>Eggysodon</i>	–	Hyracodontidae	Brunet (1979); Becker (2009)	–
	<i>Ronzotherium</i>	–	Rhinocerotidae	Brunet (1979); Tissier <i>et al.</i> (2021)	–
	<i>Epiaceratherium</i>	–	Rhinocerotidae	Pandolfi <i>et al.</i> (2016)	–
	<i>Entelodon antiquus</i>	–	Entelodontidae	Brunet (1979)	–
	<i>Feliformia</i>	–	Carnivora	Solé <i>et al.</i> (2022)	–
	<i>Caniformia</i>	–	Carnivora	Solé <i>et al.</i> (2022)	–
	<i>Eucricetodon</i>	–	Muroidea	Dienemann (1987); Freudenthal (1988)	–
	<i>Pseudocricetodon</i>	–	Muroidea	Dienemann (1987)	–
	<i>Eomys</i>	–	Eomyidae	Bosma <i>et al.</i> (2023)	–
	<i>Steneofiber</i>	–	Castoriidae	Smith (2003)	–
	<i>Ephemerolagus</i>	–	Lagomorpha	Vianey-Liaud & Lebrun (2013)	–
	<i>Percygale</i>	–	Talpidae	Huguéney & Maridet (2018)	–
<i>Oligonyctia</i>	–	Nyctitheriidae	Smith & van den Hoek Ostende (2006)	–	

TABLE 3. — Continuation.

Phase	First appearance		Family	Reference	Locality
	Western Europe	Balkanatolia			
	<i>Belgicasorex</i>	–	Heterosoricidae	Smith & van den Hoek Ostende (2006)	–
	<i>Butselia</i>	–	Plesiosoricidae	Smith & Smith (2012)	–
	<i>Tetracus</i>	–	Erinaceidae	Smith (2004)	–
Phase 3					–
MP22 (early Oligocene)	<i>Lophiomeryx</i>	–	Lophiomerycidae	Brunet & Sudre (1987); Fejfar (1987)	–
	<i>Paenanthracotherium</i>	–	Anthracotheeriidae	Scherler <i>et al.</i> (2018)	–
Phase 4					–
MP23 (early Oligocene)	<i>Doliochoerus</i>	–	Suidae	Sudre (1995)	–
	<i>Iberomeryx</i>	–	Tragulidae	Mennecart <i>et al.</i> (2011)	–
	<i>Bachitherium</i>	–	Bachitheriidae	Sudre (1986); Blondel (2001)	–

Sahni 1998) and Thailand (Marivaux *et al.* 2004), but which is lacking from Oligocene faunas from the Thrace Basin of Turkey and from Kazakhstan. Likewise, diatomyids are absent from the abundant rodent fauna recovered from Süngülü, suggesting either taphonomic/paleoecological biases within Balkanatolia or relatively late colonization of this terrane by diatomyids during the early Oligocene (after the time of deposition of Süngülü).

Although paleobiogeographic obstacles remained, a leaky Balkanatolian holding pen may be preferable to dispersal across high latitudes for those Asian mammal taxa that colonized western Europe during the Grande Coupure. From a purely paleontological perspective, there is virtually no record of late Eocene mammals across the broad geographic expanse from western Europe to China and Mongolia, making this dispersal pathway theoretically possible but entirely conjectural. This dearth of paleontological data contrasts with the broad range of late Eocene Asian mammals in the Balkanatolian holding pen (Table 3). Moreover, recent geological assessments of Eurasian paleogeography highlight the precocious (Priabonian) drying out of the Turgai Strait (Kaya *et al.* 2019), an event that has been considered pivotal for establishing a northern dispersal pathway for Asian mammals to colonize western Europe at the Grande Coupure (Mennecart *et al.* 2018b). As Licht *et al.* (2022) emphasized, if high-latitude dispersal between Asia and western Europe was insignificant during the late Eocene when climate was relatively mild, it is difficult to conceive how dispersal across this pathway would have been significantly enhanced during the Grande Coupure, when central Asia would have been cooler and drier (Oi-1, Zachos *et al.* 2001; Barbolini *et al.* 2020). Likewise, regardless of the timing of regression of the Turgai Strait, a second marine barrier to dispersal persisted between western Europe and central Asia in the form of an epicontinental seaway that connected the North Sea with Paratethys across what is now the Baltic region (see Kaya *et al.* 2019: fig. 12).

The Süngülü fauna lacks the endemic elements known from older (Lutetian) sites of central Anatolia, where the most iconic Balkanatolian mammals have been described

(Métais *et al.* 2018); and the site postdates the proposed age for the end of Balkanatolian endemism (*c.* 40–38 Ma; Licht *et al.* 2022). The Süngülü fauna shows clear Asian affinities and corroborates regional exchanges between East and southern Asia and eastern Anatolia during the late Eocene. This assemblage therefore corroborates the existence of a dispersal route between East Asia and Europe along the Neotethys via the Middle East and Balkanatolia, although late Eocene mammals from Balkanatolia remain highly dissimilar at the generic level from those known in MP 21 in Western Europe (Table 3). This dispersal route has previously been invoked to explain the appearance of Asian taxa in the Balkans prior to the Grande Coupure (Heissig 1979; Böhme *et al.* 2013). Several large or medium-sized mammal families that arrived in Central or Western Europe after the Grande Coupure were already present in Balkanatolia during the late Eocene, including ruminants (Mennecart *et al.* 2018a, b), rhinocerotoids (Nikolov & Heissig 1985; Tissier *et al.* 2018; Licht *et al.* 2022), anthracotheriids (Böhme *et al.* 2013), and Hyaenodonta (Solé *et al.* 2022). This is also true for small mammals, especially muroids which are known from Süngülü and various late Eocene localities in the Balkans (Baciu & Hartenberger 2001; de Bruijn *et al.* 2018, 2019).

Finally, the presence of an anthropoid primate in the Süngülü fauna, showing affinities with *Phileosimias* and related taxa from the Eocene and Oligocene of South Asia, indicates some degree of continuity for forested ecosystems along the Eurasian Neotethyan margin. This continuity is consistent with palynological data (Kayseri Özer 2017), but at odds with late Eocene climate simulations indicating desertic conditions along a significant part of the Neotethyan domain (Tardif *et al.* 2021). We suggest that coastal mangrove forests, which can develop despite dry conditions inland, could have played a significant role in favoring the dispersal of arboreal mammals from East Asia to Anatolia. This is also consistent with the occurrence in Süngülü of small tragulids which are currently restricted to tropical forested environments (Meijaard & Groves 2004). Interestingly, the end of Balkanatolian endemism, dated at 40–38 Ma, is roughly synchronous with the arrival of eosimiiform primates in

Africa (Jaeger *et al.* 2010). We thus suggest that eosimiiform primates might have arrived in Anatolia as early as 40–38 Ma and used Balkanatolia as a stepping stone to pursue their dispersal to Africa. Precisely how and when they reached Africa from Balkanatolia remains unknown.

CONCLUSION

New paleontological data from the latest Eocene-earliest Oligocene of Balkanatolia reveal a complex biogeographic province that remained separated from Europe until the later Paleogene while being connected to Asia. Regional small mammal assemblages indicate that major differences in faunal composition between western Europe and Balkanatolia persisted until the early part of the late Oligocene (van der Weerd *et al.* 2021, 2023), suggesting the persistence of a substantial geographic (and possibly climatic) barrier, related to the persistence of several seaways in eastern Europe and/or the early growth of the Alpine orogeny. The assembly of the Balkanatolian paleobiogeographic province during the early Paleogene remains an enigma and its demise is barely better documented. In that regard, the fauna from Süngülü offers an important snapshot for documenting the Eocene-Oligocene transition in Eastern Anatolia, along a strategic corridor for faunal exchanges between Europe, South and East Asia. Intensive screen washing efforts carried out at Süngülü has produced remains of additional and so far unstudied taxa that may provide supplementary biochronologic or paleoenvironmental data. According to the data provided by de Bruijn *et al.* (2003), including the nature of the lithological sequence containing the fossil locality, the application of modern geochronological methods to this section may help constrain the age of Süngülü.

Acknowledgements

We thank Hans de Bruijn, who invited the first author to Utrecht in 2017 to study the artiodactyl material from Süngülü, which also led to the recognition of the primate tooth. The fossils described in this work were collected by Hans de Bruijn, Gerçek Saraç, Engin Ünay, Ali Yılmaz, Fadime Suata-Alpaslan, Jes de Bruijn and Wilma Wessels. Funding has been provided by National Geographic CRE GRANT #9215-12 to GM, National Science Foundation grant #2141115 to KCB, and the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 101043268) to AL. GM warmly thanks Wilma Wessels (University of Utrecht) for various help including access to specimens. We thank F. Sénégas, N. Poulet-Crovoisier, F. Goussard, and L. Cazes (all from CR2P) for photos and surface scans. We also thank the editor-in-chief, Michel Laurin, the associated editor, Lorenzo Rook, and referees, including P.-O. Antoine and L. Marivaux (ISEM, Montpellier), B. Mennecart (Basel Museum) and an anonymous reviewer for their useful comments that helped improve the manuscript.

REFERENCES

- ABBASSI N., ALINASIRI S. & LUCAS S. G. 2016. — New localities of Late Eocene vertebrate footprints from the Tarom Mountains, Northwestern Iran. *Historical Biology* 29 (7): 1–20. <https://doi.org/10.1080/08912963.2016.1267162>
- ADAMIA S., ALANIA V., CHABUKIANI A., CHICHUA G., ENUKIDZE O. & SADRADZE N. 2010. — Evolution of the Late Cenozoic basins of Georgia (SW Caucasus): a review. *Geological Society, London, Special Publications* 340 (1): 239–259. <https://doi.org/10.1144/SP340.11>
- BACIU C. & HARTENBERGER J.-L. 2001. — Un exemple de corrélation marin-continental dans le Priabonien de Roumanie. Remarques sur la Grande Coupure. *Comptes Rendus de l'Académie des Sciences - Series IIA - Earth and Planetary Science* 333 (8): 441–446. [https://doi.org/10.1016/S1251-8050\(01\)01659-7](https://doi.org/10.1016/S1251-8050(01)01659-7)
- BARBOLINI N., WOUTERSEN A., DUPONT-NIVET G., SILVESTRO D., TARDIF D., COSTER P. M. C., MEIJER N., CHANG C., ZHANG H. X., LICHT A., RYDIN C., KOUTSODENDRIS A., HAN F., ROHRMANN A., LIU X. J., ZHANG Y., DONNADIEU Y., FLUTEAU F., LADANT J. B., LE HIR G. & HOORN C. 2020. — Cenozoic evolution of the steppe-desert biome in Central Asia. *Science Advances* 6 (41): eabb8227. <https://doi.org/10.1126/sciadv.abb8227>
- BARMANN E. V. & RÖSSNER G. E. 2011. — Dental nomenclature in Ruminantia: Towards a standard terminological framework. *Mammalian Biology* 76 (6): 762–768. <https://doi.org/10.1016/j.mambio.2011.07.002>
- BARRIER E., VRIELYNCK B., BROUILLET J.-F. & BRUNET M.-F. 2018. — *Palaeotectonic maps of the Middle East. Tectono-sedimentary-palaeospastic maps from Late Norian to Pliocene. Atlas of 20 maps.* CCGM/CGMW, Paris.
- BEARD K. C. & WANG J. 2004. — The eosimiid primates (Anthropoidea) of the Heti Formation, Yuanqu Basin, Shanxi and Henan Provinces, People's Republic of China. *Journal of Human Evolution* 46 (4): 401–432. <https://doi.org/10.1016/j.jhevol.2004.01.002>
- BEARD K. C., QI T., DAWSON M. R., WANG B. & LI C. 1994. — A diverse new primate fauna from middle Eocene fissure-fillings in southeastern China. *Nature* 368: 604–609. <https://doi.org/10.1038/368604a0>
- BEARD K. C., TONG Y., DAWSON M. R., WANG J. & HUANG X. 1996. — Earliest complete dentition of an anthropoid primate from the late middle Eocene of Shanxi Province, China. *Science* 272 (5258): 82–85. <https://www.jstor.org/stable/2890776>
- BEARD K. C., MÉTAIS G., OCAKOĞLU F. & LICHT A. 2021. — An omomyid primate from the Pontide microcontinent of north-central Anatolia: implications for sweepstakes dispersal of terrestrial mammals during the Eocene. *Geobios* 66–67: 143–152. <https://doi.org/10.1016/j.geobios.2020.06.008>
- BEARD K. C., COSTER P. M. C., OCAKOĞLU F., LICHT A. & MÉTAIS G. 2023. — Dental anatomy, phylogenetic relationships and paleoecology of *Orhaniyeia nauta* (Metatheria, Anatoliadelphidae), a Gondwanan component of the insular Eocene mammal fauna of Balkanatolia (north-central Turkey). *Journal of Mammalian Evolution* 30: 859–872. <https://doi.org/10.1007/s10914-023-09680-6>
- BECKER D. 2009. — Earliest record of rhinocerotoids (Mammalia: Perissodactyla) from Switzerland: systematics and biostratigraphy. *Swiss Journal of Geosciences* 102: 489–504. <https://doi.org/10.1007/s00015-009-1330-4>
- BIOCHROM'97 1997. — Tableaux de corrélations et synthèses, in AGUILAR J.-P., LEGENDRE S. & MICHAUX J. (eds), *Actes du congrès Biochrom'97*. École Pratique des Hautes Études, Montpellier: 769–805.
- BLONDEL C. 2001. — The Eocene-Oligocene ungulates from Western Europe and their environment. *Palaeogeography, Palaeoclimatology, Palaeoecology* 168 (1–2): 125–139. [https://doi.org/10.1016/S0031-0182\(00\)00252-2](https://doi.org/10.1016/S0031-0182(00)00252-2)

- BÖHME M., AIGSTORFER M., ANTOINE P.-O., APPEL E., HAVLIK P., MÉTAIS G., PHUC L. T., SCHNEIDER S., SETZER F., TAPPERT R., TRAN D. N., UHL D. & PRIETO J. 2013. — Na Duong (Northern Vietnam) – an exceptional window into Eocene ecosystems from South-East Asia. *Zitteliana* A53: 120-167. <https://doi.org/10.5282/ubm/epub.19019>
- BONIS L. DE 1964. — Étude de quelques mammifères du Ludien de la Débruge (Vaucluse). *Annales de Paléontologie* 50: 119-154.
- BOSMA A. A., BRUIJN H. DE & WESSELS W. 2023. — A large *Eomys antiquus* (Aymard, 1853) (Mammalia, Rodentia) from the early Oligocene sedimentary deposits at Bouldnor Cliff (Isle of Wight, England, UK). *Palaeobiodiversity and Palaeoenvironments* 103 (2): 447-464. <https://doi.org/10.1007/s12549-022-00535-1>
- BRUIJN H. DE, ÜNAY E., SARAÇ G. & YILMAZ A. 2003. — A rodent assemblage from the Eo/Oligocene boundary interval near Süngülü, Lesser Caucasus, Turkey. *Coloquios de Paleontología* Volumen Extraordinario, 1: 47-76.
- BRUIJN H. DE, MARKOVIĆ Z., WESSELS W., MILIVOJEVIĆ M. & VAN DE WEERD A. A. 2018. — Rodent faunas from the Paleogene of south-east Serbia. *Palaeobiodiversity and Palaeoenvironments* 98 (3): 441-458. <https://doi.org/10.1007/s12549-017-0305-0>
- BRUIJN H. DE, MARKOVIĆ Z., WESSELS W. & VAN DE WEERD A. A. 2019. — Pappocricetodontinae (Rodentia, Muridae) from the Paleogene of south-east Serbia. *Palaeobiodiversity and Palaeoenvironments* 99 (3): 511-526. <https://doi.org/10.1007/s12549-018-0343-2>
- BRUIJN H. DE, MARKOVIĆ Z., WESSELS W. & VAN DE WEERD A. A. 2023. — On the antiquity and status of the Spalacidae, new data from the late Eocene of south-East Serbia. *Palaeobiodiversity and Palaeoenvironments* 103 (2): 433-445. <https://doi.org/10.1007/s12549-022-00529-z>
- BRUNET M. 1979. — *Les grands mammifères chefs de file de l'immigration oligocène et le problème de la limite Eocène-Oligocène en Europe*. Éditions de la Fondation Singer-Polignac, Paris, 281 p.
- BRUNET M. & SUDRE J. 1987. — Évolution et systématique du genre *Lophiomeryx* Pomel 1853 (Mammalia, Artiodactyla). *Münchener Geowissenschaftliche Abhandlungen* (A) 10: 225-242.
- BRUNET M. & VIANEY-LIAUD M. 1987. — Mammalian reference levels MP 21-30. *Münchener Geowissenschaftliche Abhandlungen* 10: 30-31.
- DIENEMANN A. 1987. — Die Gattungen *Eucrietodon* und *Pseudocricetodon* (Rodentia, Mammalia) aus dem Oligozän Süddeutschlands. *Abhandlungen-Bayerische Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Klasse* 165: 1-158.
- FEJFAR O. 1987. — A lower Oligocene mammalian fauna from Detan and Dvorce, NW Bohemia, Czechoslovakia. *Münchener Geowissenschaftliche Abhandlungen* (A) 10: 253-264.
- FREUDENTHAL M. 1988. — Redescription of *Eucrietodon atavus* (Misonne, 1957) (Rodentia) from Hoogbutsel (Belgium). *Bulletin de l'Institut royal des sciences naturelles de Belgique* 58: 199-207.
- GABUNIA L. K. 1964. — Benara fauna of Oligocene vertebrates. "Metsniereba" Press: 1-267.
- GERAADS D., BOUVRAIN G. & SUDRE J. 1987. — Relations phylétiques de *Bachitherium* Filhol, ruminant de l'Oligocène d'Europe occidentale. *Palaeovertebrata* 17 (2): 43-73.
- GODINOT M. 2014. — Fossil record of the primates from the Paleocene to the Oligocene, in HENKE W. & TATTERSALL I. (eds), *Handbook of paleoanthropology*. Springer, Heidelberg, Berlin: 1137-1259. https://doi.org/10.1007/978-3-642-27800-6_68-1
- GRANDI F. & BONA F. 2017. — *Prominatherium dalmatinum* from the late Eocene of Grancona (Vicenza, NE Italy). The oldest terrestrial mammal of the Italian peninsula. *Comptes Rendus Palevol* 16 (7): 738-745. <https://doi.org/10.1016/j.crpv.2017.04.002>
- HEISSIG K. 1979. — Die hypothetische Rolle Südosteuropas bei den Säugetierwanderungen im Eozän und Oligozän. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1979: 83-96.
- HEISSIG K. 1990. — Ein Oberkiefer von *Anthracoehyus* (Mammalia, ?Artiodactyla) aus dem Eozän Jugoslawiens. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 30: 57-64.
- HUGUENEY M. & MARIDET O. 2018. — Evolution of Oligo-Miocene talpids (Mammalia, Talpidae) in Europe: focus on the genera Myxomysgale and Percymysgale n. gen. *Historical Biology* 30 (1-2): 267-275. <https://doi.org/10.1080/08912963.2017.1282477>
- JAEGER J.-J., THEIN T., BENAMMI M., CHAIMANEE Y., SOE A. N., LWIN T., TUN T., WAI S. & DUCROCQ S. 1999. — A new Primate from the middle Eocene of Myanmar and the early origin of anthropoids. *Science* 286 (5439): 528-530. <https://doi.org/10.1126/science.286.5439.528>
- JAEGER J.-J., BEARD K. C., CHAIMANEE Y., SALEM M., BENAMMI M., HLAL O., COSTER P., BILAL A. A., DURINGER P., SCHUSTER M., VALENTIN X., MARANDAT B., MARIVAUX L., MÉTAIS E., HAMMUDA O. & BRUNET M. 2010. — Late middle Eocene epoch of Libya yields earliest known radiation of African anthropoids. *Nature* 467 (7319): 1095-1098. <https://doi.org/10.1038/nature09425>
- KAPPELMAN J. W., MAAS M. C., SEN S., ALPAGUT B., FORTELIUS M. & LUNKKA J.-P. 1996. — A new early tertiary mammalian fauna from Turkey and its paleobiogeographic significance. *Journal of Vertebrate Paleontology* 16 (3): 592-595. <https://www.jstor.org/stable/4523748>
- KAYA M. Y., DUPONT-NIVET G., PROUST J.-N., ROPERCH P., BOUGEUIS L., MEIJER N., FRIELING J., FIORONI C., ÖZKAN ALTINER S., VARDAR E., BARBOLINI N., STOICA M., AMINOV J., MAMTIMIN M. & ZHAOJIE G. 2019. — Paleogene evolution and demise of the proto-Paratethys Sea in Central Asia (Tarim and Tajik basins): Role of intensified tectonic activity at ca. 41 Ma. *Basin Research* 31 (3): 461-486. <https://doi.org/10.1111/bre.12330>
- KAYSERİ ÖZER M. S., SANÇAY R. H., ŞEN Ş., SÖZERİ K., MÉTAIS G., AYYILDIZ T. & VAROL B. 2017. — Paleoenvironment of the Late Oligocene from the Kağızman-Tuzluca Basin (northeastern Anatolia) based on the micro- and macrofloras. *Turkish Journal of Earth Sciences* 26 (3): 227-248. <https://doi.org/10.3906/yer-1612-26>
- LE VERGER K., SOLÉ F. & LADEVÈZE S. 2020. — Description of a new species of *Cynodictis* Bravard & Pomel, 1850 (Carnivora, Mammalia) from the Quercy Phosphorites with comments on the use of skull morphology for phylogenetics. *Geodiversitas* 42 (16): 239-255. <https://doi.org/10.5252/geodiversitas2020v42a16>
- KÖHLER M. & MOYÀ-SOLÀ S. 1999. — A finding of Oligocene primates on the European continent. *Proceedings of National Academy of Science* 96 (25): 14664-14667. <https://doi.org/10.1073/pnas.96.25.14664>
- LICHT A., MÉTAIS G., COSTER P., İBİLİOĞLU D., OCAKOĞLU F., WESTERWEEL J., MUELLER M., CAMPBELL C., MATTINGLY S., WOOD M. C. & BEARD K. C. 2022. — Balkanatolia: The insular mammalian biogeographic province that partly paved the way to the Grande Coupure. *Earth-Science Reviews* 226: 103929. <https://doi.org/10.1016/j.earscirev.2022.103929>
- LUCAS S. G. & EMRY R. J. 1999. — Taxonomy and biochronological significance of *Paraentelodon*, a giant entelodont (Mammalia, Artiodactyla) from the Late Oligocene of Eurasia. *Journal of Vertebrate Paleontology* 19 (1): 160-168. <https://doi.org/10.1080/002724634.1999.10011130>
- MARIVAUX L. & WELCOMME J.-L. 2003. — New diatomyid and baluchimyine rodents from the Oligocene of Pakistan (Bugti Hills, Balochistan): systematic and paleobiogeographic implications. *Journal of Vertebrate Paleontology* 23 (2): 420-434. <https://doi.org/10.1016/j.jvpa.2003.03.002>
- MARIVAUX L., CHAIMANEE Y., YAMEE C., SRISUK P. & JAEGER J.-J. 2004. — Discovery of *Fallomus ladakhensis* Nanda & Sahni, 1998 (Mammalia, Rodentia, Diatomyidae) in the lignites of Nong Ya Plong (Phetchaburi Province, Thailand): systematic, biochronological and paleoenvironmental implications. *Geodiversitas* 26 (3): 493-507.

- MARIVAUX L., ANTOINE P.-O., BAQRI S. R. H., BENAMMI M., CHAIMANEE Y., CROCHET J.-Y., FRANCESCHI D. DE, IQBAL N., JAEGER J.-J., MÉTAIS G., ROOHI G. & WELCOMME J.-L. 2005. — Anthropoid primates from the Oligocene of Pakistan (Bugti Hills): Data on early anthropoid evolution and biogeography. *Proceedings of National Academy of Science, USA* 102 (24): 8436-8441. <https://doi.org/10.1073/pnas.0503469102>
- MARIVAUX L., ESSID E. M., MARZOUGUI W., KHAYATI AMMAR H., ADNEN S., MARANDAT B., MERZERAUD G., RAMDARSHAN A., TABUCE R., VIANEY-LIAUD M. & YANS J. 2014. — A morphological intermediate between eosimiiform and simiiform primates from the late middle Eocene of Tunisia: Macroevolutionary and paleobiogeographic implications of early anthropoids. *American Journal of Physical Anthropology* 154 (3): 387-401. <https://doi.org/10.1002/ajpa.22523>
- MARKOVIĆ Z., WESSELS W., VAN DE WEERD A. A. & BRUIJN H. DE 2018. — On a new diatomyid (Rodentia, Mammalia) from the Paleogene of south-east Serbia, the first record of the family in Europe. *Palaeobiodiversity and Palaeoenvironments* 98 (3): 459-469. <https://doi.org/10.1007/s12549-017-0301-4>
- MARTINEZ J.-N. & SUDRE J. 1995. — The astragalus of Paleogene artiodactyls: comparative morphology, variability and prediction of body mass. *Lethaia* 28 (3): 197-209. <https://doi.org/10.1111/j.1502-3931.1995.tb01423.x>
- MEIJAARD E. & GROVES C. P. 2004. — A taxonomic revision of the Tragulus mouse-deer (Artiodactyla). *Zoological Journal of the Linnean Society* 140 (1): 63-102. <https://doi.org/10.1111/j.1096-3642.2004.00091.x>
- MENNECART B. & MÉTAIS G. 2015. — *Mosaicomeryx quercyi*, new comb.: ruminant from the Oligocene of Europe, and the significance of “gelocids”. *Journal of Systematic Palaeontology* 13 (7): 581-600. <https://doi.org/10.1080/14772019.2014.948505>
- MENNECART B., BECKER D. & BERGER J.-P. 2011. — *Iberomeryx minor* (Mammalia, Artiodactyla) from the Early Oligocene of Soulce (Canton Jura, NW Switzerland): systematics and palaeodiet. *Swiss Journal of Geosciences* 104 (Suppl. 1): 115-132. <https://doi.org/10.1007/s00015-010-0034-0>
- MENNECART B., GERAADS D., SPASSOV N. & ZAGORCHEV I. 2018a. — Discovery of the oldest European ruminant in the late Eocene of Bulgaria: did tectonics influence the diachronic development of the Grande Coupure? *Palaeogeography, Palaeoclimatology, Palaeoecology* 498: 1-8. <https://doi.org/10.1016/j.palaeo.2018.01.011>
- MENNECART B., RADOVIĆ P. & MARKOVIĆ Z. 2018b. — New data on the earliest European ruminant (Mammalia, Artiodactyla): A revision of the fossil mandible from Rusce in the Pčinja basin (late Eocene, Southeastern Serbia). *Palaeontologia Electronica* 21.3.38A: 1-12. <https://doi.org/10.26879/883>
- MENNECART B., AIGLSTORFER M., LI Y. & WANG S.-Q. 2021. — Ruminants reveal Eocene Asiatic palaeobiogeographical provinces as the origin of diachronous mammalian Oligocene dispersals into Europe. *Scientific Report* 11: 17710. <https://doi.org/10.1038/s41598-021-96221-x>
- MÉTAIS G. & VISLOBOKOVA I. A. 2007. — Basal ruminants, in PROTHERO D. R. & FOSS S. E. (eds), *The evolution of artiodactyls*. The Johns Hopkins University Press, Baltimore: 189-212.
- MÉTAIS G., CHAIMANEE Y., JAEGER J.-J. & DUCROCQ S. 2001. — New remains of primitive ruminants from Thailand: evidence of the early evolution of the Ruminantia in Asia. *Zoologica Scripta* 30 (4): 231-249. <https://doi.org/10.1046/j.0300-3256.2001.00071.x>
- MÉTAIS G., WELCOMME J.-L. & DUCROCQ S. 2009. — Lophiomerycid ruminants from the Oligocene of the Bugti Hills (Balochistan, Pakistan). *Journal of Vertebrate Paleontology* 29 (1): 1-12. <https://doi.org/10.1671/039.029.0115>
- MÉTAIS G., SEN S., SÖZERI K., PEIGNÉ S. & VAROL B. 2015. — Late Paleogene terrestrial fauna and paleoenvironments in Eastern Anatolia: New insights from the Kağızman-Tuzluca Basin. *Journal of Asian Earth Sciences* 107: 96-109. <https://doi.org/10.1016/j.jseas.2015.03.048>
- MÉTAIS G., ALBAYRAK E., ANTOINE P.-O., ERDAL O., KARADENIZLI L., OYAL N., SARAÇ G., İSLAMOĞLU Y. & SEN S. 2016. — Oligocene ruminants from the Kızılırmak Formation, Çankırı-Çorum Basin, Central Anatolia, Turkey. *Palaeontologia Electronica* 19.3.37A: 1-23. <https://hal.umontpellier.fr/hal-01922849>
- MÉTAIS G., ERDAL O., ERTURAC K. & BEARD K. C. 2017. — Tarsal morphology of the pleuraspidotheriid mammal *Hilalia* from the middle Eocene of Turkey. *Acta Palaeontologica Polonica* 62 (1): 173-179. <https://doi.org/10.4202/app.00314.2016>
- MÉTAIS G., COSTER P. M., KAPPELMAN J. R., LICHT A., OCAKOĞLU F., TAYLOR M. H. & BEARD K. C. 2018. — Eocene metatherians from Anatolia illuminate the assembly of an island fauna during Deep Time. *PLoS ONE* 13 (11): e0206181. <https://doi.org/10.1371/journal.pone.0206181>
- MILNE-EDWARDS A. 1864. — Recherches anatomiques, zoologiques, et paléontologiques sur la famille des chevrotains. *Annales des Sciences Naturelles, Paris* 5 (2): 1-56. <https://doi.org/10.5962/bhl.title.82492>
- MIVART S. G. 1864. — Notes on the crania and dentition of the Lemuridae. *Proceedings of the Zoological Society London* 1865: 836-844.
- MOTTL M. 1961. — Die Dorcatherien der steiermark. *Mitteilungen, Museum Bergbau, Geologie und Technik, Graz* 22: 21-71.
- NANDA A. C. & SAHNI A. 1998. — Ctenodactyloid Rodent assemblage from Kargil Formation, Ladakh molasse Group: age and palaeobiogeographic implications for the Indian Subcontinent in the Oligo-Miocene. *Geobios* 31 (4): 533-544. [https://doi.org/10.1016/S0016-6995\(98\)80124-1](https://doi.org/10.1016/S0016-6995(98)80124-1)
- NI X., LI Q., LI L. & BEARD K. C. 2016. — Oligocene primates from China reveal divergence between African and Asian primate evolution. *Science* 352 (6286): 673-677. <https://doi.org/10.1126/science.aaf2107>
- NIKOLOV I. 1967. — Neue obereozäne Arten der Gattung *Elomeryx*. *Neues Jahrbuch für Geologie und Paläontologie* 128 (2): 205-214.
- NIKOLOV I. & HEISSIG K. 1985. — Fossile Säugetiere aus dem Obereozän und Unteroligozän Bulgariens und ihre Bedeutung für die Palaeogeographie. *Mitteilungen des Bayerische Staatssammlung für Paläontologie und historische Geologie* 25: 61-79.
- OWEN R. 1848. — Description of teeth and portions of jaws of two extinct anthracotherioid quadrupeds (*Hyopotamus vectianus* and *Hyop. bovinus*) discovered by the Marchioness of Hastings in the Eocene deposits on the N.W. coast of the Isle of Wight: with an attempt to develop Cuvier's idea of the classification of pachyderms by the number of their toes. *Quarterly Journal of the Geological Society of London* 4: 103-141. <https://doi.org/10.1144/GSL.JGS.1848.004.01-02.21>
- PALCU D. V. & KRIJGSMAN W. 2023. — The dire straits of Paratethys: gateways to the anoxic giant of Eurasia. *Geological Society, London, Special Publications* 523 (1): SP523-2021-2073. <https://doi.org/10.1144/SP523-2021-73>
- PANDOLFI L., CARNEVALE G., COSTEUR L., FAVERO L. D., FORNASIERO M., GHEZZO E., MAIORINO L., MIETTO P., PIRAS P., ROOK L., SANSALONE G. & KOTSAKIS T. 2016. — Reassessing the earliest Oligocene vertebrate assemblage of Monteviale (Vicenza, Italy). *Journal of Systematic Palaeontology* 15 (2): 1-45. <https://doi.org/10.1080/14772019.2016.1147170>
- POPOV S. V., RÖGL F., ROZANOV A. Y., STEININGER F. F., SHCHERBA I. G. & KOVAC M. 2004. — Lithologic-Paleogeographic maps of Paratethys. *Courier Forschungsinstitut Senckenberg* 250: 1-46.
- ROSE K. D. & BOWN T. M. 1991. — Additional fossil evidence on the differentiation of the earliest euprimates. *Proceedings of the National Academy of Sciences* 88 (1): 98-101. <https://doi.org/10.1073/pnas.88.1.98>
- RÖSSNER G. E. 2007. — Family Tragulidae, in PROTHERO D. R. & FOSS S. E. (eds), *The Evolution of Artiodactyls*. Johns Hopkins University Press, Baltimore: 213-220.

- SCHERLER L., LIHOREAU F. & BECKER D. 2018. — To split or not to split *Anthracotherium*? A phylogeny of Anthracotheriinae (Cetartiodactyla: Hippopotamoidea) and its palaeobiogeographical implications. *Zoological Journal of the Linnean Society* 185 (2): 487-510. <https://doi.org/10.1093/zoolinnean/zly052>
- SCOPOLI G. A. & GERLE W. A. 1777. — *Introductio ad historiam naturalem sistens genera lapidum, plantarum et animalium hactenus detecta, caracteribus essentialibus donata, in tribus divisa, subinde ad leges naturae*. Apud Wolfgangum Gerie, Prague, 506 p. <https://doi.org/10.5962/bhl.title.10827>
- SEIFFERT E. R. 2012. — Early primate evolution in Afro-Arabia. *Evolutionary Anthropology* 21 (6): 239-253. <https://doi.org/10.1002/evan.21335>
- SEIFFERT E. R., SIMONS E. L., CLYDE W. C., ROSSIE J. B., ATTIA Y., BOWN T. M., CHATRATH P. & MATHISON M. E. 2005. — Basal anthropoids from Egypt and the antiquity of Africa's higher primate radiation. *Science* 310 (5746): 300-304. <https://doi.org/10.1126/science.1116569>
- SEN S., ANTOINE P.-O., VAROL B., AYYILDIZ T. & SÖZERI K. 2011. — Giant rhinoceros *Paraceratherium* and other vertebrates from Oligocene and middle Miocene deposits of the Kağızman-Tuzluca Basin, Eastern Turkey. *Naturwissenschaften* 98 (5): 407-423. <https://doi.org/10.1007/s00114-011-0786-z>
- ŞENGÖR A. M. C., ÖZEREN M. S., KESKIN M., SAKINÇ M., ÖZBAKIR A. D. & KAYAN I. 2008. — Eastern Turkish high plateau as a small Turkic-type orogen: Implications for post-collisional crust-forming processes in Turkic-type orogens. *Earth-Science Reviews* 90 (1-2): 1-48. <https://doi.org/10.1016/j.earsci-rev.2008.05.002>
- SMITH R. 2003. — Les vertébrés terrestres de l'Oligocène inférieur de Belgique (Formation de Borgloon, MP 21): inventaire et interprétation des données actuelles. *Coloquios de Paleontología* Vol. Ext. 1: 647-657.
- SMITH R. 2004. — Insectivores (Mammalia) from the earliest Oligocene (MP 21) of Belgium. *Netherlands Journal of Geosciences* 83 (3): 187-192. <https://doi.org/10.1017/S0016774600020254>
- SMITH R. & VAN DEN HOEK OSTENDE L. W. 2006. — A new heterosoricid shrew from the lowermost Oligocene of Europe. *Acta Palaeontologica Polonica* 51 (2): 381-384.
- SMITH R. & SMITH T. 2012. — The carnivoran-like insectivore *Butselia biveri* Quinet & Misonne, 1965 (Mammalia, Plesiosoricidae) from the lowermost Oligocene of Europe. *Spanish Journal of Palaeontology* 27 (2): 105-116.
- SOLÉ F., FISCHER V., LE VERGER K., MENNECART B., SPEIJER R. P., PEIGNÉ S. & SMITH T. 2022. — Evolution of European carnivorous mammal assemblages through the Palaeogene. *Biological Journal of the Linnean Society* 135 (4): 734-753. <https://doi.org/10.1093/biolinnean/blac002>
- STEHLIN H. G. 1909. — Remarques sur les faunules de mammifères de l'Eocène et de l'Oligocène du Bassin de Paris. *Bulletin de la Société géologique de France* 4 (9): 488-520.
- SUDRE J. 1986. — Le genre *Bachitherium* Filhol 1882 (Mammalia, Artiodactyla): diversité spécifique, phylogénie, extension chronologique. *Comptes Rendus de l'Académie des Sciences de Paris, série IIa* 303: 749-754.
- SUDRE J. 1995. — Le Garouillas et les sites contemporains (Oligocène, MP 25) des phosphorites du Quercy (Lot, Tarn-et-Garonne, France) et leurs faunes de vertébrés. *Palaeontographica* 236 (1-6): 205-256. <https://doi.org/10.1127/pala/236/1995/205>
- TARDIF D., TOUMOULIN A., FLUTEAU F., DONNADIEU Y., LE HIR G., BARBOLINI N., LICHT A., LADANT J.-B., SEPULCHRE P., VIOVY N., HOORN C. & DUPONT-NIVET G. 2021. — Orbital variations as a major driver of climate and biome distribution during the greenhouse to icehouse transition. *Science Advances* 7 (43): eabh2819. <https://doi.org/10.1126/sciadv.abh2819>
- TISSIER J., BECKER D., CODREA V., COSTEUR L., FÂRÇAŞ C., SOLOMON A., VENCZEL M. & MARIDET O. 2018. — New data on Aymnodontidae (Mammalia, Perissodactyla) from Eastern Europe: Phylogenetic and palaeobiogeographic implications around the Eocene-Oligocene transition. *PLoS ONE* 13 (4): e0193774. <https://doi.org/10.1371/journal.pone.0193774>
- TISSIER J., ANTOINE P.-O. & BECKER D. 2021. — New species, revision, and phylogeny of *Ronzotherium* Aymard, 1854 (Perissodactyla, Rhinocerotidae). *European Journal of Taxonomy* 753: 1-80. <https://doi.org/10.5852/ejt.2021.753.1389>
- VAN HINSBERGEN D. J. J., TORSVIK T. H., SCHMID S. M., MAÏENCO L. C., MAFFIONE M., VISSERS R. L. M., GÜRER D. & SPAKMAN W. 2020. — Orogenic architecture of the Mediterranean region and kinematic reconstruction of its tectonic evolution since the Triassic. *Gondwana Research* 81: 79-229. <https://doi.org/10.1016/j.gr.2019.07.009>
- VAN DE WEERD A. A., BRUIJN H. DE & WESSELS W. 2021. — New rodents from the late Oligocene site of Gözüyükzilli in Anatolia (Turkey). *Historical Biology* 33 (10): 2406-2431. <https://doi.org/10.1080/08912963.2020.1800682>
- VAN DE WEERD A. A., BRUIJN H. DE & WESSELS W. 2023. — A small assemblage of early Oligocene rodents and insectivores from the Sivas basin, Turkey. *Palaeobiodiversity and Palaeoenvironments* 103: 609-632. <https://doi.org/10.1007/s12549-022-00563-x>
- VIANEY-LIAUD M. & LEBRUN R. 2013. — New data about the oldest European lagomorpha: description of the new genus *Ephemeralagus nievae* gen. nov. et sp. nov. *Spanish Journal of Palaeontology* 28 (1): 3-16. <https://doi.org/10.7203/sjp.28.1.17812>
- WESSELS W., VAN DE WEERD A. A., BRUIJN H. DE & MARKOVIĆ Z. 2018. — New Melissodontinae (Mammalia, Rodentia) from the Paleogene of south-east Serbia. *Palaeobiodiversity and Palaeoenvironments* 98 (3): 471-487. <https://doi.org/10.1007/s12549-017-0311-2>
- WESSELS W., VAN DE WEERD A. A., BRUIJN H. DE & MARKOVIĆ Z. 2020. — Dipodidae (Mammalia, Rodentia) from the Paleogene of south-east Serbia. *Palaeobiodiversity and Palaeoenvironments* 100 (3): 841-848. <https://doi.org/10.1007/s12549-019-00392-5>
- WOODBURNE M. O. 2010. — The Great American Biotic Interchange: Dispersals, Tectonics, Climate, Sea Level and Holding Pens. *Journal of Mammalian Evolution* 17 (4): 245-264. <https://doi.org/10.1007/s10914-010-9144-8>
- YILMAZ A., ENGIN T., ADAMIA S. & LAZARASHVILI T. 2001. — *Geological Studies of the Area Along Turkish-Georgian Border*. MTA, Ankara, Turkey: 1-338.
- ZACHOS J. C., SHACKLETON N. J., REVENAUGH J. S., PALIKE H. & FLOWER B. P. 2001. — Climate Response to Orbital Forcing Across the Oligocene-Miocene Boundary. *Science* 292 (5515): 274-278. <https://doi.org/10.1126/science.1058288>

Submitted on 4 April 2023;
accepted on 11 July 2023;
published on 11 December 2023.