



HAL
open science

SINGULARITIES OF HOLOMORPHIC CODIMENSION ONE FOLIATIONS OF THE COMPLEX PROJECTIVE PLANE

Dominique Cerveau, Julie Déserti

► **To cite this version:**

Dominique Cerveau, Julie Déserti. SINGULARITIES OF HOLOMORPHIC CODIMENSION ONE FOLIATIONS OF THE COMPLEX PROJECTIVE PLANE. Boletim da Sociedade Brasileira de Matemática / Bulletin of the Brazilian Mathematical Society, 2023, 54 (2), 10.1007/s00574-023-00332-0 . hal-04335451

HAL Id: hal-04335451

<https://hal.science/hal-04335451>

Submitted on 11 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

SINGULARITIES OF HOLOMORPHIC CODIMENSION ONE FOLIATIONS OF THE COMPLEX PROJECTIVE PLANE

DOMINIQUE CERVEAU AND JULIE DÉSERTI

ABSTRACT. We prove that any holomorphic codimension 1 foliation on the complex projective plane has at most one singularity up to the action of an ad-hoc birational map. Consequently, any algebraic foliation on the affine plane has no singularities up to the action of a suitable birational self map of the complex projective plane into itself.

Let \mathcal{F} be a codimension one holomorphic foliation of degree N on $\mathbb{P}_{\mathbb{C}}^2$. Denote by $\pi: \mathbb{C}^3 \setminus \{0\} \rightarrow \mathbb{P}_{\mathbb{C}}^2$ the canonical projection. The homogeneous foliation $\pi^{-1}\mathcal{F}$ extends to \mathbb{C}^3 and is defined by a 1-form

$$\omega = a(x, y, z) dx + b(x, y, z) dy + c(x, y, z) dz$$

where a, b, c are homogeneous polynomials of degree $N + 1$ without common component satisfying the Euler identity: $ax + by + cz = 0$; it is the Chow theorem for foliations. The singular set $\text{Sing}(\mathcal{F})$ of \mathcal{F} is given by

$$\pi(\{a = b = c = 0\} \setminus \{0\}).$$

Let us recall what is the Milnor number $\mu(\mathcal{F}, m)$ of a foliation \mathcal{F} at a singular point m . Let us fix a local chart (u, v) such that $m = (0, 0)$. The germ of \mathcal{F} at m is defined, up to multiplication by a unit at 0, by a 1-form $E du + F dv$. Denote by $\langle E, F \rangle$ the ideal generated by E and F , then

$$\mu(\mathcal{F}, m) = \dim \frac{\mathbb{C}\{u, v\}}{\langle E, F \rangle};$$

it is also the multiplicity of intersection of the germs of curves $(E = 0)$ and $(F = 0)$. Let us recall that ([Bru15, Chapter 2, Section 3, Example 2])

$$\sum_{m \in \text{Sing}(\mathcal{F})} \mu(\mathcal{F}, m) = N^2 + N + 1;$$

in particular, \mathcal{F} has a least one singular point.

2020 *Mathematics Subject Classification.* 32S65.

Key words and phrases. Singularities of holomorphic foliations.

Example. Let us consider the diagonal linear foliation \mathcal{F} given by the 1-form

$$\omega = \lambda yz dx + xz dy - (1 + \lambda)xy dz$$

with $\lambda \in \mathbb{C} \setminus \{0, -1, -2\}$. Note that $\text{Sing}(\mathcal{F}) = \{(0:0:1), (1:0:0), (0:1:0)\}$; moreover, $x^\lambda yz^{-(1+\lambda)}$ is a first integral of \mathcal{F} .

The birational involution from $\mathbb{P}_{\mathbb{C}}^2$ into itself given by

$$\mathcal{I}_1: (x:y:z) \dashrightarrow (xy:y^2:x^2-yz),$$

defines an isomorphism of $\{y \neq 0\}$. Remark that the foliation $\mathcal{I}_1^{-1}\mathcal{F}$ is described by the 1-form

$$-y((2+\lambda)x^2 + \lambda yz) dx + x((2+\lambda)x^2 - yz) dy + (1+\lambda)xy^2 dz;$$

in particular, $\text{Sing}(\mathcal{I}_1^{-1}\mathcal{F}) = \{(0:0:1), (0:1:0)\}$.

Let us now consider the birational involution \mathcal{I}_2 from $\mathbb{P}_{\mathbb{C}}^2$ into itself given by

$$\mathcal{I}_2: (x:y:z) \dashrightarrow (x^2:-xy+z^2:xz);$$

note that \mathcal{I}_2 defines an isomorphism of $\{x \neq 0\}$. Furthermore, the restriction of $\mathcal{I}_2^{-1}\mathcal{I}_1^{-1}\mathcal{F}$ on $(x=0)$ is described by the 1-form $(1-\lambda)z^5 dx$; consequently, $\text{Sing}(\mathcal{I}_2^{-1}\mathcal{I}_1^{-1}\mathcal{F}) = \{(0:1:0)\}$.

Can we generalize this to all holomorphic foliations on $\mathbb{P}_{\mathbb{C}}^2$? The answer is yes:

Theorem 1. *Let \mathcal{F} be a holomorphic foliation on $\mathbb{P}_{\mathbb{C}}^2$. There exists a birational self map ϕ from $\mathbb{P}_{\mathbb{C}}^2$ into itself such that $\phi^{-1}\mathcal{F}$ has at most one singular point.*

Note that many properties are preserved by conjugacy by birational maps, for instance the existence of invariant algebraic curves of genus ≥ 2 , but also the existence of dense leaves.

Corollary 2. *Let \mathcal{F} be an algebraic foliation on \mathbb{C}^2 . There exists a birational map ϕ from \mathbb{C}^2 into itself such that $\phi^{-1}\mathcal{F}$ has no singular point.*

In a certain sense Corollary 2 indicates that algebraic foliations of \mathbb{C}^2 have no special properties.

Note that the situation is completely different on $\mathbb{P}_{\mathbb{C}}^1 \times \mathbb{P}_{\mathbb{C}}^1$; in that space there exist foliations without singularities: a foliation on $\mathbb{P}_{\mathbb{C}}^1 \times \mathbb{P}_{\mathbb{C}}^1$ birationally conjugate to a foliation without singularities has a rational first integral ([Bru15]).

To prove Theorem 1 we will use the following statement:

Lemma 3. *Let \mathcal{F} be a holomorphic codimension one foliation on $\mathbb{P}_{\mathbb{C}}^2$. Assume that \mathcal{F} has N singular points, and among them $n \geq 2$ singular points on a line L . There exists a birational self map ϕ from $\mathbb{P}_{\mathbb{C}}^2$ into itself such that $\phi^{-1}\mathcal{F}$ has $N - n + 1$ singular points, one on L and $N - n$ on $\mathbb{P}_{\mathbb{C}}^2 \setminus L$.*

Proof. \diamond Assume first that L is not invariant by \mathcal{F} . Let us choose coordinates such that

$$\begin{cases} L = (z = 0) \\ (0 : 1 : 0) \notin \text{Sing}(\mathcal{F}) \\ L \text{ is not tangent to the leaf of } \mathcal{F} \text{ through } (0 : 1 : 0) \end{cases} \quad (\star)$$

The foliation \mathcal{F} is defined by

$$\omega = A(x, y, z) dx + B(x, y, z) dy + C(x, y, z) dz.$$

Let us recall the Euler condition: $xA + yB + zC = 0$; in particular note that $B(0, 1, 0) = 0$. The tangency condition (\star) says that $A(0, 1, 0) \neq 0$.

Let us consider the birational involution ϕ of $\mathbb{P}_{\mathbb{C}}^2$ into itself given by

$$\phi : (x : y : z) \dashrightarrow (xz : -yz + x^2 : z^2);$$

note that

$$\text{Ind}(\phi) = \{(0 : 1 : 0)\}, \quad \text{Exc}(\phi) = \{(z = 0)\}$$

where $\text{Ind}(\phi)$ (resp. $\text{Exc}(\phi)$) denotes the set of indeterminacy (resp. the exceptional set) of ϕ . In particular, ϕ defines an isomorphism of $\{z \neq 0\}$. One has

$$\begin{aligned} \phi^* \omega &= A(xz, -yz + x^2, z^2)(x dz + z dx) \\ &\quad + B(xz, -yz + x^2, z^2)(-y dz - z dy + 2x dx) + 2C(xz, -yz + x^2, z^2)z dz, \end{aligned}$$

and

$$\phi^* \omega|_{z=0} = xA(0, x^2, 0) dx + \underbrace{B(0, x^2, 0)}_{=0} (-y dz + 2x dx) = xA(0, x^2, 0) dx;$$

in particular $\phi^* \omega|_{z=0}$ vanishes only at $x = 0$ since $A(0, 1, 0) \neq 0$. As a consequence, $\#(\text{Sing}(\phi^{-1} \mathcal{F}) \cap (z = 0)) = 1$ and $\#\text{Sing}(\phi^{-1} \mathcal{F}) = N - n + 1$.

\diamond Suppose now that L is invariant by \mathcal{F} . Let us choose coordinates such that $\begin{cases} L = (z = 0) \\ (0 : 1 : 0) \notin \text{Sing}(\mathcal{F}) \end{cases}$

The foliation \mathcal{F} is defined by

$$\omega = zA(x, y, z) dx + zB(x, y, z) dy + C(x, y, z) dz.$$

Let us still consider the birational involution ϕ of $\mathbb{P}_{\mathbb{C}}^2$ into itself given by

$$\phi : (x : y : z) \dashrightarrow (xz : -yz + x^2 : z^2)$$

that defines an isomorphism of $\{z \neq 0\}$. Then

$$\begin{aligned} \phi^* \omega &= z \left(zA(xz, -yz + x^2, z^2)(x dz + z dx) \right. \\ &\quad \left. + zB(xz, -yz + x^2, z^2)(-y dz - z dy + 2x dx) + 2C(xz, -yz + x^2, z^2) dz \right), \end{aligned}$$

and the restriction of $\phi^{-1}\mathcal{F}$ on $(z=0)$ is described by the 1-form $2C(0, x^2, 0) dz$. But $C(0, 1, 0) \neq 0$ (because $(0:1:0) \notin \text{Sing}(\mathcal{F})$) so $C(0, x^2, 0) = 0$ if and only if $x = 0$. In particular, $\#(\text{Sing}(\phi^{-1}\mathcal{F}) \cap (z=0)) = 1$ and $\#\text{Sing}(\phi^{-1}\mathcal{F}) = N - n + 1$. □

Proof of Theorem 1. We get the result by iteration using Lemma 3. □

Assume that \mathcal{F} is an algebraic foliation of any codimension of \mathbb{C}^n ; is it true that there exists a birational self map of $\mathbb{P}_{\mathbb{C}}^n$ into itself such that $\phi^{-1}\mathcal{F}$ has no singularity ? In the real case (for example in $\mathbb{P}_{\mathbb{R}}^2$) what is the best result we can expect ?

Remark that following [CDGBM10] in which we classify, up to automorphisms of $\mathbb{P}_{\mathbb{C}}^2$, the quadratic foliations having a unique singularity there is a lot of activity around the foliations of $\mathbb{P}_{\mathbb{C}}^2$ having a unique singularity (for instance [FPR22, Alc22, Cou22, CJ21, CF21, CF20]).

REFERENCES

- [Alc22] C. R. Alcántara. Special foliations on $\mathbb{C}\mathbb{P}^2$ with a unique singular point. *Res. Math. Sci.*, 9(1):Paper No. 15, 11, 2022.
- [Bru15] M. Brunella. *Birational geometry of foliations*, volume 1 of *IMPA Monographs*. Springer, Cham, 2015.
- [CDGBM10] D. Cerveau, J. Déserti, D. Garba Belko, and R. Mezzani. Géométrie classique de certains feuilletages de degré deux. *Bull. Braz. Math. Soc. (N.S.)*, 41(2):161–198, 2010.
- [CF20] S. C. Coutinho and F. R. Ferreira. A family of foliations with one singularity. *Bull. Braz. Math. Soc. (N.S.)*, 51(4):957–974, 2020.
- [CF21] S. C. Coutinho and F. R. Ferreira. Correction to: A family of foliations with one singularity. *Bull. Braz. Math. Soc. (N.S.)*, 52(3):767–769, 2021.
- [CJ21] S. C. Coutinho and L. F. G. Jales. Foliation with one singularity and finite isotropy group. *Bull. Sci. Math.*, 169:Paper No. 102988, 18, 2021.
- [Cou22] S. C. Coutinho. On the classification of foliations of degree three with one singularity. *J. Symbolic Comput.*, 112:62–78, 2022.
- [FPR22] P. Fernández, L. Puchuri, and R. Rosas. Foliation on \mathbb{P}^2 with only one singular point. *Geom. Dedicata*, 216(5):Paper No. 59, 17, 2022.

UNIV. RENNES, CNRS, IRMAR-UMR 6625, F-35000 RENNES, FRANCE

Email address: dominique.cerveau@univ-rennes1.fr

UNIVERSITÉ D'ORLÉANS, INSTITUT DENIS POISSON, ROUTE DE CHARTRES, 45067 ORLÉANS CEDEX 2, FRANCE

Email address: deserti@math.cnrs.fr