# A note on the Casas-Alvero Conjecture

<span id="page-0-0"></span>Daniel Schaub, Univ Angers, CNRS, LAREMA, SFR MATHSTIC F-49000 Angers, France email: daniel.schaub@univ-angers.fr

Mark Spivakovsky, Univ Paul Sabatier, CNRS, IMT UMR 5219 F-31062 Toulouse, France and CNRS, LaSol UMI 2001, UNAM. email: mark.spivakovsky@math.univ-toulouse.fr

November 7, 2024

#### Abstract

The Casas–Alvero conjecture predicts that every univariate polynomial  $f$  over a field K of characteristic zero having a common factor with each of its derivatives  $H_i(f)$  is a power of a linear polynomial. Let  $f = x^d + a_1 x^{d-1} + \cdots + a_1 x \in K[a_1, \ldots, a_{d-1}][x]$  and let  $R_i = Res(f, H_i(f)) \in K[a_1, \ldots, a_{d-1}]$  be the resultant of f and  $H_i(f), i \in \{1, \ldots, d-1\}.$ The Casas-Alvero Conjecture for algebraically closed fields is equivalent to saying that  $R_1, \ldots, R_{d-1}$  are "independent" in a certain sense, namely that the height  $ht(R_1, \ldots, R_{d-1}) =$  $d-1$  in  $K[a_1,\ldots,a_{d-1}]$ . In this paper we prove a partial result in this direction: if  $i \in \{d-3, d-2, d-1\}$  then  $R_i \notin \sqrt{(R_1, \ldots, R_i, \ldots, R_{d-1})}$ .

# 1 Introduction

In the year 2001 Eduardo Casas–Alvero published a paper on higher order polar germs of plane curve singularities [\[1\]](#page-3-0). His work on polar germs inspired him to make the following conjecture (according to the testimony of José Manuel Aroca, E. Casas communicated the problem orally well before 2001).

A lot of work on the Casas-Alvero conjecture has been done since 2001. Some of the most important partial results can be found in [\[1–](#page-3-0)[7\]](#page-4-0).

Let K be a field, d a strictly positive integer and  $f = x^d + a_1 x^{d-1} + \cdots + a_{d-1} x + a_d$  a monic univariate polynomial of degree  $d$  over  $K$ . Let

$$
H_i(f) = {d \choose i} x^{d-i} + {d-1 \choose i} a_1 x^{d-i-1} + \dots + {i \choose i} a_{d-i}
$$

be the *i*-th Hasse derivative of  $f$ .

**Definition 1** The polynomial f is said to be a Casas–Alvero polynomial if for each  $i \in$  $\{1,\ldots,d-1\}$  it has a non-constant common factor with its i-th Hasse derivative  $H_i(f)$ .

Note that, by definition, a Casas-Alvero polynomial f has a common root with  $H_{d-1}(f)$ . In particular, if char  $K = 0$ , it has at least one root  $b \in K$ , regardless of whether or not K is algebraically closed. Making the change of variables  $x \rightsquigarrow x - b$ , we may assume that 0 is a root of f, in other words,  $a_d = 0$ . In the sequel, we will always make this assumption without mentioning it explicitly.

Conjecture 1 (Casas–Alvero) Assume that char  $K = 0$ . If  $f \in K[x]$  is a Casas-Alvero polynomial of degree d with  $a_d = 0$ , then  $f(x) = x^d$ .

For  $i \in \{1, \ldots, d-1\}$ , let  $R_i = \text{Res}(f, H_i(f)) \in K[a_1, \ldots, a_{d-1}]$  be the resultant of f and  $H_i(f)$ . The polynomials f and  $H_i(f)$  have a common factor if and only if  $R_i = 0$ . Thus f is Casas–Alvero if and only if the point  $(a_1, \ldots, a_{d-1}) \in K^{d-1}$  belongs to the algebraic variety  $V(R_1, \ldots, R_{d-1}) \subset K^{d-1}$ . In those terms the Conjecture can be reformulated as follows:

Conjecture 2 Let  $V = V(R_1, ..., R_{d-1})$  ⊂  $K^{d-1}$ . Then  $V = \{0\}$ .

When  $K$  is algebraically closed, this is also equivalent to

#### <span id="page-1-0"></span>Conjecture 3

<span id="page-1-2"></span>
$$
\sqrt{(R_1, \ldots, R_{d-1})} = (a_1, \ldots, a_{d-1})
$$
\n(1)

or, equivalently,

<span id="page-1-1"></span> $a_i^N \in (R_1, ..., R_{d-1})$  for all  $i \in \{1, ..., d-1\}$  and some  $N \in \mathbb{N}$ . (2)

**Remark 2** (i) When K is not algebraically closed, Conjecture [3](#page-1-0) is a priori stronger. (ii) Let  $K \subset K'$  be a field extension. The induced extension

$$
K[a_1,\ldots,a_{d-1}]\subset K'[a_1,\ldots,a_{d-1}]
$$

is faithfully flat. Since the polynomials  $R_1, \ldots, R_{d-1}$  have coefficients in  $\mathbb{Z}$ , [\(2\)](#page-1-1) holds in  $K[a_1, \ldots, a_{d-1}]$  if and only if it holds in  $K'[a_1, \ldots, a_{d-1}]$ . Hence the truth of Conjecture 5 depends only on the characteristic of  $K$  but not on the choice of the field  $K$  itself. Because of this, we will take  $K = \mathbb{C}$  in the sequel.

<span id="page-1-3"></span>**Remark 3** Formulae [\(1\)](#page-1-2) and [\(2\)](#page-1-1) can be interpreted in terms of Gröbner bases. Namely, (1) and  $(2)$  are equivalent to saying that for any choice of monomial ordering and Gröbner basis  $(f_1, \ldots, f_s)$  of  $(R_1, \ldots, R_{d-1})$ , after renumbering the  $f_j$ , the leading monomial of  $f_j$  is a power of  $a_j$  for  $j \in \{1, ..., d-1\}$ .

<span id="page-1-4"></span>**Remark 4** Conjecture [3](#page-1-3) and Remark 3 say that, as polynomials in  $K[a_1, \ldots, a_{d-1}]$ , the resultants  $R_1, \ldots, R_{d-1}$  are "independent" in a certain sense.

Each of the following statements is also equivalent to the Casas-Alvero Conjecture.

- (a) For each  $i \in \{1, \ldots, d-2\}$ , the element  $R_{i+1}$  is not a zero divisor modulo  $(R_1, \ldots, R_i)$  (in other words,  $R_1, \ldots, R_{d-1}$  form a regular sequence in  $K[a_1, \ldots, a_{d-1}].$
- (b) For each  $i \in \{1, ..., d-2\}$ ,

$$
R_{i+1} \not\in \bigcup_{\mathfrak{p} \in Ass((R_1, \ldots, R_i))} \mathfrak{p}.
$$

where  $Ass((R_1, \ldots, R_i))$  is the set of associated primes of the ideal  $(R_1, \ldots, R_i)$ .

Moreover, the above statements (a) and (b) are independent of the numbering of the  $R_i$  (since the  $R_i$  are quasi-homogeneous); a permutation of the  $R_i$  yields equivalent statements.

**Notation.** We will denote by  $(R_1, R_2, \ldots, R_{i}, \ldots, R_{d-1})$  the ideal of  $K[a_1, \ldots, a_{d-1}]$  generated by the set  $\{R_1, R_2, \ldots, R_{d-1}\} \setminus \{R_i\}.$ 

The main theorem of this paper is the following partial result in the direction of Conjecture [3](#page-1-0) and statements (a) and (b) of Remark [4.](#page-1-4)

<span id="page-2-0"></span>**Theorem 5** Take an element  $i \in \{d-3, d-2, d-1\}$ . We have

$$
R_i \not\in \sqrt{(R_1, R_2, \ldots, R_i, \ldots, R_{d-1})}.
$$

### 2 Ideals generated by all the resultants but one

In this section we prove Theorem [5](#page-2-0) after recalling some preliminary results.

<span id="page-2-1"></span>**Proposition 6** Let f be a polynomial of degree d with real roots  $\beta_1 \leq \beta_2 \leq \ldots \leq \beta_d$ . Then  $H_1(f)$  has real roots  $\gamma_1 \leq \gamma_2 \leq \ldots \leq \gamma_{d-1}$ , counted with multiplicity, where  $\gamma_i \in ]\beta_i, \beta_{i+1}[$  is  $\beta_i < \beta_{i+1}$  and  $\gamma_i = \beta_i$  if  $\beta_i = \beta_{i+1}$ .

Proof: Assume that f has s distincts roots  $\delta_1 < \delta_2 < \cdots < \delta_s$  of multiplicities  $m_1, \ldots, m_s$ , respectively. Then  $\delta_j$  is a root of  $H_1(f)$  of multiplicity  $m_j - 1$ , where we say that  $\delta_j$  is a root of multiplicity 0 if it is not a root of  $H_1(f)$ .

By Rolle's theorem, there is at least one root of  $H_1(f)$  in each of the  $s-1$  open intervals  $]\delta_1, \delta_2[,\ldots,]\delta_{s-1}, \delta_s[$ .

**Notation.** Let  $\text{Int}(\beta_i, \beta_{i+1}) := ]\beta_i, \beta_{i+1}[$  if  $\beta_i < \beta_{i+1}$  and  $\text{Int}(\beta_i, \beta_{i+1}) := \{\beta_i\}$  if  $\beta_i = \beta_{i+1}$ .

According to the above, there is at least one real root of  $H_1(f)$  in each of  $Int(\beta_i, \beta_{i+1}),$  $i \in \{1, \ldots, d-1\}$ , where  $\gamma_1 \in \text{Int}(\beta_1, \beta_2), \ldots, \gamma_{m_1-1} \in \text{Int}(\beta_{m_1-1}, \beta_{m_1})$  are the first  $m_1-1$  roots of  $H_1(f)$  (in fact, the same root counted with multiplicity  $m_1 - 1$ ) and similarly for the other multiple roots of  $f$ .

We have accounted for a total of  $s-1+(m_1-1)+\cdots+(m_s-1)=m_1+\cdots+m_s-1=d-1$ real roots of  $H_1(f)$  counted with multiplicities. Hence  $H_1(f)$  has no roots, real or complex, other than the ones listed above, and the result follows.  $\square$ 

Corollary 7 Let f be a polynomial of degree d with d real roots, counted with multiplicities. Then each of the  $H_i(f)$ ,  $i \in \{1, \ldots, d-1\}$ , has  $d-i$  real roots, counted with multiplicity. In other words, all the roots of  $H_i(f)$  are real.

Next, we recall a result from [\[4\]](#page-3-1) on almost counterexamples to the Casas-Alvero conjecture.

Definition 8 Fix an  $i \in \{1, \ldots, d-1\}$ . An almost counterexample to the Casas-Alvero conjecture of level i is a polynomial f that has a common root with  $H_i(f)$  for all  $j \in \{1, \ldots, d-\}$  $1\} \setminus \{i\}$  but is not a power of a linear polynomial.

**Notation.** Given a polynomial f of degree d with d real roots, for a pair  $(k, m)$  of integers with  $1 \leq k \leq d-1$  and  $1 \leq m \leq d-k$  we write  $\alpha_{k,m}(f)$  for the m-th root of  $H_k(f)$ , where the roots of  $H_k(f)$  are ordered (weakly) increasingly.

We state the next theorem in a somewhat stronger form than in  $[4]$ : the extra information about the recycled roots  $\alpha_{k_j,m_j}(f)$  does not appear explicitly in the statement of the result in [\[4\]](#page-3-1), but is shown in the course of its proof.

Theorem 9 (J. Draisma–J. P. de Jong [\[4\]](#page-3-1), Theorem 5) Fix  $d-2$  pairs of integers

$$
(k_j, m_j), \quad j \in \{1, \ldots, d-2\},\
$$

with

$$
1 \le k_1 < k_2 < \dots < k_{d-2} \le d-1
$$

and  $1 \leq m_i \leq d - k_i$ . There exists a polynomial  $f \in \mathbb{R}[x]$  with  $f(0) = f(1) = 0$ , all of whose roots are real and lie in [0,1], such that  $\alpha_{k_j,m_j}(f)$  is a root of f for all  $j \in \{1,\ldots,d-2\}$  (in particular, f is an almost counterexample to the Casas-Alvero conjecture of level i, where i is the unique element of the set  $\{1,\ldots,d-1\} \setminus \{k_1,\ldots,k_{d-2}\}$ .

We also recall the following result, Theorem 13 of [\[2\]](#page-3-2):

<span id="page-3-4"></span>Theorem 10 Assume that f is a counterexample to the Casas-Alvero Conjecture. Then f has at least five distinct roots.

Proof of Theorem [5:](#page-2-0) We argue by contradiction. Assume that

<span id="page-3-3"></span>
$$
R_i \in \sqrt{(R_1, \ldots, \breve{R}_i, \ldots, R_{d-1})}.
$$
\n(3)

Let f be an almost counterexample of level i to the Casas-Alvero conjecture with  $m_i = 1$  for all  $j \in \{1, \ldots, d-2\}$ , given by Theorem [9.](#page-0-0) By [\(3\)](#page-3-3), f is a Casas-Alvero polynomial that is a counterexample to the Casas-Alvero conjecture. By definition of f,  $\alpha_{k,i}(f)$  is the first root of  $H_{k_j}(f)$  and also a root of f, for all  $j \in \{1, \ldots, d-2\}$ . In particular, since  $i \in \{d-3, d-2, d-1\}$ ,  $\alpha_{\ell_1}(f)$  is the first root of  $H_{\ell}(f)$  and a root of f for all  $\ell \in \{1, \ldots, d-4\}.$ 

By Proposition [6,](#page-2-1)  $\alpha_{11}(f)$  is also the first root of f (so  $\alpha_{11}(f) = 0$ ). Let m denote the multiplicity of this root of f. By Theorem [10,](#page-3-4) f has at least 5 distinct roots, hence  $m \leq d-4$ . Let  $\beta$  denote the first strictly positive root of f.

By Proposition [6,](#page-2-1) there is a unique root  $\beta^{(1)}$  of  $H_1(f)$  with  $\beta^{(1)} \in ]0, \beta[$ . If  $m > 2$ , again by Proposition [6,](#page-2-1) there is a unique root  $\beta^{(2)}$  of  $H_2(f)$  with  $\beta^{(2)} \in ]0, \beta^{(1)}[\subset]0, \beta[$ . We continue like this recursively until  $H_{m-1}(f)$ , to show that there is a unique root  $\beta^{(m-1)}$  of  $H_{m-1}(f)$  with  $\beta^{(m-1)} \in ]0, \beta^{(m-2)}[\subset]0, \beta[$ . Now,  $H_m(f)$  has no root at 0, hence its first root  $\alpha_{m1}(f)$  belongs to the open interval  $]0, \beta^{(m-1)}[\subset]0, \beta[$ .

This contradicts the fact that  $\alpha_{m1}(f)$  is a root of f. This completes the proof.  $\Box$ 

## References

- <span id="page-3-0"></span>[1] Eduardo Casas–Alvero, Higher Order Polar Germs, Journal of Algebra, Volume 240, Issue 1, 1 June 2001, pages 326–337
- <span id="page-3-2"></span>[2] W. Castryck, R. Laterveer, M. Ounaïes, Constraints on counterexamples to the Casa-Alvero conjecture and a verification in degree 12, arXiv:1208.5404v1, 27/08/2018.
- [3] M. Chellali, A. Salinier, La conjecture de Casas-Alvero pour les degrés  $5p^e$ , hal-00748843, 2012.
- <span id="page-3-1"></span>[4] J. Draisma and J.P. de Jong, On the Casas-Alvero conjecture, Newsletter of the EMS 80 (June 2011) 29–33
- [5] R.M. de Frutos Marín, Perspectivas Aritméticas para la Conjectura de Casas-Alvero, PhD thesis, Universidad de Valladolid, 2012.
- [6] S. Ghosh, A finiteness result towards the Casas-Alvero conjecture, preprint, arXiv:2402.18717v2, [math.AG], 2024.
- <span id="page-4-0"></span>[7] H.-C. Graf von Bothmer, O. Labs, J. Schicho and C. Van de Woestline, The Casas-Alvero conjecture for infinitely many degrees Journal of Algebra, Vol. 316(1),224-230, 2007.