

Climate and agricultural productivity in southern Roman Gaul: the contribution of agent-based and agrosystemic modelling

Nicolas Bernigaud, Alberte Bondeau, Joel Guiot, Frédérique Bertoncello,

Marie-Jeanne Ouriachi

▶ To cite this version:

Nicolas Bernigaud, Alberte Bondeau, Joel Guiot, Frédérique Bertoncello, Marie-Jeanne Ouriachi. Climate and agricultural productivity in southern Roman Gaul: the contribution of agent-based and agrosystemic modelling. EAA 2020, Virtual Meeting 27 August, Session #415: From Abacus to Calculus. Computational Approaches to Roman Economy, Philipp Verhagen; Iza Romanowska, Aug 2020, Budapest (HO), Hungary. hal-04342118

HAL Id: hal-04342118 https://hal.science/hal-04342118

Submitted on 13 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



EAA 2020, Virtual Meeting 27 August Session #415: From Abacus to Calculus. Computational Approaches to Roman Economy



Climate and agricultural productivity in southern Roman Gaul: the contribution of agent-based and agrosystemic modelling¹

N. Bernigaud*, A. Bondeau**, J. Guiot*, F. Bertoncello***, M.-J. Ouriachi***

¹ RDMed project

*CEREGE, Aix-en-Provence, **IMBE, Aix-en-Provence, ***CEPAM, Nice



Goals and Objectives Presentation

What was the impact of climate change on agriculture and agricultural yields in the south of France between the Iron Age and the Late Roman antiquity ?

Contribution of Agrosystemic and Agent-Based models to this question.

Roman Climate Optimum (RCO) or Roman Warm Period (RWP)




Figure 2 | Eurasian summer temperature variability. a,b, Reconstructed temperatures from the Russian Altai (a) and the European Alps⁵ (b), with blue and red squares indicating (51/47) positive and (53/57) negative annual extremes (>2 standard deviations), respectively. Grey background shadings denote reconstruction uncertainty after 80-year low-pass filtering (Supplementary Fig. 9). c,d, Ice-core-derived global estimates of volcanic¹⁴ (GVF; dark grey; c) and total solar irradiance¹⁷ (TSI; light dashed grey; d) forcing. Black dashed lines refer to the long-term reconstruction mean of the Common Era.


• III c. BC – II c. AD

- High summer temperatures
- High precipitations

Büntgen et al. 2011

"THE CLIMATE WAS THE ENABLING BACKGROUND OF THE ROMAN MIRACLE. THE RCO TURNED THE LANDS RULED BY ROME INTO A GIANT GREENHOUSE"

- "The RCO was a boon to the fecundity of Mediterranean wheat farming"
- The RCO "fueled the agricultural engine of the economy" (Harper 2017, 52; Cf. McCormick et al. 2012; Manning 2013, 169).



Did really the RCO makes the prosperity of the Roman Empire ?

Climate change and the productive landscape in the Mediterranean region in the Roman Period (draft paper by P. Erdkamp, Vrije Universiteit Brussel)

- "Many recent studies claim that climate change and its impact on agriculture determined the fate of past societies. The prosperity of the Roman Empire at its height is attributed to the *Roman Climatic Optimum*, which "fueled the agricultural engine of the economy" (Harper 2017, 52; Cf. McCormick et al. 2012; Manning 2013, 169)"
- « It seems very unlikely that the impact on agriculture in the Roman world was as dramatic as often assumed »
- "However, a thorough analysis of the causal links between climate change and agricultural productivity is generally missing"

How to link climate and agricultural productivity ?


Lund-Potsdam-Jena managed Land (LPJmL) **Process-based** agriculture, vegetation and hydrology model

• MODEL INPUTS

- Monthly climate : temperatures, precipitations, rainy days, cloud cover
- Global atm. CO2concentrations
- Soil structure
- Land use dataset

• MODEL OUTPUTS

- Yields (t/ha), production (t), growing period
- Carbon cycle
- Soils evaporation, plant transpiration,...



Modeling ancient yields with paleoclimatic data






FIGURE 1 | Map of the 295 sites used for the reconstruction; the color indicates the elevation of the site.


 Joël Guiot, David Kaniewski. The Mediterranean Basin and Southern Europe in a warmer world: what can we learn from the past ?. Frontiers in Earth Science, Frontiers Media, 2015, 3


Using LPJmL for reconstructing agricultural yields of the past





Temperatures






Cloud cover






Reconstruction of potential yields in Ist c. BC




Wheat yield per T/ha (2000 BP)

LPJmL results

Centennial variations of WHEAT 's potential yields in t/HA 6th c. BC - 7th c. AD









Centennial variations of WHEAT 's potential yields in t/HA 6th c. BC - 7th c. AD

- Increase of yields during Iron Age
- Noticeable peak in 3rd c. BC
- High yields during Roman period
- Decrease during early Middle age





Centennial average of potential yields of WHEAT in tons/HA






Centennial variations of GRAPES's potential yields in hl/HA 6th c. BC - 7th c. AD







1st c. BC (Maximum)




Centennial variations of GRAPES's potential yields in hl/HA 6th c. BC - 7th c. AD


- 1st peak : 3rd c. BC 1 st. c. BC
- 2nd peak : 4th c. AD 6th c. AD
- Highest value : 1st c. BC = 32 hl/HA
- Lowest value : 6th c. BC = 24 hl/HA
- Yields of Roman period more stable than others






Centennial average of potential yields of rainfed **OLIVES** in t/HA 6th c. BC - 7th c. AD






Centennial average of potential yields of rainfed **OLIVES** in t/HA 6th c. BC - 7th c. AD

- Irregular yields during Iron Age with noticeable peak in 3rd
 c. BC
- Relative high yields during Roman period
- Irregular yields during early Middle age




Comparison of centennial averages of POTENTIAL yields

Climate change does not have the same effect on the yields of different plant species




From POTENTIAL YIELDS to POTENTIAL PRODUCTION : working with agent-based modeling




Agent-based Model "LPJmLvine" (NETLOGO)


- Integrate LPJmL results + archeological data (sites, roads, towns,...)
- Agents are wineries attested by archaeological data
- Allows to calculate a global wine production according to centennial potential yields


6th c. BC







5th c. BC







4th c. BC







3rd c. BC







2nd c. BC







1st c. BC







1st c. AD







2nd c. AD








3nd c. AD



























POTENTIAL wine production (according to NETLOGO model) 6th c. BC – 7th c. AD

- Progressive increase of 18 % during Iron Age until the 1st c.
 BC
- Decrease after the 1st c. BC but stabilization of production level during Roman period, with a new increase in the Late Antiquity (5th c. AD)
- Slight decrease during early Middle Age, but no collapse of the production



Conclusion

• Did the Roman Climate Optimum have had a significant effect on agricultural production ?

According to the results of our models, the potential yields and production was indeed higher in the Roman period than Iron Age and early Middle Age.

- \succ Increase of 7% of wheat potential yields
- > Increase of 18 % of potential wine production