



HAL
open science

Electrophilicity, Mechanism and Structure-Reactivity Relationships of Cyclic Secondary Amines Addition to 2-Methoxy-3,5-dinitropyridine

Takwa Slama, Ons Amamou, Amel Hedhli, Jean-claude Guillemin, Sahbi Ayachi, Taoufik Boubaker

► **To cite this version:**

Takwa Slama, Ons Amamou, Amel Hedhli, Jean-claude Guillemin, Sahbi Ayachi, et al.. Electrophilicity, Mechanism and Structure-Reactivity Relationships of Cyclic Secondary Amines Addition to 2-Methoxy-3,5-dinitropyridine. *Journal of Molecular Structure*, 2024, 1302, pp.137258. 10.1016/j.molstruc.2023.137258 . hal-04344339

HAL Id: hal-04344339

<https://hal.science/hal-04344339v1>

Submitted on 7 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Highlights

- SET mechanism has been suggested to describe the nucleophilic aromatic substitution reactions of 2-methoxy-3,5-dinitropyridine by secondary cyclic amines in acetonitrile solution.
- E values of six anisoles have been predicted according to the E vs ω correlation (Eq.I) demonstrated in this work and found to agree with those reported in the literature.
- $E = -93.41 + 23.73 \omega$ ($R^2 = 0.9997$) (I)
- The expansion of the applicability of Mayr's equation (II) to these types of processes further demonstrates the generality of this relationship.
- $\log k (20\text{ }^\circ\text{C}) = s_N (E + N)$ (II)

Journal Pre-proof

Electrophilicity, Mechanism and Structure-Reactivity Relationships of Cyclic Secondary Amines Addition to 2-Methoxy-3,5-dinitropyridine

Takwa Slama,^a Ons Amamou,^a Amel Hedhli,^a Jean-Claude Guillemin,^b Sahbi Ayachi,^c and Taoufik Boubaker^a

^aLaboratoire de Chimie Hétérocyclique, produits Naturels et Réactivité (LR11SE39), Faculté des Sciences de Monastir, Université de Monastir, Avenue de l'Environnement, 5019 Monastir, Tunisie.

^bUniv Rennes, Ecole Nationale Supérieure de Chimie de Rennes, CNRS, ISCR – UMR6226, F-35000 Rennes, France

^cLaboratoire de physico-chimie des matériaux (LR01ES19), Faculté des Sciences, Université de Monastir, Avenue de l'Environnement, 5019 Monastir, Tunisie.

*Corresponding author: Prof. Taoufik Boubaker
e-mail: boubaker_toufik@yahoo.fr

Phone : 00 216 98 65 21 21


ABSTRACT

The second-order rate constants (k) for the nucleophilic aromatic substitution reactions of 2-methoxy-3,5-dinitropyridine **1** with secondary cyclic amines **2a-c** in acetonitrile solution at 20 °C are reported. The logarithms of these rate constants are particularly significant as possible measures of the electrophilicity parameters E for 2-methoxy-3,5-dinitropyridine **1** according to the linear free energy relationship $\log k (20\text{ °C}) = s_N(N + E)$. Additional kinetic data for reactions of 2-methoxy-3,5-dinitropyridine **1** with a series of nitroalkyl anions in DMSO solution are also measured and found to agree with those calculated by Mayr's approach from the known two solvent-dependent parameters N and s_N of nitroalkyl anions and the electrophilicity parameter E measured in this work. A Single Electron Transfer (SET) mechanism is proposed for the S_NAr reactions, on the basis of the linear Brønsted plot with the β_{nuc} value of 0.94. The validity of this mechanism is confirmed by the agreement between the rate constants, k , and the oxidation potentials E° of these series of secondary cyclic amines. Also, reactivity descriptors such as the electronic chemical potential (μ), and the chemical hardness (η) for a series of pyridine derivatives were calculated by Density Functional Theory (DFT), and it is shown that the global electrophilicity index ($\omega = \mu^2/\eta$) correlate significantly with the electrophilicity parameters E .

Keywords : Substituted Pyridine Derivatives / Kinetics / Equation of Mayr / Electrophilicity / Single Electron Transfer Mechanism / DFT Analysis.

1. Introduction


Pyridine derivatives have been widely investigated for their importance in biological and industrial processes and synthetic applications.[1-7] In terms of reactivity, it should be noted that isomeric addition at substituted and unsubstituted carbons of some pyridine derivatives has been the subject of extensive experimental and theoretical studies. [8-10] Reactions of 2-methoxy-3-nitropyridine **3** with a series of secondary cyclic amines have generally been understood to proceed through a polar S_NAr mechanism on the basis of the fact that Brønsted plot was found to be linear with $\beta_{nuc} = 0.52$. [11] Similar conclusion has also been drawn for the corresponding reactions with 2-methoxy-5-nitropyridine **4** ($\beta_{nuc} = 0.55$). [11] While, Mahdhaoui and coworkers have recently reported that the reactions of 2,6-dimethoxy-3,5-dinitropyridine **5** with weakly basic 4-X-substituted phenoxide anions (e.g., X = Cl and CN) proceeds through a polar σ -complexation mechanism, while the corresponding reaction with strongly basic phenoxide anions (e.g., X = Me and OMe) proceeds through a single electron transfer mechanism (SET). [12] Such single electron transfer mechanism in σ -complexation reactions is considered to be the only time reported in literature but have been observed in some S_NAr reactions. [13-15]



To get more information on reactivity and mechanism, our kinetic study has been extended to reactions of 2-methoxy-3,5-dinitropyridine **1** with the secondary cyclic amines **2a-c** as reference nucleophiles in acetonitrile at 20 °C (Scheme 1). As will be seen, the effect of the nucleophile nature on reactivity and mechanism have been examined quantitatively and discussed on the basis of linear free-energy relationships (LFER). On the other hand, the second-order rate constants (k) of these reactions have been described by Equation (1) (where N and s_N are nucleophile-specific parameters) [16] in order to evaluate the electrophilicity parameter E of the electrophile **1** and to include it into the comprehensive electrophilicity scale of Mayr.

$$\log k (20\text{ }^\circ\text{C}) = s_N (E + N) \quad (1)$$

In addition, theoretical study based on density functional theory has been performed at the B3LYP/6-311g(d,p) level [17] in order to establish a correlation between the calculated global electrophilicity index (ω) values for a series of pyridines and their electrophilicity parameter E measured using Mayr's approach.



Scheme 1. S_NAr reactions of the 2-methoxy-3,5-dinitropyridine **1** with various secondary cyclic amines in CH_3CN at $20\text{ }^\circ C$.

2. Results and Discussion

2.1. Structural studies of the substitution products **6a-c**

Addition of dilute solutions of amines **2a-c** to a colorless solution of 2-methoxy-3,5-dinitropyridine **1** in acetonitrile resulted in the formation of yellow species exhibiting intense absorption maxima at 356, 358 and 331 nm, respectively. These λ_{max} are characteristic for the formation of substitution products. Figure 3 illustrates the UV-vis absorption spectra describing the progressive conversion of the 2-methoxy-3,5-dinitropyridine **1** to the substitution product **6b**. A clear isosbestic point is observed at 312 nm, indicating the involvement of two species in an equilibrium.






Figure 1. Time dependence of the electronic absorption spectrum of 2-methoxy-3,5-dinitropyridine **1** ($5 \times 10^{-5}\text{ mol L}^{-1}$) in the presence of piperidine **2b** ($2 \times 10^{-2}\text{ mol L}^{-1}$) in CH_3CN solution at $20\text{ }^\circ C$.

It is interesting to note that pyridine **1** does not give σ -adduct of C-H attack with any of the investigated nucleophiles under the reaction conditions; only attack at C-OMe to afford the substitution products **6** is observed. Substitution products **6a-c** have also been characterized through analysis of ^1H NMR spectra. In the case of substitution product **6a**, all the chemical shifts and coupling constants data are summarized in Table 1. Interestingly, we have succeeded in isolating **6a**. Dissolution of this compound in CD_3CN afforded NMR spectra which were identical to those recorded in the *in situ* generation of **6a** in this solvent.

Table 1. ^1H NMR shifts values for 2-methoxy-3,5-dinitropyridine **1** and the substitution product (3,5-dinitro-2-(pyrrolidin-1-yl)pyridine) **6a** (300 MHz, $\text{CD}_3\text{CN-d}_3$).

	δ (ppm)					Coupling constants (Hz)
	H₄	H₆	CH₃	H₇₋₁₀	H₈₋₉	
 <p>1</p>	9.27	8.99	4.20	–	–	$J_{4-6} = 2.6$
 <p>6a</p>	8.78 ^a (8.78) ^b	9.11 (9.11)	–	3.47 (3.49)	1.97 – 2.01 (1.97 – 2.01)	$J_{4-6} = 2.6$ ($J_{4-6} = 2.4$)

^aProduct *in situ*.

^bSynthesized product.

2.2. Kinetics of the reactions of 2-methoxy-3,5-dinitropyridine **1** with secondary cyclic amines **2a-c**

The kinetics of the reactions of 2-methoxy-3,5-dinitropyridine **1** with secondary amines **2a-c** in acetonitrile at 20 °C were monitored photometrically by following the appearance of the absorbances of substitution products **6a-c** at or close to their absorption maxima ($330 \text{ nm} < \lambda_{\text{max}} < 360 \text{ nm}$).

Our results can be interpreted on the basis of Scheme 1, in which the nucleophilic attack by amine **2** is the rate-limiting step. Under these experimental conditions, the commonly accepted general expression for the observed first-order rate constants, k_{obsd} , pertaining to the process is given by Eq. (2).^[18-22] All k_{obsd} values obtained are summarized in Supplementary Table S1 and displayed in Figure 2.

$$k_{\text{obsd}} = \frac{k_1(k_2 + k_3[\text{NuH}])}{k_{-1} + k_2 + k_3[\text{NuH}]}[\text{NuH}] \quad (2)$$

Figure 2 shows that in all cases the plots of k_{obsd} against the concentrations of the amine **2** were linear with negligible intercepts. This result confirmed the first-order of the reactions of **1** with **2a-c**, and indicated that the formation of the zwitterionic intermediate T^\ddagger is the rate-limiting step of the uncatalysed reaction and the pseudo-first-order rate constants (k_{obsd}), for

all the reactions can be readily derived from Eq. (3) ($k_{-1} \ll k_2 + k_3 [\text{NuH}]$). The values of the second-order rate constants, k_1 , determined from the slopes of k_{obsd} vs. $[\text{NuH}]$ plots are collected in Table 2.







Figure 2. Influence of the concentration of amine **2a-c** on the observed first-order rate constant (k_{obsd}) for addition to 2-methoxy-3,5-dinitropyridine **1** in acetonitrile solution at 20 °C.

Table 2. Summary of second-order rate constants for addition reactions of secondary amines **2a-c** with 2-methoxy-3,5-dinitropyridine **1** in acetonitrile solution at 20 °C.

Amine			
	2a	2b	2c
k_1^a ($\text{M}^{-1} \text{s}^{-1}$)	6.84×10^{-3}	7.55×10^{-4}	1.06×10^{-5}
pK_{aH}	19.62 ^b	19.35 ^b	16.61 ^c
N / s_N^d	18.64 / 0.60	17.35 / 0.68	15.65 / 0.74
E_p^{ox} (V vs NHE)	1.09 ^e	1.36 ^e	1.48 ^f

^aValues of k_1 are determined in this work from the slope of plots of k_{obsd} vs. $[\text{Amine}]$.

^bpK_{aH} values for conjugated acid of amines **2a** and **2b** in CH₃CN solution taken from ref [23a].

^cpK_{aH} value for conjugated acid of amine **2c** in CH₃CN solution taken from ref [23b].

^dN and *S_N* values in CH₃CN solution were taken from ref [30].

^eE_{p^{ox}} values for amines **2a** and **2b** in CH₃CN solution were taken from ref [28].

^fE_{p^{ox}} value for amine **2c** in CH₃CN solution was taken from ref [29].


2.3. Effect of amine basicity and mechanism

The effect of amine basicity on reactivity is illustrated in Figure 3 for the reactions of 2-methoxy-3,5-dinitropyridine **1** with secondary amines **2a-c**. As seen from Figure 3, the Brønsted plot of log *k*₁ in function of the pK_{aH} values of the conjugate acids of the amines **2a-c** in acetonitrile solution [23] listed in Table 2 is in good agreement and the corresponding relationship is the following

$$\log k_1 = -20.54 + 0.94 \text{ pK}_{\text{aH}} \quad (R^2 = 0.9958) \quad (2)$$

The β_{nuc} value for the reactions of **1** (i.e., β_{nuc} = 0.94) is much larger than that reported previously for the corresponding reactions of 2-methoxy-3-nitropyridine **4** (i.e., β_{nuc} = 0.52) and 2-methoxy-5-nitropyridine **5** (i.e., β_{nuc} = 0.55) performed in acetonitrile, which were proposed to proceed through a S_NAr mechanism.[11] Such a large β_{nuc} value is typical for reactions reported previously to proceed through a single electron transfer (SET) process. [12-15, 24-27]

In our study, the fact that the log *k*₁ is linearly related to the oxidation peak potentials E_{p^{ox}} (V vs NHE) values of the amine radical cations measured in acetonitrile [28, 29] (Figure 4) confirms the possibility that our S_NAr reactions proceed via electron transfer mechanism (Scheme 2) in which the formation of the substitution products **6a-c** will then proceed through initial (fast) electron-transfer from the amines **2a-c** to the 2-methoxy-3,5-dinitropyridine **1** and subsequent (slow) coupling of the resulting cation and anion radicals within the solvent cage via the transition state **T[±]**. [10]



Scheme 2. Proposed mechanistic pathway for the reactions of 2-methoxy-3,5-dinitropyridine **1** with **2a-c** in acetonitrile at 20 °C.




Figure 3. Effect of the amine basicity on the rate of the reaction of 2-methoxy-3,5-dinitropyridine **1** with secondary amines **2a-c** in CH_3CN solution at 20 °C. The pK_{aH} values were taken from ref [23].




Figure 4. The influence of the oxidation peak potential E_p^{ox} of amines on the rate constants of reactions of 2-methoxy-3,5-dinitropyridine **1** with **2a-c** in acetonitrile at 20 °C. The identity of points is given in Table 2.

2.4. Quantification of electrophilicity E of 2-methoxy-3,5-dinitropyridine **1**

In order to measure the electrophilicity parameters E describing the reactivity at C-OMe position of 2-methoxy-3,5-dinitropyridine **1**, we used the nucleophilicity parameters N and s_N for the three secondary amines **2a-c** determined by Kanzian and co-workers [30] in CH_3CN solution at 20 °C (Table 2). As required by Eq. (1), a linear plot of $(\log k_1)/s_N$ versus

nucleophilicity parameters N was obtained with slope close to 1, indicating that Eq. (1) is applicable to these S_NAr reactions (Figure 5).[16, 31-36] From the intercept of this line with the ordinate axis, the reactivity of pyridine **1** is derived ($E = -21.99$).




Figure 5. Plot of the rate constants $(\log k_1)/s_N$ for the reaction of the 2-methoxy-3,5-dinitropyridine **1** with the reference nucleophiles **2a-c** versus their nucleophilicity parameters N in CH_3CN solution at 20°C .

Figure 6 compares the reactivity of the 2-methoxy-3,5-dinitropyridine **1** ($E = -21.99$) with other pyridines, namely 2-methoxy-3-nitropyridine **3** ($E = -26.18$), and 2-methoxy-5-nitropyridine **4** ($E = -27.14$) recently reported by Mahdaoui and co-workers.[11] Also given for purposes of comparison are the E values reported for 2,6-dinitroanisole **7**, ($E = -20.55$), 2,4-dinitroanisole **8** ($E = -20.82$) and 2,4,6-trinitroanisole **9** ($E = -13.92$).[11]

As can be seen in Figure 6, the electrophilicity parameter E remains essentially unchanged on going from **3** to **4**, while the difference in reactivity between 2-methoxy-3,5-dinitropyridine **1** and 2-methoxy-5-nitropyridine **4**, the weakest ones electrophiles known to date is approximately 5 units of E .

Also to be noted is that the electrophilicity E values measured for pyridines **3** and **4** are 5.50-6.50 orders of magnitude lower than those reported for anisoles **7** and **8**.[11] This important decreases is therefore of particular interest in the understanding of the factors affecting the reactivity of the aza group.




Figure 6. Comparison of the electrophilicities of pyridine **1** with those of a few pyridines and anisoles.

It is interesting to note that applicability of the electrophilicity parameter E of our 2-methoxy-3,5-dinitropyridine **1** for predicting the rate constants of their reactions with other types of nucleophiles has been confirmed. In fact, the kinetics data obtained for the reactions of 2-methoxy-3,5-dinitropyridine **1** with a series of nitroalkyl anions **10a-c** in DMSO solution at 20 °C (the detailed results are collected in Table 3) indicates that the predicted (k_1^{calcd}) and experimental (k_1^{exp}) rate constants differ only by a factor of 4-5. This result agrees within the confidence limit of Mayr's approach (factor 10-100) [38-40] with the predicted (k_1^{calcd}) by Mayr's equation.

Most importantly, we demonstrated that the ($\log k_1^{\text{exp}}$) values are linearly related to the calculated rate constants ($\log k_1^{\text{calcd}}$) values (see Figure 7 and Eq. 3).

$$\log k_1^{\text{exp}} = 0.65 + 0.98 \log k_1^{\text{calcd}} \quad (R^2 = 0.9922) \quad (3)$$

The deviations between calculated rate constants from Eq. (3) and experimental rate constants have been observed by many authors.[41-45] Necibi and co-workers, for example, have observed a linear correlation between the experimental rate constants ($\log k_1^{\text{exp}}$) and

calculated rate constants ($\log k_1^{\text{calcd}}$) for reactions of benzoselenadiazolium cation **11** with the nitroalkyl anions **10a-c** in methanol ($\log k_1^{\text{exp}} = -0.88 + 1.14 \log k_1^{\text{calcd}}$ ($R^2 = 0.9875$)).[45]

Raissi and co-workers have demonstrated that the ($\log k_1^{\text{exp}}$) values for reactions of 4-nitrobenzofurazan **12** with a series of secondary cyclic amines in acetonitrile solution at 20 °C also correlate well with ($\log k_1^{\text{calcd}}$) values ($\log k_1^{\text{exp}} = -1.60 + 0.98 \log k_1^{\text{calcd}}$ ($R^2 = 0.9860$)).[41]



As discussed in a previous paper,[43] the reasons for such linear correlations are still not fully understood. Therefore, a more detailed examination of the results is necessary to confirm this approach.

Table 3. Experimental and calculated second-order rate constants for the reactions of 2-methoxy-3,5-dinitropyridine **1** with a series of nitroalkyl anions **10a-c** in DMSO solution at 20 °C.

Electrophile	Nucleophile	N/s_N ^a	k_1^{exp} ^b ($\text{M}^{-1} \text{s}^{-1}$)	k_1^{calcd} ^c ($\text{M}^{-1} \text{s}^{-1}$)	$k_1^{\text{exp}}/k_1^{\text{calcd}}$
<p>1 $E = -21.99$</p>	10a $\text{MeCH}=\text{NO}_2^-$	21.54/0.62	2.48	5.26×10^{-1}	4.71
	10b $\text{C H}_2=\text{NO}_2^-$	20.71/0.60	7.12×10^{-1}	1.71×10^{-1}	4.16
	10c $\text{Me}_2\text{C}=\text{NO}_2^-$	20.61/0.69	5.23×10^{-1}	1.12×10^{-1}	4.67

^a N and s_N values in DMSO were taken from ref. [37].

^bThis work.

^c k_1^{calcd} calculated by Eq. (1) with the N and s_N parameters for the nitroalkyl anions in DMSO indicated in this Table and the electrophilicity parameter E for pyridine **1**.




Figure 7. Linear correlation between $\log k_1^{\text{exp}}$ and $\log k_1^{\text{calcd}}$ for the reactions of 2-methoxy-3,5 dinitropyridine **1** with a series of nitroalkyl anions **10a-c** in DMSO solution at 20 °C.

2.5. Correlation between electrophilicity parameter (E) and global electrophilicity index (ω)

In order to establish a relationship between the electrophilicity parameters E proposed by Mayr [16] and the global electrophilicity index (ω) well-known of Parr,[46] an theoretical study has been performed using density functional theory (DFT). In this context, electronic chemical potential (μ) and chemical hardness (η) for the three nitropyridines **1**, **3** and **4** have been calculated at B3LYP/6-311g(d,P) basis set, in acetonitrile, as follow

$$\mu = (\epsilon_{\text{HOMO}} + \epsilon_{\text{LUMO}})/2 \quad (5)$$

$$\eta = (\epsilon_{\text{LUMO}} - \epsilon_{\text{HOMO}}) \quad (6)$$

where ϵ_{HOMO} and ϵ_{LUMO} are energies of the frontier molecular orbitals HOMO and LUMO, respectively.[46]

The obtained values of μ and η were considered for the calculation of the global electrophilicity index (ω) for these series of electrophiles **1**, **3**, and **4** according to Eq. (7) developed by Parr and Yang.[46a] Table 4 reports the calculated quantum chemical parameters which are related to the molecular electronic structure of the molecules under consideration, including their experimentally measured electrophilicity parameters E .

$$\omega = \mu^2/2\eta \quad (7)$$

Analysis of the data summarized in Table 4 revealed that the values of electrophilicity (E) proposed by Mayr [16] are linearly related to the global electrophilicity index (ω) defined by Parr and Yang [46a] As can be seen in Figure 8, excellent straight line is obtained on plotting

the measured E values of these series of pyridines against their global electrophilicity ω calculated in this work. The following equation (8) describes this linear dependence.

$$E = -93.41 + 23.73 \omega \quad (R^2 = 0.9997) \quad (8)$$

Relationships between parameters E and ω have also been reported by many authors.[47-54] More recently, Rammah and co-workers have also observed a satisfactory correlation between E and ω parameters of a series of benzotriazoles ($E = -50.25 + 4.09 \omega$, $R^2 = 0.9788$).[53]







Figure 8. Correlation between experimentally determined electrophilicities (E) and global electrophilicity index (ω) calculated at B3LYP/6-311g(d,p) level for the pyridines **1**, **3** and **4** listed in Table 4.

Table 4. Global properties for the whole series of pyridines **1**, **3** and **4**.

Electrophile	ϵ_{HOMO} (AU)	ϵ_{LUMO} (AU)	μ (eV)	η (eV)	ω (eV)	E
 1	-0.28318	-0.10838	-5.39	4.660	3.010	-21.99 ^b
 3	-0.26390	-0.10248	-4.985	4.390	2.830	-26.18 ^c
 4	-0.26393	-0.10133	-4.970	4.420	2.794	-27.14 ^c

^a ϵ_{HOMO} , ϵ_{LUMO} , μ , η and ω values determined in this work at the B3LYP/6-311g(d,p) level of theory.

^b This work.

^c From ref [11].

Interestingly, correlation (8) can be used as a good model to predict the unknown electrophilicity parameters E at C-OMe position of several neutral electron-deficient structures. At this point, we decided to use Eq. 8 to recalculate the E values of six anisoles **7-9** and **13-15** whose experimental data are available.[11] In fact, using their global electrophilicity ω determined in the present study by quantum chemical calculations at the B3LYP/6-311g(d,p) level of theory using Gaussian 09, we predict the electrophilicity parameters E' of these six anisoles. The results are compared in Table 5. It can be seen that these theoretical E' values vary significantly from -21.67 to -12.56. It is interestingly to note that these results from the Eq. (8) are in reasonable agreement with those reported by Mahdhaoui and coworkers.[11]

Plotting the electrophilicity parameters E of anisoles **7-9** and **13-15** and the three pyridines **1**, **3** and **4** against their ϵ_{LUMO} calculated at B3LYP/6-311g(d,p) level listed in Tables 4 and 5, we obtained a nice correlation as shown in Figure 9. This behavior indicates that Equation (8) can successfully predict the values of this series of anisoles **7-9** and **13-15**.

The results from the present study should represent one of the first confirmation for the applicability of the reactivity parameters E reported in Table 5 for predicting rate constants with various types of nucleophiles.










Figure 9. Correlation between experimentally determined electrophilicities (E) and lowest unoccupied molecular orbital energies (ϵ_{LUMO}) calculated at B3LYP/6-311g(d,p) level for the nine electrophiles listed in Tables 4 and 5.

Table 5. Global properties for the whole series of anisoles **7-9, 13-15**.

Electrophile	ϵ_{HOMO} (AU)	ϵ_{LUMO} (AU)	μ (eV)	η (eV)	ω (eV)	E' ^b	E ^c
 7	-0.27067	-0.10896	-5.165	4.410	3.025	-21.63	-20.82
 8	-0.27145	-0.10889	-5.175	4.430	3.023	-21.67	-20.55
 9	-0.28675	-0.11916	-5.52	4.560	3.341	-14.13	-13.92
 13	-0.28418	-0.11703	-5.4550	4.550	3.270	-15.81	-15.34
 14	-0.29234	-0.12107	-5.6200	4.660	3.389	-12.99	-11.94
 15	-0.29468	-0.12146	-5.6650	4.710	3.407	-12.56	-13.56

^a ϵ_{HOMO} , ϵ_{LUMO} , μ , η and ω values calculated in this work at the B3LYP/6-311g(d,p) level of theory.

^b E' values calculated in this work from Eq. (8).

^c E values taken from ref [11].

3. Conclusion

On the basis of the large Brønsted coefficients ($\beta_{\text{nuc}} = 0.94$), a single electron transfer mechanism has been suggested to describe the nucleophilic aromatic substitution reactions of 2-methoxy-3,5-dinitropyridine by secondary cyclic amines in acetonitrile solution. The SET mechanism has also been supported by the satisfactory linear correlation obtained between ($\log k_1$) and the oxidation potentials of amines (E°). Using Mayr's approach, the electrophilicity parameter E for the 2-methoxy-3,5-dinitropyridine has been determined ($E = -21.99$) and integrated into a comprehensive electrophilicity scale of Mayr. From global electrophilicity ω determined in the present study employing density functional (DFT) theory, the E values of six anisoles have been predicted according to the E vs ω correlation (Eq.8) and found to agree with those reported in the literature.

4. Experimental Section

4.1. Materials

The 2-methoxy-3,5-dinitropyridine was prepared, as previously described.^[8, 9] All the secondary cyclic amines **2** (morpholine, piperidine and pyrrolidine) were commercial products and were redistilled before use whenever necessary. Commercially available nitromethane (99%), nitroethane (98%), and 2-nitropropane (99%) were used as received. Acetonitrile (Aldrich, HPLC: 99.9%) was available of the highest quality and used without further purification.

All spectrophotometric measurements were carried out using a spectrophotometer (UV-1650 Shimadzu, Japan) equipped with a Peltier temperature controller (model TCC-240 A), which is able to keep to temperature within 0.1 K.

4.2. Kinetic Measurements

All kinetic investigations were performed in acetonitrile solution at 20 °C by following the disappearance of the UV/vis absorptions of the 2-methoxy-3,5-dinitropyridine **1** (284 nm) under pseudo-first-order conditions (2-methoxy-3,5-dinitropyridine concentration of approximately 3×10^{-5} mol L⁻¹ and nucleophile concentrations in the range 2×10^{-3} to 8×10^{-2} mol L⁻¹).

The observed first-order rate constants, k_{obsd} , were evaluated from the correlation $\ln(A_{\infty} - A_t)$ as a function of time, where A_{∞} and A_t are the values of equilibrium absorbance and the absorbance after time t , respectively. Measured first-order rate constants values, k_{obsd} , together with detailed reaction conditions are summarized in Tables S1 and S2 in the Supporting Information.

4.3. Methods and Computational Aspects

The optimized geometries for the ground (S_0) states of investigated compounds resulting from the S_NAr reactions of the 2-methoxy-3,5-dinitropyridine with various secondary cyclic amines were obtained using density functional theory (DFT) at the B3LYP/6-311g(d,p) level of theory,^[55, 56] with the Gaussian 09 quantum software package developed by Frisch and coworkers.^[57] Theoretical calculations were performed in polar acetonitrile (CH₃CN) using the conductor-like polarizable continuum model (CPCM).^[58] The Gaussian view 05 software^[59] was used to read Gaussian output files and visualize the obtained results.

The extracted highest occupied molecular orbital (HOMO) and lowest unoccupied molecular orbital (LUMO) energy levels are used to calculate the electronic chemical potential (μ), chemical hardness (η), as well as the global electrophilicity index (ω), as described in Equations 5-7.

Supporting Information Available

Concentrations and the measured rate constants, k_{obsd} , for the individual kinetic experiments values for reactions the 2-methoxy-3,5-dinitropyridine **1** with secondary cyclic amines **2a-c** in CH_3CN solution (Table S1).

Concentrations and the measured rate constants, k_{obsd} , for the individual kinetic experiments values for reactions of the 2-methoxy-3,5-dinitropyridine **1** with nitroalkyl anions in DMSO (Table S2).

Conflicts of interest statement

The authors hereby confirm they have no conflict of interest.

Authors contribution

TS, OA and AH achieved her experimental works with the help of TB and J-CG. Theoretical calculations were conducted by SA. The manuscript was jointly written and arranged under TB, SA and JCG guidance.

References

- 1 L. Lian, S. Yan, B. Yao, S. A. Chan, W. Son, Photochemical transformation of nicotine in wastewater effluent, *Environ. Sci. Technol.* 51 (2017) 11718-11730.
- 2 Y. Kitade, Y. Akao, MicroRNAs and Their Therapeutic Potential for Human Diseases: MicroRNAs, miR-143 and -145, Function as Anti-oncomirs and the Application of Chemically Modified miR-143 as an Anti-cancer Drug, *J. Pharm. Sc.* 114 (2010) 276-280.
- 3 R. K. Anchoori, M. Susanne, M. Hidalgo, T. Sarkav, G. Nalluv, R. Bai, P. J. V. Diest, E. Hamel, S. R. Khan, Novel microtubule-interacting phenoxy pyridine and phenyl sulfanyl pyridine analogues for cancer therapy, *J. Med. Chem.* 51 (2008) 5953-5957.
- 4 A. Abbottot, E. H. Calderon, M. S. Dangate, F. Angelist, N. Manfredit, G. M. Marit, C. Marinziti, E. Mosconis, M. Muccini, R. Ruffort, M. Seris, Pyridine- EDOT heteroarylene – vinylene donor-acceptor polymers, *Macromol.* 43 (2010) 9698-9713.
- 5 F. A. Egbewande, M. J. Coster, I. D. Jenkins, R. A. Davis Reaction of Papaverine with Baran Diversinates, *Mol.* 24 (2019) 3938-3951.
- 6 M. Sargolzaei, M. Afshar, H. Nikoofard, Solvent Effect on The Equilibrium and Rate Constant of the Tautomeric Reaction in Nexium, Skelaxin, Aldara and Efavirenz Drugs: A DFT Study, *J. Struc. Chem.* 59 (2018) 297-305.
- 7 S. Johanson, S. Lindstedt, H. G. Tiselius, Metabolic interconversions of different forms of vitamin B₆, *J. Bio. Chem.* 249 (1974) 6040-6046.

- 8 H. C. Van der Plas, H. Jongejan, A. Koudijs, Ring transformations in reactions of heterocyclic compounds with nucleophiles. Conversion of 5-nitropyrimidine into 3,5-dinitropyridine. A novel ring transformation, *J. Heterocycl. Chem.* 15 (1978) 485-487.
- 9 (a) F. Terrier, A. P. Chatrousse, R. Schaal, Meisenheimer complexes Stopped-flow study of the interaction of 3,5-dinitro-4-methoxypyridine with methoxide ion in methanol and methanol-dimethyl sulfoxide mixtures, *J. Org. Chem.* 37 (1972) 3010-3014. (b) P. G. Farrell, F. Terrier, H-Q. Xie, T. Boubaker, Reversal of the Order of Catalytic Efficiency of Primary and Secondary Amines in the Ionization of a Sterically Hindered Carbon Acid, *J. Org. Chem.* 55 (1990) 2546-2548.
- 10 F. Terrier, *Modern Nucleophilic Aromatic Substitution*; John Wiley & Sons: Weinheim. (2013).
- 11 F. Mahdhaoui, R. Zaier, N. Dhahri, S. Ayachi, T. Boubaker, S_NAr reactions of substituted pyridines with secondary amines in aqueous solution: Kinetic and reactivity indices in density functional theory, *Int. J. Chem. Kinet.* 51 (2019) 249-257.
- 12 F. Mahdhaoui, N. Dhahri, T. Boubaker, Single Electron Transfer in σ -complexation Reactions of 2,6-dimethoxy-3,5-dinitropyridine with Para-X-phenoxide Anions in Aqueous Solution, *Int. J. Chem. Kinet.* 48 (2016) 523-530.
- 13 I. Jamaoui, T. Boubaker, R. Goumont, Nonlinear Brønsted and Hammett Correlations Associated to Reactions of 4-chloro-7-nitrobenzofurazan with Anilines in dimethyl sulfoxide solution, *Int. Chem. Kinet.* 45 (2013) 152-160.
- 14 S. Ben Salah, T. Boubaker, R. Goumont, Kinetic studies on Nucleophilic addition reaction of 4-x-substituted anilines to N1-methyl-4-nitro-2,1,3-benzothiadiazolium tetrafluoroborate in acetonitrile: Reaction mechanism and Mayr's approach, *Int. J. Chem. Kinet.* 47 (2015) 461-470.
- 15 H. Raissi, I. Jamaoui, R. Goumont, T. Boubaker, Kinetic Studies on S_NAr Reactions of Substituted Benzofurazan Derivatives: Quantification of the Electrophilic Reactivities and Effect of Amine Nature on Reaction Mechanism, *Int. J. Chem. Kinet.* 49 (2017) 835-846.
- 16 H. Mayr, M. Patz, Scales of Nucleophilicity and Electrophilicity: A System for Ordering Polar Organic and Organometallic Reactions, *Angew. Chem. Int. Ed. Engl.* 33 (1994) 938-957.
- 17 I. Chérif, H. Raissi, K. Abiedh, B. Gassoumi, M. T. Caccamo, S. Magazu, A. Haj Said, F. Hassen, T. Boubaker, S. Ayachi, Photophysical and nonlinear optical properties of para-substituted nitrobenzofurazan: A comprehensive DFT investigation, *J. Photochem. Photobiol.* 443 (2023) 114850.

- 18 A. Echaieb, W. Gabsi, T. Boubaker, Nucleophilic Substitution Reaction of 2-methoxy-3-X-5-nitrothiophene: Effect of Substituents and Structure-Reactivity Correlation, *Int. J. Chem. Kinet.* 46 (2014) 470-476.
- 19 S. Souissi, W. Gabsi, A. Echaieb, J-C. Hierso, P. Fleurat-Lessard, T. Boubaker, Influence of solvent mixture on nucleophilicity parameters: The case of pyrrolidine in methanol-acetonitrile, *RSC Advances.* 10 (2020) 28635-28643.
- 20 E. Buncel, F. Terrier, Assessing the superelectrophilic dimension through sigma-complexation, S_NAr and Diels-Alder reactivity, *Org. Biomol. Chem.* 10 (2010) 2285-2308.
- 21 S. Souissi, W. Gabsi, A. Echaieb, T. Boubaker, Kinetics and Quantification of the Electrophilic Reactivities of Substituted Thiophenes and Structure-Reactivity Relationships, *Int. J. Chem. Kinet.* 50 (2018) 582-590.
- 22 (a) J. F. Bunnett, R. E. Zahler, Aromatic Nucleophilic Substitution Reactions, *Chem. Rev.* 49 (1951) 273-412.
(b) J. F. Bunnett, S. Sekiguchi, L. Smith, Kinetics of Reactions of Cyclic Secondary Amines with 2,4-Dinitro-1-naphthyl Ethyl Ether in Dimethyl Sulfoxide Solution. Spectacular Difference between the Behavior of Pyrrolidine and Piperidine, *J. Am. Chem. Soc.* 103 (1981) 4865-4871.
- 23 (a) S. Tshepelevitsh, A. Kütt, M. Lõkov, I. Kaljurand, J. Saame, A. Heering, P. Plieger, R. Vianello, I. Leito, On the Basicity of Organic Bases in Different Media, *Eur. J. Org. Chem.* 2019 (2019) 6735-6748. (b) J. F. Coetzee, G. R. Padmanabhan, Properties of Bases in Acetonitrile as Solvent. IV. Proton Acceptor Power and Homoconjugation of Mono- and Diamines, *J. Am. Chem. Soc.* 87 (1965) 5005-5010.
- 24 H. Raissi, H. Ayachi, F. Mahdhaoui, S. Ayachi, T. Boubaker, Relationships between experimental and theoretical scales of electrophilicity of 7-L-4-nitrobenzofurazans, *J. Mol. Struct.* 1224 (2021) 128843-128853.
- 25 A. Hunter, M. Renfrew, D. Rettura, J. A. Taylor, J. M. Whitmore, A. Williams, Stepwise versus Concerted Mechanisms at Trigonal Carbon: Transfer of the 1,3,5-Triazinyl Group between Aryl Oxide Ions in Aqueous Solution, *J. Am. Chem. Soc.* 117 (1995) 5484-5491.
- 26 (a) N. Guesmi, T. Boubaker, R. Goumont, Single electron transfer in S_NAr process: Reactions of 2,4,6-tris(trifluoromethanesulfonyl)anisole with anilines in dimethyl sulfoxide, *Prog. React. Kinet. Mech.* 38 (2013) 130-142. (b) N. El Guesmi, T. Boubaker, R. Goumont, F. Terrier, The ambident reactivity of 2,4,6-tris(trifluoromethanesulfonyl)anisole in methanol: Using the SO_2CF_3 group as a tool to reach the superelectrophilic dimension in σ -complexation processes, *Chem. Eur. J.* 15 (2009) 12018-12029. (c) N. El

- Guesmi, T. Boubaker, R. Goumont, Activation of the aromatic system by the SO_2CF_3 group: Kinetics study and structure-reactivity relationships, *Int. J. Chem. Kinet.* 42 (2010) 203-210.
- 27 F. Terrier, M. Mokhtari, R. Goumont, J. C. Halle, E. Buncel, High Bronsted β_{nuc} values in $\text{S}_{\text{N}}\text{Ar}$ displacement. An indicator of the set pathway, *Org. Biomol. Chem.* 1 (2003) 1757-1763.
- 28 M. Jonsson, D. Danial, M. Wayner, J. Luszyk, Redox and Acidity Properties of Alkyl- and Arylamine Radical Cations and the Corresponding Aminyl Radicals, *J. Phys. Chem.* 100 (1996) 17539-17543.
- 29 (a) E. L. Dolengovski, T. V. Gryaznova, O. G. Sinyashin, E. L. Gavrilova, K. V. Kholin, Y. H. Budnikova, Morpholine Radical in the Electrochemical Reaction with Quinoline N-Oxide, *Catalyst.* 13 (2023) 1279-1291. (b) V. Pavlishchuk, A. W. Addison, Conversion constants for redox potentials measured versus different reference electrodes in acetonitrile solutions at 25°C, *Inorg. Chim. Acta.* 298 (2000) 97-102.
- 30 T. Kanzian, T. A. Nigst, A. Maier, S. Pichl, H. Mayr, Nucleophilic Reactivities of Primary and Secondary Amines in Acetonitrile, *Eur. J. Org. Chem.* 36 (2009) 6379-6385.
- 31 D. S. Timofeeva, A. R. Ofial, H. Mayr, Nucleophilic reactivities of Schiff base derivatives of amino acids, *Tetrahedron.* 75 (2019) 459-463.
- 32 F. Necibi, S. B. Salah, J. Roger, J-C. Hierso, T. Boubaker, Ambident electrophilicity of 4-nitrobenzochalcogenadiazoles: Kinetic studies and structure-reactivity relationships, *Int. J. Chem. Kinet.* 52 (2020) 669-680.
- 33 L. Li, J-R. Hsu, H. Zhao, A. R. Ofial, Nucleophilicities of Cyclic α -Diazo Carbonyl Compounds, *Eur. J. Org. Chem.* 26 (2023) 1-6.
- 34 S. Ben Salah, F. Necibi, R. Goumont, T. Boubaker, Electrophilicities of 4-Nitrobenzochalcogenadiazoles, *Chem. Select.* 5 (2020) 7648-7657.
- 35 A. Eitzinger, R. J. Mayer, N. Hampel, P. Mayer, M. Waser, A. R. Ofial, Electrophilic Reactivities of Vinyl *p*-Quinone Methides, *Org. Lett.* 22 (2020) 2182-2186.
- 36 M. Rammah, F. Mahdhaoui, W. Gabsi, T. Boubaker, Quantification of the Electrophilic Reactivities of Benzotriazoles and Structure-Reactivity Relationships, *Chem. Select.* 6 (2021) 4424-4431.
- 37 T. Bug, T. Lemek, H. Mayr, Nucleophilicities of Nitroalkyl Anions, *J. Org. Chem.* 69 (2004) 7565-7576.
- 38 Q. Chen, P. Mayer, H. Mayr, Ethenesulfonyl Fluoride: The Most Perfect Michael Acceptor Ever Found, *Angew. Chem. Int. Ed.* 55 (2016) 12664-12667.


- 39 H. Mayr, A. R. Ofial, Do General Nucleophilicity Scales Exist, *J. Phys. Org. Chem.* 21 (2008) 584-595.
- 40 E. Follet, P. Mayer, H. Mayr, Lewis Acidities of Indol-3-Ylmethylum Ions and Intrinsic Barriers of their Reactions with Phosphines and Pyridines, *Eur. J. Org. Chem.* 23 (2016) 4050-4058.
- 41 S. Ben Salah, T. Boubaker, R. Goumont, Reactions of N1-methyl-4-nitro-2,1,3-benzoselenadiazolium Tetrafluoroborate with Aliphatic and Cyclic Amines in Acetonitrile: Kinetic and Structure-Reactivity Correlations, *Int. J. Chem. Kinet.* 49 (2017) 576-583.
- 42 K. Azaiez, N. Dhahri, T. Boubaker, Reactivity of Nitroalkyl Anions Addition to Substituted Benzylidenecyanoacetate: Electrophilicity Parameters and Free Energy Relationships, *Int. Chem. Kinet.* 49 (2017) 847-858.
- 43 N. Dhahri, T. Boubaker, R. Goumont, Kinetic of Alkaline Hydrolysis of p-Substituted Benzylidenemalonitrile in 50% Aqueous Acetonitrile: Substituent Effects and Quantification of the Electrophilic Reactivity, *J. Phys. Org. Chem.* 27 (2014) 484-489.
- 44 H. Raissi, F. Mahdhaoui, S. Ayachi, T. Boubaker, Solvent effect, quantification and correlation analysis of the nucleophilicities of cyclic secondary amines, *Chem. Papers.* 77 (2023) 307-319.
- 45 F. Necibi, S. Ben Salah, J-C. Hierso, P. Fleurat-Lessard, S. Ayachi, T. Boubaker, Nucleophilicity Parameters for Nitroalkyl Anions in Methanol and Structure-Reactivity Analysis, *Chem. Select.* 8 (2023) 1-8.
- 46 (a) R. G. Parr, W. Yang Density Functional Theory of Atoms and Molecules, Oxford University Press. New York. (1989). (b) R. G. Parr, R. G. Pearson, Absolute hardness: companion parameter to absolute electronegativity, *J. Am. Chem. Soc.* 105 (1983) 7512-7516.
- 47 (a) P. Pérez, Global and local electrophilicity patterns of diazonium ions and their reactivity toward π -nucleophiles, *Org. Chem.* 68 (2003) 5886-5889. (b) P. Pérez, A. Labbé, A. Aizman, R. Contreas, Comparison between experimental and theoretical scales of electrophilicity in benzhydryl cations, *J. Org. Chem.* 67 (2002) 4747-4752.
- 48 L. R. Domingo, P. Pérez, R. Contreras, Reactivity of the carbon-carbon double bond towards nucleophilic additions. A DFT analysis, *Tetrahedron.* 60 (2004) 6585-6591.
- 49 D. S. Allgäuer, H. Jangra, H. Asahara, Z. Li, Q. Chen, H. Zipse, A. R. Ofial, H. Mayr, Quantification and Theoretical Analysis of the Electrophilicities of Michael Acceptors, *J. Am. Chem. Soc.* 139 (2017) 13318-13329.

- 50 H. Raissi, I. Chérif, H. Ayachi, A. Haj Said, F. Hassen, S. Ayachi, T. Boubaker, Structure-Property Relationships in Benzofurazan Derivatives: A Combined Experimental and DFT/TD-DFT Investigation, *IntechOpen Book Series Chapter 17*. (2021)
- 51 E. Chamorro, M. Duque-Noreña, R. Notario, P. Pérez, Intrinsic Relative Scales of Electrophilicity and Nucleophilicity, *J. Phys. Chem.* 12 (2013) 2636-2643.
- 52 W. Gabsi, T. Boubaker, R. Goumont, Azo-coupling Reactions of para-X-benzenediazonium cations with 3-ethoxythiophene in Acetonitrile, *Int. J. Chem. Kinet.* 48 (2016) 266-273.
- 53 (a) M. Rammah, F. Mahdhaoui, S. Ayachi, T. Boubaker, Exploring the reactivity of benzotriazole derivatives: Mayr's approach and density functional theory analysis, *J. Mol. Struct.* 1247 (2022) 131310. (b) A. Laouiti, F. Couty, J. Marrot, T. Boubaker, M. M. Rammah, M. B. Rammah, G. Evano, Exploring the anionic reactivity of ynimines, useful precursors of metalated ketenimines, *Org. Lett.* 16 (2014), 2252-2255.
- 54 E. Chamorro, M. Duque-Noreña, P. Pérez, A comparison between theoretical and experimental models of electrophilicity and nucleophilicity, *J. Mol. Struct.* 896 (2009) 73-79.
- 55 M. Govindarajan, S. Periandy, K. Carthigayen, FT-IR and FT-Raman spectra, thermo dynamical behavior, HOMO and LUMO, UV, NLO properties, computed frequency estimation analysis and electronic structure calculations on α -bromotoluene, *Spectrochim. Acta Part A, Mol. Biomol. Spectro.* 97 (2012) 411-422.
- 56 V. Mishra, A. Raghuvanshi, A. K. Saini, S. M. Mobin, Anthracene derived dinuclear gold(I) diacetylide complexes: Synthesis, photophysical properties and supramolecular interactions, *J. Organomet. Chem.* 813 (2016) 103-109.
- 57 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, D. J. Fox, *Gaussian 09*, Revision A.01, Gaussian: Wallingford, CT., 2009.

- 58 A. D. Becke, Density-functional thermochemistry. III. The role of exact exchange, J. Chem. Phys. 98 (1993) 5648-5652.
- 59 C. Lee, W. Yang, R. G. Parr, Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density, Phys. Rev. B. 37 (1988) 785-789.

Graphical Abstract

The E parameters of various anisoles (see Figure) were derived from the correlation between experimentally measured electrophilicities (E) and global electrophilicity index (ω) calculated at B3LYP/6-311g(d,p) level for a series of pyridines.



AUTHORS CONTRIBUTION

TS, OA and AH achieved her experimental works with the help of TB and J-CG. Theoretical calculations were conducted by SA. The manuscript was jointly written and arranged under TB, SA and JCG guidance.

DECLARATION-OF-COMPETING-INTERESTS

The authors hereby confirm they have no conflict of interest.