



HAL
open science

Two-term large-time asymptotic expansion of the value function for dissipative nonlinear optimal control problems

Veljko Askovic, Emmanuel Trélat, Hasnaa Zidani

► **To cite this version:**

Veljko Askovic, Emmanuel Trélat, Hasnaa Zidani. Two-term large-time asymptotic expansion of the value function for dissipative nonlinear optimal control problems. 2023. hal-04345255

HAL Id: hal-04345255

<https://hal.science/hal-04345255>

Preprint submitted on 14 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Two-term large-time asymptotic expansion of the value function for dissipative nonlinear optimal control problems

Veljko Ašković* Emmanuel Trélat[†] Hasnaa Zidani[‡]

Abstract

Considering a general nonlinear dissipative finite dimensional optimal control problem in fixed time horizon T , we establish a two-term asymptotic expansion of the value function as $T \rightarrow +\infty$. The dominating term is T times the optimal value obtained from the optimal static problem within the classical turnpike theory. The second term, of order unity, is interpreted as the sum of two values associated with optimal stabilization problems related to the turnpike.

1 Introduction

The long-term asymptotic properties of the value function have been extensively explored from the perspective of partial differential equations (PDE), primarily within the framework of Hamilton-Jacobi-Bellman (HJB) equations ([1], [2], [3]) and ergodic theory ([4]). For instance, in [1, Chapter VII], the authors investigate the optimal control problem with discounted Lagrange cost over an infinite time horizon. To characterize the ergodic behavior of the value function, the authors take the limit as the discount factor tends to zero and identify the limit value function as the viscosity solution of the limiting equation. In [2], assuming suitable conditions, including periodicity assumptions on the Hamiltonian, the authors characterize the large-time behavior of the solution to the first-order HJB equation as the solution of a stationary equation. The extension of these results to deterministic zero-sum differential games with two conflicting controllers has been studied, as seen in [3]. It is worth noting that more general results are available, such as those presented in [5], where the authors demonstrate that, under appropriate assumptions, there exists at most one potential accumulation point (in the uniform convergence topology) of the

*MBDA, France (veljkoaskovic@hotmail.com).

[†]Sorbonne Université, CNRS, Université Paris Cité, Inria, Laboratoire Jacques-Louis Lions (LJLL), F-75005 Paris, France (emmanuel.trelat@sorbonne-universite.fr).

[‡]Insa Rouen Normandie, LMI, Saint-Étienne-du-Rouvray, France (hasnaa.zidani@insa-rouen.fr).

values. This occurs when the time horizon of Cesaro means converges to infinity or the discount factor of Abel means converges to zero.

From the classical optimal control perspective, the large-time behavior of the value function is typically inferred as a consequence of a property satisfied by a broad class of optimal control problems when the time horizon is sufficiently large. This property is known as the *turnpike property*, indicating that, for certain optimal control problems with a sufficiently large time horizon, any optimal solution tends to remain close to the optimal solution of an associated *static optimization* problem for the majority of the time. This optimal static solution is referred to as the turnpike (the term originates from the concept that a turnpike represents the fastest route between two distant points, even if it is not the most direct route; see Figure 1).




Figure 1: Turnpike illustration: (a) small time case (b) large time case

The turnpike phenomenon was initially observed and investigated by economists for discrete-time optimal control problems (see [6], [7]). Various notions of turnpike properties exist, with some being stronger than others (see [8]). Exponential turnpike properties have been established in [9], [10], [11], [12] and [13] for the optimal triple resulting of the application of Pontryagin’s Maximum Principle (PMP), ensuring that the extremal solution (state, adjoint and control) remains exponentially close to an optimal solution of the corresponding static controlled problem, except at the beginning and at the end of the time interval, when the time horizon T is sufficiently large. As unravelled in [12] this phenomenon is closely related to hyperbolicity properties of the Hamiltonian flow. For discrete-time problems it has been shown for instance in [14], [15] that the exponential turnpike property is also closely related to a strict dissipativity property. Measure-turnpike is a weaker notion of turnpike, meaning that any optimal solution, along the time frame, remains close to an optimal static solution except during a subset of times with small Lebesgue measure. It has been proved in [16], [17] that measure turnpike follows from strict dissipativity or from strong duality properties.

Based on the turnpike property, an equivalent as $T \rightarrow +\infty$ of the value

function has been derived in [11] or [12], (see Figure 2):

$$\frac{1}{T} \cdot v(T, x, z) \underset{T \rightarrow +\infty}{\sim} \bar{v} \text{ if } \bar{v} \neq 0 \quad (1)$$

where $v(T, x, z)$ is the optimal cost to steer the system from x to z in time T and \bar{v} is the “steady” cost at the turnpike.




Figure 2: Turnpike phenomenon

Significant insights have been achieved in [18] and [19], where the authors explore variations of the linear quadratic (LQ) problem. Assuming certain controllability conditions, they derive the large-time expansion of the value function at the order of one in $1/T$. In [18], the various aspects are identified, and some are interpreted within the framework of the Hamilton-Jacobi-Bellman (HJB) theory. Meanwhile, in [19], all the terms are characterized as functions of the solutions of the Riccati algebraic equation, initial/final states, and the Lagrange multiplier of the static optimization problem.

In this manuscript, we extend the results of [18, 19] to the class of *dissipative* nonlinear systems. The concept of (strict) dissipativity, as introduced in [21], is defined in a broad context, accompanied by related notions like the available storage function and the supply rate function. In cases where a system exhibits dissipativity with a specified supply rate function, the inquiry into identifying a suitable storage function has been extensively investigated (refer to, e.g., [32]). This inquiry bears resemblance to the task of determining an appropriate Lyapunov function in the Lyapunov second method, which ensures the stability of a system. A precise mathematical definition will be provided later in this paper.

2 Setting of the problem

We consider the finite-dimensional optimal control problem, consisting of minimizing the cost functional

$$J_{T,x,z}(u) = \int_0^T f^0(y(t), u(t)) dt \quad (2)$$

over the time interval $[0, T]$, $T > 0$ being fixed under the constraints

$$\dot{y}(t) = f(y(t), u(t)), \quad (3a)$$

$$y(0) = x, \quad y(T) = z, \quad (3b)$$

where $f : \mathbb{R}^n \times \mathbb{R}^p \rightarrow \mathbb{R}^n$ and $f^0 : \mathbb{R}^n \times \mathbb{R}^p \rightarrow \mathbb{R}$ are of class C^1 . The associated value function defined by

$$v(T, x, z) := \min_{u(\cdot) \in \mathcal{U}_T^\Omega} J_{T,x,z}(u) \quad (4)$$

where $u \in \mathcal{U}_T^\Omega := L^\infty([0, T], \Omega)$, Ω being a fixed compact subset of \mathbb{R}^p . In the sequel, we assume that $\Omega \subset \bar{B}(0, c)$ for some $c > 0$.

Remark 2.1. Since Ω does not depend on T , the space of controls \mathcal{U}^Ω is independent of T . This is because, beyond the time horizon T , the controls can be trivially extended within Ω and consequently, the space of controls will be denoted by \mathcal{U}^Ω .

We assume that for T large enough, (2)-(3)-(4) admits an optimal solution, denoted by $(\hat{y}_T(\cdot), \hat{u}_T(\cdot))$. The conditions ensuring the existence of such a solution are well known (see for instance [20], [37]). For instance, if the set of velocities $\{f(y, u) \mid u \in \Omega\}$ is a convex subset of \mathbb{R}^n for any $y \in \mathbb{R}^n$, with mild growth at infinity and if the epigraph of f^0 is convex, then there exists at least one optimal solution. These conditions are for example satisfied if the dynamics are control-affine and if the cost functional is convex with respect to u .

By the Pontryagin maximum principle ([20]-[35]), there exist $\lambda^0 \leq 0$ and an absolutely continuous mapping $\hat{\lambda}_T : [0, T] \rightarrow \mathbb{R}^n$ (called adjoint vector) satisfying $(\hat{\lambda}_T(\cdot), \lambda^0) \neq (0, 0)$ such that:

$$\begin{aligned} \dot{\hat{y}}_T(t) &= \frac{\partial H}{\partial \lambda} \left(\hat{y}_T(t), \hat{\lambda}_T(t), \lambda^0, \hat{u}_T(t) \right) \\ \dot{\hat{\lambda}}_T(t) &= -\frac{\partial H}{\partial y} \left(\hat{y}_T(t), \hat{\lambda}_T(t), \lambda^0, \hat{u}_T(t) \right) \\ \frac{\partial H}{\partial u} \left(\hat{y}_T(t), \hat{\lambda}_T(t), \lambda^0, \hat{u}_T(t) \right) &= 0 \end{aligned} \quad (5)$$

for almost every $t \in [0, T]$, where the Hamiltonian H is defined by

$$H(y, \lambda, \lambda^0, u) := \langle \lambda, f(y, u) \rangle + \lambda^0 f^0(y, u) \quad (6)$$

and $\langle \cdot, \cdot \rangle$ is the Euclidean scalar product in \mathbb{R}^n . We assume throughout that the abnormal case does not occur, and we set $\lambda^0 = -1$. Moreover, we assume the existence and uniqueness of the solution (denoted by (\bar{y}, \bar{u})) to the *static optimization problem*

$$\bar{v} := \min_{f(y,u)=0} f^0(y, u). \quad (7)$$

This is a usual nonlinear constrained optimization problem settled in $\mathbb{R}^n \times \mathbb{R}^p$. Note that the minimizer exists and is unique in the linear quadratic case.

By the Karush Kuhn Tucker (KKT) optimality conditions, assuming that the abnormal case does not occur, there exists $\bar{\lambda} \in \mathbb{R}^n$ such that

$$\begin{aligned} f(\bar{y}, \bar{u}) &= 0 \\ -\frac{\partial f^0}{\partial y}(\bar{y}, \bar{u}) + \langle \bar{\lambda}, \frac{\partial f}{\partial y}(\bar{y}, \bar{u}) \rangle &= 0 \\ -\frac{\partial f^0}{\partial u}(\bar{y}, \bar{u}) + \langle \bar{\lambda}, \frac{\partial f}{\partial u}(\bar{y}, \bar{u}) \rangle &= 0 \end{aligned} \quad (8)$$

In the Hamiltonian formalism, the first-order optimality system (8) is equivalent to

$$\begin{aligned} \frac{\partial H}{\partial \lambda}(\bar{y}, \bar{\lambda}, -1, \bar{u}) &= 0 \\ -\frac{\partial H}{\partial y}(\bar{y}, \bar{\lambda}, -1, \bar{u}) &= 0 \\ \frac{\partial H}{\partial u}(\bar{y}, \bar{\lambda}, -1, \bar{u}) &= 0 \end{aligned} \quad (9)$$

Because of the turnpike property (see further), since it is expected that the dominating term in the asymptotic expansion of $v(\cdot)$ is $T \cdot \bar{v}$, we "absorb" it by subtracting it to the cost $J_{T,x,z}$ and consider the "shifted" optimal control problem defined as

$$\min_{u(\cdot) \in \mathcal{U}^\Omega} C_T(u, x, z) := \min_{u(\cdot) \in \mathcal{U}^\Omega} \int_0^T w(y(t), u(t)) dt \quad (10a)$$

$$\left(\mathcal{P}_{[0,T]}^{x,z} \right) \quad \dot{y}(t) = f(y(t), u(t)), \quad \forall t \in [0, T] \quad (10b)$$

$$y(0) = x, \quad y(T) = z \quad (10c)$$

where $w(\cdot)$ is the "shifted cost" defined by

$$w(y, u) := f^0(y, u) - f^0(\bar{y}, \bar{u}). \quad (11)$$

The readers acquainted with the notion of dissipativity in nonlinear optimal control will recognize in (11) a classical storage function. This is where the link with dissipativity appears.

We introduce also the shifted infinite-time optimal control problems

$$\begin{aligned} (\mathcal{P}_{\infty}^x) \quad v_f(x) &:= \min_{u(\cdot) \in \mathcal{U}^\Omega} \int_0^{+\infty} w(y(t), u(t)) dt \\ \dot{y}(t) &= f(y(t), u(t)), \quad y(0) = x \end{aligned} \quad (12)$$

and

$$\begin{aligned}
(\mathcal{P}_{\infty b}^z) \quad v_b(z) &:= \min_{u(\cdot) \in \mathcal{U}^\Omega} \int_0^{+\infty} w(y(t), u(t)) dt \\
\dot{y}(t) &= -f(y(t), u(t)), \quad y(0) = z
\end{aligned} \tag{13}$$

In our notations the index “f” stands for “forward” while the index “b” stands for “backward”. We assume the existence of optimal solutions to $(\mathcal{P}_{\infty f}^x)$ and $(\mathcal{P}_{\infty b}^z)$ with a finite cost.

Remark 2.2. The function $v_f(\cdot)$ (resp. $v_b(\cdot)$) is the optimal cost of the infinite-time optimal control problem consisting in steering the system from x (resp. z) forward (resp. backward) in time and minimizing the cost functional $\int_0^{+\infty} w(y(t), u(t)) dt$.

3 Assumptions

Assumptions of global nature:

(A₁): (*Regularity of f and f^0*): We assume that f and f^0 are of class C^1 .

(A₂): (*Existence and uniqueness of optimal solutions*): There exists $T_0 > 0$ such that for any $T \geq T_0$ each of the optimal control problems $(\mathcal{P}_{[0,T]}^{x,z})$, $(\mathcal{P}_{\infty f}^x)$ and $(\mathcal{P}_{\infty b}^z)$ admits a unique optimal solution denoted respectively by $(\hat{y}_T(\cdot), \hat{u}_T(\cdot))$, $(\hat{y}_{\infty f}(\cdot), \hat{u}_{\infty f}(\cdot))$ and $(\hat{y}_{\infty b}(\cdot), \hat{u}_{\infty b}(\cdot))$.

(A₃): (*Boundedness of the optimal trajectories*): The optimal trajectories of $(\mathcal{P}_{[0,T]}^{x,z})$, $(\mathcal{P}_{\infty f}^x)$ and $(\mathcal{P}_{\infty b}^z)$ are bounded uniformly with respect to $T \geq T_0$:

$$\exists b > 0, \quad | \quad \forall t \geq 0, \quad \|\hat{y}(t)\| \leq b. \tag{14}$$

(A₄): (*Boundedness of the optimal costs*): The optimal costs of $(\mathcal{P}_{[0,T]}^{x,z})$, $(\mathcal{P}_{\infty f}^x)$ and $(\mathcal{P}_{\infty b}^z)$ are bounded uniformly with respect to $T \geq T_0$.

(A₅): The minimizer (\bar{y}, \bar{u}) of (7) is unique and there exists a unique Lagrange multiplier $\bar{\lambda}$ satisfying (8). Moreover, we assume that $\bar{u} \in \overset{\circ}{\Omega}$ (interior of Ω).

In the sequel, we will also need a concept of *strict dissipativity*. We recall that (10a)-(10b) is *dissipative* at (\bar{y}, \bar{u}) with respect to the supply rate function w if there exists a bounded function $S : \mathbb{R}^n \rightarrow \mathbb{R}$, called *storage function* such that for any admissible pair $(y(\cdot), u(\cdot))$ and any $T > 0$:

$$S(y(0)) + \int_0^T w(y(t), u(t)) dt \geq S(y(T)). \tag{15}$$

The system is *strictly dissipative* if, in addition, there exists some function $\alpha(\cdot)$ of class \mathcal{K} (i.e., $\alpha : [0, +\infty) \rightarrow [0, +\infty)$ continuous, increasing, and such that

$\alpha(0) = 0$) and for any $T > 0$ we have:

$$S(y(0)) + \int_0^T w(y(t), u(t)) dt \geq S(y(T)) + \int_0^T \alpha \left(\left\| \begin{array}{c} y(t) - \bar{y} \\ u(t) - \bar{u} \end{array} \right\| \right) dt. \quad (16)$$

In this work we assume the following.

(A₆): (*Strict dissipativity property*): The family of optimal control problems (10a)-(10b) indexed by T is strictly dissipative at (\bar{y}, \bar{u}) with respect to the supply rate function w defined by (11) with a storage function S .

The notion of strict dissipativity was introduced in [21] and already used in [14], [22], [16] and [15] to derive turnpike properties.

Assumption of local nature:

(A₇): Setting $A := \frac{\partial f}{\partial y}(\bar{y}, \bar{u})$, $B := \frac{\partial f}{\partial u}(\bar{y}, \bar{u})$, the pair (A, B) satisfies the Kalman rank condition, i.e., the linearized control system at (\bar{y}, \bar{u}) is controllable.

(A₈): (*Local boundedness of the minimum time trajectories and controls near the turnpike*): There exists $r > 0$ such that for any $x \in \bar{B}(\bar{y}, r)$ the minimum time trajectory to \bar{y} starting from x , denoted by $y_x^{\tau_f}(\cdot)$ and the associated control, denoted by $u^{\tau_f}(\cdot)$ remain in the neighbourhood of respectively \bar{y} and \bar{u} uniformly with respect to x , i.e.,

$$\exists r, K_r > 0 \mid \forall x \in \bar{B}(\bar{y}, r) \quad \forall t \in [0, \tau_f(x)] \quad \|y_x^{\tau_f}(t) - \bar{y}\| + \|u^{\tau_f}(t) - \bar{u}\| \leq K_r \quad (17)$$

where $\tau_f(\cdot)$ is the minimum time function to reach \bar{y} with the dynamics f .

4 Comments

The assumptions (A₁) and (A₇) together imply that there exists $r > 0$ such that for any $x \in B(\bar{y}, r)$, there exists an admissible trajectory steering the control system from x to \bar{y} in finite time. Thus the minimum time function $\tau_f(\cdot)$ is well defined on $B(\bar{y}, r)$ and is continuous at \bar{y} . This is classical result that can be found, for instance, in [1]. The result remains true for the minimum time function associated to the backward-in-time dynamics $-f$, denoted by $\tau_{-f}(\cdot)$.

The assumption (A₈) requires the local boundedness of the minimum time trajectories and controls for any trajectory starting in the previously defined neighbourhood of the turnpike. This assumption is satisfied if the minimum time function is C^1 in the neighbourhood of \bar{y} . We highlight here that the regularity of the minimum time function has been widely studied in the literature: it is well known that under appropriate controllability type conditions the minimum time function has an open domain of definition and is locally Lipschitz on it, see for instance [1]-[23]. It is thus differentiable almost everywhere on its domain. The value function fails in general to be differentiable at points that are reached by at least two minimum time trajectories and its differentiability at a point does not guarantee continuous differentiability around this

point. In [24], the authors show that, under some assumptions on the regularity and target smoothness (which excludes the singleton case), the nonemptiness of the proximal subdifferential of the minimum time function at a point implies its continuous differentiability in a neighborhood of this point. An analogous result has been proved for the value function of the Bolza problem in [25] in the case where the initial state is a prescribed point and the final state is let free. In [26], the author gives a survey of results on the regularity of the minimum time map for control-affine systems with prescribed initial and final points. Finally, for results on the set where the value function is differentiable we refer the reader to [27], [28], [29], [30], [31] and references therein.

The (strict) dissipativity property (A_6) is certainly the less intuitive assumption to check in practice. In general, when the system is dissipative, storage functions are closely related to some viscosity sub-solutions of partial differential inequalities called Hamilton-Jacobi inequalities. We refer the reader to [32, Chapter 4] for more details on this subject. One can remark that, under suitable regularity and boundedness assumptions on the dynamics and the cost, the value function (its opposite more precisely) can be taken as a storage function, and the dissipativity inequality is then deduced from the Dynamic Programming Principle (DPP). Note also that the infinitesimal form of the (non-strict) dissipativity inequality (15), with $\alpha = 0$, is the Hamilton-Jacobi inequality:

$$H_1(y, \nabla S(y), u) \leq -f^0(\bar{y}, \bar{u})$$

where H_1 is the maximized normal Hamiltonian (indeed, divide by $t_1 - t_0 > 0$ and take the limit $t_1 - t_0 \rightarrow 0$). The existence of C^1 solutions is therefore related to the so-called weak KAM theory (see [38]). In this context, the singleton $\{(\bar{y}, \bar{u})\}$ is the Aubry set and $f^0(\bar{y}, \bar{u})$ is the Mañé critical value.

Remark 4.1. The strict dissipativity inequality (16) remains true for the dynamics $-f$, provided that one switches the initial and final states. The corresponding storage function is $-S(\cdot)$.

5 Main result

Theorem 5.1. *Under Assumptions (A_1)–(A_8), the value function (4) satisfies*

$$v(T, x, z) = T\bar{v} + v_f(x) + v_b(z) + o(1) \tag{18}$$

as $T \rightarrow +\infty$.

In order to prove Theorem 5.1, we need some preliminary lemmas.

5.1 Some useful lemmas

Lemma 5.2 (Barbalat's lemma). *Assume that $f : [0, +\infty) \rightarrow \mathbb{R}$ is uniformly continuous and that $\lim_{t \rightarrow +\infty} \int_0^t f(\tau) d\tau$ exists and is finite, then $\lim_{t \rightarrow +\infty} f(t) = 0$.*

Proof. A proof can be found for instance in [36]. By contradiction, take $\epsilon > 0$ and assume that $f(t)$ does not converge to 0 as $t \rightarrow +\infty$. In this case, there exists an increasing sequence $(t_n)_{n \in \mathbb{N}}$ in \mathbb{R}^+ such that $|f(t_n)| > \epsilon$. By the uniform continuity of f there exists $\delta > 0$ such that, for any $n \in \mathbb{N}$, and any $t \in \mathbb{R}$

$$|t - t_n| \leq \delta \implies |f(t) - f(t_n)| \leq \frac{\epsilon}{2}.$$

So for any $t \in [t_n, t_n + \delta]$ and any $n \in \mathbb{N}$, one has

$$|f(t)| = |f(t_n) - (f(t_n) - f(t))| \geq |f(t_n)| - |f(t_n) - f(t)| \geq \frac{\epsilon}{2}.$$

Therefore,

$$\left| \int_0^{t_n+\delta} f(t)dt - \int_0^{t_n} f(t)dt \right| = \left| \int_{t_n}^{t_n+\delta} f(t)dt \right| = \int_{t_n}^{t_n+\delta} |f(t)|dt \geq \frac{\delta \epsilon}{2}.$$

The latter inequality contradicts the convergence of $\int_0^t f(\tau)d\tau$ as $t \rightarrow +\infty$ and the lemma follows. \square

Lemma 5.3. *The optimal trajectory $\hat{y}_T(\cdot)$ of $(\mathcal{P}_{0,T}^{x,z})$ satisfies*

$$\exists t(T) \in [0, T] \quad | \quad \hat{y}_T(t(T)) \longrightarrow \bar{y} \text{ as } T \rightarrow +\infty. \quad (19)$$

Proof. For $T \geq T_0$, the strict dissipativity inequality applied to the optimal pair $(\hat{y}_T(\cdot), \hat{u}_T(\cdot))$ implies

$$\begin{aligned} f^0(\bar{y}, \bar{u}) &\leq \frac{1}{T} \int_0^T f^0(\hat{y}_T(s), \hat{u}_T(s))ds + \frac{S(x) - S(z)}{T} \\ &\quad - \frac{1}{T} \int_0^T \alpha \left(\left\| \begin{array}{c} \hat{y}_T(s) - \bar{y} \\ \hat{u}_T(s) - \bar{u} \end{array} \right\| \right) ds. \end{aligned} \quad (20)$$

Let us prove that

$$\frac{1}{T} \int_0^T \alpha \left(\left\| \begin{array}{c} \hat{y}_T(s) - \bar{y} \\ \hat{u}_T(s) - \bar{u} \end{array} \right\| \right) ds \longrightarrow 0 \text{ as } T \rightarrow +\infty. \quad (21)$$

Let us assume by contradiction that this is not true: then there exists $\eta > 0$ and a sequence $T_k \rightarrow +\infty$ such that

$$\frac{1}{T_k} \int_0^{T_k} \alpha \left(\left\| \begin{array}{c} \hat{y}_{T_k}(s) - \bar{y} \\ \hat{u}_{T_k}(s) - \bar{u} \end{array} \right\| \right) ds \geq \eta. \quad (22)$$

By multiplying the inequality (20) by T_k one gets

$$\int_0^{T_k} \alpha \left(\left\| \begin{array}{c} \hat{y}_{T_k}(s) - \bar{y} \\ \hat{u}_{T_k}(s) - \bar{u} \end{array} \right\| \right) ds \leq \int_0^{T_k} w(\hat{y}_{T_k}(s), \hat{u}_{T_k}(s))ds + S(x) - S(z) \quad (23)$$

which implies from (22)

$$T_k \cdot \eta \leq \int_0^{T_k} w(\hat{y}_{T_k}(s), \hat{u}_{T_k}(s)) ds + S(x) - S(z). \quad (24)$$

Assumption (A₄) leads to a contradiction in the above inequality when $k \rightarrow +\infty$.

Then we have $\frac{1}{T} \int_0^T \alpha \left(\left\| \begin{array}{c} \hat{y}_T(t) - \bar{y} \\ \hat{u}_T(t) - \bar{u} \end{array} \right\| \right) ds \geq \frac{1}{T} \int_0^T \alpha(\|\hat{y}_T(t) - \bar{y}\|) ds \rightarrow 0$ as $T \rightarrow +\infty$ which implies, using the mean value theorem, that

$$\exists t(T) \in [0, T] \quad | \quad \alpha(\|\hat{y}_T(t(T)) - \bar{y}\|) \rightarrow 0 \text{ as } T \rightarrow +\infty. \quad (25)$$

From the properties of $\alpha(\cdot)$, this leads to

$$\hat{y}_T(t(T)) \rightarrow \bar{y} \text{ as } T \rightarrow +\infty. \quad (26)$$

The lemma is proved. \square

Remark 5.4. As noted in [33], if one makes the change of variable $s = \frac{t}{T}$, in (21) then

$$\frac{1}{T} \int_0^T \alpha \left(\left\| \begin{array}{c} \hat{y}_T(t) - \bar{y} \\ \hat{u}_T(t) - \bar{u} \end{array} \right\| \right) dt = \int_0^1 \alpha \left(\left\| \begin{array}{c} \hat{y}_T(T \cdot s) - \bar{y} \\ \hat{u}_T(T \cdot s) - \bar{u} \end{array} \right\| \right) ds \rightarrow 0 \text{ as } T \rightarrow +\infty$$

which implies, by the converse Lebesgue theorem (see [34, Theorem IV.9]) that there exists an increasing sequence of time horizons $(T_k)_{k \in \mathbb{N}}$ such that $\hat{y}_{T_k}(T_k \cdot s) \rightarrow \bar{y}$ and $\hat{u}_{T_k}(T_k \cdot s) \rightarrow \bar{u}$ as $k \rightarrow +\infty$ for almost every $s \in [0, 1]$. The latter looks like a measure turnpike result. However we did not exploit this result in our paper.

Lemma 5.5. *The optimal trajectory $\hat{y}_{\infty f}(\cdot)$ of $(\mathcal{P}_{\infty f}^x)$ satisfies*

$$\hat{y}_{\infty f}(t) \rightarrow \bar{y} \text{ as } t \rightarrow +\infty \quad (27)$$

Proof. From (16) we have

$$\int_0^T \alpha \left(\left\| \begin{array}{c} \hat{y}_{\infty f}(t) - \bar{y} \\ \hat{u}_{\infty f}(t) - \bar{u} \end{array} \right\| \right) dt \leq \int_0^T w(\hat{y}_{\infty f}(t), \hat{u}_{\infty f}(t)) dt + S(x) - S(\hat{y}_{\infty f}(T)) \quad (28)$$

the right-hand side of the inequality being bounded uniformly with respect to T one gets

$$\int_0^T \alpha(\|\hat{y}_{\infty f}(t) - \bar{y}\|) dt \leq \int_0^T \alpha \left(\left\| \begin{array}{c} \hat{y}_{\infty f}(t) - \bar{y} \\ \hat{u}_{\infty f}(t) - \bar{u} \end{array} \right\| \right) dt \underset{T \rightarrow +\infty}{=} O(1) \quad (29)$$

which implies

$$\Phi(T) := \int_0^T \alpha(\|\hat{y}_{\infty f}(t) - \bar{y}\|) dt \underset{T \rightarrow +\infty}{=} O(1). \quad (30)$$

First, we remark that (30) and the positivity of $\alpha(\cdot)$ imply the convergence of $\Phi(T)$ as $T \rightarrow +\infty$.

On the other hand, from (A_1) and (A_3) we know that f is of class C^1 on the compact set $\overline{B}(0, b) \times \overline{B}(0, c)$ thus bounded by a global constant, denoted by $k > 0$. Therefore $\hat{y}_{\infty f}(\cdot)$ is globally Lipschitz continuous in time t , and consequently $t \mapsto \|\hat{y}_{\infty f}(t) - \bar{y}\|$ as well.

From the boundedness of $\hat{y}_{\infty f}(\cdot)$ (see (A_3)) and the continuity of $\alpha(\cdot)$, we deduce the uniform continuity of $t \mapsto \alpha(\|\hat{y}_{\infty f}(t) - \bar{y}\|)$ over $[0 + \infty)$. Indeed, $\alpha(\cdot)$ being continuous on a compact set, it is uniformly continuous (Heine theorem).

Applying Barbalat's lemma (see 5.2), one has $\alpha(\|\hat{y}_{\infty f}(t) - \bar{y}\|) \rightarrow 0$ as $t \rightarrow +\infty$, which implies $\hat{y}_{\infty f}(t) \rightarrow \bar{y}$ as $t \rightarrow +\infty$ and the proof is over. \square

Corollary 5.6. *Lemma 5.5 remains true for the optimal trajectory of $(\mathcal{P}_{\infty b}^z)$ problem (with the backward-in-time dynamics $-f$).*

5.2 Proof of the main result

Proof. For $T \geq T_0$, let us consider the respective optimal solutions $(\hat{y}_T(\cdot), \hat{u}_T(\cdot))$, $(\hat{y}_{\infty f}(\cdot), \hat{u}_{\infty f}(\cdot))$ and $(\hat{y}_{\infty b}(\cdot), \hat{u}_{\infty b}(\cdot))$ of $(\mathcal{P}_{[0, T]}^{x, z})$, $(\mathcal{P}_{\infty f}^x)$ and $(\mathcal{P}_{\infty b}^z)$.

We split the optimal cost $C_T(\hat{u}_T, x, z)$ as

$$C_T(\hat{u}_T, x, z) = \int_0^T w(\hat{y}_T(t), \hat{u}_T(t)) dt \quad (31)$$

$$= \underbrace{\int_0^{t(T)} w(\hat{y}_T(t), \hat{u}_T(t)) dt}_{C_T^f} + \underbrace{\int_{t(T)}^T w(\hat{y}_T(t), \hat{u}_T(t)) dt}_{C_T^b} \quad (32)$$

where $t(T)$ is defined by (19).

We perform the proof in two steps. We first prove that

$$\textbf{Step 1:} \quad v_f(x) + v_b(z) \leq \liminf_{T \rightarrow +\infty} C_T(\hat{u}_T, x, z) \quad (33)$$

Then we prove that:

$$\textbf{Step 2:} \quad \limsup_{T \rightarrow +\infty} C_T(\hat{u}_T, x, z) \leq v_f(x) + v_b(z) \quad (34)$$

which will prove the required result.

The real number r being defined in Assumption (A_8) , we first remark that $(y, u) \mapsto w(y, u)$ is continuous on $\Omega_r := \overline{B}(\bar{y}, K_r) \times \overline{B}(\bar{u}, K_r)$ which is a compact set of $\mathbb{R}^n \times \mathbb{R}^p$. Consequently,

$$\exists M_r > 0 \quad | \quad \forall (y, u) \in \Omega_r, \quad |w(y, u)| \leq M_r. \quad (35)$$

Let $\epsilon > 0$. The continuity of $\tau_f(\cdot)$ at \bar{y} gives:

$$\exists \eta > 0 \text{ s.t. } \|x - \bar{y}\| \leq \eta \Rightarrow |\tau_f(x)| \leq \frac{\epsilon}{2.M_r}. \quad (36)$$

The continuity of $\tau_{-f}(\cdot)$ at \bar{y} gives:

$$\exists \nu > 0 \text{ s.t. } \|x - \bar{y}\| \leq \nu \Rightarrow |\tau_{-f}(x)| \leq \frac{\epsilon}{2.M_r}. \quad (37)$$

We set $\gamma := \min(\eta, \nu, r) > 0$, and we denote by $B := B(\bar{y}, \gamma)$.

▷ **Step 1:** From Lemma 5.3, we know that:

$$\exists T_1 \geq 0 \text{ s.t. } \forall T \geq T_1, \hat{y}_T(t(T)) \in \bar{B}. \quad (38)$$

We select a time horizon T such that $T \geq \max(T_0, T_1)$ and construct $\check{u}(\cdot)$ admissible control for the (\mathcal{P}_{∞}^x) problem as follows:

$$\check{u}(t) := \begin{cases} \hat{u}_T(t) & \text{if } t \in [0, t(T)] \\ \hat{u}_0(t) & \text{if } t \in [t(T), t(T) + \tau_0] \\ \bar{u} & \text{if } t \geq t(T) + \tau_0 \end{cases}$$

where $\tau_0 := \tau_f(\hat{y}_T(t(T)))$ is the minimum time to reach \bar{y} from $\hat{y}_T(t(T))$ and $\hat{u}_0(\cdot)$ the associated optimal control.

We infer from (36) and (38) that

$$\begin{aligned} v_f(x) &\leq \int_0^{+\infty} w(\check{y}(t), \check{u}(t)) dt \\ &\leq \int_0^{t(T)} w(\hat{y}_T(t), \hat{u}_T(t)) dt + \int_{t(T)}^{t(T)+\tau_0} \underbrace{w(\hat{y}_0(t), \hat{u}_0(t))}_{\in \Omega_r} dt + \int_{t(T)+\tau_0}^{+\infty} \underbrace{w(\bar{y}, \bar{u})}_0 dt \\ &\leq C_T^f + \tau_0.M_r \\ &\leq C_T^f + \frac{\epsilon}{2} \end{aligned} \quad (39)$$

For the second term C_T^b , we first remark that

$$C_T^b = \int_{t(T)}^T w(\hat{y}_T(t), \hat{u}_T(t)) dt = \int_0^{T-t(T)} w(\tilde{y}_T(t), \tilde{u}_T(t)) dt \quad (40)$$

where $(\tilde{y}_T(t), \tilde{u}_T(t)) := (\hat{y}_T(T-t), \hat{u}_T(T-t))$ is such that

$$\dot{\tilde{y}}_T(t) = -f(\tilde{y}_T(t), \tilde{u}_T(t)) \text{ with } \tilde{y}_T(0) = z \quad (41)$$

Noting that the $\tilde{y}_T(T-t(T)) = \hat{y}_T(t(T))$, we construct $\check{u}(\cdot)$ the admissible control for the (\mathcal{P}_{∞}^z) problem as follows

$$\check{u}(t) := \begin{cases} \tilde{u}_T(t) & \text{if } t \in [0, T-t(T)] \\ \hat{u}_1(t) & \text{if } t \in [T-t(T), T-t(T) + \tau_1] \\ \bar{u} & \text{if } t \geq T-t(T) + \tau_1 \end{cases}$$

where $\tau_1 := \tau_{-f}(\tilde{y}_T(T - t(T)))$ is the minimum time to reach \bar{y} from $\tilde{y}_T(T - t(T))$ and $\hat{u}_1(\cdot)$ the associated optimal control.

We infer from (37) and (38) that

$$\begin{aligned}
v_b(z) &\leq \int_0^{+\infty} w(\check{y}(t), \check{u}(t)) dt \\
&\leq \int_0^{T-t(T)} w(\tilde{y}_T(t), \tilde{u}_T(t)) dt \\
&\quad + \int_{T-t(T)}^{T-t(T)+\tau_1} \underbrace{w(\hat{y}_1(t), \hat{u}_1(t))}_{\in \Omega_r} dt + \int_{T-t(T)+\tau_1}^{+\infty} \underbrace{w(\bar{y}, \bar{u})}_0 dt \\
&\leq C_T^b + \tau_1 \cdot M_r \\
&\leq C_T^b + \frac{\epsilon}{2}
\end{aligned} \tag{42}$$

Combining (39) and (42), we obtain

$$v_f(x) + v_b(z) \leq C_T^f + C_T^b + \epsilon = C_T(\hat{u}_T, x, z) + \epsilon \tag{43}$$

and thus

$$v_f(x) + v_b(z) \leq \liminf_{T \rightarrow +\infty} C_T(\hat{u}_T, x, z) \tag{44}$$

▷ **Step 2:** From Lemma 5.5 and Corollary 5.6, we have $\hat{y}_{\infty f}(t) \rightarrow \bar{y}$ and $\hat{y}_{\infty b}(t) \rightarrow \bar{y}$ as $t \rightarrow +\infty$. Consequently:

$$\exists T_2 > 0 \text{ s.t. } \forall T \geq T_2, \left\| \hat{y}_{\infty f} \left(\frac{T}{2} - 1 \right) - \bar{y} \right\| \leq \gamma \tag{45}$$

and

$$\exists T_3 > 0 \text{ s.t. } \forall T \geq T_3, \left\| \hat{y}_{\infty b} \left(\frac{T}{2} - 1 \right) - \bar{y} \right\| \leq \gamma \tag{46}$$

Take $T \geq \max(T_2, T_3)$ and denote:

- $\tau_3 := \tau_f \left(\hat{y}_{\infty f} \left(\frac{T}{2} - 1 \right) \right)$ and $\hat{u}_3(\cdot)$ the associated optimal control;
- $\tau_4 := \tau_{-f} \left(\hat{y}_{\infty b} \left(\frac{T}{2} - 1 \right) \right)$ and $\hat{u}_4(\cdot)$ the associated optimal control.

We construct the admissible control $\underline{u}_T(\cdot)$ for $(\mathcal{P}_{0,T})_{x,z}$ as follows (see Figure




Figure 3: Construction of an admissible trajectory for $(\mathcal{P}_{[0,T]}^{x,z})$

3):

$$\underline{u}_T(t) := \begin{cases} \hat{u}_{\infty f}(t) & \text{if } t \in \left[0, \frac{T}{2} - 1\right] \\ \hat{u}_3(t) & \text{if } t \in \left[\frac{T}{2} - 1, \frac{T}{2} - 1 + \tau_3\right] \\ \bar{u} & \text{if } t \in \left[\frac{T}{2} - 1 + \tau_3, \frac{T}{2} + 1 - \tau_4\right] \\ \hat{u}_4(T-t) & \text{if } t \in \left[\frac{T}{2} + 1 - \tau_4, \frac{T}{2} + 1\right] \\ \hat{u}_{\infty b}(T-t) & \text{if } t \in \left[\frac{T}{2} + 1, T\right] \end{cases}$$

We have then the upper bound for the optimal cost:

$$C_T(\hat{u}_T, x, z) \leq \int_0^T w(\underline{y}_T(t), \underline{u}_T(t)) dt \leq A + B + C + D + E \quad (47)$$

where

$$A := \int_0^{\frac{T}{2}-1} w(\hat{y}_{\infty f}(t), \hat{u}_{\infty f}(t)) dt$$

$$E := \int_{\frac{T}{2}+1}^T w(\hat{y}_{\infty b}(T-t), \hat{u}_{\infty b}(T-t)) dt = \int_0^{\frac{T}{2}-1} w(\hat{y}_{\infty b}(t), \hat{u}_{\infty b}(t)) dt$$

$$\begin{aligned}
B &:= \int_{\frac{T}{2}-1}^{\frac{T}{2}-1+\tau_3} \underbrace{w(\hat{y}_3(t), \hat{u}_3(t))}_{\in \Omega_r} dt \leq \tau_3 \cdot M_r \leq \frac{\epsilon}{2} \\
C &:= \int_{\frac{T}{2}-1+\tau_3}^{\frac{T}{2}+1-\tau_4} \underbrace{w(\bar{y}, \bar{u})}_0 dt = 0 \\
D &:= \int_{\frac{T}{2}+1-\tau_4}^{\frac{T}{2}+1} w(\hat{y}_4(T-t), \hat{u}_4(T-t)) dt = \int_{\frac{T}{2}-1}^{\frac{T}{2}-1+\tau_4} \underbrace{w(\hat{y}_4(t), \hat{u}_4(t))}_{\in \Omega_r} dt \leq \tau_4 \cdot M_r \leq \frac{\epsilon}{2}
\end{aligned}$$

Finally, we obtain

$$C_T(\hat{u}_T, x, z) \leq \int_0^{\frac{T}{2}-1} w(\hat{y}_{\infty f}(t), \hat{u}_{\infty f}(t)) dt + \int_0^{\frac{T}{2}-1} w(\hat{y}_{\infty b}(t), \hat{u}_{\infty b}(t)) dt + \epsilon \tag{48}$$

Noting that the integrals A and E converge, we take the limit superior as $T \rightarrow +\infty$ of the above inequality and we obtain:

$$\limsup_{T \rightarrow +\infty} C_T(\hat{u}_T, x, z) \leq v_f(x) + v_b(z). \tag{49}$$

▷ **Conclusion:** By combining (44) and (49), we obtain the required result:

$$\lim_{T \rightarrow +\infty} C_T(\hat{u}_T, x, z) = v_f(x) + v_b(z). \tag{50}$$

□

Remark 5.7. In the LQ case, a two-term large-time expansion of the value function has been derived in [19] solely under the Kalman condition. Assumptions such as the strict dissipativity property and the existence/uniqueness of the solution of the static optimization problem are automatically satisfied in this case. For the proof, see [19].

6 Conclusions

Considering a general nonlinear dissipative optimal control problem in finite dimension and fixed time T , we have established a two-term asymptotic expansion of the value function as $T \rightarrow +\infty$. Essentially based on the strict dissipativity property, the result is a generalization of the expansion established in the variants of the linear quadratic case treated in [18] and [19]. We highlight here that a generalization of [19] to the infinite dimensional case is currently being finalized. The infinite dimensional nonlinear case remains open.

References

- [1] M. Bardi, I. Capuzzo-Dolcetta, Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman Equations, Modern Birkhäuser Classics, 1997
- [2] G. Barles, P. Souganidis, On the Large Time Behavior of Solutions of Hamilton-Jacobi Equations, *SIAM J. Mathematical Analysis*, 31(4), pp. 925-939, 2000
- [3] O. Alvarez, M. Bardi, Ergodic Problems in Differential Games, in *Advances in Dynamic Game Theory: Numerical Methods, Algorithms, and Applications to Ecology and Economics*. Editors: S. Jørgensen, M. Quincampoix, and T.L Vincent. pp. 131-152, 2007
- [4] M. Arisawa, Ergodic problem for the Hamilton Jacobi-Bellman equation i. existence of the ergodic attractor, *Annales de l'Institut Henri Poincaré, Nonlinear Analysis*, vol. 14, pp. 415-438, 1997
- [5] R. Buckdahn, M. Quincampoix, J. Renault, On Representation Formulas for Long Run Averaging Optimal Control Problem, *J. Differential Equations*, 259(11), pp. 5554-5581, 2015
- [6] P.A. Samuelson, R. Dorfman and R.M. Solow, *Linear Programming and Economic Analysis*, McGraw-Hill, New York, 1958
- [7] L. W. McKenzie, Turnpike theorems for a generalized leontief model, *Econometrica*, 31(1/2) 165-180, 1963
- [8] A. J. Zaslavski, *Turnpike theory of continuous-time linear optimal control problems*, Springer Optimization and Its Applications, 2015
- [9] L. Grune, M. Schaller, A. Schiela, Exponential sensitivity and turnpike analysis for linear quadratic optimal control of general evolution equations, *J. Differential Equations*, 268(12), 7311-7341, 2020
- [10] A. Porretta, E. Zuazua, Remarks on long time versus steady state optimal control, in *Mathematical paradigms of climate science, volume 15 of Springer INdAM Series*, pp. 67-89, 2016
- [11] A. Porretta, E. Zuazua, Long time versus steady state optimal control, *SIAM J. Control Optim*, 51(6), pp. 4242-4273, 2013
- [12] E. Trélat, E. Zuazua, The turnpike property in finite-dimensional nonlinear optimal control, *Journal of Differential Equations*, 258 no.1, pp. 81-114, 2015
- [13] E. Trélat, C. Zhang, E. Zuazua, Steady-state and periodic exponential turnpike property for optimal control problems in Hilbert spaces, *SIAM J. Control Optim* 56 (2), pp. 1222-1252, 2018

- [14] T. Damm, L. Grüne, M. Stieler, K. Worthmann, An exponential turnpike theorem for dissipative discrete time optimal control problems, *SIAM J. Control Optim.* 52, pp. 1935–1957, 2014
- [15] L. Grüne, R. Guglielmi, Turnpike properties and strict dissipativity for discrete time linear quadratic optimal control problems, *SIAM J. Control Optim.* 56(2), pp. 1282-1302, 2018
- [16] T. Faulwasser, M. Korda, C. Jones, D. Bonvin, On turnpike and dissipativity properties of continuous time optimal control problems, *Automatica J. IFAC*, 81, pp. 297-304, 2017
- [17] E. Trélat, C. Zhang, Integral and measure-turnpike properties for infinite-dimensional optimal control systems, *Math. Control Signals Systems*, 30(1), pp. 30:3, 2018
- [18] C. Esteve, H. Kouhkouh, D. Pighin, E. Zuazua, On the turnpike property and the long time behavior of the Hamilton-Jacobi equation, *Math. Control Signals Systems*, 34, pp. 819–853, 2022
- [19] V. Askovic, E. Trélat, H. Zidani, On the asymptotic behavior of the value function in large time optimal control problems, *IFAC Papers 55(16)*, pp. 38–43, 2022
- [20] E. Trélat, *Contrôle optimal: théorie et applications*, Mathématiques Concrètes, 2008
- [21] J.C. Willems, Dissipative dynamical systems, Part I: General Theory, *Arch. Ration. Mech. Anal.* , 45, 321–351, 1972
- [22] L. Grüne, M.A. Müller, On the relation between strict dissipativity and turnpike properties, *Systems Control Lett.*, 90, pp. 45-53, 2016
- [23] H. Frankowska, Value Function in Optimal Control, *Summer School on Mathematical Control Theory*, 2001
- [24] H. Frankowska, L. N. Nguyen, Local Regularity of the Minimum Time Function, *Journal of Optimal Theory applications*, 164, pp. 68–91, 2015
- [25] P. Cannarsa, H. Frankowska, Local regularity of the value function in optimal control, *System and Control Letters*, 62, pp. 791–794, 2013
- [26] G. Stefani, Regularity properties of the minimum time map, *Nonlinear Synthesis*, Birkhauser, 1991
- [27] J-P. Aubin, H. Frankowska, *Set-Valued Analysis*, Birkhauser, Boston, Inc., Boston, MA, 1990
- [28] F.H. Clarke, R.B. Vinter, The relationship between the maximum principle and dynamic programming, *SIAM J. Control Optim.*, 25, pp. 1291–1311, 1987

- [29] L.Rifford, E. Trélat, Morse-Sard type results in sub-Riemannian geometry, *Math. Ann.*, 332, pp. 145–149, 2005
- [30] L.Rifford, E. Trélat, On the stabilization problem for nonholonomic distributions, *J. Eur. Math. Soc (JEMS)*, 11, pp. 223–255, 2009
- [31] F.H. Clarke, R.B. Vinter, Semiconcave Functions, Hamilton-Jacobi Equations and Optimal Control, *Progr. Nonlinear Differential Equations Appl.* 58, Birkhauser Boston, 25, pp. 1291–1311, 2004
- [32] B. Brogliato, R. Lozano, B. Maschke and O. Egeland, *Dissipative systems analysis and control*, Springer-Verlag London, Ltd, 2006
- [33] E. Trélat, Linear turnpike theorem, *Math. Control Signals Systems*, 35, no. 3, pp. 685–739, 2023
- [34] H. Brezis, *Functional analysis, Theory and applications*, Dunod, 2005
- [35] L.S. Pontryagin, V.G. Boltyanskii, R.V. Gamkrelidze, E.F. Mishchenko, *The mathematical theory of optimal processes*, J. Wiley & Sons, 1962
- [36] H. K. Khalil, *Nonlinear Systems*, Prentice Hall, Upper Saddle River, 2016
- [37] L. Cesari, *Optimization-theory and applications. Problems with ordinary differential equations*, Applications of Mathematics, Springer-Verlag, 1983
- [38] A. Fathi, Weak KAM theorem in Lagrangian dynamics, *Cambridge Studies in Advanced Mathematics, Hardcover, 2016*