



HAL
open science

A total variation version of Breuer–Major Central Limit Theorem under $D^{1,2}$ assumption

Jürgen Angst, Federico Dalmao, Guillaume Poly

► To cite this version:

Jürgen Angst, Federico Dalmao, Guillaume Poly. A total variation version of Breuer–Major Central Limit Theorem under $D^{1,2}$ assumption. 2023. hal-04349914

HAL Id: hal-04349914

<https://hal.science/hal-04349914>

Preprint submitted on 18 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A total variation version of Breuer–Major Central Limit Theorem under $\mathbb{D}^{1,2}$ assumption

Jürgen Angst*, Federico Dalmao† and Guillaume Poly*

September 13, 2023

Abstract

In this note, we establish a qualitative total variation version of Breuer–Major Central Limit Theorem for a sequence of the type $\frac{1}{\sqrt{n}} \sum_{1 \leq k \leq n} f(X_k)$, where $(X_k)_{k \geq 1}$ is a centered stationary Gaussian process, under the hypothesis that the function f has Hermite rank $d \geq 1$ and belongs to the Malliavin space $\mathbb{D}^{1,2}$. This result in particular extends the recent works of [NNP21], where a quantitative version of this result was obtained under the assumption that the function f has Hermite rank $d = 2$ and belongs to the Malliavin space $\mathbb{D}^{1,4}$. We thus weaken the $\mathbb{D}^{1,4}$ integrability assumption to $\mathbb{D}^{1,2}$ and remove the restriction on the Hermite rank of the base function. While our method is still based on Malliavin calculus, we exploit a particular instance of Malliavin gradient called the sharp operator, which reduces the desired convergence in total variation to the convergence in distribution of a bidimensional Breuer–Major type sequence.

1 Framework and main result

Let us consider $X = (X_n)_{n \geq 1}$ a real-valued centered stationary Gaussian sequence with unit variance, defined on an abstract probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $\rho : \mathbb{N} \rightarrow \mathbb{R}$ be the associated correlation function, in other words $\rho(|k - \ell|) = \mathbb{E}[X_k X_\ell]$, for all $k, \ell \geq 1$. We will also classically denote by $\mathcal{N}(0, \sigma^2)$ the law of a centered normal variable with variance σ^2 . Set $\gamma(dx) := (2\pi)^{-1/2} e^{-x^2/2} dx$ the standard Gaussian measure on the real line and $\gamma_d = \otimes_{k=1}^d \gamma$ its analogue in \mathbb{R}^d . We then denote by $(H_m)_{m \geq 0}$ the family of Hermite polynomials which are orthogonal with respect to γ , namely $H_0 \equiv 1$ and

$$H_m(x) := (-1)^m e^{\frac{x^2}{2}} \frac{d^m}{dx^m} e^{-\frac{x^2}{2}}, \quad m \geq 1.$$

We denote by $L^2(\mathbb{R}, \gamma)$ the space of square integrable real functions with respect to the Gaussian measure. Recall that a real function $f \in L^2(\mathbb{R}, \gamma)$ is said to have Hermite rank

*Univ Rennes, CNRS, IRMAR - UMR 6625, F-35000 Rennes, France.

jurgen.angst@univ-rennes1.fr, guillaume.poly@univ-rennes1.fr

†DMEL, Cenur LN, Universidad de la República, Salto, Uruguay. fdalmao@unorte.edu.uy

This work was supported by the ANR grant UNIRANDOM, ANR-17-CE40-0008.

$d \geq 0$ if it can be decomposed as a sum of the form

$$f(x) = \sum_{m=d}^{+\infty} c_m H_m(x), \quad c_d \neq 0.$$

For integers $k, p \geq 1$, we further denote by $\mathbb{D}^{k,p}(\mathbb{R}, \gamma)$ the Malliavin–Sobolev space consisting of the completion of the family of polynomial functions $q : \mathbb{R} \rightarrow \mathbb{R}$ with respect to the norm

$$\|q\|_{k,p} := \left| \int_{\mathbb{R}} \left(|q(x)|^p + \sum_{\ell=1}^k |q^{(\ell)}(x)|^p \right) \gamma(dx) \right|^{1/p},$$

where $q^{(\ell)}$ is the ℓ -th derivative of q . Given a real function f , let us finally set

$$S_n(f) := \frac{1}{\sqrt{n}} \sum_{k=1}^n f(X_k).$$

In this framework, the celebrated Central Limit Theorem (CLT) by Breuer and Major gives sufficient conditions on ρ and f so that the sequence $S_n(f)$ satisfies a CLT.

Theorem 1 (Theorem 1 in [BM83]). *If the function f belongs to $L^2(\mathbb{R}, \gamma)$ with Hermite rank $d \geq 1$ and if $\rho \in \ell^d(\mathbb{N})$, i.e. $\sum_{\mathbb{N}} |\rho(k)|^d < +\infty$, then the sequence $(S_n(f))_{n \geq 1}$ converges in distribution as n goes to infinity to a normal distribution $\mathcal{N}(0, \sigma^2)$, where the limit variance is given by*

$$\sigma^2 := \sum_{m=d}^{\infty} m! c_m^2 \sum_{k \in \mathbb{Z}} \rho(k)^m,$$

with c_m being the coefficients appearing in the Hermite expansion of f .

Recently, under mild additional assumptions, a series of articles has reinforced the above convergence in distribution into a convergence in total variation, with polynomial quantitative bounds, see e.g. [KN19, NPY19, NZ21, NNP21]. Recall that the total variation distance between the distributions of two real random variables X and Y is given by

$$d_{\text{TV}}(X, Y) := \sup_{A \in \mathcal{B}(\mathbb{R})} |\mathbb{P}(X \in A) - \mathbb{P}(Y \in A)|,$$

where the supremum runs over $\mathcal{B}(\mathbb{R})$, the Borel sigma field on the real line. To the best of our knowledge, the best statement so far in this direction is the following

Theorem 2 (Theorem 1.2 in [NNP21]). *Assume that $f \in L^2(\mathbb{R}, \gamma)$ has Hermite rank $d = 2$ and that it belongs to $\mathbb{D}^{1,4}(\mathbb{R}, \gamma)$. Suppose that $\rho \in \ell^d(\mathbb{N})$ and that the variance σ^2 of Theorem 1 is positive. Then, there exists a constant $C > 0$ independent of n such that*

$$d_{\text{TV}} \left(\frac{S_n(f)}{\sqrt{\text{var}(S_n(f))}}, \mathcal{N}(0, 1) \right) \leq \frac{C}{\sqrt{n}} \left[\left(\sum_{|k| \leq n} |\rho(k)| \right)^{\frac{1}{2}} + \left(\sum_{|k| \leq n} |\rho(k)|^{\frac{4}{3}} \right)^{\frac{3}{2}} \right].$$

The goal of this note is to establish that the convergence in total variation in fact holds as soon as the function f is in the Malliavin–Sobolev space $\mathbb{D}^{1,2}(\mathbb{R}, \gamma)$ and has Hermite rank $d \geq 1$.

Theorem 3. *Suppose that $f \in \mathbb{D}^{1,2}(\mathbb{R}, \gamma)$ has Hermite rank $d \geq 1$. Suppose moreover that $\rho \in \ell^d(\mathbb{N})$ and that the variance σ^2 of Theorem 1 is positive. Then, as n goes to infinity*

$$d_{\text{TV}} \left(\frac{S_n(f)}{\sqrt{\text{var}(S_n(f))}}, \mathcal{N}(0, 1) \right) \xrightarrow{n \rightarrow +\infty} 0.$$

Note that, for the sake of simplicity, we only consider here a real Gaussian sequence $(X_n)_{n \geq 1}$ and a real function f but our method is robust and would yield, under similar covariance and rank assumptions, a convergence in total variation for a properly renormalized sequence of the type $\sum_{k=1}^n f(X_k^1, \dots, X_k^d)$ associated with a sequence of Gaussian vectors $(X_n)_{n \geq 1}$ with values in \mathbb{R}^d and a function f in the corresponding Malliavin–Sobolev space $\mathbb{D}^{1,2}(\mathbb{R}^d, \gamma_d)$.

The detailed proof of Theorem 3 is the object of the next section and the rest of the paper. Unsurprisingly, we use the Malliavin–Stein approach to establish the CLT in total variation. However, our approach differs from the other works mentioned above in that we make use of the so called “sharp gradient”, whose definition and main properties are recalled in Section 2.2. With this tool at hand and in view of using Malliavin–Stein equation to characterize the proximity to the normal distribution, we shall see that the convergence in total variation in fact reduces to two rather simple steps

- i*) a two-dimensional version of the classical Breuer–Major CLT (i.e. in distribution not in total variation), see Section 2.3 ;
- ii*) some elementary uniform integrability estimates, allowing to pass from a convergence in probability to a convergence in L^1 , see Section 2.4.

2 Proof of the main result

As mentioned just above, the setting of the proof of Theorem 3 is the one of Malliavin–Stein calculus. Note that for each fixed $n \geq 1$, the quantity of interest $S_n(f)$ involves only a finite number of Gaussian coefficients. So let us sketch the framework of Malliavin–Stein method in the finite dimensional setting, and we refer to [Nua09] or [NP12] for a more general introduction.

2.1 A glimpse of Malliavin calculus

Let us fix an integer $n \geq 1$ and let us place ourselves in the product probability space $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n), \gamma_n)$ with $\gamma_n := \otimes_{k=1}^n \gamma$, the n -dimensional standard Gaussian distribution on \mathbb{R}^n . Consider the classical *Ornstein–Uhlenbeck* operator $\mathcal{L}_n := \Delta - \vec{x} \cdot \nabla$ which is

symmetric with respect to γ_n . We have then the standard decomposition of the L^2 -space in Wiener chaoses, namely

$$L^2(\gamma_n) = \bigoplus_{k=0}^{\infty} \text{Ker}(\mathcal{L}_n + k\text{I}), \quad \text{with}$$

$$\text{Ker}(\mathcal{L}_n + k\text{I}) = \text{Vect} \left(\prod_{i=0}^n H_{k_i}(x_i) \middle| \sum_{i=0}^n k_i = n \right) := \underbrace{\mathcal{W}_k}_{k\text{-th Wiener chaos}}.$$

The square field or “carré du champ” operator Γ_n is then defined as the bilinear operator $\Gamma_n := [\cdot, \cdot] = \nabla \cdot \nabla$. As a glimpse of the power of Malliavin–Stein approach in view of establishing total variation estimates, recall that if $F \in \text{Ker}(\mathcal{L}_n + k\text{I})$ is such that $\mathbb{E}[F^2] = 1$, then for some constant C_k only depending on k , the total variation distance between the variable F and a standard Gaussian can be upper bounded by

$$d_{TV}(F, \mathcal{N}(0, 1)) \leq C_k \sqrt{\text{var}(\Gamma[F, F])}.$$

Via the notion of isonormal Gaussian process, the finite dimensional framework for Malliavin–Stein method sketched above can in fact be extended to the infinite dimensional setting giving rise to an Ornstein–Uhlenbeck operator \mathcal{L} and an associated “carré du champ” Γ , see e.g. Chapter 2 in [NP12].

2.2 The sharp gradient

A detailed introduction to the sharp gradient can be found in Section 4.1 of the reference [AP20]. We only recall here the basics which will be useful to our purpose. Let us assume that $(N_k)_{k \geq 1}$ is an i.i.d. sequence of standard Gaussian variables on $(\Omega, \mathcal{F}, \mathbb{P})$ which generate the first Wiener chaos. Without loss of generality, we shall assume that $\mathcal{F} = \sigma(N_k, k \geq 1)$. We will also need a copy $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$ of this probability space as well as $(\hat{N}_i)_{i \geq 1}$ a corresponding i.i.d. sequence of standard Gaussian variables such that $\hat{\mathcal{F}} = \sigma(\hat{N}_k, k \geq 1)$. We will denote by $\hat{\mathbb{E}}$ the expectation with respect to the measure $\hat{\mathbb{P}}$. For any integer $m \geq 1$ and any function Φ in the space $\mathcal{C}_b^1(\mathbb{R}^m, \mathbb{R})$ of continuously differentiable functions with a bounded gradient, we then set

$$\sharp \Phi(N_1, \dots, N_m) := \sum_{i=1}^m \partial_i \Phi(N_1, \dots, N_m) \hat{N}_i. \quad (1)$$

In Sections 4.1.1 and 4.1.2 of [AP20], it is shown that this *gradient* is closable and extends to the Malliavin space $\mathbb{D}^{1,2}$, where

$$\mathbb{D}^{1,2} := \left\{ F \in \mathbb{L}^2(\Omega, \mathcal{F}, \mathbb{P}), \mathbb{E}[F^2] + \mathbb{E}[\sharp F]^2 < +\infty \right\}.$$

The last space $\mathbb{D}^{1,2}$ is naturally the infinite dimensional version of the Malliavin–Sobolev space $\mathbb{D}^{1,2}(\mathbb{R}, \gamma)$ introduced in Section 1 in the one-dimensional setting. In particular, Proposition 8 in the latter reference shows that

$$\forall F \in \mathbb{D}^{1,2}, \forall \phi \in \mathcal{C}_b^1(\mathbb{R}, \mathbb{R}) : \sharp \phi(F) = \phi'(F) \sharp F.$$

Given $F \in \mathbb{D}^{1,2}$, taking first the expectation $\hat{\mathbb{E}}$ with respect $\hat{\mathbb{P}}$ and using Fubini inversion of sums yields the following key relation, for all $\xi \in \mathbb{R}$

$$\mathbb{E} \left(\exp \left(-\frac{\xi^2}{2} \Gamma[F, F] \right) \right) = \hat{\mathbb{E}} \mathbb{E} \left(\exp \left(i\xi \sharp F \right) \right). \quad (2)$$

By essence, via their Laplace/Fourier transforms, this key equation allows to relate the asymptotic behavior in distribution (or in probability if the limit is constant) of the carré du champ $\Gamma[F, F]$ with the one of the sharp gradient $\sharp F$.

Finally, let us remark that by definition, the image $(\sharp X_k)_{k \geq 1}$ of our initial stationary sequence $(X_k)_{k \geq 1}$ by the sharp gradient is an independent copy of $(X_k)_{k \geq 1}$. We will write $(\sharp X_k)_{k \geq 1} = (\hat{X}_k)_{k \geq 1}$ in the sequel.

2.3 Convergence in probability via a two dimensional CLT

Let us suppose that f satisfies the assumptions of Theorem 3, namely $f \in \mathbb{D}^{1,2}(\mathbb{R}, \gamma)$ with Hermite rank $d \geq 1$, so that it can be decomposed as $f = \sum_{m=d}^{\infty} c_m H_m$ in $L^2(\mathbb{R}, \gamma)$. Let \mathcal{L}^{-1} denote the pseudo-inverse of the Ornstein–Uhlenbeck operator and consider the pre-image

$$g(x) := -\mathcal{L}^{-1}[f](x) = \sum_{m=d}^{\infty} \frac{c_m}{m} H_m(x).$$

To simplify the expressions in the sequel, we set

$$F_n := S_n(f) = \frac{1}{\sqrt{n}} \sum_{k=1}^n f(X_k), \quad \text{and} \quad G_n := S_n(g) = -\mathcal{L}^{-1} F_n = \frac{1}{\sqrt{n}} \sum_{k=1}^n g(X_k).$$

Now, take $(s, t, \xi) \in \mathbb{R}^3$ and let us apply the above key relation (2) with the random variable $tF_n + sG_n$, we get

$$\mathbb{E} \left[\exp \left(-\frac{\xi^2}{2} \Gamma[tF_n + sG_n, tF_n + sG_n] \right) \right] = \hat{\mathbb{E}} \mathbb{E} \left[\exp \left(i\xi \left(t \sharp F_n + s \sharp G_n \right) \right) \right]. \quad (3)$$

On the one hand, by bilinearity of the carré du champ operator, we have

$$\Gamma[tF_n + sG_n, tF_n + sG_n] = t^2 \Gamma[F_n, F_n] + s^2 \Gamma[G_n, G_n] + 2ts \Gamma[F_n, -\mathcal{L}^{-1} F_n]. \quad (4)$$

On the other hand, the right hand side of Equation (3) is simply the characteristic function under $\mathbb{P} \otimes \hat{\mathbb{P}}$ of the couple $(\sharp F_n, \sharp G_n)$ where

$$\left(\sharp F_n, \sharp G_n \right) = \frac{1}{\sqrt{n}} \sum_{k=1}^n \left(f'(X_k) \hat{X}_k, g'(X_k) \hat{X}_k \right),$$

is a “Breuer–Major type” sequence with respect to the \mathbb{R}^2 -valued centered stationary Gaussian process $(\hat{X}_k, X_k)_{k \geq 1}$ and the \mathbb{R}^2 -valued functional

$$(x, y) \mapsto \Psi(x, y) := (f'(x)y, g'(x)y).$$

Since f is in $\mathbb{D}^{1,2}(\mathbb{R}, \gamma)$, its derivative f' is in $L^2(\mathbb{R}, \gamma)$ and $(\hat{X}_k)_{k \geq 1}$ and $(X_k)_{k \geq 1}$ are independent, therefore the functional Ψ is in $L^2(\mathbb{R}^2, \gamma_2)$ and the multivariate counterpart of the classical Breuer–Major Theorem applies, see Theorem 4 of [Arc94].

As a result, the bidimensional sequence $(\#F_n, \#G_n)$ converges in distribution, under $\mathbb{P} \otimes \hat{\mathbb{P}}$, towards a bidimensional centered Gaussian vector with a symmetric semi-positive covariance matrix Σ . Therefore, from Equations (3) and (4) and via the characterization of convergence in distribution in terms of Fourier transform, there exists real numbers λ, μ, ν (depending on the limit covariance matrix Σ) such that for any $(s, t, \xi) \in \mathbb{R}^3$, as n goes to infinity, we have

$$\mathbb{E} \left[e^{-\frac{\xi^2 t^2}{2} \Gamma[F_n, F_n] - \frac{\xi^2 s^2}{2} \Gamma[G_n, G_n] - \xi^2 t s \Gamma[F_n, -\mathcal{L}^{-1} F_n]} \right] \xrightarrow[n \rightarrow \infty]{} e^{-\frac{\xi^2}{2} (\lambda t^2 + \mu s^2 + 2\nu t s)}.$$

Since the above convergence is valid for any $\xi \in \mathbb{R}$, this shows in particular that for any fixed $(s, t) \in \mathbb{R}^2$, the sequence $\Gamma[tF_n + sG_n, tF_n + sG_n]$ converges in distribution (and thus in probability) towards the constant variable $(\lambda t^2 + \mu s^2 + 2\nu t s)$. Choosing $s = t = 1$, we thus get that $\Gamma[F_n + G_n, F_n + G_n]$ converges in probability towards $(\lambda + \mu + 2\nu)$. Choosing $s = 0$ and $t = 1$, then $t = 0$ and $s = 1$, one deduce in the same manner that $\Gamma[F_n, F_n]$ and $\Gamma[G_n, G_n]$ both converge in probability towards λ and μ respectively. Finally, by Equation (4), one can conclude that the cross term

$$\Gamma[F_n, G_n] = \Gamma(F_n, -\mathcal{L}^{-1} F_n) = \hat{\mathbb{E}} \left[\#F_n \#G_n \right]$$

also converges in probability towards the constant limit variable ν .

2.4 Gaining some uniform integrability

Since our goal is to derive convergence in total variation of $F_n = S_n(f)$, the convergence in probability of the term $\Gamma[F_n, -\mathcal{L}^{-1} F_n]$ is not sufficient. Indeed, with Stein's Equation in mind, the lack of uniform integrability is a problem to deduce the following required asymptotic behavior for any $\phi \in \mathcal{C}_b^1(\mathbb{R})$, as n goes to infinity

$$\mathbb{E} \left[\phi'(F_n) \Gamma[F_n, -\mathcal{L}^{-1} F_n] \right] \approx \nu \mathbb{E} \left[\phi'(F_n) \right].$$

In order to bypass this problem, let us go back to the two-dimensional classical Breuer–Major theorem associated with the functional Ψ used in the last section. For any integer $p \geq 1$, let us denote by Ψ_p the projection of Ψ on the first p -th chaoses. Applying Theorem 4 and Equation (2.43) of [Arc94], we get that there exists a constant $C > 0$ (which depends only on the covariance structure of the underlying Gaussian process) such that

$$\sup_{n \geq 1} \mathbb{E} \hat{\mathbb{E}} \left[\left| \frac{1}{\sqrt{n}} \sum_{k=1}^n (\Psi - \Psi_p)(X_k, \hat{X}_k) \right|^2 \right] \leq C \times \int_{\mathbb{R}^2} |(\Psi - \Psi_p)(x)|^2 \gamma_2(dx).$$

Since Ψ belongs to $L^2(\mathbb{R}^2, \gamma_2)$, the last term on the right hand side goes to zero as p goes to infinity. As a result, uniformly in $n \geq 1$, the two-dimensional process

$$\left(\#F_n, \#G_n \right) = \frac{1}{\sqrt{n}} \sum_{k=1}^n \Psi(X_k, \hat{X}_k)$$

can be approximated arbitrarily closely in $L^2(\mathbb{P} \otimes \hat{\mathbb{P}})$ by the following process which is finitely expanded on the Wiener chaoses

$$Z_n^p := (Z_n^{p,1}, Z_n^{p,2}) := \frac{1}{\sqrt{n}} \sum_{k=1}^n \Psi_p(X_k, \hat{X}_k).$$

Therefore, choosing $p \geq 1$ large enough, uniformly in $n \geq 1$, the product $\#F_n \times \#G_n$ can be approximated arbitrarily closely in $L^1(\mathbb{P} \otimes \hat{\mathbb{P}})$ by $\Delta_n^p := Z_n^{p,1} \times Z_n^{p,2}$. In other words, for any $\varepsilon > 0$ and $p \geq 1$ large enough, we have

$$\sup_n \mathbb{E} \left[\left| \hat{\mathbb{E}} \left(\#F_n \times \#G_n \right) - \hat{\mathbb{E}} \left(\Delta_n^p \right) \right| \right] \leq \sup_n \mathbb{E} \hat{\mathbb{E}} \left[\left| \#F_n \times \#G_n - \Delta_n^p \right| \right] < \varepsilon.$$

But mimicking the proof detailed in the previous Section 2.3 for the convergence in probability of $\Gamma[F_n, G_n]$ towards the constant variable ν , one would then similarly get here that $\hat{\mathbb{E}}[\Delta_n^p]$ converges in probability under \mathbb{P} towards a constant random variable $\nu_p \in \mathbb{R}$, and by construction $\lim_{p \rightarrow +\infty} \nu_p = \nu$. The crucial point here is that both random variables Δ_n^p and $\hat{\mathbb{E}}[\Delta_n^p]$ are now finitely expanded on the Wiener chaoses under $\mathbb{P} \otimes \hat{\mathbb{P}}$ and \mathbb{P} respectively. Therefore, by hypercontractivity, the convergence in probability can be freely upgraded to the convergence in L^q for every $q \geq 1$. In particular, as n goes to infinity, the sequence $\hat{\mathbb{E}}[\Delta_n^p]$ converges in L^1 to the constant variable ν_p .

2.5 Conclusion

We go back to Stein's Equation. Let $\phi \in \mathcal{C}_b^1(\mathbb{R})$ and $\varepsilon > 0$. Integrating by parts, for $p \geq 1$ large enough and by the results of the last section, we have

$$\begin{aligned} & \left| \mathbb{E} [F_n \phi(F_n)] - \nu \mathbb{E} [\phi'(F_n)] \right| = \left| \mathbb{E} [\phi'(F_n) \Gamma[F_n, -\mathcal{L}^{-1} F_n]] - \nu \mathbb{E} [\phi'(F_n)] \right| \\ & = \left| \mathbb{E} [\phi'(F_n) \Gamma[F_n, G_n]] - \nu \mathbb{E} [\phi'(F_n)] \right| \\ & = \left| \mathbb{E} \left[\phi'(F_n) \left(\Gamma[F_n, G_n] - \hat{\mathbb{E}}[\Delta_n^p] \right) \right] + \mathbb{E} \left[\phi'(F_n) \left(\hat{\mathbb{E}}[\Delta_n^p] - \nu_p \right) \right] + (\nu_p - \nu) \mathbb{E} [\phi'(F_n)] \right| \\ & \leq \|\phi'\|_\infty \varepsilon + \|\phi'\|_\infty \mathbb{E} \left[\left| \hat{\mathbb{E}}[\Delta_n^p] - \nu_p \right| \right] + \|\phi'\|_\infty |\nu_p - \nu|. \end{aligned}$$

As a result, letting first n and then p go to infinity, we get that uniformly in ϕ such that $\|\phi'\|_\infty \leq C$

$$\limsup_{n \rightarrow +\infty} \left| \mathbb{E} [F_n \phi(F_n)] - \nu \mathbb{E} [\phi'(F_n)] \right| = 0.$$

One can then classically conclude using Stein's approach for the convergence in total variation.

References

- [AP20] Jürgen Angst and Guillaume Poly. On the absolute continuity of random nodal volumes. *The Annals of Probability*, 48(5):2145 – 2175, 2020.
- [Arc94] Miguel A. Arcones. Limit theorems for nonlinear functionals of a stationary gaussian sequence of vectors. *The Annals of Probability*, 22(4):2242–2274, 1994.
- [BM83] Péter Breuer and Péter Major. Central limit theorems for non-linear functionals of gaussian fields. *Journal of Multivariate Analysis*, 13(3):425–441, 1983.
- [KN19] Sefika Kuzgun and David Nualart. Rate of convergence in the Breuer–Major theorem via chaos expansions. *Stochastic Analysis and Applications*, 37(6):1057–1091, 2019.
- [NNP21] Ivan Nourdin, David Nualart, and Giovanni Peccati. The Breuer–Major theorem in total variation: Improved rates under minimal regularity. *Stochastic Processes and their Applications*, 131:1–20, 2021.
- [NP12] Ivan Nourdin and Giovanni Peccati. *Normal approximations with Malliavin calculus*, volume 192 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2012. From Stein’s method to universality.
- [NPY19] Ivan Nourdin, Giovanni Peccati, and Xiaochuan Yang. Berry–Esseen bounds in the Breuer–Major CLT and Gebelein’s inequality. *Electronic Communications in Probability*, 24(none):1 – 12, 2019.
- [Nua09] David Nualart. *Malliavin calculus and its applications*, volume 110 of *CBMS Regional Conference Series in Mathematics*. Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2009.
- [NZ21] David Nualart and Hongjuan Zhou. Total variation estimates in the Breuer–Major theorem. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 57(2):740 – 777, 2021.